Published in final edited form as:

Gut. 2014 February ; 63(2): . doi:10.1136/gutjnl-2012-304331.

Functional effects of 13 rare *PRSS1* variants presumed to cause chronic pancreatitis

Andrea Schnúr^{1,2}, Sebastian Beer¹, Heiko Witt^{3,4}, Péter Hegyi², and Miklós Sahin-Tóth^{1,*}

¹Department of Molecular and Cell Biology, Henry M. Goldman School of Dental Medicine, Boston University, Boston, MA 02118

²First Department of Medicine, University of Szeged, Szeged, Hungary

³Else Kröner-Fresenius-Zentrum für Ernährungsmedizin (EKFZ) & Zentralinstitut für Ernährungs- und Lebensmittelforschung (ZIEL), Technische Universität München (TUM), 85350 Freising, Germany

⁴Department of Pediatrics, Klinikum rechts der Isar (MRI), Technische Universität München (TUM), 80804 Munich, Germany

Abstract

Objective—Hereditary pancreatitis is caused by mutations in human cationic trypsinogen (*PRSS1*) which lead to increased autoactivation by altering chymotrypsin C (CTRC)-dependent trypsinogen activation and degradation. Exceptions are some cysteine mutations which cause misfolding, intracellular retention and endoplasmic reticulum stress. Clinical relevance of many *PRSS1* variants found in patients with sporadic chronic pancreatitis is unknown but often assumed by analogy with known disease-causing mutations. Functional comparison of *PRSS1* variants found in sporadic and hereditary cases is needed to resolve this dilemma.

Design—Here, we investigated the functional phenotype of 13 published *PRSS1* variants with respect to autoactivation in the presence of CTCRC and cellular secretion.

Results—Only mutation p.D100H increased trypsinogen autoactivation, but this gain in function was offset by a marked reduction in secretion. Five mutants (p.P36R, p.G83E, p.I88N, p.V123M, p.S124F) showed decreased autoactivation due to increased degradation by CTCRC. Five mutants exhibited strongly (p.D100H, p.C139F) or moderately (p.K92N, p.S124F, p.G208A) reduced secretion, whereas mutant p.K170E showed slightly increased secretion. Mutant p.I88N was also secreted to higher levels but was rapidly degraded by CTCRC. Finally, three mutants (p.Q98K, p.T137M, p.S181G) had no phenotypic alterations relative to wild-type trypsinogen.

Conclusion—Rare *PRSS1* variants found in sporadic chronic pancreatitis do not stimulate autoactivation but may cause increased degradation, impaired secretion or no functional change. Variants with reduced secretion are likely pathogenic due to mutation-induced misfolding and consequent endoplasmic reticulum stress.

*Correspondence to Miklós Sahin-Tóth, 72 East Concord Street, Evans-433; Boston, MA 02118; Tel: (617) 414-1070; Fax: (617) 414-1041; miklos@bu.edu.

CONTRIBUTIONS

The study was designed by A.S., P.H. and M.S.-T. The experiments were performed by A.S., S.B. and H.W.. The manuscript was written by A.S., S.B., P.H. and M.S.-T.

COMPETING INTEREST STATEMENT

No competing interest to declare.

Keywords

chronic pancreatitis; trypsinogen; trypsin; chymotrypsin C; misfolding

Hereditary pancreatitis is an autosomal dominant disorder caused by mutations in the serine protease 1 (*PRSSI*) gene that codes for cationic trypsinogen [1]. Causative mutations are p.R122H (~65%), p.N29I (~25%) and less frequently p.A16V, p.D22G, p.K23R, p.K23_I24insIDK, p.N29T, p.V39A, p.R116C and p.R122C [1–5]. To date, 21 additional rare missense *PRSSI* variants have been reported, the majority of which were found in patients with sporadic chronic pancreatitis with no family history [5], www.pancreasgenetics.org). The mechanism of action of hereditary pancreatitis-associated mutations involves increased autoactivation of mutant trypsinogens resulting in elevated intrapancreatic trypsin activity levels [6] (Figure 1). Recent studies uncovered that *PRSSI* mutations alter the regulation of activation and degradation of cationic trypsinogen by chymotrypsin C (CTRC). The digestive enzyme CTRC stimulates trypsinogen activation by processing the activation peptide to a shorter form, which is easier cleaved by trypsin [7]. Somewhat paradoxically, CTRC also promotes degradation of trypsinogen by cleaving the calcium binding loop [6, 8]. This cleavage in combination with a trypsin-mediated autolytic cleavage results in inactivation of trypsinogen during autoactivation and lower trypsin levels attained. Pancreatitis-associated mutations render trypsinogen resistant to CTRC-dependent degradation and/or increase N-terminal processing by CTRC and thereby elevate trypsin levels generated through autoactivation [6] (Figure 1).

The unifying pathological mechanism described above does not seem to apply to some mutations that alter the number of cysteine residues in cationic trypsinogen. Hereditary-pancreatitis associated mutation p.R116C was shown to induce protein misfolding with intracellular retention and degradation, which may represent an alternative disease-causing mechanism unrelated to trypsinogen activation and trypsin activity [9]. Mutation p.C139S, which was reported in sporadic cases of chronic pancreatitis, exhibits similar properties [9]. Mutation-dependent misfolding can elicit endoplasmic reticulum (ER) stress, which might be responsible for increased pancreatitis risk, although the mechanism remains unclear (Figure 1).

In the present study we surveyed the functional properties of 13 rare missense *PRSSI* variants found in patients with sporadic chronic pancreatitis. Our primary objective was to test whether these variants also exhibit increased activation in the presence of CTRC as previously seen with disease-causing mutants in hereditary pancreatitis. A second objective of the study was to assess cellular secretion of the mutants to determine whether mutation-induced changes in folding and secretion may be a more common phenotype of *PRSSI* variants than previously appreciated.

EXPERIMENTAL PROCEDURES

Nomenclature

Amino acid residues in human cationic trypsinogen (serine protease 1, *PRSSI*) are numbered starting with the initiator methionine of the primary translation product, in accordance with the recommendations of the Human Genome Variation Society.

Plasmid construction and mutagenesis

The pTrapT7 intein-*PRSSI*, pcDNA3.1(–) *PRSSI* and pcDNA3.1(–) *CTRC* 10His expression plasmids were constructed previously [7, 8, 10]. Missense mutations were

introduced by overlap extension PCR mutagenesis, cloned into the expression plasmids and verified by DNA sequencing.

Expression and purification of trypsinogen

Wild-type and mutant trypsinogens were expressed in the aminopeptidase P deficient LG-3 *E. coli* strain as fusions with a self-splicing mini-intein, as described in [10, 11]. This expression system was developed to produce recombinant trypsinogen with uniform, authentic N termini. Isolation of cytoplasmic inclusion bodies, *in vitro* refolding and purification with ecotin affinity chromatography were carried out according to published protocols [10, 11]. Mutant p.C139F could not be purified by this method, as it misfolded during *in vitro* refolding. Concentrations of trypsinogen preparations were calculated from their UV absorbance at 280 nm using the extinction coefficient $37,525 \text{ M}^{-1} \text{ cm}^{-1}$.

Cell culture and transfection

Human embryonic kidney 293T (HEK 293T) cells were cultured and transfected as described previously [12]. Transfections were performed using 1 μg expression plasmid and 2.5 μL Lipofectamine 2000 (Invitrogen, Carlsbad, CA) in 2 mL Dulbecco's Modified Eagle Medium medium (DMEM). After overnight incubation, cells were washed and the transfection medium was replaced with 2 mL OPTI-MEM I Reduced Serum Medium (Invitrogen) containing 1 mM benzamidine (final concentration) to inhibit autoactivation of secreted trypsinogen. Conditioned media were harvested 24 h after addition of OPTI-MEM.

Expression and purification of human CTRC

Large-scale expression of human CTRC in HEK 293T cells and purification from the conditioned medium using nickel-affinity chromatography were performed as reported previously [6]. CTRC was activated with human cationic trypsin and active CTRC concentrations were determined by active site titration with ecotin, as described [13].

Trypsinogen autoactivation in the presence of CTRC

Trypsinogen at 1 μM concentration was incubated with 5 nM human CTRC and 10 nM cationic trypsin in 0.1 M Tris-HCl (pH 8.0), 1 mM CaCl_2 and 0.05% Tween-20 (final concentrations) at 37 °C. At given times, 2 μL aliquots were withdrawn and mixed with 48 μL assay buffer containing 0.1 M Tris-HCl (pH 8.0), 1mM CaCl_2 , and 0.05% Tween-20. Trypsin activity was measured by adding 150 μL 200 μM N-CBZ-Gly-Pro-Arg-p-nitroanilide substrate and following the release of the yellow p-nitroanilin at 405 nm in a SpectraMax plus384 microplate reader (Molecular Devices, Sunnyvale, CA) for 1 min. Reaction rates were calculated from fits to the initial linear portions of the curves. The trypsin substrate was dissolved in 0.1 M Tris-HCl (pH 8.0), 1 mM CaCl_2 and 0.05% Tween-20.

Measurement of trypsin activity in conditioned media

Aliquots (50 μL) of conditioned media were supplemented with 5 μL 1 M Tris-HCl (pH 8.0) and 1 μL 0.5 M CaCl_2 and trypsinogens were activated by adding 1 μL 1.4 $\mu\text{g/mL}$ human enteropeptidase (R&D Systems, Minneapolis, MN). After incubation for 1 h at 37 °C, a 50 μL aliquot was removed and mixed with 150 μL 200 μM N-CBZ-Gly-Pro-Arg-p-nitroanilide substrate. Activity was determined as described above.

SDS-PAGE and Western blotting

Conditioned media (200 μL) was precipitated with 10% trichloroacetic acid (final concentration); resuspended in 20 μL Laemmli sample buffer containing 100 mM

dithiothreitol, heat-denatured at 95 °C for 5 min and run on 15% SDS-polyacrylamide gels. The gels were stained with Coomassie Blue R-250. For Western blotting, 5 µL conditioned media were directly mixed with sample buffer and electrophoresed as described above. Proteins were transferred onto an Immobilon-P membrane (Millipore Corporation, Bedford, MA) at 300 mA for 1.5 h. The membrane was blocked with 5% milk powder dissolved in phosphate-buffered saline supplemented with 0.1% Tween-20 (final concentration), at 4 °C overnight. Trypsinogen was detected with a sheep polyclonal antibody (R&D Systems, #AF3848) used at a dilution of 1:5000 followed by horse-radish peroxidase (HRP)-conjugated donkey polyclonal anti-sheep IgG (R&D Systems, #HAF016) used at 1:2000 dilution. Incubations with primary and secondary antibodies were performed at room temperature for 1 h each. HRP was detected using the SuperSignal West Pico Chemiluminescent Substrate (Thermo Scientific). Quantitation of bands was carried out with the Image Lab 3.0 (Bio-Rad) software.

RESULTS

***PRSS1* variants found in subjects with sporadic chronic pancreatitis**

We studied here 13 missense *PRSS1* variants, found in patients with idiopathic chronic pancreatitis or recurrent acute pancreatitis with no family history (Table 1). The variants were detected in the heterozygous state, with the exception of p.G208A, which was also identified in a homozygous individual. As indicated, the variants were found only in one to three cases each, therefore, the genetic information was insufficient to determine whether these variants are pathogenic or neutral. Unaffected carrier parent was reported in two cases, for variant p.C139F and p.S181G. Note that variants p.T137M, p.K170E and p.G208A were from subjects of Asian origin, whereas all other variants were from subjects of European origin.

To determine whether these 13 variants are prevalent in the general population, we consulted the NHLBI Exome Sequencing Project Exome Variant Server (<http://evs.gs.washington.edu>) which lists aggregate exome-sequencing data for approximately 4300 European-American and 2200 African-American individuals. We found *PRSS1* variants p.V123M and p.T137M in one European-American subject each and variant p.S181G in one African-American subject. However, interpretation of these findings is difficult, because the database also lists the disease-causing mutation p.R122C found in two European-American individuals, suggesting that the studied cohort may have contained pancreatitis patients. Therefore, we sequenced exon-3 of the *PRSS1* gene in 1000 German subjects without any pancreatic disease. Exon-3 was selected because most of the studied variants (9 of 13) were in this region. With the exception of a novel c.367G>T (p.V123L) variant found in one subject, no other alterations were identified. This observation is in agreement with published sequencing data showing the absence of these 13 *PRSS1* variants in 200 French [14], 82 German [15], 420 Chinese [16, 17] and 28 Korean [18] control subjects. We conclude that the 13 *PRSS1* variants presumed to cause chronic pancreatitis (Table 1) are not generally found in a healthy population.

Autoactivation and catalytic activity of *PRSS1* variants

First, trypsinogen variants were characterized for their ability to autoactivate in the absence of CTTC. When measured at pH 8.0, in 1 mM Ca²⁺, only one mutant, p.D100H, exhibited increased autoactivation, which was about 2.5-fold faster than wild type (Supplementary Figure S1A). Five mutants (p.Q98K, p.T137M, p.K170E, p.S181G, and p.G208A) autoactivated as well as wild-type cationic trypsinogen (Supplementary Figure S1B). Mutants p.K92N and p.S124F autoactivated at a decreased rate but reached essentially the same final trypsin levels as wild-type trypsinogen (Supplementary Figure S1C). In contrast,

four mutants exhibited decreased rates of autoactivation (p.P36R, p.G83E, p.I88N, and p.V123M) with reduced final trypsin levels, suggesting that these mutants may become partially degraded during autoactivation (Supplementary Figure S1D).

Catalytic activity of the activated trypsin variants was measured with the synthetic peptide substrate *N*-CBZ-Gly-Pro-Arg-p-nitroanilide. As shown in Supplementary Table S1, enzyme kinetic parameters of trypsin variants were comparable to those of wild-type cationic trypsin. This finding is consistent with previous observations that natural *PRSS1* variants almost never affect catalytic function of trypsin.

Note that mutant p.C139F could not be purified in sufficient quantities due to misfolding, therefore it was not tested in the autoactivation and catalytic activity assays.

Autoactivation of *PRSS1* variants in the presence of CTTC

To determine the effect of CTTC on the trypsinogen variants, we measured autoactivation in the presence of 5 nM human CTTC. This low CTTC concentration slightly stimulates the rate of autoactivation but causes a reduction in final active trypsin levels due to trypsinogen degradation during activation [6]. In the experiments presented, wild-type cationic trypsinogen reached about 50% activity in the presence of CTTC, relative to trypsin activity observed in the absence of CTTC (not shown, also see Figure 1A in ref [6]). Under these conditions, both increases and decreases in autoactivation are readily detectable.

Only one of the 12 variants tested, p.D100H exhibited increased activation in the presence of CTTC (Figure 2A). A similar phenotype was observed with this mutant when autoactivation was performed in the absence of CTTC (see above), indicating that the mutation increases autoactivation independent of CTTC. Consistent with this interpretation, degradation of p.D100H mutant trypsinogen and trypsin by CTTC was unchanged (not shown). Six mutants (p.K92N, p.Q98K, p.T137M, p.K170E, p.S181G, and p.G208A) autoactivated in the presence of CTTC in a manner that was comparable to wild type (Figure 2B). Surprisingly, however, five mutants (p.P36R, p.G83E, p.I88N, p.V123M and p.S124F) reached markedly reduced trypsin activity levels during autoactivation, suggesting increased susceptibility to CTTC-dependent degradation (Figure 2C). This notion was confirmed by direct degradation experiments which demonstrated that these five mutants were degraded by CTTC at increased rates (Supplementary Figure S2).

Secretion of *PRSS1* variants from HEK 293T cells

To identify trypsinogen variants defective in folding, we evaluated their secretion from transfected cells to the conditioned medium using SDS-PAGE with Coomassie Blue staining, Western blot analysis and trypsin activity measurements after activation with enteropeptidase. The use of HEK 293T cells for cellular secretion studies is a compromise, as efficient transfection of pancreatic acinar cells is not feasible. Despite its constitutive secretion, this cell line remains a good model to study ER related disturbances such as misfolding, retention and degradation of secretory proteins.

Six of 13 mutants tested (p.P36R, p.G83E, p.Q98K, p.V123M, p.T137M, and p.S181G) showed trypsinogen secretion close to wild-type levels (~70–85%) (Figure 3AB, 3C, black bars). Severe secretion defects (~20% of wild type) were observed with mutants p.D100H and p.C139F. Moderate reduction in secretion (~40–50% of wild type) was noted with three mutants (p.K92N, p.S124F and p.G208A). Two mutants (p.I88N and p.K170E) showed increased secretion levels, 140% and 130% of wild type, respectively. The higher secretion of mutant p.I88N was due to the appearance of a second trypsinogen band which represents an aberrantly glycosylated trypsinogen form (Figure 3AB). This band was also observed in a

much fainter form with wild-type trypsinogen and all other variants and may be an artifact of the heterologous cell line used. Although increased secretion of variant p.I88N may represent a gain of function, the rapid degradation of this variant by CTTC (see above) would cancel out this effect and result in a loss-of-function phenotype. On the other hand, it seems probable that the slightly more efficient secretion of mutant p.K170E may confer increased risk for pancreatitis.

For the majority of the variants, trypsin activity levels in the conditioned medium correlated well with the protein levels secreted (Figure 3C, gray bars). Four variants (p.P36R, p.G83E, p.K92N, and p.C139F) had considerably lower enzyme activity relative to their protein levels. Variant p.C139F may be misfolded and catalytically defective, which explains its lower activity, whereas mutants p.P36R and p.G83E suffer some degradation during activation, as seen in the autoactivation experiments in Supplementary Figure S1D. The lower activity of mutant p.K92N is unexplained and may be related to increased sensitivity to some inhibitory component in the cell culture medium, because when this mutant was purified either from *E. coli* or from the conditioned medium of transfected HEK 293T cells, wild-type trypsin activity levels were reached after activation with enteropeptidase.

DISCUSSION

In this study we investigated the functional properties of 13 rare *PRSSI* variants detected in patients with sporadic chronic pancreatitis. The common biochemical phenotype of *PRSSI* mutations associated with hereditary pancreatitis is the generation of greatly increased trypsin levels during autoactivation in the presence of CTTC [6]. This phenomenon is due to increased sensitivity of trypsinogen mutants to CTTC-mediated stimulation of autoactivation and/or resistance to CTTC-dependent trypsinogen degradation (Figure 1). Unexpectedly, only one of the investigated 13 mutants exhibited a similar phenotype when autoactivation was tested in the presence of CTTC. Mutant p.D100H autoactivated at a faster rate and reached higher trypsin levels than wild-type trypsinogen, however, this was independent of CTTC as the same phenotype was observed in its absence. More importantly, further studies, discussed below, demonstrated that mutant p.D100H was poorly secreted, which would completely negate any increased propensity for autoactivation and result in diminished trypsin levels upon activation.

Five mutants were degraded by CTTC at an increased rate resulting in lower trypsin levels during autoactivation, relative to wild-type trypsinogen. This biochemical phenotype is inconsistent with the trypsin-dependent model of hereditary pancreatitis and suggests that these variants are not pathogenic (Table 2). Previous studies found that *PRSSI* mutations p.R116C and p.C139S resulted in reduced trypsinogen secretion [9]. These mutations alter the number of Cys residues and thus may interfere with correct disulfide formation during folding, which results in intracellular retention and degradation. A similar secretion defect, likely caused by mutation-induced misfolding, was observed in the present study for variants p.D100H and p.C139F, and to a smaller extent, variants p.K92N, p.S124F and p.G208A, suggesting that these variants are pathogenic (Table 2). Trypsinogen misfolding underlying the reduced secretion may elicit ER stress and thereby increase the risk for pancreatitis, as previously shown for mutants p.R116C and p.C139S [9] (Figure 1). Secretion defect and ER stress was also observed for a subset of CTTC mutants associated with chronic pancreatitis [12, 19]. Increased secretion of mutant trypsinogens may result in higher trypsinogen levels in the pancreatic juice with consequently increased risk for autoactivation (Figure 1). Copy number mutations in trypsinogen genes may exert their pathogenic effect via this mechanism [20, 21]. Conversely, decreased trypsinogen expression due to the c.1–408C>T polymorphism in the 5' region is associated with decreased pancreatitis risk [22]. In this study we observed slightly increased secretion with two mutants, p.I88N and p.K170E.

While the rapid degradation of mutant p.I88N by CTRC would counteract the effect of higher trypsinogen concentrations, the phenotype of mutant p.K170E may indicate a true gain of function with potentially elevated risk of pancreatitis (Table 2). Finally, three trypsinogen variants (p.Q98K, p.T137M, and p.S181G) proved functionally neutral in this study, indicating that *PRSSI* variants of no clinical significance may be incidental findings when subjects with chronic pancreatitis are screened for underlying genetic defects (Table 2).

In summary, functional analysis of 13 *PRSSI* variants found in patients with chronic pancreatitis demonstrated that these rare *PRSSI* variants do not phenocopy the disease-causing hereditary-pancreatitis associated *PRSSI* mutations. Instead, increased degradation or reduced secretion was observed mostly. Variants with decreased secretion may increase pancreatitis risk through an alternative pathological pathway related to mutation-induced misfolding rather than trypsin activity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

These studies were supported by NIH grants R01DK058088, R01DK082412, R01DK082412-S2 and R01DK095753 (to M.S.-T.). A.S. was also supported by a fellowship from the Rosztoczy Foundation in Boston and A.S. and P.H. were supported by the TÁMOP-4.2.2/B-10/1-2010-0012 grant in Szeged. The authors thank Vinciane Rebours (Beaujon Hospital, Clichy, France) and Jian-Min Chen (Université de Bretagne Occidentale, Brest, France) for providing unpublished information.

References

- Whitcomb DC, Gorry MC, Preston RA, et al. Hereditary pancreatitis is caused by a mutation in the cationic trypsinogen gene. *Nat Genet.* 1996; 14:141–145. [PubMed: 8841182]
- Howes N, Lerch MM, Greenhalf W, et al. Clinical and genetic characteristics of hereditary pancreatitis in Europe. *Clin Gastroenterol Hepatol.* 2004; 2:252–261. [PubMed: 15017610]
- Teich N, Rosendahl J, Tóth M, et al. Mutations of human cationic trypsinogen (*PRSS1*) and chronic pancreatitis. *Hum Mutat.* 2006; 27:721–730. [PubMed: 16791840]
- Rebours V, Boutron-Ruault MC, Schnee M, et al. The natural history of hereditary pancreatitis: a national series. *Gut.* 2009; 58:97–103. [PubMed: 18755888]
- Szmola R, Sahin-Tóth M. Uncertainties in the classification of human cationic trypsinogen (*PRSS1*) variants as hereditary pancreatitis-associated mutations. *J Med Genet.* 2010; 47:348–350. [PubMed: 20452997]
- Szabó A, Sahin-Tóth M. Increased activation of hereditary pancreatitis-associated human cationic trypsinogen mutants in presence of chymotrypsin C. *J Biol Chem.* 2012; 287:20701–20710. [PubMed: 22539344]
- Nemoda Z, Sahin-Tóth M. Chymotrypsin C (caldecrin) stimulates autoactivation of human cationic trypsinogen. *J Biol Chem.* 2006; 281:11879–11886. [PubMed: 16505482]
- Szmola R, Sahin-Tóth M. Chymotrypsin C (caldecrin) promotes degradation of human cationic trypsin: identity with Rinderknecht's enzyme Y. *Proc Natl Acad Sci USA.* 2007; 104:11227–11232. [PubMed: 17592142]
- Kereszturi E, Szmola R, Kukor Z, et al. Hereditary pancreatitis caused by mutation-induced misfolding of human cationic trypsinogen: a novel disease mechanism. *Hum Mutat.* 2009; 30:575–582. [PubMed: 19191323]
- Király O, Guan L, Szepessy E, et al. Expression of human cationic trypsinogen with an authentic N terminus using intein-mediated splicing in aminopeptidase P deficient *Escherichia coli*. *Protein Expr Purif.* 2006; 48:104–111. [PubMed: 16542853]

11. Király O, Guan L, Sahin-Tóth M. Expression of recombinant proteins with uniform N-termini. *Methods Mol Biol.* 2011; 705:175–194. [PubMed: 21125386]
12. Beer S, Zhou J, Szabó A, et al. Comprehensive functional analysis of chymotrypsin C (CTRC) variants reveals distinct loss-of-function mechanisms associated with pancreatitis risk. *Gut.* 2012 Sep 1. [Epub ahead of print].
13. Szabó A, Héja D, Szakács D, et al. High affinity small protein inhibitors of human chymotrypsin C (CTRC) selected by phage display reveal unusual preference for P4' acidic residues. *J Biol Chem.* 2011; 286:22535–22545. [PubMed: 21515688]
14. Chen JM, Piepoli Bis A, Le Bodic L, et al. Mutational screening of the cationic trypsinogen gene in a large cohort of subjects with idiopathic chronic pancreatitis. *Clin Genet.* 2001; 59:189–193. [PubMed: 11260229]
15. Teich N, Bauer N, Mössner J, et al. Mutational screening of patients with nonalcoholic chronic pancreatitis: identification of further trypsinogen variants. *Am J Gastroenterol.* 2002; 97:341–346. [PubMed: 11866271]
16. Chang YT, Wei SC, LPC, et al. Association and differential role of PRSS1 and SPINK1 mutation in early-onset and late-onset idiopathic chronic pancreatitis in Chinese subjects. *Gut.* 2009; 58:885. [PubMed: 19433603]
17. Zeng K, Liu QC, Lin JH, et al. Novel mutations of *PRSS1* gene in patients with pancreatic cancer among Han population. *Chin Med J (Engl).* 2011; 124:2065–2067. [PubMed: 22088471]
18. Lee YJ, Kim KM, Choi JH, et al. High incidence of PRSS1 and SPINK1 mutations in Korean children with acute recurrent and chronic pancreatitis. *J Pediatr Gastroenterol Nutr.* 2011; 52:478–481. [PubMed: 21415673]
19. Szmola R, Sahin-Tóth M. Pancreatitis-associated chymotrypsinogen C (CTRC) mutant elicits endoplasmic reticulum stress in pancreatic acinar cells. *Gut.* 2010; 59:365–372. [PubMed: 19951900]
20. Le Maréchal C, Masson E, Chen JM, et al. Hereditary pancreatitis caused by triplication of the trypsinogen locus. *Nat Genet.* 2006; 38:1372–1374. [PubMed: 17072318]
21. Masson E, Le Maréchal C, Chandak GR, et al. Trypsinogen copy number mutations in patients with idiopathic chronic pancreatitis. *Clin Gastroenterol Hepatol.* 2008; 6:82–88. [PubMed: 18063422]
22. Whitcomb DC, Larusch J, Krasinskas AM, et al. Common genetic variants in the CLDN2 and PRSS1-PRSS2 loci alter risk for alcohol-related and sporadic pancreatitis. *Nat Genet.* 2012; 44:1349–1354. [PubMed: 23143602]
23. Keiles S, Kammesheidt A. Identification of *CFTR*, *PRSS1* and *SPINK1* mutations in 381 patients with pancreatitis. *Pancreas.* 2006; 33:221–227. [PubMed: 17003641]
24. Tautermann G, Ruebsamen H, Beck M, et al. R116C mutation of cationic trypsinogen in a Turkish family with recurrent pancreatitis illustrates genetic microheterogeneity of hereditary pancreatitis. *Digestion.* 2001; 64:226–232. [PubMed: 11842279]
25. Rosendahl J, Landt O, Bernadova J, et al. CFTR, SPINK1, CTRC and PRSS1 variants in chronic pancreatitis: is the role of mutated CFTR overestimated? *Gut.* 2012 Mar 17. [Epub ahead of print]. 10.1136/gutjnl-2011-300645
26. Sultan M, Werlin S, Venkatasubramani N. Genetic prevalence and characteristics in children with recurrent pancreatitis. *J Pediatr Gastroenterol Nutr.* 2012; 54:645–650. [PubMed: 22094894]
27. Corleto VD, Gambardella S, Gullotta F, et al. New PRSS1 and common CFTR mutations in a child with acute recurrent pancreatitis, could be considered an “Hereditary” form of pancreatitis? *BMC Gastroenterol.* 2010; 10:119. [PubMed: 20950468]

Summary box

What is already known about this subject

- Hereditary pancreatitis is caused by mutations in human cationic trypsinogen (*PRSSI*) that increase autoactivation.
- Some pathogenic *PRSSI* variants misfold, suffer intracellular retention and cause endoplasmic reticulum stress.
- Clinical relevance of rare *PRSSI* variants found in subjects with sporadic chronic pancreatitis is uncertain but frequently assumed.

What are the new findings

- Rare *PRSSI* variants found in sporadic chronic pancreatitis do not increase trypsinogen activation.
- Rather they cause increased degradation, loss of secretion or no functional change.
- Reduced secretion is indicative of misfolding that may increase pancreatitis risk through endoplasmic reticulum stress.

How might it impact on clinical practice in the foreseeable future?

- Functional classification of *PRSSI* variants found in patients with chronic pancreatitis is required for determination of clinical relevance.

Figure 1.

Pathological pathways associated with *PRSS1* mutations in hereditary and sporadic chronic pancreatitis. Mutations in *PRSS1* can increase autoactivation of cationic trypsinogen by different mechanisms: increased trypsinogen expression or secretion; inhibition of chymotrypsin C (CTRC)-dependent trypsinogen degradation, stimulation of N-terminal processing of the trypsinogen activation peptide by CTRC; and direct stimulation of autoactivation. Alternatively, *PRSS1* mutations can cause misfolding and endoplasmic reticulum (ER) stress. See text for further details.

Figure 2.

Autoactivation of cationic trypsinogen variants in the presence of chymotrypsin C (CTRC). Trypsinogen was incubated at 1 μ M concentration with 5 nM CTRC and 10 nM initial trypsin in 0.1 M Tris-HCl (pH 8.0), 1 mM CaCl_2 and 0.05% Tween-20 (final concentrations) at 37° C. At the indicated times, 2 μ L aliquots were removed and trypsin activity was measured as described in *Experimental Procedures*. Trypsin activity was expressed as percent of the maximal wild-type activity in the absence of CTRC. **A**, mutant with increased autoactivation, **B**, mutants with unchanged autoactivation; **C**, mutants with decreased autoactivation.

Figure 3.

Secretion of cationic trypsinogen variants from HEK 293T cells. Cells were transiently transfected and conditioned media were collected after 24 h. Trypsinogen protein levels were qualitatively analyzed on Coomassie Blue stained SDS-polyacrylamide gels (**A**) and then quantitatively determined by Western blots (**B**) and densitometry (**C**, black bars). Trypsin activity was measured after activation with enteropeptidase (**C**, gray bars). See *Experimental Procedures* for details. Representative gels and blots of four independent experiments are shown. Average of four experiments with SD was plotted. Trypsinogen protein and trypsin activity levels were expressed as percent of wild-type values.

Table 1

Rare *PRSS1* variants in pancreatitis.

Region	Nucleotide change	Amino acid change	Carriers reported	Clinical and genetic information	Citations
Exon 2	c.107C>G	p.P36R	2	French m and fm, ICP	[14] [#]
Exon 3	c.248G>A	p.G83E	1	Italian, ICP	[14] [#]
Exon 3	c.263T>A	p.I88N	1	18 y white fm, CP, no <i>SPINK1</i> or <i>CFTR</i> mut	[23]
Exon 3	c.276G>T	p.K92N	1	Italian, ICP	[14] [#]
Exon 3	c.292C>A	p.Q98K	1	55 y French m, CP, no <i>SPINK1</i> , <i>CTRC</i> or <i>CFTR</i> mut	[4] [*]
Exon 3	c.298G>C	p.D100H	1	German m, ICP	[24]
Exon 3	c.367G>A	p.V123M	1	French fm, ICP	[14] [#]
Exon 3	c.371C>T	p.S124F	1	46 y German fm, ICP, no <i>SPINK1</i> , <i>CTRC</i> or <i>CFTR</i> mut	[25]
Exon 3	c.410C>T	p.T137M	3	13 y Asian fm, AP, no <i>SPINK1</i> or <i>CFTR</i> mut	[23]
Exon 3	c.416G>T	p.C139F	2	Chinese, late onset ICP, no <i>SPINK1</i> mut	[16]
				28 y Chinese, pancreatic cancer, also carried p.T135A <i>PRSS1</i>	[17]
				14 y German m, RAP, unaffected mother carrier	[15]
Exon 4	c.508A>G	p.K170E	1	8 y fm, CP with chronic pain, no <i>SPINK1</i> or <i>CFTR</i> mut	[26]
				30 y Indian m, CP, no <i>SPINK1</i> , <i>CTRC</i> or <i>CFTR</i> mut	[4] [*]
				6 y Italian m, RAP, p.F508del <i>CFTR</i> , unaffected mother carrier	[27]
Exon 5	c.623G>C	p.G208A	3	12 y Asian m, pancreatitis, hm p.F508del and p.Q1352H <i>CFTR</i>	[23]
				Korean child, RAP, no <i>SPINK1</i> mut	[18]
				7 y Korean, necrotizing AP	[18]

AP, acute pancreatitis, CP, chronic pancreatitis; ICP, idiopathic chronic pancreatitis, RAP, recurrent acute pancreatitis; m, male; fm, female; hm, homozygous, y, year, mut, mutation. Unless indicated otherwise, mutations are heterozygous. The age indicates the time of diagnosis.

* Vinciane Rebours, personal communication.

[#] Jian-Min Chen, personal communication.

Table 2

Clinical relevance of *PRSSI* variants classified on the basis of functional phenotype.

Region	Nucleotide change	Amino acid change	Relevant phenotype
Pathogenic variants, strong			
Exon 3	c.298G>C	p.D100H	severe secretion defect
Exon 3	c.416G>T	p.C139F	severe secretion defect
Pathogenic variants, mild			
Exon 3	c.276G>T	p.K92N	moderate secretion defect
Exon 3	c.371C>T	p.S124F	moderate secretion defect
Exon 4	c.508A>G	p.K170E	moderate secretion increase
Exon 5	c.623G>C	p.G208A	moderate secretion defect
Non-pathogenic variants, functionally neutral			
Exon 3	c.292C>A	p.Q98K	none
Exon 3	c.410C>T	p.T137M	none
Exon 4	c.541A>G	p.S181G	none
Non-pathogenic variants, loss of function			
Exon 2	c.107C>G	p.P36R	degradation by CTRC
Exon 3	c.248G>A	p.G83E	degradation by CTRC
Exon 3	c.263T>A	p.I88N	degradation by CTRC
Exon 3	c.367G>A	p.V123M	degradation by CTRC