ERRATUM

Erratum to: Risk Factors for Nosocomial Gastrointestinal Bleeding and Use of Acid-Suppressive Medication in Non-Critically Ill Patients

Shoshana J. Herzig, MD, MPH¹, Michael B. Rothberg, MD, MPH², David B. Feinbloom, MD¹, Michael D. Howell, MD, MPH³, Kalon K. L. Ho, MD, MSc⁴, Long H. Ngo, PhD¹, and Edward R. Marcantonio, MD, SM^{1,5}

¹Division of General Medicine and Primary Care, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA, USA;

²Department of Internal Medicine, Medicine Institute, Cleveland Clinic, Cleveland, OH, USA; ³Division of Pulmonary and Critical Care, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA, USA; ⁴Division of Cardiology, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA, USA; ⁵Division of Gerontology, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA, USA.

J Gen Intern Med 28(7):978

DOI: 10.1007/s11606-013-2425-1

© Society of General Internal Medicine 2013

Erratum to: J Gen Intern Med

DOI: 10.1007/s11606-012-2296-x

In Table 4 of the original publication, the third column of the first row of the table (number with no bleeding amongst those patients in the “Low risk” category) should read 57,317 rather than 5,737.

The authors regret this error.

Corresponding Author: Shoshana J. Herzig, MD, MPH; Division of General Medicine and Primary Care, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA, USA (e-mail: sherzig@bidmc.harvard.edu)

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11606-012-2296-x>.

Published online April 16, 2013