Published in final edited form as:

World Health Popul. 2013 ; 14(2): 12–23.

Cancer Registration Needs Assessment at a Tertiary Medical Center in Kilimanjaro, Tanzania

Leah L. Zullig, MPH¹, Charles Muiruri, MPH^{1,2}, Amy Abernethy, MD³, Bryan J. Weiner, PhD, MA¹, John Bartlett, MD², Olola Oneko, MD⁴, and S. Yousuf Zafar, MD, MS^{2,3}

¹Department of Health Policy and Management, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC

²Duke Global Health Institute, Duke University, Durham, NC

³Duke Cancer Institute, Duke University, Durham, NC

⁴Kilimanjaro Christian Medical Centre, Moshi, Tanzania

Abstract

Cancer burden is increasing in Africa more any other continent, but population-based tracking of cancer incidence is incomplete. Cancer registries can improve understanding of cancer incidence.

To assess organizational readiness to sustain registry development, we conducted a survey assessing change efficacy, resource availability, and change commitment at an academic hospital in Moshi, Tanzania. Fifty-two surveys were returned (80% response rate). There was strong reliability among change efficacy and commitment survey items with Cronbach's alphas of 0.93 and 0.77, respectively.

Clinicians, nurses, and administrators conveyed similar responses regarding change efficacy. Clinicians had similar responses for change commitment. Echoing many low- and middle-income countries, approximately one-third of respondents indicated there were no funds to maintain the registry and funds were not obtainable. For most resources respondents felt that resources were sufficient or attainable. Respondents were generally confident and committed to registry implementation. Lessons learned at KCMC may be more broadly relevant.

Keywords

Tanzania; needs assessment; registries; neoplasms

INTRODUCTION

Cancer and cancer registration in Africa

The burden of cancer in sub-Saharan Africa is growing. Over the next ten years, the burden of cancer and chronic disease will increase in Africa more than in than any other continent (de-Graft Aikins et al., 2010). By best estimates, the cumulative incidence of cancer (excluding Kaposi's sarcoma and non-melanoma skin cancer) in Zimbabwe, for example, is only about 30% less than that of France, but the cancer mortality is almost double the cancer mortality in France (Parkin et al., 2008). These data suggest that, in Africa, both the

Corresponding Author: S. Yousuf Zafar, MD, MHS, yousuf.zafar@duke.edu, Phone: 919.684.0138, Address: Duke Cancer Institute | DUMC 3505, Durham, NC 27710.

Conflicts of interest: The authors have no conflicts of interest to disclose.

incidence of and mortality due to cancer are severely underestimated. An important first step in improving these estimates involves the establishment of reliable, population-based cancer registries. Unfortunately, in most of Africa the population-based tracking of cancer incidence has been incomplete; resources are limited, knowledge of cancer trends in this area is inadequate, and trends are based mostly on case series (Orem & Wabinga, 2009; Parkin, et al., 2008).

Models of capacity building to support cancer registry development are a chief issue for low- and middle-income countries. In fact, the International Agency for Research on Cancer (IARC) has a Global Initiative for Cancer Registry Development in low- and middle-income countries that aims to create capacity to produce reliable and high quality information on the burden of cancer to inform effective cancer control policies. The U.S.-based National Cancer Institute (NCI) recently issued a report seeking to facilitate lessons learned from IARC and other groups to provide technical assistance and training to foster development of cancer registries in low- and middle-income countries (National Cancer Institute, 2012). Our approach for the assessment of organizational readiness to change in preparation for cancer registry development supports this aim.

Assessing the perceptions of key stakeholders presents a critical foundation for establishing a sustainable population-based cancer registry. In fact, experts in organizational change have found that readiness to change is an important predictor of successful implementation (Kotter, 1996; Weiner, Amick, & Lee, 2008). Without a clear understanding of organizational perceptions and readiness to change, the development of a cancer registry might not be aligned with the organization's values and needs. Organizational readiness to change is a multifaceted issue comprised of two chief constructs –organizational members' change commitment and change efficacy to implement organizational change. Change is comprised of intertwining levels, including both individual and organizational levels. When an individual believes that they can successfully make a change that change is personally beneficial, they may be more likely to support an organizational change (Holt, et al., 2010). Moreover, assessing individuals' readiness to change and their perceived needs are critical for a sustainable foundation for population-based cancer registry development. Our objective was to assess shared beliefs among stakeholders in the organizational ability to initiate and maintain cancer registration activities at a tertiary referral medical center in Moshi, Tanzania. In doing so we present a model for ensuring sustainable cancer registry development in low- and middle-income countries (LMICs).

METHODS

Kilimanjaro Christian Medical Centre

Located in northeastern Tanzania, Kilimanjaro Christian Medical Centre (KCMC) was founded in 1971 by the Good Samaritan Foundation and has been in continuous operation for over four decades. KCMC is a tertiary referral center, with approximately 560 beds, serving a patient population of over 11 million people living in the Kilimanjaro region (Kilimanjaro Christian Medical Centre, 2012). In addition to providing patient care, in 1996 KCMC was expanded to include medical training with the establishment of Tumaini University/Kilimanjaro Christian Medical College (Kilimanjaro Christian Medical Centre, 2012).

KCMC has a history of cancer registration, although data collection has not been systematic and the quality and completeness of registry data have not been documented. At the time of manuscript preparation, KCMC cancer registration activities were inactive. Because of the increasing cancer burden in eastern Africa and the need for high quality, comprehensive data

to inform resource planning, there has been renewed interest in expanding KCMC cancer registration activities.

Survey Design

We conducted a self-administered needs assessment survey at KCMC over a two-week period in June 2012. The survey contained 41 items assessing change commitment, change efficacy (Gist & Mitchell, 1992; Weiner, 2009), and determinants of change efficacy such as resource availability. The survey contained items assessing change commitment. Survey items included, “How committed is your facility to implementing this change?” and “How willing is your facility to work hard to implement this change?” among others. Response options included: “not at all,” “a little,” “fairly,” “very,” and “don’t know.”

The survey also assessed change efficacy. Change efficacy refers to organizational members’ shared organizational confidence in the communal ability to implement a change effectively – in this case further development of a cancer registration program. Examples of change efficacy items are, “Can the organization assign the right tasks to the right people in implementing this change?” and “Can the organization effectively use the resources that are currently available to implement the program?” Change efficacy items used in the survey were rated on a 5-point Likert scale: “not at all confident,” “a little confident,” “fairly confident,” “very confident,” and “don’t know.”

The survey addressed two determinants of change efficacy – resource availability and situational factors (Weiner, 2009). Resource availability refers to organizational members’ perceptions about the availability and obtainability of resources. This includes an assessment of material, financial, and human resources needed to advance the cancer registry. Respondents were asked about resources including pathologists, clinical staff to collect patient data, computers, and software, among others. Response options for resource availability items included: “have none and resource is not available,” “have insufficient, but resource is attainable,” “already have sufficient resource,” “not necessary,” and “don’t know.” In addition to these discrete items, respondents were also given the opportunity to answer open-ended questions with free responses. Regarding situational factors, respondents were asked whether hospital staff has sufficient time and political support to make needed changes.

Sampling and Recruitment

Because our goal was to obtain a comprehensive, organizational perspective we conducted non-probability, purposive sampling (Trochim & Donnelly, 2006). To maximize organizational leaders’ support we initially met with the hospital’s executive director and subsequently contacted the department chairs of obstetrics and gynecology, pediatrics, urology, surgery, internal medicine, medical records, and nursing. Department chairs provided additional contacts and facilitated in obtaining responses from their unit. We attended clinical meetings such as morning report to increase survey awareness and participation. In accordance with respondents’ preference, surveys were administered on paper as well as electronically via email.

Statistical Analysis

Frequencies were tabulated and, where appropriate, inconsistencies were checked against the original surveys for manual correction if required. Missing variables and incomplete questions were coded “missing.” Missing values were examined for patterns in reporting and correlations with professional role. Data were first analyzed for descriptive outcomes, including proportions, means, medians, and ranges. Chronbach’s alphas were examined as a measure of reliability among the change efficacy and change commitment survey items.

Bivariate associations among clinical role, resource availability, and change efficacy survey items were reported using chi-squared and Fisher's exact tests as appropriate.

After examining raw change efficacy survey responses, survey responses were collapsed (e.g., "not at all confident" and "a little confident" were merged) and re-examined descriptively. Survey items were then classified into two categories – organizational management versus motivation. Within each category, survey items were ordered from highest percentage of confidence to lowest percentage of confidence.

All analyses were conducted using Stata (StataCorp. 2011. *Stata Statistical Software: Release 12*. College Station, TX: StataCorp LP). Duke University Health System's Institutional Review Board reviewed and exempted this study. The Research Ethics Committee at Kilimanjaro Christian Medical Centre and Tumaini University Kilimanjaro Christian Medical College approved this study.

RESULTS

Response Rate and Respondent Characteristics

A total of 64 surveys were administered (57 paper-based surveys delivered to possible participants; 7 electronic surveys emailed). Fifty-two surveys were returned for a response rate of approximately 80%. Respondent characteristics are presented in Table 1.

Missing Values

Data were examined for systematic patterns of missing values. No patterns were detected. No association was found between professional role and missing value. We examined associations between year of birth, a proxy measure for years of professional experience, and missing values; no association was found.

Change Commitment

In addition to examining individuals' confidence in the organization's readiness to change, respondents were asked about the organization's commitment, motivation, and willingness to work hard to implement the cancer registration program (Table 3). Cronbach's alpha for the change commitment items were 0.77, indicating acceptable reliability. Regarding both commitment to the cancer registry and willingness to work hard to implement the registry, approximately half of respondents reported that the organization is very committed/willing. However, only 37% (n=19) reported that the organization was very motivated to implement the cancer registry.

Change Efficacy

Cronbach's alpha for the change efficacy items was 0.93, indicating strong reliability. A comprehensive list of the survey items assessing change efficacy are described in Table 2. Forty percent (n=21) of respondents indicated that they were very confident that their organization could effectively use the resources that are currently available in order to implement the cancer registry. Over half of respondents (56%, n=19) indicated that they were very confident that clinicians could be encouraged to use the cancer registry. Similarly, 42% (n=22) reported that their organization could effectively coordinate the efforts of those involved in implementing the cancer registry. Respondents were divided regarding their perceptions of the organization's ability to communicate clearly regarding implementation of a cancer registry. Slightly fewer than half (44%, n=23) of respondents believed that the organization could maintain the momentum going in order to implement the cancer registry. Approximately one-third of respondents were very confident and fairly confident (35%,

n=18; 37%, n=19, respectively) that the organization can keep track of progress in implementing the registry.

Perceptions of Organizational Motivation versus Management

The change efficacy survey items were collapsed and categorized (e.g., “fairly confident” and “very confident” were merged; classified into motivation versus management domains). There was a clear threshold of confidence/agreement between the organizational motivation versus management domains (Table 4). For example, being fairly or very confident within the management domain survey items ranged from 64% to 77%. The mean percentage response of fairly or very confident across these survey items was 73%. In contrast, respondents reported being fairly or very confident (or similar) for between 81% and 89% of motivational survey items.

Resource Availability

Respondents were asked to report about perceived availability of resources needed for development of the cancer registry (Table 5). When asked about the availability of pathologists, approximately half of respondents indicated they “have none and resource is not attainable.” Respondents were asked about the availability of staff to collect, enter, follow-up on, and audit cancer registry patient data. For each of these items, most respondents reported that the organization has insufficient staff, but the resource is attainable; specifically, staff to collect patient data (44%, n=23), staff to enter patient data (46%, n=24), staff to follow-up on registered patients (35%, n=18), and staff to audit the registry (44%, n=23). The resource assessment also inquired about the availability of computers, software, and staff with computing expertise. In each of these categories, the most common response category was that the resource was insufficient, but attainable. Respondents were also asked about the availability of funds to maintain the registry. Interestingly, answers were more evenly distributed across response categories for this survey item than for any other.

Bivariate Analysis

Due to the small sample size, we utilized Fisher’s exact test to examine whether professional roles (e.g., physician, administrator) was associated with differences in perception regarding availability of resources. For the majority of resources, there was no association between professional role and perception of resource availability. However, there was an association between professional role and availability of pathology technicians ($p=0.04$). Further, we assessed differences in reported change efficacy and professional role. For each of the 13 change efficacy questions, there was no association between reported confidence in an organization’s ability to implement the registry and professional role.

We then examined whether specialty (e.g., internal medicine, surgery) was associated with reported change efficacy. Out of 13 items, differences between specialties were found for only 4 measures. Significant differences were found for the following items: support clinicians as they adjust clinical practice in response to the program ($p=0.01$), keep the momentum going in implementing this change ($p=0.00$), assign the right tasks to the right people ($p=0.04$), and coordinate tasks so that implementation goes smoothly ($p=0.02$).

DISCUSSION

Overall, respondents reported a high degree of change commitment and change efficacy, suggesting that KCMC clinicians and administrators expressed confidence in their shared ability to organize and implement the process needed to develop and implement the cancer registry. Furthermore, responses were consistent among people of varied professional roles;

clinicians, nurses, trainees, and administrators all conveyed similar responses regarding their organization's readiness and ability to implement systematic cancer registration. In general there was also agreement between professionals of different medical specialties. Readiness to change and organizational endorsement are strong predictors of success in making a change (Kotter, 1996; Sweeney & Whitaker, 1994; Weiner, 2009). This consensus among a multidisciplinary team suggests that KCMC is an organization with a drive to change. While each institution in low- and middle-income countries is unique, identifying strengths and concerns at KCMC may inform other future cancer registration projects in similar environments.

The two measures with the highest reported levels of confidence were to encourage clinicians to try using the registry (e.g., refer patients to the registry) and, secondly, to support clinicians as they adjust their clinical practice in response to the registry. Comparatively lower levels of confidence were reported for three measures: 1) effectively solve problems that might arise, 2) keep track of how things are going, and 3) manage the politics of implementing this program. This suggested that there was perceived support for clinicians, but respondents were not as confident about the organization's ability to manage change throughout implementation. When we collapsed and classified the change efficacy items into two categories – organizational motivation versus management – a clear threshold presented itself. Over 80% responded positively (e.g. “fairly confident” or “very confident”) for each of the motivational survey items; respondents indicated that KCMC was very committed to registry implementation, very willing to work hard to implement it, and wanted very much to implement the registry. However, when asked about issues of management such as task coordination and problem solving, respondents responded positively far less frequently.

This distinction between organizational motivation and task management is important to understand, particularly in the context of cancer registry development in low- and middle-income countries. In learning healthcare organizations, managers must extend beyond traditional management roles and must serve as designers and teachers (Hernandez, Kaluzny, & Haddock, 2000). This may be particularly true in the resource-limited settings where managers must be innovative. Organizations must communicate goals and values to their employees in an effort to build a sense of organizational belonging. To that end, it is essential those internal and external stakeholders are involved early in the implementation process and that there is organizational endorsement. Relevant stakeholders within low- and middle-income countries must be identified and their attitudes toward the change, such as cancer registry implementation, must be assessed (Hernandez, et al., 2000). Moreover, it is critical that the political climate both internal and external to the organization is aligned well to support cancer registration activities. There must be clear delineation of responsibilities, an understanding of decision-making patterns, and a balance of tasks between employees (Charns & Gittel, 2000; Hernandez, et al., 2000).

Despite the high level of change efficacy, limited resources are available for all human resources except pathologists (e.g., staff to collect and enter patient data), the largest percentage of respondents indicated that currently the resource was insufficient but that it was obtainable. We surveyed a wide variety of professionals ranging from medical records technicians to hospital administrators. Respondents' professional role may impact their knowledge of certain resources. For example, a medical records technician may not be as knowledgeable about the availability of funds to maintain the registry compared to the hospital administrator. Interestingly, we found similar responses across professional roles for most items. From a change management perspective, the assessment of perception of resource availability may be particularly important. If resources are truly unavailable, change efficacy may be undermined. Nearly half of respondents reported that they had no

pathologists and none were obtainable. However, pathology residents from KCMC are currently undergoing training to return as staff pathologists; clearly, a communication gap exists between perception and actual availability of resources. Despite this communication gap, the assumption of KCMC staff is likely based on the known shortage of trained pathologists in sub-Saharan Africa (Awadelkarim, Mohamedani, & Barberis, 2012). For example, the entire country of Tanzania has only 15 pathologists; this equates to one pathologist per 2.5 million people (Okesina, 2009; Rambau, 2011). This shortage of pathologists needs to be addressed on a broader scale. A potential solution to shortage of trained pathologists involves telepathology, where pathologic slides are prepared and images are transmitted to pathologists located off-site (Hazin & Qaddoumi, 2010), and working partnerships with high-income countries (Knaul et al., 2012).

Slightly more than one-third of respondents indicated that there were no funds to maintain the registry and funds were not obtainable. This is a sentiment that is echoed in many LMICs, so lessons learned at KCMC may be more broadly relevant. Allocating scarce resources to a cancer registry can create additional strain in a system already stretching funds to provide appropriate health care. However, an initial investment in cancer registration is critical to understanding the disease burden, and without understanding disease burden, appropriate resources cannot be allocated to cancer control. In order to ensure sustainable funding for population-based cancer registries, those leading cancer registration should early on involve health ministry officials in their respective countries (Anderson et al., 2011). Furthermore, long-term partnerships between high- and low-income countries have led to successful implementation of cancer control and treatment programs (Carlson et al., 2010). In order to bridge the resource gap, smaller-scale, hospital-based cancer registries can be initially developed and later scaled-up for more comprehensive population coverage. Information obtained in our survey support the aims of NCI and other organizations aimed at facilitating lessons learned in low- and middle-income settings for broader dissemination in the implementation of cancer registry development (National Cancer Institute, 2012). To that end, we posit that conducting a needs assessment is a critical early-stage component when beginning or expanding a cancer registry program, particularly in LMIC's where resources are scarce. Perspectives gained from a survey like ours may be useful in conducting intra-institutional educational programs and/or prioritizing resource acquisition. While each country and institution is unique, our survey design and methodology may be suitable for other contexts.

There are a number several successful cancer registries in LMICs that serve as an exemplar for aspiring cancer registration program like KCMC. An example of one such registry is the Gharbiah regional cancer registry in Egypt. For well over a decade the Gharbiah registry has achieved over 90% population coverage. Data from the Gharbiah registry have informed several publications describing patterns of cancer in the region, including a monograph describing cancer incidence in several Middle Eastern countries and comparing that information with the U.S.-based Surveillance Epidemiology and End Results (SEER) cancer registry (Freedman, Edwards, Ries, & Young, 2006). There are two unique attributes of the Gharbiah registry that may have facilitated its success. First, in 1996 Ministers of Health in Egypt, Israel, Jordan, Cyprus, and the Palestinian Authority established the Middle East Cancer Consortium (Middle East Cancer Consortium, 2010). These registries use common software called CANREG, which was developed by IARC. Secondly, the registry is affiliated with Cairo University. These strategic alliances bolster resources and provide collaborative partnerships, which will help developing cancer registry programs in low- and middle-incomes, succeed. KCMC is uniquely positioned in its affiliations with Tumaini University and Kilimanjaro Christian Medical College. Despite this, respondents reported a perception that their organization is generally supportive of change regarding the cancer registry program, but lacks many tools needed to expand. This information shapes our

approach in terms of interaction with KCMC staff and registry expansion. Though not geographically proximal to KCMC, Dar es Salaam in Tanzania is also home to referral cancer center. KCMC may partner with local institutions and those in higher income countries to increase its capacity for cancer registration.

This study has several limitations. KCMC is a single institution, and survey results from KCMC may not be generalizable to other organizations. Surveys were self-administered. While no identifying information was collected, it is possible that respondents were hesitant to disclose responses that might reflect negatively on their organization. Given the consistency of responses across surveys, this is not a significant concern. We utilized a cross-sectional survey technique. Change efficacy and readiness to change will evolve longitudinally. It is possible that, even within a single institution, results would differ if measured over time. Because of the two-week survey administration time period and the sampling strategy, it is possible that important potential survey participants were not contacted. Effort was made to mitigate this limitation by sending electronic, emailed surveys.

In conclusion, administrators and medical staff at a large referral center in sub-Saharan Africa believed strongly in the utility of cancer registration. However, they expressed concerns regarding registry management and sustainability. These are valid concerns that likely translate across borders to other institutions or regions in LMICs looking to develop cancer registries. Identifying such strengths and concerns prior to initiating cancer registration will help focus resources and improve chances of long-term success.

Acknowledgments

Funding: Ms. Zullig is funded by the National Cancer Institute (5R25CA116339). Dr. Zafar is supported by the American Cancer Society Mentored Research Scholar Grant and the Duke Cancer Institute Cancer Prevention, Detection and Control Research Pilot Studies Award. Dr. Bartlett is supported by HRSA grant T84HS21123 and NIH grants P30AI064518, D43CA153722, U01AI069484 and D43TW06732. This project was also supported by the HIV Associated Malignancies Training and Research Program (D43 CA153722) and Medical Education Partnership Initiative (T84HA21123).

References

- Anderson BO, Cazap E, El Saghir NS, Yip CH, Khaled HM, Otero IV, Harford JB. Optimisation of breast cancer management in low-resource and middle-resource countries: executive summary of the Breast Health Global Initiative consensus, 2010. *Lancet Oncol.* 2011; 12(4):387–398.10.1016/S1470-2045(11)70031-6 [PubMed: 21463833]
- Awadelkarim KD, Mohamedani AA, Barberis M. Role of pathology in sub-Saharan Africa: An example from Sudan. *Pathology and Laboratory Medicine International.* 2012; 2:49–57.
- Carlson JW, Lyon E, Walton D, Foo WC, Sievers AC, Shulman LN, Milner DA Jr. Partners in pathology: a collaborative model to bring pathology to resource poor settings. *Am J Surg Pathol.* 2010; 34(1):118–123.10.1097/PAS.0b013e3181c17fe6 [PubMed: 19898229]
- Charns, MP.; Gittel, JH. Work Design. In: Shortell, SM.; Kaluzny, AD., editors. *Health Care Management: Organization Design and Behavior.* 4. Albany, NY: Delmar; 2000.
- de-Graft Aikins A, Unwin N, Agyemang C, Allotey P, Campbell C, Arhinful D. Tackling Africa's chronic disease burden: from the local to the global. *Global Health.* 2010; 6:5.10.1186/1744-8603-6-5 [PubMed: 20403167]
- Freedman, LS.; Edwards, BK.; Ries, LAG.; Young, JL. NIH Pub No 06-5873. Bethesda, MD: 2006. Cancer Incidence in Four Member Countries (Cyprus, Egypt, Israel, and Jordan) of the Middle East Cancer Consortium (MECC) Compared with US SEER.
- Gist ME, Mitchell TR. Self-Efficacy - a Theoretical-Analysis of Its Determinants and Malleability. *Academy of Management Review.* 1992; 17:183–211.

- Hazin R, Qaddoumi I. Teleoncology: current and future applications for improving cancer care globally. *Lancet Oncol.* 2010; 11(2):204–210.10.1016/s1470-2045(09)70288-8 [PubMed: 20152772]
- Hernandez, SR.; Kaluzny, AD.; Haddock, CC. Organizational Innovation, Change, and Learning. In: Shortell, SM.; Kaluzny, AD., editors. *Health Care Management: Organization Design and Behavior.* 4. Albany, NY: Delmar; 2000.
- Holt DT, Helfrich CD, Hall CG, Weiner BJ. Are you ready? How health professionals can comprehensively conceptualize readiness for change. *J Gen Intern Med.* 2010; 25(Suppl 1):50–55.10.1007/s11606-009-1112-8 [PubMed: 20077152]
- Kilimanjaro Christian Medical Centre. Kilimanjaro Christian Medical Centre; 2012. Retrieved July 7, 2012, from <http://www.kcmc.ac.tz/index.html>
- Knaul FM, Shulman LN, Gralow J, Quick J, Bhadelia A, Frenk J. Improving pathology for better cancer care and control in countries of low and middle income. *Lancet.* 2012; 379(9831): 2052.10.1016/s0140-6736(12)60894-0 [PubMed: 22656884]
- Kotter, JP. *Leading change.* Boston: 1996.
- Middle East Cancer Consortium, T. M. E. C. Welcome to the Middle East Cancer Consortium (MECC). 2010. Retrieved Oct. 12, 2012, from <http://mecc.cancer.gov>
- National Cancer Institute. *Setting Priorities for Global Cancer Research: NCI Programs in Global Health 2010–2011.* Bethesda, MD: 2012.
- Okesina A. Pathology in West Africa: problems of training and practice. *The Bulletin of The Royal College of Pathologists.* 2009:306–308.
- Orem J, Wabinga H. The roles of national cancer research institutions in evolving a comprehensive cancer control program in a developing country: experience from Uganda. *Oncology.* 2009; 77(5): 272–280.10.1159/000259258 [PubMed: 19923865]
- Parkin DM, Sitas F, Chirenje M, Stein L, Abratt R, Wabinga H. Part I: Cancer in Indigenous Africans--burden, distribution, and trends. *Lancet Oncol.* 2008; 9(7):683–692.10.1016/s1470-2045(08)70175-x [PubMed: 18598933]
- Rambau PF. Pathology practice in a resource-poor setting: Mwanza, Tanzania. *Arch Pathol Lab Med.* 2011; 135(2):191–193.10.1043/1543-2165-135.2.191 [PubMed: 21284436]
- Sweeney YT, Whitaker C. Successful change: renaissance without revolution. *Semin Nurse Manag.* 1994; 2(4):196–202. [PubMed: 7704648]
- Trochim, W.; Donnelly, JP. *The Research Methods Knowledge Base.* 3. Cincinnati, OH: Atomic Dog Publishing; 2006.
- Weiner BJ. A theory of organizational readiness for change. *Implement Sci.* 2009; 4:67.10.1186/1748-5908-4-67 [PubMed: 19840381]
- Weiner BJ, Amick H, Lee SY. Conceptualization and measurement of organizational readiness for change: a review of the literature in health services research and other fields. *Med Care Res Rev.* 2008; 65(4):379–436.10.1177/1077558708317802 [PubMed: 18511812]

Table 1

Respondent Characteristics (n=52)

	Percent/Mean	SD	Minimum, Maximum
Male	61.5%	--	--
Age	38.3	9.9	24, 66
Professional Role			
Physician	57.7%	--	--
Nurse	17.3%	--	--
Resident	9.6%	--	--
Administrator	7.7%	--	--
Medical student	7.7%	--	--
Medical Department			
Internal medicine	32.7	--	--
Surgery	13.5	--	--
Pediatrics	11.5	--	--
Urology	7.7	--	--
Medical records	7.7	--	--
Gynecology	5.8	--	--
Dermatology	3.9	--	--
Pathology	1.9	--	--
Not reported *	15.4	--	--

* Many respondents interpreted the medical department/specialty question as though it applied to physicians only. Therefore, medical department was not reported by many nurse-respondents.

Table 2

Change Commitment (n=52)*

	Not At All Committed % (n)	A Little Committed % (n)	Fairly Committed % (n)	Very Committed % (n)	Don't Know % (n)	Missing % (n)
How committed is your facility to implementing this change?	3.9% (2)	5.8% (3)	28.9% (15)	51.9% (27)	9.6% (5)	0% (0)
	Not At All Motivated % (n)	A Little Motivated % (n)	Fairly Motivated % (n)	Very Motivated % (n)	Don't Know % (n)	Missing % (n)
How motivated is your facility to implementing this change?	1.9% (1)	9.6% (5)	44.2% (23)	36.5% (19)	7.7% (4)	0% (0)
	Not At All Willing % (n)	A Little Willing % (n)	Fairly Willing % (n)	Very Willing % (n)	Don't Know % (n)	Missing % (n)
How willing is your facility to work hard to implement this change?	0.0% (0)	5.8% (3)	36.5% (19)	50.0% (26)	7.7% (4)	0% (0)
	Not At All % (n)	A Little % (n)	Somewhat % (n)	Very Much % (n)	Don't Know % (n)	Missing % (n)
How much does your facility want to implement this change?	1.9% (1)	3.9% (2)	30.8% (16)	55.8% (29)	7.7% (4)	0% (0)

Notes:

1) Respondents were instructed that when an organization makes a change, like expanding the cancer registry program, there is a core group of people who implement the change and use the program. This group probably includes administrators, managers, physicians, nurses, or other staff. Respondents were advised that the survey items assessed their perception and opinion, not factual knowledge. They were then asked to indicate which response best reflected their understanding and confidence in their facility and its leadership regarding expanding the cancer registry.

2) Cronbach's alpha for the change commitment items was 0.77.

Table 3

Change Efficacy (n=52)

	Not At All Confident % (n)	A Little Confident % (n)	Fairly Confident % (n)	Very Confident % (n)	Don't Know % (n)	Missing % (n)
Effectively use the resources that are currently available to implement the program?	9.6% (5)	7.7% (4)	36.5% (19)	40.4% (21)	5.8% (3)	0% (0)
Encourage clinicians to try using this program (e.g., refer patients)?	0% (0)	9.6% (5)	32.7% (17)	55.9% (29)	1.9% (1)	0% (0)
Effectively coordinate the efforts of those involved in implementing this program?	5.8% (3)	13.5% (7)	42.3% (22)	34.6% (18)	3.9% (2)	0% (0)
Support clinicians as they adjust their clinical practice in response to this program?	1.9% (1)	11.5% (6)	36.5% (19)	44.2% (23)	3.9% (2)	1.9% (1)
Effectively solve problems that might arise in implementing this program?	7.7% (4)	17.3% (9)	42.3% (22)	26.9% (14)	3.9% (2)	1.9% (1)
Communicate clearly why the facility is implementing this change?	1.9% (1)	15.4% (8)	38.5% (20)	38.5% (20)	5.8% (3)	0% (0)
Keep the momentum going in implementing this change?	1.9% (1)	21.2% (11)	44.2% (23)	28.9% (15)	1.9% (1)	1.9% (1)
Keep track of how things are going in implementing this change?	1.9% (1)	19.2% (10)	36.5% (19)	34.6% (18)	3.9% (2)	3.9% (2)
Assign the right tasks to the right people in implementing this change?	5.8% (3)	15.4% (8)	32.7% (17)	44.2% (23)	1.9% (1)	0% (0)
Coordinate tasks so that implementation goes smoothly?	3.9% (2)	21.2% (11)	28.9% (15)	44.2% (23)	1.9% (1)	0% (0)
Support people as they adjust to this change?	1.9% (1)	17.3% (9)	50.0% (26)	23.1% (12)	5.8% (3)	1.9% (1)
Get people involved and interested in implementing this change?	9.6% (5)	9.6% (5)	42.3% (22)	34.6% (18)	1.9% (1)	1.9% (1)
Manage the politics of implementing this change?	7.7% (4)	17.3% (9)	32.7% (17)	30.8% (16)	7.7% (4)	3.9% (2)

Notes:

¹⁾ Respondents were instructed that when an organization makes a change, like expanding the cancer registry program, there is a core group of people who implement the change and use the program. This group probably includes administrators, managers, physicians, nurses, or other staff. Respondents were advised that the survey items assessed their perception and opinion, not factual knowledge. They were then asked to indicate which response best reflected their understanding and confidence in their facility and its leadership regarding expanding the cancer registry.

²⁾ Cronbach's alpha for the change efficacy items was 0.93.

Table 4

Collapsed Change Efficacy and Change Commitment Responses (n=52) *

	Not At All or A Little Confident % (n)	Fairly or Very Confident % (n)	Don't Know or Missing % (n)
Management Tasks			
Communicate clearly why the facility is implementing this change?	17.3% (9)	76.9% (40)	5.8% (3)
Effectively use the resources that are currently available to implement the program?	17.3% (9)	76.9% (40)	5.8% (3)
Effectively coordinate the efforts of those involved in implementing this program?	19.3% (10)	76.9% (40)	3.9% (2)
Get people involved and interested in implementing this change?	19.3% (10)	76.9% (40)	3.9% (2)
Assign the right tasks to the right people in implementing this change?	21.2% (11)	76.9% (40)	1.9% (1)
Support people as they adjust to this change?	19.2% (10)	73.1% (38)	7.7% (4)
Keep the momentum going in implementing this change?	22.2% (12)	73.1% (38)	3.8% (2)
Coordinate tasks so that implementation goes smoothly?	25.1% (13)	73.1% (38)	1.9% (1)
Keep track of how things are going in implementing this change?	21.1% (11)	71.1% (37)	7.8% (4)
Effectively solve problems that might arise in implementing this program?	25% (13)	69.2% (36)	3.9% (2)
Manage the politics of implementing this change?	25.0% (13)	63.5% (33)	11.6% (6)
Organizational Motivation			
Encourage clinicians to try using this program (e.g., refer patients)?	9.6% (5)	88.6% (46)	1.9% (1)
Support clinicians as they adjust their clinical practice in response to this program?	13.4% (7)	80.7% (42)	5.8% (3)
	Not At All or A Little % (n)	Somewhat or Very Much % (n)	Don't Know or Missing % (n)
How much does your facility want to implement this change?	5.8% (3)	86.6% (45)	7.7% (4)
	Not At All or A Little Willing % (n)	Fairly or Very Willing % (n)	Don't Know or Missing % (n)
How willing is your facility to work hard to implement this change?	5.8% (3)	86.5% (45)	7.7% (4)
	Not At All or A Little Committed % (n)	Fairly or Very Committed % (n)	Don't Know or Missing % (n)
How committed is your facility to implementing this change?	9.7% (5)	80.8% (42)	9.6% (5)
	Not At All or A Little Motivated % (n)	Fairly or Very Motivated % (n)	Don't Know or Missing % (n)
How motivated is your facility to implementing this change?	11.5% (6)	80.7% (42)	7.7% (4)

* Survey responses presented in Table 2 were collapsed (e.g., "not at all confident" and "a little confident" were merged). Survey items were reorganized into two categories – organizational motivation and management tasks. Within each category, survey items were then ordered from highest percentage of confidence to lowest percentage of confidence.

Table 5

Resource Availability (n=52)

	Have None and Resource is Not Attainable % (n)	Have Insufficient, But Resource is Attainable % (n)	Already Have Sufficient Resource % (n)	Not Necessary % (n)	Don't Know % (n)	Missing % (n)
Pathologists	48.1% (25)	38.5% (20)	5.8% (3)	0.0% (0)	1.9% (1)	5.8% (3)
Pathology technicians	9.6% (5)	63.5% (33)	13.5% (7)	1.9% (1)	3.9% (2)	7.7% (4)
Staff to collect patient data	11.5% (6)	44.2% (23)	28.9% (15)	0.0% (0)	5.8% (3)	9.6% (5)
Staff to enter patient data	13.5% (7)	46.2% (24)	26.9% (14)	0.0% (0)	5.8% (3)	7.7% (4)
Staff to follow-up on registered patients	17.3% (9)	34.6% (18)	25.0% (13)	1.9% (1)	13.5% (7)	7.7% (4)
Staff to audit the registry	19.2% (10)	44.2% (23)	15.4% (8)	0.0% (0)	13.5% (7)	7.7% (4)
Computers	9.6% (5)	57.7% (30)	17.3% (9)	0.0% (0)	7.7% (4)	7.7% (4)
Software	17.3% (9)	48.1% (25)	11.5% (6)	0.0% (0)	15.4% (8)	7.7% (4)
Staff with computing expertise	17.3% (9)	44.2% (23)	17.3% (9)	0.0% (0)	13.5% (7)	7.7% (4)
Funds to maintain the registry	34.6% (18)	19.2% (10)	5.8% (3)	0.0% (0)	32.7% (17)	7.7% (4)