Published in final edited form as:

Bioorg Med Chem Lett. 2007 July 15; 17(14): 4062–4065. doi:10.1016/j.bmcl.2007.04.070.

Synthesis and characterization of a perchlorotriphenylmethyl (trityl) triester radical: a potential sensor for superoxide and oxygen in biological systems

Vinh Dang^a, Jinhua Wang^b, Song Feng^b, Christophe Buron^a, Frederick A. Villamena^a, Peng George Wang^b, and Periannan Kuppusamy^{a,*}

^aCenter for Biomedical EPR Spectroscopy and Imaging, Davis Heart and Lung Research Institute, and Division of Cardiovascular Medicine, Department of Internal Medicine, The Ohio State University, Columbus, OH 43210, USA

^bDepartment of Chemistry, The Ohio State University, Columbus, OH 43210, USA

Abstract

Synthesis and characterization of an inert perchlorotriphenylmethyl triester radical, PTM-TE, are reported. PTM-TE was prepared by a facile 3-step synthesis using Friedel-Crafts reaction of tetrachlorobenzene with chloroform followed by ethoxycarbonylation and subsequent oxidation. PTM-TE is paramagnetic, and detected as a single sharp peak by EPR spectroscopy. In solution, the EPR linewidth of PTM-TE is highly sensitive to the dissolved oxygen content, thus enabling accurate measurement of oxygen concentration (oximetry). In addition, the radical also shows high reactivity to superoxide. The ester radical has the potential for use as a high-sensitive probe for determination of oxygen concentration and superoxide in biological systems.

Keywords

Perchlorotriphenylmethyl radical; perchlorotriphenylmethyl triester; superoxide; free radicals; EPR; oximetry; spectrophotometry

In 1900, Moses Gomberg reported the existence of a stable, trivalent organic free radical which challenged the prevailing belief that carbon could only have four chemical bonds.¹ As this report provided new possibilities to carbon chemistry, research aimed at understanding and developing these radicals continued to thrive and permeated through various fields of science including medicine and industrial applications. Most notably, Gomberg's work paved the way for future research on using persistent trityl radicals in clinical applications.²⁻⁴ However, it is known that some radicals and diradicals are short-lived and very reactive - the half-life of triphenylmethyl radicals in aerated solution may be as low as a fraction of a second. Studies have also shown that these reactive radicals become stable and chemically inert upon perchlorination.⁵⁻⁸ In light of this endeavor, highly chlorinated mono-, di-, and triarylmethanes were developed and since then they have become the most valuable chemical precursors of inert free radicals with estimated half-lives in the order of

© 2007 Elsevier Ltd. All rights reserved.

*Corresponding author. Tel: +1 614 292 8998; fax +1 614 292 8454; kuppusamy.1@osu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

100 years.⁶ Together, these studies have been regarded as both promising and significant.⁵ As a result, the synthesis of perchlorinated trityl radical has become an immense undertaking with promising implications.

The role of reactive oxygen species (ROS) such as superoxide, hydroxyl, and alkylperoxyl radicals has been linked to a variety of pathophysiological processes.⁹ Since, oxidative stress leads to an increased production of ROS, especially superoxide, accurate determination of these free radicals can help us gain a better understanding of oxidative stress-related diseases. Of the many techniques currently available for the detection of superoxide in biological systems, electron paramagnetic resonance (EPR) spectroscopy has the distinct capability for direct and real-time detection. However, the EPR technique requires the use of high concentrations (10–100 mM) of spin-trap molecules to maximize the efficiency of trapping, which may alter the redox environment of the system.^{10,11} The poor efficiency of the spin-trap molecules for superoxide detection are attributed to small bimolecular reaction rate constants (typically about $10^2 \text{ M}^{-1} \text{ s}^{-1}$). Thus, there is a great need to develop probes with orders of magnitude increased reactivity with superoxide. To this end, we have developed a perchlorotriphenylmethyl triester radical (PTM-TE) that can overcome this limitation. PTM-TE has a high detection-sensitivity due to sharp single-line EPR spectrum. In addition, PTM-TE is an attractive candidate for measurement of oxygen concentration (oximetry). Furthermore, the ester nature of the PTM-TE radical is expected to facilitate its penetration into cells, followed by hydrolysis by cellular enzymes such as esterases into carboxylate radical and thus can be trapped in the cells. This will enable the detection of superoxide radicals produced inside the cells. As an oxygen probe, PTM-TE can be utilized for measurement as well as mapping (imaging) of oxygen distribution in tissues.¹² In this communication, we report the synthesis, characterization, and preliminary evaluation of PTM-TE for oximetry and detection of superoxide.

The synthesis of PTM-TE radical **4** (scheme 1) used tris(2,3,5,6-tetrachlorophenyl)methane generated from the Friedel-Craft reaction of 1,2,4,5-tetrachlorobenzene and chloroform.¹³ Reaction of compound **2** with slight excess of butyllithium and TMEDA in THF at low temperature yielded the corresponding tri-anion. This tri-anion was reacted with ethylchloroformate *in situ* to yield compound **3**.¹⁴ Compound **3** was then converted into **4** by a two-step process: conversion into the triphenylcarbanion with NaOH in a mixture solution of DMSO and Et₂O, followed by oxidation with I₂.¹⁵ Since compound **3** has the ethoxycarbonyl functional group, it can be easily modified to afford more variants for future research.

PTM-TE was characterized by EPR spectroscopy (Figure 1). The solid showed a broad EPR signal with a peak-to-peak width (ΔB_{pp}) of 5.650 G in room air. However, a solution of PTM-TE (100 μM) in dimethylsulfoxide (DMSO) at ambient conditions showed a single sharp peak with a linewidth of 0.540 G. No self-induced spin-spin broadening was observed for up to 12 mM concentration of PTM-TE. The linewidth of PTM-TE in DMSO, however, was dependent on the concentration of dissolved molecular oxygen in solution. We measured the linewidth of PTM-TE (100 μM) in DMSO as well as in hexafluorobenzene (HFB) as a function of partial pressure of equilibrating oxygen ($p\text{O}_2$). As shown in Figure 2, the linewidth increased proportionally to $p\text{O}_2$. The oxygen-induced broadening is attributed to the paramagnetic nature of molecular oxygen, which upon collision with PTM-TE molecule can undergo Heisenberg spin exchange interaction resulting in broadening of the observed EPR signal. It should be noted that this exchange interaction does not chemically alter the two molecules involved. Further, the oxygen-induced broadening has no effect on the EPR signal intensity (double-integral of the EPR spectrum). Hence, the width of the EPR signal can be used to measure the concentration of O₂ in the system. The calibration curves

of EPR linewidth versus pO_2 can be generated for any solvent by equilibrating it with known concentrations of oxygen.

The results shown in Figure 2 indicate a significant contrast between the slopes of the calibration curves of PTM-TE in DMSO (1.6 mG/mmHg) and HFB (17.71 mG/mmHg). The higher sensitivity in HFB is largely attributed to higher solubility of oxygen in HFB. For example, the solubility of oxygen at atmospheric pressure (21% of oxygen) at 25°C is 4400 μM in HFB as compared to 478 μM in DMSO.¹⁶ In addition, viscosity and polarity of solvent molecule may also contribute to the observed difference.

The reactivity of PTM-TE to superoxide was determined. PTM-TE in DMSO (0.8 mM) was treated with a solution of potassium superoxide (KO_2) in DMSO (2.4 mM). As shown in Figure 3, the EPR spectrum of PTM-TE was completely quenched by superoxide suggesting that PTM-TE radical reacts with superoxide to form a diamagnetic product. Although the mechanism by which PTM-TE reacts with superoxide is unclear, previous reports suggested that the neutral protonated form of superoxide, $\bullet\text{OOH}$, rather than $\text{O}_2^{\bullet-}$ is involved in the reaction, as illustrated in the following equation.

It is likely that the superoxide anion or $\bullet\text{OOH}$ radical interacts with PTM-TE to form a covalent bond between the methyl radical epicenter and the oxygen atom of $\bullet\text{OOH}$.¹⁰ The resulting adduct is diamagnetic and hence, EPR silent. This observation is confirmed by ^1H NMR spectroscopy where only the triplet and quartet splittings for the ethoxycarbonyl substituent were identified. The absence of a singlet peak at δ 7 (ppm) corresponding to the epicenter methine proton (data not shown) indicates that the resulting compound appears most likely to be the proposed adduct, $[\text{R}_3\text{C} - \text{OOH}]$.

PTM-TE was specifically designed with the ethoxycarbonyl substituent attached to each of the phenyl ring for a specific purpose, i.e., cellular internalization. The hydrophobic character of the molecule permits it to easily diffuse across the membrane lipid bilayers. Then, by the nonspecific action of intracellular esterases, the molecule is expected to be cleaved to yield the corresponding tricarboxylate radical, PTM-TC. The resulting PTM-TC is another stable “inert” radical which also gives a single sharp EPR peak in aqueous solutions.¹⁷ PTM-TC gives a characteristic UV-VIS absorption at 380 nm. Our recent study has shown that PTM-TC, on reaction with superoxide, induces a decrease in EPR signal or optical absorption at 380 nm with high specificity and sensitivity.¹⁸ Thus, PTM-TE can be used to monitor and quantify the production of intracellular superoxide.

In summary, we have developed an efficient synthetic route for the synthesis of PTM-TE. Additionally, we have demonstrated the effect of molecular oxygen and superoxide on the EPR spectrum of PTM-TE in solution. Unlike molecular oxygen, superoxide induces a signal loss. Thus, PTM-TE potentially can be used to determine oxygen and/or superoxide generation in biological applications.

Acknowledgments

This study was supported by NIH grants EB004031.

References and notes

1. Gomberg M. J Am Chem Soc. 1900; 22:757.

2. Golman K, Zandt RI, Lerche M, Pehrson R, Ardenkjaer-Larsen JH. *Cancer Res.* 2006; 66:10855. [PubMed: 17108122]
3. Ardenkjaer-Larsen JH, Fridlund B, Gram A, Hansson G, Hansson L, Lerche MH, Serving R, Thaning M, Golman K. *Proc Natl Acad Sci USA.* 2003; 100:10158. [PubMed: 12930897]
4. Golman K, Ardenkjaer-Larsen JH, Petersson JS, Mansson S, Leunbach I. *Proc Natl Acad Sci USA.* 2003; 100:10435. [PubMed: 12930896]
5. Ballester M, Pascual I, Carreras C, Vidal-Gancedo J. *J Am Chem Soc.* 1994; 116:4205.
6. Ballester M. *Pure Appl Chem.* 1967; 15:123.
7. Ballester M. *Adv Phys Org Chem.* 1989; 25:267.
8. Ballester M, Riera J, Castañer J, Badía C, Monsó JM. *J Am Chem Soc.* 1971; 93:2215.
9. Sies H. *Int J Vitam Nutr Res Suppl.* 1989; 30:215. [PubMed: 2507700]
10. Rizzi C, Samouilov A, Kutala VK, Parinandi NL, Zweier JL, Kuppusamy P. *Free Radic Biol Med.* 2003; 35:1608. [PubMed: 14680684]
11. Kutala VK, Parinandi NL, Zweier JL, Kuppusamy P. *Arch Biochem Biophys.* 2004; 424:81. [PubMed: 15019839]
12. Bratasz A, Kulkarni AC, Kuppusamy P. *Biophys J.* 2007; 92:2918. [PubMed: 17259268]
13. Ballester M, Riera J, Castañer J, Rovira C, Armet O. *Synthesis.* 1986; 1:64.
14. Tris(2,3,5,6-tetrachlorophenyl)methane (486 mg, 0.74 mmol) and TMEDA (377 μ l, 2.4 mmol) were dissolved in dry THF (50 ml) under dry nitrogen atmosphere and cooled to -78°C . A solution of 1.6 M butyllithium in hexanes (1.6 ml, 2.5 mmol) was added in one portion and the mixture stirred at low temperature for 1 hour. The reaction was quenched with ethyl chloroformate and the mixture was warmed up to room temperature overnight. Solvent was evaporated and the residue dissolved in dichloromethane. The organic layer was washed with water and dried over sodium sulfate. Solvent was evaporated under vacuum. Silica gel chromatography (hexanes/AcOEt:12/1) yielded 500 mg (77%) of tris(2,3,5,6-tetrachloro-4-ethoxycarbonyl-phenyl) methane **3** as a colorless solid. ^1H NMR (400 MHz, CDCl_3) δ 7.00 (s, 1H); 4.47 (q, $J = 7.1$ Hz, 6H); 1.41 (t, $J = 7.1$ Hz, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 163.2, 138.5, 135.5, 135.0, 134.0, 130.5, 129.5, 63.1, 56.3, 14.0. UV-vis (hexanes): λ_{max} (e) 292 (2220), 302 (2480). IR: ν 2982, 2936, 1743, 1556, 1465, 1446, 1371, 1341, 1329, 1299, 1263, 1225, 1208, 1121, 1096, 1019, 859 cm^{-1} . HRMS $[\text{M} + \text{Na}]^+$ calcd. for $\text{C}_{28}\text{H}_{16}\text{Cl}_{12}\text{O}_6\text{Na}$ 896.7018, found 896.7041.
15. Tris(2,3,5,6-tetrachloro-4-ethoxycarbonyl-phenyl) methane (1.49 g, 1.78 mmol) was dissolved in the mixture of DMSO/ Et_2O (75/425). To the solution powdered NaOH was added, the mixture was stirred in the dark for 24h. The mixture was filtered through a dry funnel into a solution of iodine (1.9 g) in Et_2O (100 ml). The solution was left undisturbed in the dark for 24h. The resulting solution was then washed with concentrated NaHCO_3 (39%, 50 ml) brine, then water (50 ml $\times$ 2). Solvent was dried over Na_2SO_4 and evaporated. The residue was purified by silica gel chromatography (Hexane/AcOEt: 80/20) to give a red solid (1.378 g, 92.5%). EPR (solid, 298 K): single peak, $\Delta B_{\text{pp}} = 5.650$. UV-vis (hexane): λ_{max} (e) 289 (5010), 378 (8830). IR: ν 2938, 1740, 1536, 1445, 1380, 1329, 1284, 1222, 1136, 1007, 858, 720 cm^{-1} . HRMS $[\text{M} + \text{Na}]^+$ calculated for $\text{C}_{28}\text{H}_{15}\text{Cl}_{12}\text{O}_6\text{Na}$ 895.694, found 895.692.
16. Murov, SL.; Hug, GL.; Carmichael, I. *Hand Book of Photochemistry.* New York: M. Dekker; 1993.
17. Maspoch D, Gerbier P, Catala L, Ruiz-Molina D, Vidal-Gancedo J, Wurst K, Rovira C, Veciana J. *Chem Eur J.* 2002; 8:3635. [PubMed: 12203290]
18. Kutala VK, Villamena FA, Ilangoan G, Maspoch D, Roques N, Veciana J, Rovira C, Kuppusamy P. *Biochem J.* in review.

Figure 1.

EPR spectra of PTM-TE. The spectra were measured from a sample of crystalline powder and in a solution of PTM-TE (100 μ M) in DMSO at ambient conditions using an X-band EPR spectrometer. The EPR acquisition settings were: modulation amplitude, 160 mG; microwave power, 1 mW; time constant, 40 ms; scan time, 15 s. The first derivative peak-to-peak widths were 5.650 G for solid and 0.540 G for solution. The small satellite peaks indicated with * on both sides of the main peak are due to hyperfine coupling with ^{13}C nuclei (natural abundance).

Figure 2.

Effect of oxygen on the EPR linewidth of PTM-TE in solution. Measurements were made by equilibrating a solution of PTM-TE (100 μ M) in DMSO or HFB with pre-calibrated gas mixtures of known O₂ concentration. The data show a linear dependence of linewidth with pO₂. The oxygen-sensitivity was 17.71 mG/mmHg for HFB and 1.6 mG/mmHg for DMSO.

Figure 3. The effect of superoxide on the EPR spectrum of PTM- TE radical in DMSO. **(A)** EPR spectrum of PTM-TE in DMSO (0.8 mM). **(B)** EPR spectrum of PTM-TE in DMSO following addition of solid KO₂ (2.4 mM). The PTM-TE signal is quenched by O₂•⁻.

Synthesis of PTM-TE radical. Reagents and conditions: (a) CHCl_3 , AlCl_3 ; (b) *n*-BuLi, TMEDA, ethyl chloroformate, 77%; (c) (1) NaOH, DMSO/ Et_2O , (2) I_2 , Et_2O , 93%.