Published in final edited form as:

J Am Chem Soc. 2005 October 26; 127(42): 14538–14539. doi:10.1021/ja052401q.

The Reactant State for Substrate-Activated Turnover of Acetylthiocholine by Butyrylcholinesterase is a Tetrahedral Intermediate

Jose R. Tormos[†], Kenneth L. Wiley^{†,¶}, Javier Seravalli^{†,‡}, Florian Nachon[‡], Patrick Masson[‡], Yvain Nicolet[§], and Daniel M. Quinn[†]

Department of Chemistry, The University of Iowa, Iowa City, Iowa 52242; Centre de Recherches du Service de Santé Armées, Unité d'Enzymologie, BP 87-38702 La Tronche CEDEX, France; and the Institut de Biologie Structurale, Laboratoire de Cristallographie et Cristallogénèse des Protéines, 38027 Grenoble, France

Daniel M. Quinn: daniel-quinn@uiowa.edu

Abstract

Secondary β -deuterium kinetic isotope effects have been measured as a function of substrate concentration for recombinant human butyrylcholinesterase-catalyzed hydrolysis of acetyl-L₃-thiocholine ($L = {}^1\text{H}$ or ${}^2\text{H}$). The isotope effect on V/K is inverse, $D^3V/K = 0.93 \pm 0.03$, which is consistent with conversion of the sp^2 hybridized carbonyl carbon of the scissile ester bond of the E + A reactant state to a quasi-tetrahedral structure in the acylation transition state. In contrast, the isotope effect on V_{max} under conditions of substrate activation is markedly normal, $D^3(\beta V_{\text{max}}) = 1.29 \pm 0.06$, an observation that is consistent with accumulation of a tetrahedral intermediate as the reactant state for catalytic turnover. Generally, tetrahedral intermediates for nonenzymic ester hydrolyses are high-energy steady-state intermediates. Apparently, butyrylcholinesterase displays an unusual ability to stabilize such intermediates. Hence, the catalytic power of cholinesterases can largely be understood in terms of their ability to stabilize tetrahedral intermediates in the multi-step reaction mechanism.

Acetylcholinesterase (AChE¹) and butyrylcholinesterase (BuChE¹) catalyze the hydrolysis of choline esters with very high catalytic efficiency. At neutral pH cholinesterase reactions are $\sim 10^{13}$ fold faster than spontaneous substrate hydrolysis, a factor that corresponds to 74 kJ mol⁻¹ of transition state stabilization.² Though many structural features that contribute to the catalytic mechanism have been illuminated^{3,4} – the Ser-His-Glu catalytic triad, the oxyanion hole, the acyl binding site, the quaternary ammonium binding site – a decisive understanding of cholinesterase catalytic power is yet to be had.

Recently, Nicolet *et al.*⁵ reported crystal structures of human BuChE and complexes of the enzyme. Surprisingly, in the supposed unliganded enzyme structure, determined at a

Correspondence to: Daniel M. Quinn, daniel-quinn@uiowa.edu.

[†]The University of Iowa

[‡]Centre de Recherches du Service de Santé des Armées

[§]Institut de Biologie Structurale.

[¶]Current address: Harvard University, Department of Molecular and Cellular Biology, Cambridge, MA 02138.

[#]Current address: University of Nebraska, Department of Biochemistry, Lincoln, NE 68588-0664.

Supporting Information Available: Kinetic parameters (V_{max} , V/K) from fits of time courses to the integrated Michaelis-Menten equation for BuChE-catalyzed hydrolyses of acetyl-L₃-thiocholines. Derivations of equations 1 and 2.

¹Abbreviations: AChE, acetylcholinesterase; ATCh, acetylthiocholine; BuChE, butyrylcholinesterase; BSA, bovine serum albumin; DTNB, 5,5'-dithiobis(2-nitrobenzoic acid); mAU, milli-absorbance unit.

resolution of 2.0 Å, electron density that was proximal to the nucleophilic oxygen of the active site Ser198 was interpreted as a bound butyrate. This ligand could be displaced with 3-bromopropionate, which placed the structural assignment on a firm footing. However, it was not possible to differentiate cleanly between a complex in which the ligand is bound as a tetrahedral intermediate and one which is a BuChE-carboxylate complex. Though angular distortions toward tetrahedrality were noted about the carbonyl carbon of the bound butyrate, the resolution of the structure and the rather long Ser198 γO to butyrate carbonyl carbon distance (2.16 Å) suggest that the complex may be a mixture of tetrahedral intermediate and EP complexes, and perhaps also a covalent acylenzyme.

These unusual observations beg for an experimental probe of the catalytic behavior of BuChE that can address the nature of substrate hybridization changes that accompany catalytic turnover in the steady-state. Herein we report on the measurements of secondary β-deuterium kinetic isotope effects with isotopomers of acetylthiocholine (*i.e.* acetyl-L₃-thiocholine, L = ¹H or ²H) that provide such a probe.⁶ Secondary deuterium isotope effects are inverse (< 1.00) when a sp²-hybridized reactant is converted to a quasi-tetrahedral transition state, and normal (> 1.00) when an sp³ reactant state is converted to a transition state that has greater trigonal planar character.⁷

BuChE-catalyzed reactions often show substrate activation at high substrate concentrations.³ An interaction mechanism that accords with substrate activation is outlined in Scheme 1. E and A are the enzyme and substrate respectively, EA is the Michaelis complex, and EAA is the ternary complex of enzyme with substrate monomers bound at both the active and substrate activation sites. The active site of BuChE is at the bottom of a 20 Å deep gorge,⁵ and the substrate activation site is thought to coincide with the peripheral site at the mouth of the gorge.⁹ Substrate activation is observed because turnover of EAA via βk_{cat} (= βV_{max}/[E]_T) is faster than turnover of EA by a factor β. As Figure 1 shows, the dependence of human BuChE-catalyzed hydrolysis of acetyl-¹H₃-thiocholine on substrate concentration shows deviations from Michaelis-Menten kinetics that are consistent with substrate activation. The data in Figure 1 were fit to equation 1, derived for the kinetic mechanism of Scheme 1 (see Supporting Information for derivation of equation 1). The fit yielded the following parameters: V_{max} = 20 ± 2 mA min⁻¹, K_m = 80 ± 20 μM, K_a = 3.5 ± 0.5 mM, and β = 2.6 ± 0.2. Hence, the ternary EAA complex of Scheme 1 turns over to product 2.6 times faster than does the EA complex. Though substrate activation has long been observed for BuChE-catalyzed reactions, an understanding of its functional origins remains elusive. Fortunately, the β-secondary deuterium kinetic isotope effects detailed below shed important new light on this phenomenon.

$$v_i = \frac{V_{\max}[A](1+\beta[A]/K_A)}{K_m(1+\beta[A]/K_A)+[A](1+[A]/K_A)} \quad (1)$$

The least-squares uncertainties that result from the fit in Figure 1 preclude determination of secondary isotope effects on the kinetic parameters from such experiments. An alternate approach is to determine the secondary isotope effect on the initial rate (*i.e.* $D^3 v_i = v_i^{H^3}/v_i^{D^3}$) as a function of substrate concentration. As Figure 2 shows, the isotope effect on the initial rate increases hyperbolically as the substrate concentration increases. These data were fit to equation 2, which provides a quantitative analysis and extrapolation of the isotope effects:

$$D^3 v_i = D^3 V_{\max} \frac{1+\frac{\beta[A]}{K_A}}{1+\frac{\beta[A]}{K_A D^3 \beta}} \times \frac{[A] \left(1+\frac{[A]}{K_A}\right) + K_m \left(1+\frac{\beta[A]}{K_A D^3 \beta}\right)}{[A] \left(1+\frac{[A]}{K_A}\right) + K_m \left(1+\frac{\beta[A]}{K_A}\right)} \quad (2)$$

The isotope effect extrapolated to zero concentration is D^3V/K (the isotope effect on $V_{\max}/K_m$) and $D^3\beta$ is the asymptotic value of the isotope effect on $V_{\max}$ at infinite concentration. Figure 2 shows a least-squares fit of the isotope effects to equation 2. In determining this fit, the parameters β , K_A and K_m were constrained to the values determined in the substrate activation plot in Figure 1. The isotope effects that result from the fit in Figure 2 are $D^3V/K = 0.93 \pm 0.03$ and $D^3\beta = 1.38 \pm 0.06$. Hence, the isotope effect on $\beta V_{\max}$ is 1.29 ± 0.06 .

The experiment in Figure 2 does not provide the value of the isotope effect on $V_{\max}$, the parameter that monitors non-substrate activated turnover via the EA complex of Scheme 1. To determine $D^3V_{\max}$, and to provide a check on the value of D^3V/K determined in Figure 2, time course data for the isotopic acetyl-L₃-thiocholines were fit to the integrated form of the Michaelis-Menten equation.¹⁰ The experiments utilized an initial substrate concentration $[A]_0 = 0.2$ mM that was well below K_A , and therefore the kinetics were not complicated by substrate activation. These experiments gave the following values for the isotope effects on the Michaelis-Menten parameters: $D^3V_{\max} = 1.02 \pm 0.05$ and $D^3V/K = 0.97 \pm 0.06$. Both values are within experimental error of an isotope effect of 1.00, and the value for D^3V/K indeed agrees with that determined in Figure 2. Importantly, neither isotope effect approaches the normal isotope effect on substrate-activated $V_{\max}$. Only upon substrate activation is the marked normal isotope effect expressed.

The observation of a covalent tetrahedral adduct in the crystal structure of human BuChE⁵ suggests that turnover of the acylenzyme intermediate is rate limiting for $V_{\max}$. Therefore, the sizeable normal isotope effect on substrate-activated $V_{\max}$ suggests that in the FA complex allosteric modulation of the active site is accompanied by stabilization of the tetrahedral adduct, which then becomes the predominant accumulating species in the steady-state. The turnover of the tetrahedral adduct would be marked by rehybridization from sp^3 toward sp^2 in the developing acetate product, consistent with the observed normal isotope effect. Hence, observations from x-ray crystallography and from isotope effects conspire to suggest that an important element in the catalytic power of cholinesterases is their ability to stabilize tetrahedral intermediates that are, in corresponding nonenzymic reactions, high-energy and metastable species.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

KLW was supported by the Predoctoral Training Program in Biotechnology, NIH grant T32 GM008365.

References

2. Quinn DM. Chem Rev. 1987; 87:955–979.
3. Taylor P, Radi Z. Annu Rev Pharmacol Toxicol. 1994; 34:281–320. [PubMed: 8042853]
4. Harel M, Quinn DM, Nair HK, Silman I, Sussman JL. J Am Chem Soc. 1996; 118:2340.
5. Nicolet Y, Lockridge O, Masson P, Fontecilla-Camps JC, Nachon F. J Biol Chem. 2003; 278:41141–41147. [PubMed: 12869558]
6. Malany S, Sawai M, Sikorski RS, Seravalli J, Quinn DM, Radi Z, Taylor P, Kronman C, Velan B, Shafferman A. J Am Chem Soc. 2000; 122:2981–2987.
7. (a) Hogg, JL. Transition State of Biochemical Processes. Gandour, RD.; Schowen, RL., editors. Plenum Press; New York and London: 1978. p. 201–224.(b) Kirsch, JF. Isotope Effects on Enzyme-Catalyzed Reactions. Cleland, WW.; O'Leary, MH.; Northrop, DB., editors. University Park Press; Baltimore: 1977. p. 100–122.

8. Ellman GL, Courtney KD, Andreas V Jr, Featherstone RM. *Biochem Pharmacol.* 1961; 7:88–95. [PubMed: 13726518]
9. Masson P, Legrand P, Bartels CF, Froment MT, Schopfer LM, Lockridge O. *Biochemistry.* 1997; 36:2266–2277. [PubMed: 9047329]
10. Acheson SA, Barlow PN, Lee GC, Swanson ML, Quinn DM. *J Am Chem Soc.* 1987; 109:246–252.

Figure 1.

Dependence of initial rate on concentration of acetyl-¹H₃-thiocholine. Reactions were monitored by the coupled assay of Ellman *et al.*⁸ at 27.0 ± 0.1 °C and pH = 7.24 in 0.1 M sodium phosphate buffer that contained 0.5 mM DTNB, 0.03 mg/ml of BSA, and 93 pM recombinant human BuChE. Reactions were followed at $\lambda = 412$ nm on a Molecular Devices SPECTRA_{MAX}® PLUS³⁸⁴ UV-visible microplate spectrophotometer. The solid line is least-squares fit to equation 1 of the text. The double-reciprocal plot in the inset clearly shows substrate activation.

Figure 2.
 β -secondary deuterium kinetic isotope effects versus substrate concentration for the recombinant human BuChE-catalyzed hydrolysis of acetyl-L₃-thiocholine (L = H or D). The solid line is a least-squares fit to equation 2 of the text.

Scheme 1.

Kinetic mechanism for substrate activation of BuChE-catalyzed reactions