Published in final edited form as:

Angew Chem Int Ed Engl. 2012 May 7; 51(19): 4685–4689. doi:10.1002/anie.201200559.

Base-Catalyzed Direct Aldolization of α -Alkyl- α -Hydroxy Trialkyl Phosphonoacetates**

Michael T. Corbett,

Department of Chemistry, The University of North Carolina at Chapel Hill, Chapel Hill, NC 27599 (USA)

Daisuke Uraguchi,

Department of Applied Chemistry, Graduate School of Engineering, Nagoya University, Furo-cho B2-3(611), Chikusa, Nagoya 464-8603 (Japan)

Takashi Ooi*, and

Department of Applied Chemistry, Graduate School of Engineering, Nagoya University, Furo-cho B2-3(611), Chikusa, Nagoya 464-8603 (Japan)

Jeffrey S. Johnson*

Department of Chemistry, The University of North Carolina at Chapel Hill, Chapel Hill, NC 27599 (USA)

Abstract

Catalytic direct aldol addition of α -hydroxy trialkyl phosphonoacetates to aldehydes affords α -hydroxy- β -phosphonyloxy ester products. The fully substituted glycolate enolate intermediate is generated in situ under mild conditions via [1,2]-phosphonate-phosphate rearrangement. High enantioselectivity and dramatic enhancement of reaction diastereocontrol is realized via the application of chiral iminophosphorane catalysts. The phosphate products undergo stereoselective nucleophilic displacement reactions.

Keywords

Direct Aldol Reaction; Phosphate; Rearrangement; Enantioselective

Catalytic direct aldol reactions offer a convenient method for the rapid construction of β -hydroxy carbonyls through the coupling of carbonyl donors and aldehyde acceptors.^[1]

**The project described was supported by Award R01 GM084927 from the National Institute of General Medical Sciences, NEXT program, and a Grant-in-Aid for Scientific Research on Innovative Areas "Advanced Molecular Transformations by Organocatalysts" from MEXT. M.T.C. acknowledges a JSPS/NSF EAPSI Fellowship and the Global COE program in Chemistry of Nagoya University. Additional support from Novartis is gratefully acknowledged. X-ray crystallographic analysis was performed by Mr. Yusuke Ueki of Nagoya University.

*Prof. Dr. J. S. Johnson, Fax: (+1) 919-962-2388, jsj@unc.edu, Prof. Dr. T. Ooi, Fax: (+81) 52-789-3338, tooi@apchem.nagoya-u.ac.jp.

Generation of the reactive enolate by a basic or nucleophilic catalyst obviates the need to preform an enolate equivalent. Reaction subtypes can be broadly categorized by the oxidation state of the pro-nucleophile. Ketones and aldehydes can be used in direct aldolization because of their relatively high acidity (via Brønsted base catalysis) or their ability to form an enamine (via Lewis base catalysis).^[2] Catalytic direct aldol and Mannich reactions when the donor is in the carboxylic acid oxidation state are considerably more elusive.^[3] Mechanistic nuances are more diverse and full substitution at the α -carbon is difficult to achieve. Reported in this communication is a new base-catalyzed direct glycolate aldol addition that relies upon the strategic use of a [1,2]-phosphonate-phosphate rearrangement (Scheme 1). This new reaction platform was then deployed in the development of highly enantio- and diastereoselective variants through the application of chiral iminophosphorane catalysts. A consequence of the reaction design is that a nucleofuge is directly delivered by the aldolization, a circumstance that is favorable for subsequent nucleophilic displacement reactions.

Although analogous to the well-studied isoelectronic [1,2]-Brook rearrangement,^[4] the [1,2]-phosphonate-phosphate rearrangement of α -hydroxy phosphonates is comparatively underutilized.^[5,6] Incorporation of an adjacent electron-withdrawing group facilitates C $\rightarrow$ O dialkoxyphosphinyl migration ($2 \rightarrow 2' \rightarrow 2''$) through stabilization of the incipient negative charge. The [1,2]-phosphonate-phosphate rearrangement of α -hydroxy phosphonates has been employed in the formal racemic and enantioselective reduction of α -keto esters;^[7] however, it has seldom been harnessed as an instrument for C–C bond construction through umpolung reactivity.^[6,8] The fully substituted glycolic acid derivatives that would result from productive trapping of $2'$ feature a β -phosphonyloxy moiety that can serve as an electrophile in secondary transformations^[9] as well as intermediates in the synthesis of biologically active compounds such as tagetitoxin, leustroducsin B, and phoslactomycin A.^[10] The generation and electrophilic trapping of glycolate enolates via base-catalyzed C $\rightarrow$ O dialkoxyphosphinyl migration was accordingly undertaken.

Initial investigations examined the reaction of α -hydroxy phosphonate **1a**^[11] with various bases in the presence of 4-chlorobenzaldehyde (Scheme 2). Temperature controlled the product identity. When the reaction was conducted at room temperature, the epoxide **5** was isolated in 23% yield with 2:1 d.r.; however, at -78 °C under otherwise identical conditions, α -hydroxy- β -phosphonyloxy ester **3a** was isolated in 97% yield and 2:1 d.r. Both products are proposed to arise from the same pathway (*vide infra*): the aldolate of **3a** can undergo intramolecular S_N2 displacement of the vicinal phosphate to afford epoxide **5** through a Darzens-type mechanism at elevated temperature.^[12] At cryogenic temperatures, epoxide formation is completely suppressed. NaO^tBu resulted in lower diastereoselectivity (entry 3), while LiO^tBu had relatively small effect (entry 5). A higher reaction temperature was required for Cs₂CO₃ to initiate the [1,2]-phosphonate-phosphate rearrangement and competitive enolate quenching was observed, resulting in the formation of **4** (entry 4). Attempts to render the reaction catalytic were successful as 20 mol % of either KHMDS or KO^tBu promoted the reaction (entries 6 and 8, respectively). The phosphazene base ^tBu-P4 was also catalytically competent and delivered the opposite (*syn*) diastereoisomer (entry 7). The effect of independently increasing either the equivalents of aldehyde or the reaction concentration was marginal in minimizing formation of **4**, but simultaneously increasing both provided the best ratio of desired product to quenched enolate (entry 8). Carboxy and phosphonate ester identity had little impact on the diastereoselectivity of the reaction (not shown).^[13]

With high-yielding reaction conditions in hand (Table 1, entry 8), an examination of the scope with various aldehydes was conducted (Table 2). Electron-poor and electron-neutral aromatic aldehydes afforded excellent yields albeit with marginal diastereoselectivity

(entries 1–10). Electron-rich aromatic, alkenyl, and alkyl aldehydes afforded product with a notable enhancement in *anti* diastereoselectivity; the use of excess aldehyde (5.0 equiv) was optimal in these cases (entries 11, 13–14). Isolation typically only consisted of separation of the product from excess aldehyde. The major byproduct formed during the reaction is the quenched glycolate enolate **4**. Bulkier aldehydes such as isobutyraldehyde and pivaldehyde are not suitable electrophiles at the present level of optimization.

The scope of the α -hydroxy phosphonate coupling partner was investigated with benzaldehyde by varying the α -substituent (Table 3). The presence of an α -substituent was found to be critical: reactions where R = H suffered from poor reactivity (<5% conversion) (entry 1). In addition to the benzyl group, other alkyl substituents were well-tolerated under the reaction conditions, thus allowing for the incorporation of functional handles such as terminal alkenes and alkynes (entries 3 and 4).

Having established a baseline protocol for achieving a new direct ester aldol addition, efforts were directed at improving the modest reaction diastereoselectivity and developing an enantioselective variant of the title reaction. Catalytic aldolization using ^tBu-P4 provided the *opposite* major diastereomer relative to alkali bases (Table 1), hinting at a possibility that both relative and absolute stereocontrol issues could conceivably be addressed through the use of a chiral iminophosphorane base possessing appropriate structural features and sufficient basicity.^[14] Exploratory experiments testing this hypothesis are summarized in Table 4 and reveal that the aldolization of **1e** indeed took place under the catalysis of chiral iminophosphoranes such as L-valine-derived **6** and the product **7** was isolated with good levels of enantioselectivity. Enantiocontrol can be tuned via electronic changes at the 4-position of the four aryl groups, but reaction efficiency drops with electron-poor variants, presumably due to the reduced basicity (entries 1–5). Solvent identity also appeared important in terms of product distribution (ratio of aldol **7** to the quenched enolate **4b**) and stereoselectivity (entries 6 and 7). Optimal results were then realized using 10 mol % of **6c** in 2-MeTHF as the reaction solvent at –50 °C: the hydroxy phosphate adduct **7a** was isolated in 71% yield and 95:5 enantioselection (entry 8). Perhaps most remarkably, the chiral iminophosphorane-catalyzed aldolization proceeds with essentially complete diastereocontrol (>30:1 *syn:anti*),^[15] revealing this strategy as a powerful method for stereoselective glycolate aldolization with a fully substituted α -center.

Preliminary experiments suggest that application of the optimized reaction conditions to other aldehyde electrophiles provides access to enantiomerically enriched aldol adducts. As illustrated in Table 5,^[17] while the product yields were slightly reduced with 2-substituted aldehydes, complete diastereoselectivity is regularly observed and good enantiocontrol is maintained. The structure and absolute stereochemistry of aldol adduct **7e** was determined by X-ray crystallography,^[18] and other products were assigned by analogy. For electron-donating aromatic and aliphatic aldehydes, the relative rate of irreversible stereoselective C–C formation to afford **7** was slower than the irreversible protonation of the chiral enolate **8** to **4b** (see Scheme 2), resulting in diminished yields (R = 4-MePh: 48% yield, *syn:anti* >30:1, or 89:11). Under proton transfer conditions, minimization of undesired protonation of the chiral enolate **8**^[7d] remains a challenge despite the enhanced basicity of the chiral catalyst **6c**.^[19]

A catalytic cycle to account for the observed reactivity of the α -hydroxy phosphonate **1e** in the aldol addition is proposed in Scheme 2. Following deprotonation of the alcohol **1e** by the basic catalyst, [1,2]-phosphonate-phosphate rearrangement proceeds to generate the reactive glycolate enolate **8**. FMO considerations^[20] lead us to favor the illustrated (*Z*)-enolate as the kinetically preferred product of C→O phosphinyl migration. Enolate addition to the aldehyde would lead to the aldolate **9**. The precise structures of the enolate **8** and the aldol

transition state remain open questions. Approximately thermoneutral O→O phosphinyl migration occurs from the 3° alcohol to the vicinal 2° alkoxide, presumably to reduce steric strain at the fully substituted center, affording aldolate **10**. The reaction is rendered catalytic via proton transfer between the aldolate **10** and the α-hydroxy phosphonate starting material (**1e**), either directly or via the free base **6**.

The title reaction is especially attractive since the aldolization directly installs a leaving group that can be immediately deployed in nucleophilic displacement chemistry. The –OPO₃R₂ groups in benzylic phosphates are viable nucleofuges in acid-promoted Friedel-Crafts alkylations.^[21] Upon treatment of (±)-**3a** (dr 2:1) and anisole with TfOH, the α-hydroxy ester **11** was isolated in 91% yield as a single diastereomer. Heteroatom and heteroaromatic nucleophiles were also employed in this Friedel-Crafts alkylation in good to excellent yields (Scheme 3),^[22] providing stereogenic β-diaryl glycolates in a single step with pronounced stereoconvergence. Formation of a discrete carbenium ion explains the stereoconvergence and finds precedent in the diastereoselective Friedel-Crafts alkylations pioneered by Bach and coworkers.^[23]

In conclusion, a catalytic direct aldol addition of α-hydroxy trialkyl phosphonoacetates to aldehydes to afford α-hydroxy β-phosphonyloxy esters has been developed. A [1,2]-phosphonate-phosphate rearrangement was utilized to generate the reactive glycolate enolate in situ. The reaction works well for a variety of alkyl, alkenyl, aryl, and heteroaryl aldehydes affording the desired products in good to excellent yields in low to moderate diastereoselectivities. Iminophosphorane catalysts enabled positive outcomes in asymmetric versions of the title process providing excellent levels of stereocontrol. Stereoconvergent second stage transformations have also been developed to enhance the synthetic utility of the methodology. The applicability of this mechanistic framework in the context of other new reactions is the topic of ongoing investigations.

Experimental Section

A dried test tube was charged with the α-hydroxy phosphonoacetate **1e** (30.2 mg, 0.10 mmol, 1.0 equiv) and benzaldehyde (51 μL, 0.50 mmol, 5.0 equiv) and dissolved in 2-MeTHF (500 μL, 0.2 M) under an atmosphere of argon. The solution was cooled to –50 °C. Iminophosphorane **6c** (5.81 mg, 0.01 mmol, 0.1 equiv) was added and the reaction stirred at –50 °C for 20 h. The reaction was quenched at –50 °C with 0.5 M TFA in toluene (40 μL, 0.2 equiv). The resulting solution was diluted with 1 N HCl at 0 °C. The aqueous layer was extracted with CHCl₃ (3×). The combined organic extracts were washed with brine (1×), dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The diastereomeric ratio was determined by ¹H NMR analysis of the crude residue. The residue was purified by column chromatography on silica gel eluting with 30% acetone:hexane to afford the aldol adduct **7a** (29.0 mg, 0.07 mmol, 71% yield, >30:1 *syn:anti*) as a white solid (mp 119–127 °C).

References

1. Trost BM, Brindle CS. Chem. Soc. Rev. 2010; 39:1600–1632. [PubMed: 20419212]
2. a) Yoshikawa N, Yamada YMA, Das J, Sasai H, Shibasaki M. J. Am. Chem. Soc. 1999; 121:4168–4178. b) Yoshikawa N, Kumagai N, Matsunaga S, Moll G, Ohshima T, Suzuki T, Shibasaki M. J. Am. Chem. Soc. 2001; 123:2466–2467. [PubMed: 11456913] c) Trost BM, Ito H, Silcoff ER. J. Am. Chem. Soc. 2001; 123:3367–3368. [PubMed: 11457073] d) List B, Lerner RA, Barbas CF III. J. Am. Chem. Soc. 2000; 122:2395–2396. e) Notz W, List B. J. Am. Chem. Soc. 2000; 122:7386–7387. f) Northrup AB, MacMillan DWC. J. Am. Chem. Soc. 2002; 124:6798–6799. [PubMed: 12059180]
3. Selected examples: Evans DA, Downey CW, Hubbs JL. J. Am. Chem. Soc. 2003; 125:8706–8707. [PubMed: 12862448] Magdziak D, Lalic G, Lee HM, Fortner KC, Aloise AD, Shair MD. J. Am.

- Chem. Soc. 2005; 127:7284–7285. [PubMed: 15898756] Willis MC, Cutting GA, Piccio VJ-D, Durbin MJ, John MP. *Angew. Chem.* 2005; 117:1567–1569. *Angew. Chem. Int. Ed.* 2005; 44:1543–1545. Saito S, Kobayashi S. *J. Am. Chem. Soc.* 2006; 128:8704–8705. [PubMed: 16819844] Matsubara R, Berthiol F, Kobayashi S. *J. Am. Chem. Soc.* 2007; 130:1804–1805. [PubMed: 18198871] Saito S, Tsubogo T, Kobayashi S. *Chem. Commun.* 2007:1236–1237. Uraguchi D, Ueki Y, Ooi T. *J. Am. Chem. Soc.* 2008; 130:14088–14089. [PubMed: 18834197] Alonso DA, Kitagaki S, Utsumi N, Barbas CF III. *Angew. Chem.* 2008; 120:4664–4667. *Angew. Chem. Int. Ed.* 2008; 47:4588–4591. Utsumi N, Kitagaki S, Barbas CF III. *Org. Lett.* 2008; 10:3405–3408. [PubMed: 18630874] Van Nguyen H, Matsubara R, Kobayashi S. *Angew. Chem.* 2009; 121:6041–6043. *Angew. Chem. Int. Ed.* 2009; 48:5927–5929. Trost BM, Malhotra S, Fried BA. *J. Am. Chem. Soc.* 2009; 131:1674–1675. [PubMed: 19191696] Liu X, Deng L, Jiang X, Yan W, Liu C, Wang R. *Org. Lett.* 2010; 12:876–879. [PubMed: 20088540] Misaki T, Takimoto G, Sugimura T. *J. Am. Chem. Soc.* 2010; 132:6286–6287. [PubMed: 20408533] Melhado AD, Amarante GW, Wang ZJ, Luparia M, Toste FD. *J. Am. Chem. Soc.* 2011; 133:3517–3527. [PubMed: 21341677] Kobayashi S, Kiyohara H, Yamaguchi M. *J. Am. Chem. Soc.* 2011; 133:708–711. [PubMed: 21171607] Iwata M, Yazaki R, Chen IH, Sureshkumar D, Kumagai N, Shibasaki M. *J. Am. Chem. Soc.* 2011; 133:5554–5560. [PubMed: 21417332].
4. a) Brook AG. *J. Am. Chem. Soc.* 1958; 80:1886–1889. b) Brook AG. *Acc. Chem. Res.* 1974; 7:77–84.
 5. a) Fitch SJ, Moedritzer K. *J. Am. Chem. Soc.* 1962; 84:1876–1879. b) Hammerschmidt F, Schneyder E, Zbiral E. *Chem. Ber.* 1980; 113:3891–3897.
 6. a) Bausch CC, Johnson JS. *Adv. Synth. Catal.* 2005; 347:1207–1211. b) Demir AS, Reis Ö, dir AÇ, Esiringü , Eymur S. *J. Org. Chem.* 2005; 70:10584–10587. [PubMed: 16323878] c) Demir AS, Reis BR, Reis Ö, Eymür S, Göllü M, Tural S, Saglam G. *J. Org. Chem.* 2007; 72:7439–7442. [PubMed: 17715973] d) Demir AS, Esiringü , Göllü M, Reis Ö. *J. Org. Chem.* 2009; 74:2197–2199. [PubMed: 19199662]
 7. a) Romanov GV, Yagfavor MS, Konovalo AI, Pudovik AN, Konovalo IV, Yusupova TN. *Zh. Obshch. Khim.* 1973; 43:2378–2386. b) Pudovik AN, Zimin MG, Sobanov AA. *Zh. Obshch. Khim.* 1975; 45:1438–1444. c) Binder J, Zbiral E. *Tetrahedron Lett.* 1984; 25:4213–4216. d) Hayashi M, Nakamura S. *Angew. Chem.* 2011; 123:2297–2300. *Angew. Chem. Int. Ed.* 2011; 50:2249–2252.
 8. Ruel R, Bouvier J-P, Young RN. *J. Org. Chem.* 1995; 60:5209–5213.
 9. a) Ito M, Koyakumar K-i, Ohta T, Takaya H. *Synthesis.* 1995:376–378. b) Jiménez-Hopkins M, Hanson PR. *Org. Lett.* 2008; 10:2223–2226. [PubMed: 18447383] c) Krawczyk H, W sek K, K dzia J. *Synthesis.* 2009:1473–1476.
 10. a) Mitchell RE, Coddington JM, Young H. *Tetrahedron Lett.* 1989; 30:501–504. b) Kohama T, Enokita R, Okazaki T, Miyaoka H, Torikata A, Inukai M, Kaneko I, Kagasaki T, Sakaida Y, Satoh A, Shirashi A. *J. Antibiot.* 1993; 46:1503–1511. [PubMed: 7503975] c) Kohama T, Nakamura T, Kinoshita T, Kaneko I, Shirashi A. *J. Antibiot.* 1993; 46:1512–1519. [PubMed: 7503976] d) Fushimi S, Nishikawa S, Shimazu A, Seto H. *J. Antibiot.* 1989; 42:1019–1025. [PubMed: 2753808] e) Fushimi S, Furihata K, Seto H. *J. Antibiot.* 1989; 42:1026–1036. [PubMed: 2753809]
 11. Nakamura E. *Tetrahedron Lett.* 1981; 22:663–666.
 12. Ballester M. *Chem. Rev.* 1955; 55:283–300.
 13. See the Supporting Information for more details.
 14. a) Uruguchi D, Sakaki S, Ooi T. *J. Am. Chem. Soc.* 2007; 129:12392–12393. [PubMed: 17880221] b) Uruguchi D, Ito T, Ooi T. *J. Am. Chem. Soc.* 2009; 131:3836–3837. [PubMed: 19249866] c) Uruguchi D, Ito T, Nakamura S, Ooi T. *Chem. Lett.* 2009; 38:1052–1053. d) Uruguchi D, Ito T, Nakamura S, Ooi T. *Chem. Sci.* 2010; 1:488–490. e) Uruguchi D, Nakamura S, Ooi T. *Angew. Chem.* 2010; 122:7724–7727. *Angew. Chem. Int. Ed.* 2010; 49:7562–7565.
 15. This is a conservative estimate. In most cases, the minor diastereomer is not detectable by ¹H NMR spectroscopy; however, the lowest observed d.r. is 34:1.
 16. See the Supporting Information for detailed optimization experiments.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

17. Other pronucleophiles were employed in the reaction with 4-bromobenzaldehyde; however, at our current level of reaction and catalyst optimization both **1c** (34% yield, >30:1 *syn:anti*, 87% *ee*) and **1d** (95% yield, >30:1 *syn:anti*, 70% *ee*) provided unsatisfactory levels of either reactivity or enantioselectivity despite an observed retention in absolute diastereocontrol.
18. CCDC 854995 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
19. Competitive enolate quenching was also observed in the KO^tBucatalyzed aldolization (Table 2). Entries 11, 13, and 14 necessitated 5 equivalents of aldehyde to compensate for their reduced electrophilicity.
20. Schmitt DC, Johnson JS. *Org. Lett.* 2010; 12:944–947. [PubMed: 20143793]
21. Smith AG, Johnson JS. *Org. Lett.* 2010; 12:1784–1787. [PubMed: 20235527]
22. See the Supporting Information for the determination of the relative stereochemistry of **11**.
23. a) Mühlthau F, Schuster O, Bach T. *J. Am. Chem. Soc.* 2005; 127:9348–9349. [PubMed: 15984844] b) Mühlthau F, Stadler D, Goepfert A, Olah GA, Prakash GKS, Bach T. *J. Am. Chem. Soc.* 2006; 128:9668–9675. [PubMed: 16866519] c) Stadler D, Bach T. *Chem. Asian J.* 2008; 3:272–284. [PubMed: 17985329] d) Sun B-F, Hong R, Kang Y-B, Deng L. *J. Am. Chem. Soc.* 2009; 131:10384–10385. [PubMed: 19601599]

Scheme 1.
Base-Catalyzed Direct-Aldolization Employing Phosphonate-Phosphate Rearrangement

Scheme 2.
Mechanistic Proposal

Scheme 3.
Stereoconvergent Friedel-Crafts Alkylations

Direct Aldol Addition: Reaction Optimization^[a]

$[f]_{[1]0} = 0.20 \text{ M}$, 2.0 equiv aldehyde.

Table 2

Aldehyde Substrate Scope^[a]

entry	R	product	yield [%] ^[b]	<i>anti:syn</i> ^[c]
1	4-ClC ₆ H ₄	3a	95	2.0:1.0
2	2-FC ₆ H ₄	3b	91	2.3:1.0
3	2-NO ₂ C ₆ H ₄	3c	87	1.5:1.0
4	3-NO ₂ C ₆ H ₄	3d	92	2.4:1.0
5	4-NO ₂ C ₆ H ₄	3e	95	2.8:1.0
6	4-CF ₃ C ₆ H ₄	3f	91	2.1:1.0
7	4-CNC ₆ H ₄	3g	93	2.4:1.0
8	C ₆ H ₅	3h	97	2.1:1.0
9	4-FC ₆ H ₄	3i	98	2.1:1.0
10	4-MeC ₆ H ₄	3j	97	2.0:1.0
11 ^[d]	4-OMeC ₆ H ₄	3k	89	4.9:1.0
12	2-thienyl	3l	81 ^[e]	6.7:1.0
13 ^[d]	(<i>E</i>)-CH=CHPh	3m	89	4.4:1.0
14 ^[d]	CH ₂ CH ₂ Ph	3n	78	5.8:1.0

^[a]Unless noted, reactions were performed on 0.20 mmol scale, using 2.0 equiv of aldehyde in THF (1.0 mL) at $-78\text{ }^\circ\text{C}$ for 2 h;

^[b]Isolated yield;

^[c]Determined by ^1H NMR analysis of the crude reaction mixture;

^[d]With 5.0 equiv of aldehyde;

^[e]Product isolated as the *anti* diol following column chromatography.

Table 3

 α -Substituent Scope^[a]

entry	R	product	yield [%] ^[b]	anti:syn ^[c]
1	H (1f)	3o	trace	-
2	Me (1b)	3p	81	1.9:1.0
3	CH ₂ CH=CH ₂ (1c)	3q	89	1.2:1.0
4	CH ₂ C≡CH (1d)	3r	93	2.9:1.0
5	CH ₂ Ph (1a)	3h	97	2.1:1.0

^[a]Reactions were performed on 0.20 mmol scale, using 2.0 equiv of aldehyde in THF (1.0 mL) at -78 °C for 2 h;

^[b]Isolated yield;

^[c]Determined by ¹H NMR analysis of the crude reaction mixture.

Table 4

Development of an Asymmetric Aldolization^[16]

entry	6	solvent	T [°C]	conv. [%] ^[b]	7:4b <i>syn:anti</i> ^[b]	e.r. ^[c]
1	6a	THF	-40	<5	-	-
2	6b	THF	-40	53	1:0.6	>30:1
3	6c	THF	-40	76	1:0.8	>30:1
4	6d	THF	-40	80	1:0.7	>30:1
5	6e	THF	-40	>95	1:1.1	>30:1
6	6c	Et ₂ O	-40	>95	1:0.6	>30:1
7	6c	2-MeTHF	-40	>95	1:0.8	>30:1
8	6c	2-MeTHF	-50	>90 (71) ^[d]	1:0.4	>30:1

^[a]Reactions were performed on 0.10 mmol scale, using 5.0 equiv of PhCHO in solvent (0.5 mL) for 20 h;

^[b]Conversions and diastereomeric ratios were determined by ¹H NMR spectroscopy of the crude reaction mixture;

^[c]Determined by chiral HPLC using a CHIRALPAK AD3 column;

^[d]Isolated yield.

Table 5

Diastereo- and Enantioselective Direct Aldol Additions^[a]

entry	R	product	yield [%] ^[b]	<i>syn:anti</i> ^[c]	e.r. ^[d]
1	C ₆ H ₅	7a	71	>30:1	95:5
2	2-naphthyl	7b	68	>30:1	94.5:5.5
3	2-FC ₆ H ₄	7c	56	>30:1	94:6
4	2-ClC ₆ H ₄	7d	64	>30:1	95:5
5	3-NCC ₆ H ₄	7e	89	>30:1	94:6
6	3-BrC ₆ H ₄	7f	87	>30:1	95:5
7	4-ClC ₆ H ₄	7g	82	>30:1	95:5
8	4-BrC ₆ H ₄	7h	91	>30:1	95.5:4.5
9	4-MeO ₂ CC ₆ H ₄	7i	89	>30:1	97:3

^[a]Reactions were performed on 0.10 mmol scale, using 5.0 equiv of aldehyde in 2-MeTHF (0.5 mL) at -50 °C for 20 h;^[b]Isolated yield;^[c]Determined by ¹H NMR analysis of the crude reaction mixture.^[d]Determined by chiral HPLC using a CHIRALPAK AD3 column;