Published in final edited form as:

Tetrahedron. 2013 April 22; 69(16): 3432–3436. doi:10.1016/j.tet.2013.02.073.

De novo asymmetric synthesis of oligo-rhamno di- and tri-saccharides related to the anthrax tetrasaccharide

Hua-Yu Leo Wang^{a,b}, Haibing Guo^{a,*}, and George A. O'Doherty^{b,*}

^aCollege of Nuclear Technology, Chemistry and Biology, Hubei University of Science and Technology, Xianning, Hubei 437100, China

^bDepartment of Chemistry and Chemical Biology, Northeastern University, 360 Huntington Avenue, Boston, MA 02115, USA

Abstract

An asymmetric synthesis of the di- and trisaccharide portion of the naturally occurring anthrax tetrasaccharide from acetylfuran has been developed. The construction of the di- and trisaccharide subunits is based upon our previously disclosed route to anthrax tetrasaccharide. The approach uses iterative diastereoselective palladium-catalyzed glycosylations, Luche reductions, diastereoselective dihydroxylations, and regioselective protections for the assembly of the rhamno- di- and tri-saccharide. The route was also modified for the preparation of the mixed D-/L-disaccharide analogue.

Keywords

Anthrax; Oligo-rhamnose; Palladium; glycosylation

1. Introduction

Until late 2001, when the unfortunate event of a powdery sample of *Bacillus anthracis* was sent through the U.S. Postal Service,¹ most cultural references to Anthrax were to the American thrash metal band from New York City. The band was formed in 1981 and named after the disease caused by *Bacillus anthracis*.² The spore-forming *Bacillus anthracis* is a Gram-positive bacterium which when inhaled causes anthrax, a fatal infectious disease in humans and other mammals.³ The mature endospores exhibit remarkable resistance to extremely harsh conditions, which allow the spores to persist for many years,⁴ and make them ideal for use as biological weapon.

While the origin of the weaponized material is still in dispute, the effects of having it sent through the post in 2001 are clear. As a result, 22 people were confirmed to be infected with *Bacillus anthracis* and only 7 survived.⁵ In fact, all the 7 survivors are believed to have the more easily treated cutaneous form of anthrax. In response, to the seriousness of the threat,

© 2013 Elsevier Ltd. All rights reserved.

*Corresponding author. g.o'doherty@neu.edu. *Corresponding author. haibingguojj@yahoo.com.

Supplementary Material

Copies of the spectral data ¹H NMR and ¹³C NMR for all the new compounds can be found in the Supporting Information.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

there have been extensive efforts aimed at the discovery of new inexpensive treatments (e.g., vaccines and antibacterials) for the disease,⁶ as well as, methods for the detection of the *Bacillus anthracis* spores (e.g., antibodies).⁷ As part of the efforts to develop methods for detecting *Bacillus anthracis*, the anthrax tetrasaccharide **1** (Figure 1) was isolated from BclA, a surface glycoprotein found on the exosporium of *Bacillus anthracis*.⁸ The unusual structure of anthrax tetrasaccharide **1** consists of unique tris-L-trisaccharide portion terminated by a never before seen D-aminosugar, called anthrose. These unique structural features in conjunction with the general resistance to evolutionary change of carbohydrates, has made the anthrax tetrasaccharide **1** an excellent target for anthrax detection and in turn total synthesis.^{9,10}

Since its discovery of anthrax tetrasaccharide **1** in 2004, there have been several syntheses of analogues of **1** with various alkyl groups at the reducing end.¹¹ The addition of functional groups at the anomeric position was made to give the anthrax tetrasaccharide a chemical handle for further conjugation and subsequent manipulations (e.g., antibody/vaccine development). In addition to analogue material with a chemical handle, our de novo approach to the anthrax tetrasaccharide **1** was the only route to prepare synthetic material identical to the isolated natural product.¹⁰ Our synthetic **1** was prepared as part of a broad search aimed at the development of a method for the selective detection of **1** and thus anthrax.⁷ To address concern that these methods might focus on the more common tris-rhamnose portion of the molecule, we needed control compounds consisting of the rhamno-di- and tri-saccharide structural motif. Thus, we targeted the all L-trisaccharide **2** and the all L-disaccharide **3** as well as the mixed D/L-disaccharide **4** for synthesis using our previously described approach to anthrax tetrasaccharide **1**.¹⁰ Herein, we describe our successful efforts toward the de novo asymmetric synthesis of anthrax-based rhamno-oligosaccharides **2**, **3** and **4** (Figure 1). This de novo asymmetric strategy features iterative use of palladium-catalyzed glycosylation and starts from a commercially available and inexpensive acetylfuran.¹²

Our de novo approach to anthrax oligosaccharides **2–4** is outlined in Scheme 1. Based on our previous route to **1**, we envisioned access the unique rhamno- oligosaccharide scaffold in both anthrax trisaccharide analogue **2** and disaccharide analogues **3** and **4** from the protected intermediates **5–7**. These precursors, in turn, could be prepared in a highly divergent manner from a glycosylation of Leyspiroketal disaccharide **7** with pyranone **8** or (*ent*)-**8**. The rapid assembly of the desired rhamno sugar relies on the iterative use of palladium-catalyzed glycosylation, Luche reduction, and Upjohn dihydroxylation. The required pyranone sugar donor in both D- and L- configuration could be obtained from the achiral starting material acetylfuran **9** through asymmetric Noyori reduction and Achmatowicz oxidative ring expansion.

2. Results/Discussion

Our approach to the monosaccharide building block began with our efforts to selectively introduce the D- or L-stereochemistry of the rhamnose, which could be derived from the furyl alcohol (*R*)-**10** or (*S*)-**10**. Previously we have shown that either (*R*) or (*S*)-furyl alcohol could be readily prepared by an asymmetric Noyori reduction of commercially available starting material acetylfuran **9** (Scheme 2).¹³ Noyori reduction of acetylfuran **9** produced the enantiomerically pure (> 96% *ee*)¹⁴ furfuryl alcohol (*R*)-**10** or (*S*)-**10**, which was exposed in the subsequent Achmatowicz oxidative rearrangement (NBS/H₂O) and *tert*-butoxyl-carbonylation to give the desired Boc-pyranone α -D-(*ent*)-**8** or α -L-**8** with good α -anomeric diastereoselectivity (α : β : γ :1).¹⁵ It is worth noting that this route provides Boc-pyranones in 60% overall yield (three steps) with only one chromatographic purification and this route is also scalable for both L- and D-enantiomers.

Using our previously described route to anthrax tetrasaccharide **1**, we prepared the desired protected known precursors **5**, **6**, and **7** (Scheme 3).¹⁰ In brief, monosaccharide **7** with the free C-2 hydroxyl group can glycosylated with Boc-pyranone α -L-**8** and in two post-glycosylation transformations (NaBH₄/CeCl₃ then OsO₄/NMO) converted into disaccharide **6**. The conversion of disaccharide **6** into trisaccharide **5** began with the selective protection of the C-2/C-4 hydroxyls via a one-pot orthoester/acylation/hydrolysis procedure to give **12**. Once again, a three-step palladium-catalyzed glycosylation/post-glycosylation sequence was used to install the final rhamnose sugar in trisaccharide **5** with excellent yield and diastereoselectivity.

The preparation of the unknown L-D-disaccharide diastereomer **4** the anthrax disaccharide analogue **3** began with the common Ley-protected rhamnose intermediate **7**. In a diastereomeric fashion, the C-2 hydroxyl in **7** was glycosylated with Boc-pyranone (*ent*)-**8** to give **14** in a nearly identical yield (85%) for the palladium-catalyzed glycosylation (Scheme 4). Similarly, the resulting enone in L-D-disaccharide **14** was reduced to give **15** (90%) and dihydroxylated to give **16** (93%), in excellent yield and stereoselectivity. Thus in a diastereomerically divergent fashion, the Ley-protected L-rhamnose **7** can be glycosylated in a three-step sequence to install either a D- or L-rhamnose sugar in nearly identical yields.

Finally, all the desired anthrax di- and tri-saccharide analogues **2**, **3** and **4** were obtained after a two- or three-step deprotection sequence. Simply treating D/L-diastereomers **6** and **16** to aqueous TFA gave disaccharides **17** and **18**, respectively. The desired fully deprotected diastereomeric disaccharides **3** and **4** were obtained by hydrogenolysis (1 atm H₂, Pd/C) in excellent yields (85 and 88% over two steps). The deprotection of trisaccharide **5** was slightly more complicated due to concomitant acetate partial hydrolysis and migration. Thus, the crude product from exposure of **5** to aqueous TFA was immediately followed by the treatment with basic methanol and hydrogenolysis to give trisaccharide **2** in excellent overall yield (81%).

3. Conclusion

In conclusion, a highly enantio- and diastereo-controlled approach to the anthrax analogues has been developed in a divergent manner using a Ley-protected rhamno-monosaccharide **7** as the common intermediate for the assembly of *rhamno*-di- and tri-saccharides (*i.e.*, **2** and **3**) and the incorporation of D-L-*bis-rhamno*-disaccharide diastereomer (*i.e.*, **4**). The feasibility of this was evident in that both the di- and tri-saccharide fragments were readily assembled from achiral starting materials by use of asymmetric catalysis. This de novo asymmetric route illustrates the utilities of Noyori reduction, palladium-catalyzed glycosylation, diastereoselective dihydroxylation, Luche reduction and selective acylation. Further use of these substrates for the screening of colorimetric assays for anthrax detection is ongoing and will be reported in due course.

4. Experimental Section

(2R,6S)-6-(((2R,3R,4aS,5S,7R,8R,8aR)-7-(benzyloxy)-2,3-dimethoxy-2,3,5-trimethylhexahydro-2H-pyrano[3,4b]-[1,4]-dioxin-8-yl)oxy)-2-methyl-2H-pyran-3(6H)-one (14)

To a solution of Boc-pyranone (*ent*)-**8** (846.6 mg, 3.713 mmol) and Ley-spiroketal rhamnose **7** (171 mg, 0.464 mmol) in dry CH₂Cl₂ (0.9 mL) was added a solution of Pd₂(dba)₃ · CHCl₃ (12.01 mg, 0.012 mmol) and PPh₃ (12.2 mg, 0.046 mmol) in 1.0 mL CH₂Cl₂ at 0 °C. The reaction mixture was stirred at 0 °C under argon gas for 4 h. The reaction mixture was quenched with 10 mL of saturated NaHCO₃ solution, followed by extraction with Et₂O (10 mL x 3). The organic layer was washed with 15 mL saturated brine solution, dried over with

Na₂SO₄ and concentrated under reduced pressure. The compound was purified using flash chromatography eluting with 25% EtOAc in hexanes to give the title compound **14** (188 mg, 0.39 mmol, 85%); a yellow oil; *R*_f 0.50 (30% EtOAc/hexanes); [α]₂₅^D = −138 (*c* 1.0, CH₂Cl₂); IR (thin film, cm^{−1}) 2995, 2940, 2909, 2836, 1698, 1459, 1374, 1114, 1035. 968; ¹H NMR (600 MHz, CDCl₃) δ 7.34–7.26 (m, 5H), 6.79 (dd, *J* = 10.2, 3.6 Hz, 1H), 6.03 (d, *J* = 10.2 Hz, 1H), 5.12 (d, *J* = 3.6 Hz, 1H), 5.02 (q, *J* = 6.6 Hz, 1H), 4.78 (d, *J* = 0.6 Hz, 1H), 4.68 (d, *J* = 12 Hz, 1H), 4.49 (d, *J* = 12 Hz, 1H), 4.06 (dd, *J* = 3.6, 1.8 Hz, 1H), 4.04 (dd, *J* = 10.2, 3.6 Hz, 1H), 3.82 (dq, *J* = 10.2, 6.6 Hz, 1H), 3.64 (dd, *J* = 10.2, 10.2 Hz, 1H), 3.25 (s, 3H), 3.17 (s, 3H), 1.31 (d, *J* = 6.6 Hz, 3H), 1.24 (d, *J* = 6.0 Hz, 3H), 1.24 (s, 3H), 1.22 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ 198.3, 143.3, 137.3, 128.6(2C), 128.2(2C), 128.1, 127.8, 100.0, 99.7, 98.3, 92.8, 75.2, 70.8, 69.3, 68.8, 67.5, 67.4, 48.1, 47.8, 18.0, 17.9, 16.8, 15.3; HRMS (CI): calcd. for [C₂₅H₃₄O₉Na⁺]: 501.20950, found: 501.20966.

(2*R*,3*S*,6*R*)-6-(((2*R*,3*R*,4*aS*,5*S*,7*R*,8*R*,8*aR*)-7-(benzyloxy)-2,3-dimethoxy-2,3,5-trimethylhexahydro-2*H*-pyrano[3,4*b*]-[1,4]-dioxin-8-yl)oxy)-2-methyl-3,6-dihydro-2*H*-pyran-3-ol (15)

A solution of disaccharide pyranone **14** (48 mg, 0.100 mmol) in dry CH₂Cl₂ (0.1 mL) and 0.4 M CeCl₃/MeOH (0.1 mL) was cooled to −78 °C. NaBH₄ (5 mg, 0.120 mmol) was added and the reaction mixture was stirred for 2 h. The reaction mixture was then diluted with Et₂O (5 mL) and quenched with saturated solution of NaHCO₃ (2 mL), then extracted with Et₂O (5 mL x 3). The organic layer was dried over with Na₂SO₄, concentrated under reduced pressure, and purified using flash chromatography with 35% EtOAc in hexanes elution to give the title compound **15** (43 mg, 0.09 mmol, 90%); colorless oil; *R*_f 0.29 (40% EtOAc/hexanes); [α]₂₅^D = −92 (*c* 1.0, CH₂Cl₂); IR (thin film, cm^{−1}) 3454, 2983, 2940, 2936, 2902, 2836, 1454, 1378, 1138, 1118, 1040, 999, 734; ¹H NMR (600 MHz, CDCl₃) δ 7.33–7.25 (m, 5H), 5.92 (d, *J* = 10.2 Hz, 1H), 5.74 (ddd, *J* = 10.2, 2.4, 1.8 Hz, 1H), 4.87 (d, *J* = 2.4 Hz, 1H), 4.75 (s, 1H), 4.67 (d, *J* = 12.0 Hz, 1H), 4.57 (d, *J* = 12.0 Hz, 1H), 4.04 (dd, *J* = 3.6, 1.8 Hz, 1H), 4.02 (dq, *J* = 9.0, 6.0 Hz, 1H), 3.99 (dd, *J* = 10.2, 3.6 Hz, 1H), 3.80 (dq, *J* = 10.2, 6.0 Hz, 1H), 3.63 (dd, *J* = 10.2, 9.6 Hz, 1H), 3.25 (s, 3H), 3.21 (s, 3H), 1.26 (d, *J* = 6.0 Hz, 3H), 1.26 (s, 3H), 1.25 (s, 3H), 1.24 (d, *J* = 6.0 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) δ 137.5, 133.9, 128.6(2C), 128.2(2C), 128.0, 126.8, 100.0, 99.8, 98.3, 93.1, 73.7, 70.1, 69.3, 68.9, 68.1, 67.2, 67.1, 48.0, 47.9, 18.3, 18.1, 17.9, 16.9; HRMS(CI): calcd. for [C₂₅H₃₆O₉Na⁺]: 503.22515, found: 503.22531.

(2*R*,3*S*,4*S*,5*S*,6*R*)-2-(((2*R*,3*R*,4*aS*,5*S*,7*R*,8*R*,8*aR*)-7-(benzyloxy)-2,3-dimethoxy-2,3,5-trimethylhexahydro-2*H*-pyrano-[3,4-*b*]-[1,4]dioxin-8-yl)oxy)-6-methyltetrahydro-2*H*-pyran-3,4,5-triol (16)

To a solution of allylic alcohol **15** (30 mg, 0.063 mmol) in *tert*-butanol/acetone (125 μ L, 1:1 (v/v), 0.5M) at 0 °C was added a solution of *N*-methymorpholine-*N*-oxide/water (50% w/v, 65 μ L, 1M). Crystalline OsO₄ (0.16 mg, 1 mol%) was added and the reaction mixture was stirred for 12 h at 0 °C. The reaction mixture was then quenched with 500 μ L of saturated Na₂S₂O₃ solution, extracted with EtOAc (5 x 2 mL), and dried over with Na₂SO₄ and concentrated under reduced pressure. The compound was purified by flash chromatography eluting with 85% EtOAc in hexanes to give the title compound **16** (29.9 mg, 0.058 mmol, 93%); colorless oil; *R*_f 0.56 (10% MeOH in EtOAc); [α]₂₅^D = −50 (*c* 1.0, CH₂Cl₂); IR (thin film, cm^{−1}) 3304, 2938, 2775, 2100, 1128, 1050; ¹H NMR (600MHz, CDCl₃) δ 7.35–7.25 (m, 5H), 4.75 (d, *J* = 0.6 Hz, 1H), 4.74 (d, *J* = 1.2 Hz, 1H), 4.67 (d, *J* = 12.0 Hz, 1H), 4.47 (d, *J* = 12.0 Hz, 1H), 4.26 (dq, *J* = 9.6, 6.0 Hz, 1H), 3.99 (dd, *J* = 10.2, 3.0 Hz, 1H), 3.96 (m, 1H), 3.95 (dd, *J* = 3.6, 1.8 Hz, 1H), 3.81 (dq, *J* = 9.6, 6.0 Hz, 1H), 3.64 (dd, *J* = 10.2, 10.2 Hz, 1H), 3.40 (dd, *J* = 9.6, 9.6 Hz, 1H), 3.23 (s, 3H), 3.22 (s, 3H), 1.25 (s, 3H), 1.24 (d, *J* = 6.0 Hz, 3H), 1.24 (s, 3H), 1.24 (d, *J* = 6.0 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) δ 137.4, 128.7(2C), 128.2(2C), 128.1, 99.9, 99.7, 98.5, 97.7, 74.00, 73.95, 72.2, 71.3, 69.4, 68.8,

68.0, 67.5, 67.1, 48.2, 47.9, 18.0, 17.9, 17.5, 16.9; HRMS (CI): calcd. for [C₂₅H₃₈O₁₁Na⁺]: 537.23063, found: 537.23074.

(2R,3S,4S,5S,6R)-2-(((2R,3R,4R,5R,6S)-2-(benzyloxy)-4,5-dihydroxy-6-methyltetrahydro-2H-pyran-3-yl)oxy)-6-methyltetrahydro-2H-pyran-3,4,5-triol (18)

To a solution of disaccharide **16** (19.3 mg, 0.0375 mmol) in CH₂Cl₂ (375 μ L) was added 75 μ L of TFA-H₂O (10:1). The reaction was stirred at room temperature for 4 h. The reaction was then quenched with saturated NaHCO₃ solution (3.0 mL), dried over with Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by flash chromatography eluting with 8% MeOH in EtOAc to give the title compound **18** (14.7 mg, 0.0367 mmol, 98%); colorless oil; R_f 0.48 (20% MeOH in EtOAc); [α]₂₅^D = +5.0 (c 1.0, MeOH); IR (thin film, cm⁻¹) 3372, 2071, 1508, 1334, 1120, 973, 786; ¹H NMR (600 MHz, CD₃OD) δ 7.35–7.28 (m, 5H), 4.87 (d, *J* = 1.8 Hz, 1H), 4.74 (d, *J* = 1.8 Hz, 1H), 4.70 (d, *J* = 12.0 Hz, 1H), 4.54 (d, *J* = 12.0 Hz, 1H), 3.98 (dq, *J* = 9.6, 6.0 Hz, 1H), 3.92 (dd, *J* = 3.6, 1.8 Hz, 1H), 3.81 (dd, *J* = 3.6, 1.8 Hz, 1H), 3.79 (dd, *J* = 9.6, 3.6 Hz, 1H), 3.74 (dd, *J* = 9.6, 3.0 Hz, 1H), 3.62 (dq, *J* = 9.0, 6.0 Hz, 1H), 3.37 (dd, *J* = 9.6, 9.6 Hz, 1H), 3.35 (dd, *J* = 9.6, 9.6 Hz, 1H), 1.25 (d, *J* = 6.6 Hz, 3H), 1.24 (d, *J* = 6.6 Hz, 3H); ¹³C NMR (150 MHz, CD₃OD) δ 139.1, 129.6(2C), 129.2(2C), 129.0, 100.1, 97.9, 76.9, 74.3, 74.1, 72.6, 72.3, 71.7, 70.5, 70.4, 70.3, 18.2, 18.0; HRMS (CI): calcd. for [C₁₉H₂₈O₉+Na⁺]: 423.16255, found: 423.16269.

(3R,4R,5R,6S)-6-methyl-3-(((2R,3S,4S,5S,6R)-3,4,5-trihydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-tetrahydro-2H-pyran-2,4,5-triol (4)

To a solution of benzyl disaccharide **18** (11.4 mg, 0.030 mmol) in MeOH (0.1 mL) was added 10% Pd/C (60 mg). The reaction suspension was degassed and replaced atmosphere with hydrogen gas three times. The reaction mixture was stirred for 24 h and filtered through a pad of Celite. The solute was concentrated under reduced pressure to give the title compound **4** (8.2 mg, 0.026 mmol, 90%); viscous oil; R_f 0.52 (40% MeOH in EtOAc); [α]₂₅^D = +16 (c 1.0, MeOH); IR (thin film, cm⁻¹) 3375, 2991, 2942, 1570, 1145, 995; ¹H NMR (600 MHz, CD₃OD) δ 5.10 (d, *J* = 1.8 Hz, 1H), 4.79 (d, *J* = 1.2 Hz, 1H), 3.98 (dq, *J* = 9.6, 6.0 Hz, 1H), 3.86 (dd, *J* = 3.0, 1.8 Hz, 1H), 3.83 (dd, *J* = 3.6, 1.8 Hz, 1H), 3.81 (dd, *J* = 9.6, 3.6 Hz, 1H), 3.80 (dq, *J* = 9.6, 6.0 Hz, 1H), 3.75 (dd, *J* = 9.6, 3.6 Hz, 1H), 3.38 (dd, *J* = 9.6, 9.6 Hz, 1H), 3.32 (dd, *J* = 9.0, 9.0 Hz, 1H), 1.25 (d, *J* = 6.0 Hz, 3H), 1.24 (d, *J* = 6.0 Hz, 3H); ¹³C NMR (150 MHz, CD₃OD) δ 100.1, 92.7, 77.8, 74.6, 74.2, 72.6, 72.3, 71.4, 70.3, 69.8, 18.4, 18.1; HRMS (CI): calcd. for [C₁₂H₂₂O₉Na⁺]: 333.11560, found: 333.11566.

(3R,4R,5R,6S)-6-methyl-3-(((2S,3R,4R,5R,6S)-3,4,5-trihydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)-tetrahydro-2H-pyran-2,4,5-triol (3)

To a solution of disaccharide **6** (26 mg, 0.05 mmol) in CH₂Cl₂ (0.5 mL), was added the 0.1 mL of 10:1 TFA-H₂O. The reaction was stirred at rt for 40 minutes. The reaction was quenched with saturated 5 mL NaHCO₃ solution, dried over with Na₂SO₄, and concentrated to remove water and solvent under reduced pressure. The crude compound was purified by passing through a pad of silica gel eluting with 10% MeOH in EtOAc to give the benzyl disaccharide **17**; To a solution of the benzyl disaccharide **17** in 0.5 mL MeOH was added 10% Pd/C (100 mg). The reaction suspension was degassed and replaced atmosphere with hydrogen gas three times. The reaction suspension was stirred with hydrogen balloon for 24 h, filtered by passing through a pad of Celite, concentrated under reduced pressure and under vacuo to give the title compound **3** (13 mg, 0.043 mmol, 85%); viscous oil; R_f 0.31 (20% Methanol in EtOAc); [α]₂₅^D = -20 (c 1.0, MeOH); IR (thin film, cm⁻¹) 3351, 2981, 2935, 1645, 1055, 1032; ¹H NMR (600 MHz, CD₃OD) δ 5.20 (d, *J* = 1.8 Hz, 1H), 4.98 (d, *J* = 1.8 Hz, 1H), 3.89 (dd, *J* = 9.6, 3.6 Hz, 1H), 3.85 (dq, *J* = 9.6, 6.0 Hz, 1H), 3.84 (dd, *J* =

3.6, 1.8 Hz, 2H), 3.78 (dq, $J = 9.6, 6.0$ Hz, 1H), 3.75 (dd, $J = 9.6, 3.6$ Hz, 1H), 3.44 (dd, $J = 9.6, 9.6$ Hz, 1H), 3.39 (dd, $J = 9.6, 9.6$ Hz, 1H), 1.31 (d, $J = 6.0$ Hz, 3H), 1.29 (d, $J = 6.0$ Hz, 3H); ^{13}C NMR (150 MHz, CD_3OD) δ 104.2, 94.6, 81.2, 74.7, 74.1, 72.4, 72.2, 71.9, 70.3, 69.5, 18.4, 18.0; HRMS (CI): calcd. for $[\text{C}_{12}\text{H}_{22}\text{O}_9\text{Na}^+]$: 333.11560, found: 333.11568.

(3*R*,4*R*,5*R*,6*S*)-3-(((2*S*,3*R*,4*R*,5*S*,6*S*)-3,5-dihydroxy-6-methyl-4-(((2*S*,3*R*,4*R*,5*R*,6*S*)-3,4,5-trihydroxy-6-methyltetrahydro-2*H*-pyran-2-yl)oxy)-tetrahydro-2*H*-pyran-2-yl)oxy)-6-methyltetrahydro-2*H*-pyran-2,4,5-triol (2)

To a solution of trisaccharide **5** (42 mg, 0.05 mmol) in CH_2Cl_2 (0.5 mL) was added 0.1 mL of 10:1 TFA- H_2O . The reaction was stirred at rt for 40 minutes. The reaction was quenched with saturated 5 mL of NaHCO_3 solution, dried over with Na_2SO_4 , and concentrated to remove water and solvent under reduced pressure. The crude compound was purified by passing through a pad of silica gel eluting with 10% MeOH in EtOAc to give diacetate; To the solution of the subsequent diacetate in MeOH (0.2 mL) was added LiOH (9.6 mg, 0.4 mmol) and stirred for 5 h. The reaction mixture was concentrated under reduced pressure and purified by passing through a pad of silica gel eluting with 15% MeOH in EtOAc to give the benzyl trisaccharide **19**; The resulting benzyl trisaccharide **19** was dissolved in 0.5 mL MeOH and added 10% Pd/C (100 mg). The reaction suspension was degassed and replaced atmosphere with hydrogen gas for three times. The reaction suspension was stirred under the hydrogen balloon for 24 h, filtered by passing through a pad of Celite, concentrated under reduced pressure and under vacuo to give the title compound **2** (18.5 mg, 0.041 mmol, 81%); viscous oil; R_f 0.19 (20% MeOH in EtOAc); $[\alpha]_{25}^{\text{D}} = -82$ (c 1.2, MeOH); IR (thin film, cm^{-1}) 3356, 2981, 2934, 1572, 1416, 1059, 986; ^1H NMR (600 MHz, CD_3OD) δ 5.18 (d, $J = 1.8$ Hz, 1H), 5.10 (d, $J = 1.8$ Hz, 1H), 4.98 (d, $J = 1.8$ Hz, 1H), 4.12 (dd, $J = 3.6, 1.8$ Hz, 1H), 4.03 (dd, $J = 3.6, 1.8$ Hz, 1H), 3.86 (m, 7H), 3.57 (dd, $J = 9.6, 9.6$ Hz, 1H), 3.45 (dd, $J = 9.6, 9.6$ Hz, 1H), 3.43 (dd, $J = 9.6, 9.6$ Hz, 1H), 1.35 (d, $J = 6.0$ Hz, 3H), 1.31 (d, $J = 6.0$ Hz, 3H), 1.29 (d, $J = 6.0$ Hz, 3H); ^{13}C NMR (150 MHz, CD_3OD) δ 104.0(2C), 94.7, 80.9, 79.4, 74.7, 74.3, 73.5, 72.4, 72.3, 72.0(2C), 70.6, 70.3, 69.5, 18.4, 18.1, 18.0; HRMS (CI): calcd. for $[\text{C}_{18}\text{H}_{32}\text{O}_{13}\text{Na}^+]$: 479.17351, found: 479.17367.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to NIH (GM090259), NSF (CHE-1213596) and NSF of China (21202136) for the support of our research programs.

References and notes

- McCarthy M. Lancet Infect Dis. 2001; 1:288–289. [PubMed: 11871791]
- www.anthrax.com
- (a) Mock M, Fouet A. Annu Rev Microbiol. 2001; 55:647–671. [PubMed: 11544370] (b) Sylvestre P, Couture-Tosi E, Mock M. Mol Microbiol. 2002; 45:169–178. [PubMed: 12100557]
- Nicholson WL, Munakata N, Horneck G, Melosh HJ, Setlow P. Microbiol Mol Biol Rev. 2000; 64:548–572. [PubMed: 10974126]
- Inglesby TV, O'Toole T, Henderson DA, Bartlett JG, Ascher MS, Eitzen E, Friedlander AM, Gerberding J, Hauer J, Hughes J, McDade J, Osterholm MT, Parker G, Perl TM, Russell PK, Tonat K. JAMA. 2002; 288:2236–2252. [PubMed: 11980524]
- (a) Bouzianan DG. J Med Chem. 2010; 53:4305–4331. [PubMed: 20102155] (b) Chitlaru T, Altboum Z, Reuveny S, Shafferman A. Immunol Rev. 2011; 239:221–236. [PubMed: 21198675]

7. (a) Tamborrini M, Werz DB, Frey J, Pluschke G, Seeberger PH. *Angew Chem Int Ed.* 2006; 45:6581–6582. (b) Tamborrini M, Holzer M, Seeberger PH, Schurch N, Pluschke G. *Clin Vaccine Immunol.* 2010; 17:1446–1451. [PubMed: 20660139] (c) Saile E, Boons GJ, Buskas T, Carlson RW, Kannenberg EL, Barr JR, Boyer AE, Gallegos-Candela M, Quinn CP. *Clin Vaccine Immunol.* 2011; 18:743–748. [PubMed: 21389148]
8. Daubenspeck JM, Zeng H, Chen P, Dong S, Steichen CT, Krishna NR, Pritchard DG, Turnbough CL Jr. *J Biol Chem.* 2004; 279:30945–30953. [PubMed: 15152001]
9. (a) Werz DB, Seeberger PH. *Angew Chem Int Ed.* 2005; 44:6315–6318. (b) Adamo R, Saksena R, Kovac P. *Carbohydr Res.* 2005; 340:2579–2582. [PubMed: 16216230] (c) Saksena R, Adamo R, Kovac P. *Bioorg Med Chem Lett.* 2006; 16:615–617. [PubMed: 16275067] (d) Adamo R, Saksena R, Kovac P. *Helv Chim Acta.* 2006; 89:1075–1089. (e) Mehta AS, Saile E, Zhong W, Buskas T, Carlson R, Kannenberg E, Reed Y, Quinn CP, Boons GJ. *Chem Eur J.* 2006; 12:9136–9149. [PubMed: 17133642] (f) Crich D, Vinogradova O. *J Org Chem.* 2007; 72:6513–6520. [PubMed: 17661521] (g) Werz DB, Adibekian A, Seeberger PH. *Eur J Org Chem.* 2007:1976–1982.
10. (a) Guo H, O'Doherty GA. *Angew Chem Int Ed.* 2007; 46:5206–5208. (b) Guo H, O'Doherty GA. *J Org Chem.* 2008; 73:5211–5220. [PubMed: 18563936]
11. Several approaches to related tri- and pentasaccharides, see: Saksena R, Adamo R, Kovac P. *Bioorg Med Chem.* 2007; 15:4283–4310. [PubMed: 17412599] Wang Y, Liang X, Wang P. *Tetrahedron Lett.* 2011; 52:3912–3915. Tamborrini M, Bauer M, Bolz M, Maho A, Oberli MA, Werz DB, Schelling E, Jakob Z, Seeberger PH, Frey J, Pluschke G. *J Bacteriol.* 2011; 193:3506–3511. [PubMed: 21571994] Milhomme O, Dhenin SGY, Djedaini-Pilard F, Moreau V, Grandjean C. *Carbohydr Res.* 2012; 356:115–131. [PubMed: 22356927]
12. (a) Babu RS, O'Doherty GA. *J Carb Chem.* 2005; 24:169–177. (b) Zhou M, O'Doherty GA. *Org Lett.* 2006; 8:4339–4342. [PubMed: 16956221] (c) Zhou M, O'Doherty GA. *Org Lett.* 2008; 10:2283–2286. [PubMed: 18461951] (d) Yu X, O'Doherty GA. *Org Lett.* 2008; 10:4529–4532. [PubMed: 18785743] (e) Yu X, Li M, O'Doherty GA. *Heterocycles.* 2011; 82:1577–1584. (f) Wang HYL, Rojanasakul Y, O'Doherty GA. *ACS Med Chem Lett.* 2011; 2:264–269. [PubMed: 21660118] (g) Babu RS, Chen Q, Kang SW, Zhou M, O'Doherty GA. *J Am Chem Soc.* 2012; 134:11952–11955. [PubMed: 22780712]
13. (a) Fujii A, Hashiguchi S, Uematsu N, Ikariya T, Noyori R. *J Am Chem Soc.* 1996; 118:2521–2522. (b) Noyori R, Ohkuma T. *Angew Chem, Int Ed.* 2001; 40:40–73.
14. The absolute stereochemistry and the level of enantioexcess of 10 and its enantiomer (ent)-10 were determined by the method of Mosher, see: Sullivan GR, Dale JA, Mosher HS. *J Org Chem.* 1973; 38:2143–2147. Yamaguchi S, Yasuhara F, Kabuto KT. *Tetrahedron.* 1976; 32:1363–1367. Bushey ML, Haukaas MH, O'Doherty GA. *J Org Chem.* 1999; 64:2984–2985. [PubMed: 11674385] For literature report of optical activity, see: Drueckhammer DG, Barbas CF III, Nozoki K, Wong C-H, Wood CY, Ciufolini MA. *J Org Chem.* 1993; 58:4511–4512.
15. The relative stereochemistry and the level of diastereoselective induction were determined by analysis of the allylic coupling constants in the ^1H NMR spectra, see: Babu RS, Guppi SR, O'Doherty GA. *Org Lett.* 2006; 8:1605–1608. [PubMed: 16597121]
16. (a) Luche JL. *J Am Chem Soc.* 1978; 100:2226–2227. (b) Abrams JN, Babu RS, Guo H, Le D, Le J, Osbourn JM, O'Doherty GA. *J Org Chem.* 2008; 73:1935–1940. [PubMed: 18237188]

Figure 1.
Anthrax tetrasaccharide and its tri- and di-saccharide fragments

Scheme 1.
Retrosynthetic approach to anthrax saccharides 1-4

Scheme 2.
Preparation of *tert*-butyl-Boc pyranone sugar donors

Scheme 3.

Previously reported approach to oligosaccharides 5, 6, and 7

Scheme 4.
Synthesis of L-D-disaccharide anthrax precursor

Scheme 5.
Preparation of anthrax saccharides **1**–**4**