Published in final edited form as:

Angew Chem Int Ed Engl. 2013 March 11; 52(11): 3237–3240. doi:10.1002/anie.201209266.

Rhodium- and Platinum-Catalyzed [4+3] Cycloaddition with Concomitant Indole Annulation: Synthesis of Cyclohepta[*b*]indoles**

Dongxu Shu

The School of Pharmacy, University of Wisconsin Madison, WI 53705-2222 (USA)

Department of Chemistry, University of Wisconsin

Wangze Song, Xiaoxun Li, and Prof. Dr. Weiping Tang*

* The School of Pharmacy, University of Wisconsin Madison, WI 53705-2222 (USA)
wtang@pharmacy.wisc.edu

Keywords

cycloaddition; heterocycles; homogeneous catalysis; transition metals

Seven-membered rings fused with an indole, cyclohepta[*b*]indoles, are present in many bioactive natural products such as silicine,^[1] ervitsine,^[2] and actinophyllic acid (Figure 1).^[3] They are also important structural motifs in numerous pharmaceuticals with various pharmacological properties such as inhibition of deacetylase SIRT1,^[4] inhibition of adipocyte fatty-acid-binding protein (A-FABP),^[5] and anti-tubercular activity.^[6] Most previous efforts have focused on building the seven-membered ring and the indole separately by cyclization reactions.^[4,5,7] Recently, an elegant three-component [4+3] cycloaddition was reported by Wu and co-workers for the synthesis of cyclohepta[*b*]indoles from indoles, aldehydes, and dienes.^[8] It represents the first example of a [4+3] cycloaddition involving an indole as the 2 π component. We herein report an efficient and versatile process that allows the simultaneous construction of both the indole and seven-membered ring through a [4+3] cycloaddition with concomitant indole annulation.^[9]

Vinyl metal carbenes derived from vinyl diazo compounds can undergo formal [4+3] cycloadditions with dienes through a cyclopropanation/Cope rearrangement sequence (Scheme 1).^[10,11] Similarly, vinyl Fisher carbenes^[12] or vinyl gold carbenes derived from propargylic esters^[13] can also react with dienes to form various seven-membered rings. It was recently reported that vinyl metal carbenes could be conveniently generated from propargylic ethers tethered with a nucleophile for a [3+2] cycloaddition^[14] and a synthesis of furans.^[15] We envisioned that the vinyl metal carbene **2**, derived from **1**, would undergo a formal [4+3] cycloaddition^[16] with diene **3** to afford the cyclohepta[*b*]indole **4** by either a cyclopropanation/Cope rearrangement sequence involving the divinyl-cyclopropane **5**, or an unusual [4+4] cycloaddition to form the eight-membered metallacycle **6** with subsequent reductive elimination. Both indole and seven-membered rings may be constructed very efficiently in this tandem process from simple building blocks.

** We thank the NIH (R01 GM088285) for funding. W.T. is grateful for a Young Investigator Award from Amgen.

© 2013 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Correspondence to: Weiping Tang.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201209266>.

The transformation from the propargylic ether **1** to the product **4** requires a metal catalyst which has enough π acidity^[17] to induce cyclization of **1**, thus forming a carbene intermediate, and the ability to promote cycloadditions. The catalyst $[\{\text{Rh}(\text{CO})_2\text{Cl}\}_2]$ can facilitate 1,3-acyloxy migration of propargylic esters, a process that is typically catalyzed by π -acidic metals,^[17] and effects cycloadditions as well.^[18,19] When a mixture of the propargylic ether **1a** and diene **3a** was treated with this catalyst at 80°C, no reaction occurred (Table 1, entry 1). We have previously found that electron-deficient phosphine or phosphite ligands often increase the acidity of rhodium catalysts and promote 1,2-acyloxy^[20] or 1,3-acyloxy^[19] migration of propargylic esters. Indeed, a mixture of the [4+3] cycloaddition product **4a** and simple indole **7** was observed when **1a** was treated with $[\{\text{Rh}(\text{CO})_2\text{Cl}\}_2]$ in the presence of such ligands (entries 2–4). The amount of **7** could be minimized by employing a greater excess of **3a** (entry 4). A 67% yield of the isolated tricyclic product **4a** could be obtained in the presence of a rhodium(I) metal complex and an electron-deficient phosphite ligand.

We also examined PtCl_2 , $\text{PtCl}_2/\text{alkene}$, and $\text{PtCl}_2/\text{PPh}_3$, all of which have been used in the generation of vinyl platinum carbenes from propargylic ethers.^[14,15] A low yield of **4a** or no product, however, was observed using these catalysts (Table 1, entries 5–8). We suspect that the coordination of the bidentate diene **3a** to PtCl_2 may reduce the acidity of the metal. Electron-deficient phosphite or phosphine ligands were then added to further enhance the reactivity of the PtCl_2 catalyst (entries 9–11). Indeed, the yield of **4a** was increased significantly. The tris(pentafluorophenyl)phosphine ligand provided the highest yield of the isolated product **4a** (entry 11). A lower catalyst loading led to lower conversion, and other metal catalysts did not afford the desired product (entries 12–14).

With the two catalysts in hand (Table 1, entries 4 and 11), we studied the scope of this tandem indole annulation/[4+3] cycloaddition with different propargylic ethers (Table 2, entries 1–7). The ketone **8a** was isolated in 82% yield after in situ hydrolysis of the silyl enol ether **4a** (entry 1). A benzyl ether could also be tolerated (entry 2), and other leaving groups (e.g. $\text{X}=\text{OH}$, OPiv , or Cl) led to a complex mixture. Electron-withdrawing or electron-donating groups on the benzene ring change the nucleophilicity of the aniline nitrogen atom, however the efficiency of the indole annulation/[4+3] cycloaddition did not change with either type of substituent (entries 3 and 4). A lower yield was observed for the substrate **1e** having a free alcohol (entry 5), and a formyl group did not interfere with the tandem reaction (entry 6). The secondary propargylic ether **1g** also participated in the tandem reaction and yielded **4g** (entry 7).

We next investigated the scope of acyclic dienes that could be used in this process (Table 2, entries 8–10). The more functionalized 2,3-disubstituted diene **3b** afforded **4ab** in high yield (entry 8, Table 2). The monosubstituted diene **3c** produced **8ac** in 59% yield in the presence of a rhodium catalyst (entry 9), and lower yields were obtained when various platinum catalysts were employed in this case. The same trend was also observed for **3d** (entry 10). When platinum catalysts were employed, the yield of **4ad** was 20–30% lower than that obtained from using the rhodium catalyst, and a diastereomeric mixture of **4gd** was isolated when **3d** was reacted with the propargylic ether **1g** (entry 11). A complex mixture was observed when substrate **1a** was treated with 2-methyl-1,3-butadiene in the presence of either the platinum or rhodium catalyst, thus suggesting that the silyloxy substituent is critical for the reactivity of acyclic dienes.

We were pleased to find that the furan **9a** participated in the tandem reaction and afforded the tetracyclic product **10a** in 71% yield (Table 3, entry 1). The arylation product **11a** was isolated in 14% yield. For the 3,4-disubstituted furan **9b**, a single product, **10b**, was observed (entry 2). The yields for the ester-substituted furans **9c** and **9d** were slightly lower

References

- [1]. Shafiee A, Ahond A, Bui AM, Langlois Y, Riche C, Potier P. *Tetrahedron Lett.* 1976; 17:921.
- [2]. Andriantsiferana M, Besselievre R, Riche C, Husson HP. *Tetrahedron Lett.* 1977; 18:2587.
- [3]. a) Carroll AR, Hyde E, Smith J, Quinn RJ, Guymer G, Forster PI. *J. Org. Chem.* 2005; 70:1096. [PubMed: 15675882] b) Taniguchi T, Martin CL, Monde K, Nakanishi K, Berova N, Overman LE. *J. Nat. Prod.* 2009; 72:430. [PubMed: 19220031]
- [4]. Napper AD, Hixon J, McDonagh T, Keavey K, Pons JF, Barker J, Yau WT, Amouzegh P, Flegg A, Hamelin E, Thomas RJ, Kates M, Jones S, Navia MA, Saunders J, DiStefano PS, Curtis R. *J. Med. Chem.* 2005; 48:8045. [PubMed: 16335928]
- [5]. Barf T, Lehmann F, Hammer K, Haile S, Axen E, Medina C, Uppenberg J, Svensson S, Rondahl L, Lundbaeck T. *Bioorg. Med. Chem. Lett.* 2009; 19:1745. [PubMed: 19217286]
- [6]. Yamuna E, Kumar RA, Zeller M, Prasad KJR. *Eur. J. Med. Chem.* 2012; 47:228. [PubMed: 22119150]
- [7]. a) Bennasar ML, Vidal B, Bosch J. *J. Am. Chem. Soc.* 1993; 115:5340. b) Bennasar ML, Vidal B, Bosch J. *J. Org. Chem.* 1997; 62:3597. c) Martin CL, Overman LE, Rohde JM. *J. Am. Chem. Soc.* 2008; 130:7568. [PubMed: 18491907] d) Vaswani RG, Day JJ, Wood JL. *Org. Lett.* 2009; 11:4532. [PubMed: 19810762] e) Amat M, Llor N, Checa B, Molins E, Bosch J. *J. Org. Chem.* 2010; 75:178. [PubMed: 19968261] f) Martin CL, Overman LE, Rohde JM. *J. Am. Chem. Soc.* 2010; 132:4894. [PubMed: 20218696] g) Zaimoku H, Taniguchi T, Ishibashi H. *Org. Lett.* 2012; 14:1656. [PubMed: 22369137]
- [8]. Han X, Li H, Hughes RP, Wu J. *Angew. Chem.* 2012; 124:10536. *Angew. Chem. Int. Ed.* 2012; 51:10390.
- [9]. For recent reviews on indole synthesis, see: Barluenga J, Rodriguez F, Fananas FJ. *Chem. Asian J.* 2009; 4:1036. [PubMed: 19360759] ; Taber DF, Tirunahari PK. *Tetrahedron.* 2011; 67:7195.; Vicente R. *Org. Biomol. Chem.* 2011; 9:6469. [PubMed: 21779596]
- [10]. For reviews on sequential cyclopropanation and Cope rearrangement, see: Davies HML. *Tetrahedron.* 1993; 49:5203.; Davies HML, Denton JR. *Chem. Soc. Rev.* 2009; 38:3061. [PubMed: 19847341]
- [11]. For selected examples of sequential cyclopropanation and Cope rearrangement, see: Davies HML, Smith HD, Korkor O. *Tetrahedron Lett.* 1987; 28:1853.; Davies HML, McAfee MJ, Oldenburg CEM. *J. Org. Chem.* 1989; 54:930.; Reddy RP, Davies HML. *J. Am. Chem. Soc.* 2007; 129:10312. [PubMed: 17685525] ; Olson JP, Davies HML. *Org. Lett.* 2008; 10:573. [PubMed: 18215048]
- [12]. a) Wulff WD, Yang DC, Murray CK. *J. Am. Chem. Soc.* 1988; 110:2653. b) Barluenga J, Aznar F, Martin A, Garciagrande S, Salvado MA, Pertierra P. *J. Chem. Soc. Chem. Commun.* 1993:319. c) Barluenga J, Aznar F, Martin A, Vazquez JT. *J. Am. Chem. Soc.* 1995; 117:9419. d) Barluenga J, Aznar F, Fernandez M. *Chem. Eur. J.* 1997; 3:1629. e) Hoffmann M, Buchert M, Reissig HU. *Chem. Eur. J.* 1999; 5:876. f) Barluenga J, Alonso J, Rodriguez F, Fananas FJ. *Angew. Chem.* 2000; 112:2555. *Angew. Chem. Int. Ed.* 2000; 39:2459. g) Barluenga J, Alonso J, Fananas FJ. *hem. Eur. J.* 2005; 11:4995. h) Aznar F, Fananas-Mastral M, Alonso J, Fananas FJ. *Chem. Eur. J.* 2008; 14:325. [PubMed: 17886851]
- [13]. a) Miki K, Ohe K, Uemura S. *J. Org. Chem.* 2003; 68:8505. [PubMed: 14575478] b) Gung BW, Bailey LN, Wonsler J. *Tetrahedron Lett.* 2010; 51:2251. [PubMed: 20352055] c) Garayalde D, Kruger K, Nevado C. *Angew. Chem.* 2011; 123:941. *Angew. Chem. Int. Ed.* 2011; 50:911.
- [14]. Saito K, Sogou H, Suga T, Kusama H, Iwasawa N. *J. Am. Chem. Soc.* 2011; 133:689. [PubMed: 21171651]
- [15]. Allegretti PA, Ferreira EM. *Org. Lett.* 2011; 13:5924. [PubMed: 21995781]
- [16]. For recent reviews on [4+3] cycloadditions, see: Harmata M. *Adv. Synth. Catal.* 2006; 348:2297.; Harmata M. *Chem. Commun.* 2010; 46:8886.; Harmata M. *Chem. Commun.* 2010; 46:8904.; Lohse AG, Hsung RP. *Chem. Eur. J.* 2011; 17:3812. [PubMed: 21384451]
- [17]. For selected reviews on π -acidic metal catalysis, see: Marion N, Nolan SP. *Angew. Chem.* 2007; 119:2806.; *Angew. Chem. Int. Ed.* 2007; 46:2750.; Fürstner A, Davies PW. *Angew. Chem.* 2007; 119:3478.; *Angew. Chem. Int. Ed.* 2007; 46:3410.; Hashmi ASK. *Chem. Rev.* 2007; 107:3180.

[PubMed: 17580975] ; Gorin DJ, Toste FD. Nature. 2007; 446:395. [PubMed: 17377576] ; Wang S, Zhang G, Zhang L. Synlett. 2010:692.; Hashmi ASK. Angew. Chem. 2010; 122:5360.; Angew. Chem. Int. Ed. 2010; 49:5232.; Shu X-Z, Shu D, Schienebeck CM, Tang W. Chem. Soc. Rev. 2012; 41:7698. [PubMed: 22895533]

- [18]. a) Shu D, Li X, Zhang M, Robichaux PJ, Tang W. Angew. Chem. 2011; 123:1382. Angew. Chem. Int. Ed. 2011; 50:1346. b) Li X, Huang S, Schienebeck CM, Shu D, Tang W. Org. Lett. 2012; 14:1584. [PubMed: 22381143]
- [19]. Huang S, Li X, Lin CL, Guzei IA, Tang W. Chem. Commun. 2012; 48:2204.
- [20]. a) Shu, X.-z.; Huang, S.; Shu, D.; Guzei, IA.; Tang, W. Angew. Chem. 2011; 123:8303. Angew. Chem. Int. Ed. 2011; 50:8153. b) Shu, X.-z.; Li, X.; Shu, D.; Huang, S.; Schienebeck, CM.; Zhou, X.; Robichaux, PJ.; Tang, W. J. Am. Chem. Soc. 2012; 134:5211. [PubMed: 22364320]
- [21]. Cramer CJ, Barrows SE. J. Org. Chem. 1998; 63:5523.
- [22]. Ito Y, Hirao T, Saegusa T. J. Org. Chem. 1978; 43:1011.

Figure 1.
Representative cyclohepta[*b*]indoles.

Previous work:

This work:

Scheme 1.

[4+3] Cycloaddition of vinyl carbenes and dienes.

Scheme 2.

Proposed mechanisms for the [4+3] cycloaddition accompanied by an indole annulation.

Table 1

Optimization for the reaction between **1a** and **3a**.^[a]

Entry	Conditions	4a Yield [%] ^[b]	4a/7
1 ^[c]	[{Rh(CO) ₂ Cl} ₂]	no reaction	–
2 ^[c]	[{Rh(CO) ₂ Cl} ₂], P[(CF ₃) ₂ C ₆ H ₃] ₃	35	1:1
3 ^[c]	[{Rh(CO) ₂ Cl} ₂], P(C ₆ F ₅) ₃	53	2:1
4	[{Rh(CO) ₂ Cl} ₂], P[OCH(CF ₃) ₂] ₃	70 (67) ^[d]	4:1
5	PtCl ₂	28	1.6:1
6	PtCl ₂ , octene	21	2:1
7	[Pt(PPh ₃) ₂ Cl ₂]	no reaction	–
8	PtCl ₂ , CO (1 atm)	18	4:1
9	PtCl ₂ , P[OCH(CF ₃) ₂] ₃	66	8:1
10	PtCl ₂ , P[(CF ₃) ₂ C ₆ H ₃] ₃	31	10:1
11	PtCl ₂ , P(C ₆ F ₅) ₃	85 ^[d]	>20:1
12	[IrCl(CO)(PPh ₃) ₂]	no reaction	–
13	PdCl ₂	no reaction	–
14	[AuClP(<i>p</i> -CF ₃ C ₆ H ₄) ₃]	no reaction	–

^[a] Reaction conditions: 10 mol% metal catalyst, 20 mol% ligand, diene **3a** (5 equiv), 80°C, 1,4-dioxane, Na₂CO₃, 12 h, unless noted otherwise.

^[b] Yields based on ¹H NMR spectroscopy.

^[c] Used 2 equiv of **3a**.

^[d] Yield of isolated product. Boc = *tert*-butoxycarbonyl, TIPS=triisopropylsilyl.

Table 2Scope of propargylic ethers and acyclic dienes.^[a]

Entry	1	3	Product	Yield [%]
1 ^[b]	1a (X=OMe)	3a	8a	82
2 ^[b]	1b (X=OBn)	3a	8a	80
3	1c (R=CO ₂ Et)	3a	4c	83
4	1d (R=OMe)	3a	4d	86
5	1e (R=CH ₂ OH)	3a	4e	52
6	1f (R=CHO)	3a	4f	83
7 ^[c]	1g	3a	4g	56
8	1a	3b	4ab	90
9 ^{[b],[d]}	1a	3c	8ac	59

Entry	1	3	Product	Yield [%]

10 ^[d]	1a	3d	4ad	66

11	1g	3d	4gd	50 (d.r.=3:1)

^[a] Reaction conditions: PtCl₂ (10 mol%), P(C₆F₅)₃ (20 mol%), diene **3** (5.0 equiv), 80°C, 1,4-dioxane, Na₂CO₃, 12 h. Yields are those of the isolated products.

^[b] The resulting product was treated with aqueous HCl (4M).

^[c] Used 1.5 equiv of **3a**.

^[d] [Rh(CO)₂Cl]₂ (5 mol%) and P[OCH-(CF₃)₂]₃ (20 mol%) were employed. TBS = *tert*-butyldimethylsilyl.

Table 3

Scope of cyclic dienes for the tandem reaction with the propargylic ether **1a**.^[a]

Entry	9	Products ^[b]

1 ^[c]	9a	10a (71%)	11a (14%)

2	9b	10b (79%)

3	9c	10c (52%)

4	9d	10d (44%)	10d' (23%)

5	9e	10e (83%) ^[d]	10e' (15%)

Entry	9	Products ^[b]

6	9f	10f (58%) 11f (30%)

7	9g	10g (63%)

8	9h	10h (61%) ^[e]

^[a] Reaction conditions: PtCl₂ (10 mol%), P(C₆F₅)₃ (20 mol%), diene **5** (1.5–5.0 equiv), 80°C, 1,4-dioxane, Na₂CO₃, 12 h.

^[b] Yield for isolated product given within parentheses.

^[c] PtCl₂ (5 mol%), P(C₆F₅)₃ (10 mol%)

^[d] Yield for **10e** and **10e'** combined.

^[e] The Boc protecting group was removed by the treatment with trifluoroacetic acid. The yield is the overall yield for two steps.