Published in final edited form as:

Pediatr Pulmonol. 2010 September ; 45(9): 919–926. doi:10.1002/ppul.21273.

EARLY WHEEZE AS REPORTED BY MOTHERS AND LUNG FUNCTION IN FOUR-YEAR-OLDS. PROSPECTIVE COHORT STUDY IN KRAKOW

Wieslaw Jedrychowski*, Umberto Maugeri**, Frederica P. Perera***, Elzbieta Mroz*, Maria Klimaszewska-Rembiasz****, Elzbieta Flak*, Susan Edwards***, Renata Majewska*, and Agata Sowa*

*Chair of Epidemiology and Preventive Medicine, Coll. Med. Jagiellonian University in Krakow. Poland **Institute for Clinical Medicine and Rehabilitation, Salvatore Maugeri Foundation, Pavia, Italy ***Columbia Center for Children's Environmental Health, Mailman School Public Health, Columbia University, NY, NY, US ****St.Louis Children's Hospital in Krakow, Poland

Abstract

The purpose of the study was to check the hypothesis that early wheezing as reported by mothers would be associated with reduced lung function in 4-year-olds. Study participants were recruited prenatally, as part of a prospective cohort study on the respiratory health of young children exposed to various ambient air pollutants. After delivery, infants were followed over four years and the interviewers visited participants at their home to record respiratory symptoms every three months in the child's first two years of life and every 6 months in the third and fourth years. In the fourth year of follow-up, children were invited for standard lung function testing by spirometry quantified by FVC, FEV₁ and FEV₀₅ levels. Out of 258 children attending spirometry testing 139 performed at least two acceptable exhalation efforts. Cohort children with acceptable spirometric measurements did not differ with respect to wheezing experience and exposure characteristics from those without. The study shows that episodic wheeze was reported in 28.1% of 4-year-olds, 6.5% had transient wheeze and 4.3% had recurrent wheeze. There was an increased frequency of wheezing symptoms and their duration in transient and recurrent wheezers. Adjusted multivariable regression models for gender and height showed that children who reported more than 2 episodes of wheezing at any point over the follow-up had FVC values lower by 120.5 mL ($p = 0.016$) and FEV₁ values lower by 98.3 mL ($p = 0.034$) compared to those who did not report any wheezing; children experiencing more than 10 wheezing days by age 4 showed FVC deficit of 87.4 mL ($p = 0.034$) and FEV₁ values of 65.7 mL ($p = 0.066$) The ratios of FEV₁/FVC% and FEV₀₅/FVC% were neither associated with wheezing episodes nor wheezing days. In recurrent wheezers lung function decrement amounted to 207 mL of FVC, 175 mL of FEV₁ and 104 mL of FEV₀₅. In conclusion, our findings show that wheezing experience during early postnatal life may be associated with lung function deficit of restrictive character in preschool children and detailed history of wheeze in early postnatal life, even though not physician-confirmed, may help define the high risk group of children for poor lung function testing.

Introduction

Wheeze is prevalent in early life, and recurrent wheezing is one of the main pathophysiological manifestations of asthma, along with increased airway responsiveness and persistent airway inflammation. Most cases of persistent wheeze and asthma begin in early childhood (1 – 4) which is often associated with increased airway responsiveness and reduced lung function (5 – 11). The clinical and epidemiologic importance of early wheezing for respiratory health in later childhood and in the course of adult life is poorly

understood (12). Wheeze originates in the narrowing of airways, either by compression or by intrabronchial or intraluminal obstruction by inflammatory mucosal edema, secretions or spasm, which causes an increase in the velocity of gas in the lungs with resultant oscillation. It is suggested that wheezing lower respiratory illness (LRI) in infants may be a consequence of anatomically small airways and is unrelated to the later development of atopic asthma (13). Assessing the association between early wheeze in early childhood and lung function is complicated by the fact that in most epidemiological studies information about wheeze was collected through parental reporting of current wheeze. There is some doubt that parents may confuse wheeze with other respiratory symptoms, and this could lead to incorrect effect estimates and overestimations of the true prevalence of wheeze.

Many studies on respiratory standard function tests and its utility in school children and adolescents have already been published (14 – 20), however, more attention has been focused recently on preschool children, who may have difficulty to fulfill precisely reproducibility criteria in performing standard spirometric tests (21, 22). Up to now, only a few studies have been concerned with measurements of lung function in preschool children and its relation with respiratory symptoms.

The main objective of this study was to find out whether the wheeze frequency and lung function of preschool children differs by wheezing phenotype reported by mothers over the first 4 years of life. We assessed the frequency of wheezing by the number of wheezing episodes (frequency) and wheezing days through regular standardized face-to-face interviews with mothers over the first four years of life. Lung function was assessed by standard spirometry and quantified by FVC, FEV₁, FEV₀₅ levels and ratios of FEV₀₅/FVC and FEV₁/FVC%.

Materials and Methods

This study uses data from an earlier established birth cohort of children in Krakow, the product of a longstanding collaboration between Jagiellonian University in Krakow and Columbia University in New York. The design of the study and the detailed selection of the population have been described previously (23). Pregnant women were recruited from ambulatory prenatal clinics in their first or second trimesters of pregnancy. Only women 18–35 years of age, who claimed to be non-smokers, with singleton pregnancies, with no history of illicit drug use and HIV infection, free from chronic diseases such as diabetes or hypertension, and who had resided in Krakow for at least one year prior to pregnancy were eligible for the study. Prior to participation, women read and signed an informed consent. The Ethical Committee of the Jagiellonian University approved the research.

Upon enrollment, a detailed questionnaire was administered to each woman to solicit information on demographic data, house characteristics, medical and reproductive history, occupational hazards, and smoking practices of others present at home. A total of 505 enrolled pregnant women gave birth between January 2001 and February 2004. After delivery, every three months in the first two years of the newborn's life, and every 6 months in their third and fourth years, a detailed standardized face-to-face interview on the infant's health and respiratory symptoms was administered to each mother by a trained interviewer. In the 4th year of follow-up, 382 mothers (76%) participated in the study. During the interview at each of the study periods at 3, 6, 9, 12, 15, 18, 21, 24, 30, 36, and 42 months after birth, a history of wheezing in children was ascertained by the following question to mothers: "Has your child had wheezing or whistling in the chest since the last interview?" Answers were to include wheeze with and without colds and was not doctor confirmed. If an affirmative response to the question on the occurrence of wheezing was recorded, the

interviewer asked about the number of wheezing episodes and its duration (in days). Medical chest auscultation was not included in the assessment.

The data collected over the course of the 12 follow-up time points was used to identify four mutually exclusive patterns of wheeze between birth and 4 years of age: 0. never wheezers (no wheezing at any of the 12 time points); 1. episodic wheeze (wheeze only in any one year of the follow-up period, 2. transient wheeze (wheeze reported in two years of the 4 year follow-up period but at non-consecutive intervals, 3. recurrent wheezing (wheeze developed in the first two years of life and continued to be present in the third or fourth year of the follow-up). The severity of respiratory symptoms was expressed by the total number of wheezing episodes and wheezing days experienced by the child recorded in the follow-up. In separate analyses, we used information on maternal report of “current wheeze”, i.e. wheeze reported over the course of the 12 month period preceding lung function testing.

Spirometry testing

Children were free of respiratory symptoms on the day of testing and none of the children had any previous experience performing spirometry. Prior to spirometric testing, standing height and weight without shoes of each child was measured using the same equipment and standardized procedures and the children were coached to engage in maximal forced expiratory efforts in a standing position without nose clip. All spirometric measurements were carried out with a PC QRS Card Spirometer with an interactive computerized incentive display software. Pulmonary function testing was performed in accordance with the ATS/ERS guidelines on lung function testing in preschool children (21) by highly experienced staff member (E.Mroz). Each day, prior to the lung function examination, the spirometer was calibrated with a one-liter syringe. Each child made at least two good forced exhalation efforts. The primary indicators of lung function are FVC (Forced Vital Capacity), FEV₁ (Forced Expiratory Volume in 1 second), and FEV_{0.5} (Forced Expiratory Volume in 0.5 second). The research worker observed the child closely in the course of the expiratory effort to ensure that the child's lips were sealed around mouthpiece, and tracked volume-time curves on the computer screen visually. The exhalation was excluded if the flow-volume curve did not demonstrate a rapid rise to peak flow, and smooth descending limb. Expiratory maneuvers were also excluded if cough or glottic closure appeared in the course of testing. Spirograms with a back-extrapolated expiratory volume greater than 100 ml or an expiration time less than 1 second were not accepted either. Expiratory volumes were calculated based on each subject's best expiratory attempt (the greatest sum of FEV₁ and FVC) and corrected to BTPS (body temperature, pressure saturated). Out of 382 children participating in the follow-up, 258 children (68%) reported for the lung function testing. The present analysis was based on 139 subjects who performed at least two acceptable expiratory lung function tests, where the second highest FVC and FEV₁ were within 100 ml. All FVC curves started from time zero, except two curves which showed a backward extrapolated volume lower than 50 ml of FVC. Of the group of 139 children with acceptable lung function tests, 114 had three exhalation efforts and 25 had two.

Statistical Analysis

The main purpose of the statistical analysis was to assess the relationship between lung function indices (outcome variables), the number of wheezing episodes, wheezing days, and wheezing phenotypes (independent variables). Lung function indices were treated as continuous variables, the severity of wheeze defined by the number wheezing episodes was transformed in a dummy variable coded as: 0 – never wheezed, 1.- one or two episodes reported over the follow-up (below median in wheezers), 2: three or more episodes of wheezing (above median in wheezers); the corresponding dummy variable for wheezing

days was coded as: 0. – never wheezed, 1. up to ten wheezing days (below median in wheezers), 2. - more than 10 wheezing days (above median in wheezers). The adjusted effect of wheeze severity was assessed in the multivariable stratified linear regression analysis where regression coefficients for wheezing strata were compared with the reference level (no wheezing). Preliminary analysis to identify potential confounders assessed associations between population characteristics and outcome variables in univariate statistical models, where Chi-square statistics (nominal variables), one way of analysis of variance (numerical variables), and nonparametric test for trend tested differences between subgroups with various wheezing experience were used. In the multivariable analyses, associations between lung function and wheezing were initially adjusted for a set of covariates (gender of child, children's anthropometry, exact age of child in months) but in the final analysis only significant confounders were taken into consideration. All statistical analyses were carried out with STATA 10 version software for Windows (24, 25).

Results

Basic characteristics of the study groups with acceptable and excluded exhalation efforts did not differ significantly from each other (Table 1). On average those with unacceptable exhalation efforts were taller (106.6 cm vs. 105.3 cm), heavier (17.76 kg vs. 17.14 kg) however, these differences were not statistically significant at the $p = 0.05$ level. Children whose exhalation efforts were not acceptable produced higher spirometric volumes (Table 2).

Wheezers reported on average 2.1 wheezing episodes (95%CI: 1.5 – 2.6) and 15.5 wheezing days (95%CI: 10.4 – 20.6) in the follow-up. There was a very strong correlation between number of wheezing episodes and wheezing days (Spearman correlation coefficient $r_s = 0.979$; 95%CI: 0.970 – 0.985). Episodic wheeze was reported in 28.1% of children (95% CI: 21.4 – 37.1%), 6.5% had transient wheeze (95% CI: 3.0 – 11.9) and 4.3% had recurrent wheeze (95% CI: 1.6 – 9.2%). There were 20 children (14.4%) where wheezing symptoms were reported for the last 12 months (current wheezers). Table 3 displays the frequency of wheezing episodes and wheezing days grouped by wheezing phenotype. It shows that the number of wheezing episodes and wheezing days was particularly high in recurrent wheezers.

Boys had a higher FVC values than girls but the difference was not statistically significant (887.6 vs. 848.4, $t = 1.375$, $df. = 137$, $p = 0.172$); similar differences were observed for FEV_1 and FEV_{05} across gender groups. Lung function indices (FVC, FEV_1 and FEV_{05}) correlated in univariate analyses with children's height ($r = 0.42 - 0.44$, $p < 0.001$) and weight ($r = 0.25 - 0.32$, $p < 0.005$) as recorded at the time of lung function testing. In the forward stepwise regression analysis of lung function indices, where gender was specified as a lockterm, only the child's height was a significant predictor of lung function in the study sample. Therefore, in the subsequent multivariable regression analyses, only children's height was considered in the regression models in addition to gender.

Controlling for gender and height, multivariable linear regression models showed an inverse but not significant association between current wheeze (the symptoms present in the last 12 months versus no wheezing over the last year) and FVC (regression coefficient = -46.93 ($p = 0.218$); current wheeze and FEV_1 -38.48 ($p = 0.216$); current wheeze and FEV_{05} -25.18 ($p = 0.433$).

We were able to show an inverse and significant association between the higher number of wheezing episodes recorded in the follow-up period and both FVC and FEV_1 indices, but not FEV_{05} (Table 4). FVC values in children who reported more than 2 episodes of

wheezing had FVC values lower by 120.5 mL ($p = 0.016$) and FEV₁ by 98.3 mL ($p = 0.034$) compared to children who did not report any wheezing; children experiencing more than 10 wheezing days showed FVC deficit of 87.4 mL ($p = 0.034$) and FEV₁ of 65.7 mL ($p = 0.066$) compared to non-wheezers. The ratios of FEV₁/FVC% and FEV_{0.5}/FVC% were associated neither with the number of wheezing episodes nor wheezing days.

There was a significant inverse trend of FVC values across wheezing phenotypes (nonparametric trend $z = -2.03$, $p = 0.042$), but lung function was noticeably decreased in recurrent wheezers among whom lung function decrement amounted to 207 mL of FVC ($p = 0.002$), 175 mL of FEV₁ ($p = 0.004$) and 105 mL of FEV_{0.5} ($p = 0.061$). Figure 1 presents the box-and-whisker plot of FVC crude values by the wheezing phenotypes. As expected, children with recurrent wheeze experienced much higher number of wheezing episodes (mean number of episodes = 6.3; 95%CI: 2.8 – 9.9) and wheezing days (mean number of days = 39.2; 95%CI: 34.3 – 44.5) than children who reported episodic or transient wheezing (mean number of episodes = 1.5; 95%CI: 1.2 – 1.91; mean number of days = 12.5; 95%CI: 11.5 – 13.6).

Discussion

Adjusting multivariable regression models for child gender and height at age of 4 years, we found that children who reported more episodes of wheezing or more wheezing days at any time point over the course of the four year follow-up, had significant lower values of FVC and FEV₁ compared to those children who did not report any wheezing. The ratios of FEV₁/FVC% and FEV_{0.5}/FVC% were neither associated with wheezing episodes nor wheezing days. The phenotype of recurrent wheezing was associated with a marked FVC decrement of restrictive pattern.

Infancy is characterized by pulmonary alveolar multiplication and extensive remodeling of the airways to accommodate growth. The recurrent wheeze may be a surrogate marker of pulmonary immune immaturity, which is manifested by an enhanced susceptibility to the effects of respiratory viruses. It has already been shown previously that frequent recurrent insults to the airway epithelium in the course of respiratory infections in wheezers may result in various inflammatory processes that induce structural airway changes leading to airways remodeling (26 – 30). In fact, recurrent wheezers in our study experienced much higher number of wheezing episodes over the follow-up than the group with episodic or transient wheezing. Moreover, in recurrent wheezers the onset of symptoms was noted in the first two years of life and continued through later age.

Our study results showing a restrictive pattern of lung function deficit in wheezers support the hypothesis that recurrent wheeze may result from early exposure to ambient insults and/or lower respiratory illness. This may prevent lungs from reaching their optimal development potential. Moreover, restricted pattern of lung function established in early childhood may be followed by accelerated lung function decline in adulthood, lead to early onset of chronic obstructive disease and increased mortality (31).

Many published papers on wheeze and lung function measured expiratory flows using various techniques. For example, Delacourt et al. (32) measured maximal flow at functional residual capacity at age 17 months and reported that persistent wheeze was associated with reduced lung function. The German Multicentre Allergy Study (33) following children from birth to 7 years of age found out that although transient early wheezers had normal or subnormal lung function, children with persistent and late-onset of symptoms at age 7 years showed significant drop of expiratory flow volumes. Also Dutch Prevention and Incidence of Asthma and Mite Allergy (34) established that resistance measured by the interrupter

technique was higher in 4-year-olds with persistent wheeze than in children who had never wheezed or had transient early wheeze. The Manchester Asthma and Allergy Study (11) has proved that reduced lung function (plethysmographic measure of sRaw) at age 3 years predicted the persistence of symptoms only in children who had wheezed over the first 3 years, but lower lung function was not associated with the onset of wheeze after age 3 if children not wheezed previously. The ALSPAC cohort study (35) also added evidence that wheezing phenotypes were associated with decrements of FEF₂₅₋₇₅ and increased airway responsiveness at 8 – 9 years of age compared with never/infrequent wheezers, and the greatest lung function decrements were associated with persistent wheezing.

Up to now there is a scarcity of reports regarding the association of wheeze with restrictive lung function pattern. The Finnish study (36) on wheezing with RSV confirmed a reduced FVC suggesting a restrictive pattern of lung function. Very recent prospective birth cohort study of Simpson et al. (37) documented that children hospitalized because of lower respiratory illness (LRI) in the first year of life had a restrictive lung function pattern without an increased airway responsiveness at age 8. In contrast to that, children who experienced LRI in the third year of life showed obstructed lung function with increased bronchial hypersensitivity. The authors concluded that severe LRI occurring at different ages in early life may have different pathogenesis and long term sequelae. This observation would be consistent with our findings, which have shown that restrictive lung function pattern was observed in recurrent wheezers with the onset of symptoms in the first two years of life.

Our study would be at variance with the results of the Childhood Allergy Study birth cohort in northern Detroit (Michigan) where wheeze from birth through the age of 6 to 7 years has been monitored annually by interviews and lung function was measured between the ages of 6 and 7 years (38). The findings indicated that the log transformed percentage of predicted FEV₁ for children who completed spirometry at the age of 6 to 7 years was not associated with wheeze present at any age. The results of the latter study, however, are not comparable with our observations since in that study detailed data were not collected on the frequency, severity, or duration of wheeze.

The strength of our study in terms of the relationship between early wheeze and lung expiratory volumes stems from the fact that we were able to monitor wheezing and its severity over many regular time points in the course of face-to-face interviews with mothers of children. Face-to-face interviews facilitated much better explanation of wheeze to parents. Other studies considered only current wheeze (over the last 12 months) and very often used self-administered questionnaires. We believe that the quarterly and semiannually collected data on wheeze offered us a much better opportunity to explore the importance of early wheeze in subsequent follow-up periods. On the other hand, we are aware of the limitations of our study which are mainly related to the relatively small sample size and the shortage of cases with recurrent wheeze. However, the results are internally consistent with the decrement of expiratory lung volumes across the wheezing strata. Because of the underpowered statistical analysis of the results the conclusions from the study should be drawn with caution. Another weak point of the study is the fact that the occurrence of wheezing over the follow up was not physician-confirmed, but the significant association between regularly collected parental data on a child's history of wheeze and the objective measurement of lung function may be an indirect evidence for treating parental reporting on children's wheeze as trustworthy.

In conclusion, our findings show that wheezing experience during early life may be associated with lung function deficit of restrictive character in preschool children and detailed history of early wheeze, even though not physician-confirmed, may help define the

high risk group of children for poor lung function testing. In order to confirm the preliminary results of this study, further monitoring of the cohort over several forthcoming years is planned.

Acknowledgments

This is part of an ongoing comparative longitudinal investigation on the health impact of prenatal exposure to outdoor/indoor air pollution in infants and children being conducted in New York City and Krakow. The study received funding from an RO1 grant entitled, "Vulnerability of the Fetus/Infant to PAH, PM_{2.5} and ETS" (5 RO1 ES10165 NIEHS; 02/01/00 - 01/31/04) and from the NIEHS (RO1 ES010165-0451) the Lundin Foundation and the Gladys T. and Roland Harriman Foundation. Principal investigator: Prof. FP Perera.

The study was also partly supported by the grant from the International Center for Research in Biomedicine. Luxembourg.

References

1. Dezateux C, Stocks J. Lung development and early origins of childhood respiratory illness. *Br Med Bull.* 1997; 53:40–57. [PubMed: 9158283]
2. Stick S. The contribution of airway development to paediatric and adult lung disease. *Thorax.* 2000; 55:587–594. [PubMed: 10856320]
3. Johnson CC, Ownby DR, Zoratti EM, Alford SH, Williams LK, Joseph CLM. Environmental epidemiology of pediatric asthma and allergy. *Epidemiol Rev.* 2002; 24:154–175. [PubMed: 12762090]
4. Bracken MB, Belanger K, Cookson WO, Triche E, Christiani DC, Leaderer BP. Genetic and perinatal risk factors for asthma onset and severity: a review and theoretical analysis. *Epidemiol Rev.* 2002; 24:176–189. [PubMed: 12762091]
5. Von Mutius E. Paediatric origins of adult lung disease. *Thorax.* 2001; 56:153–157. [PubMed: 11209107]
6. Martinez FD, Morgan WJ, Wright AL, Holberg CJ, Taussig LM. Diminished lung function as a predisposing factor for wheezing respiratory illness in infants. *N Engl J Med.* 1988; 319:1112–1117. [PubMed: 3173442]
7. Tager IB, Hanrahan JP, Tosteson TD, Castile RG, Brown RW, Weiss ST, Speizer FE. Lung function, pre- and post-natal smoke exposure, and wheezing in the first year of life. *Am Rev Respir Dis.* 1993; 147:811–817. [PubMed: 8466114]
8. Strachan D, Gerritsen J. Long-term outcome of early childhood wheezing: population data. *Eur Respir J.* 1996; 9:42–47. [PubMed: 8834332]
9. Dezateux C, Stocks J, Wade AM, Dundas I, Fletcher ME. Airway function at one year: association with premorbid airway function, wheezing and maternal smoking. *Thorax.* 2001; 56:680–686. [PubMed: 11514687]
10. Turner SW, Palmer LJ, Rye PJ, Gibson NA, Judge PK, Young S, Landau LI, Le Souef PN. Infants with flow limitation at 4 weeks: outcome at 6 and 11 years. *Am J Respir Crit Care Med.* 2002; 165:1294–1298. [PubMed: 11991882]
11. Lowe LA, Simpson A, Woodcock A, Morris J, Murray C, Custovic A. Wheeze phenotypes and lung function in preschool children. *Am J Respir Crit Care Med.* 2005; 171:231–237. [PubMed: 15502115]
12. Gappa M, Bush A. Pre-school Wheeze: More Questions Than Answers. *Pediatric Pulmonology.* 2006; 41:910–911. [PubMed: 16933219]
13. Sly, PD.; Willet, K. Developmental physiology. In: Silverman, M., editor. In the book *Childhood Asthma and Other Wheezing Disorders*. Chapman & Hall; London: 1995. p. 55–66.
14. Dockery DW, Berkey CS, Ware JH, Speizer FE, Ferris BG Jr. Distribution of forced vital capacity and forced expiratory volume in one second in children 6 to 11 years of age. *Am Rev Respir Dis.* 1983; 128:405–412. [PubMed: 6614634]

15. Kanner RE, Schenker MB, Munoz A, Speizer ER. Spirometry in children: methodology for obtaining optimal results for clinical and epidemiologic studies. *Am Rev Respir Dis.* 1983; 127:720–724. [PubMed: 6859655]
16. Strachan DP. Repeatability of ventilatory lung function measurements in a population survey of 7 year old children. *Thorax.* 1989; 44:474–479. [PubMed: 2763257]
17. Wang X, Dockery DW, Wypij D, Fay ME, Ferris BG. Pulmonary function between 6 to 18 years of age. *Pediatr Pulmonol.* 1993; 15:75–88. [PubMed: 8474788]
18. Quanjer P, Tammeling G, Cotes J, et al. Lung volume and forced ventilatory flows; 1993 update. Working Party Report “Standardisation of lung function tests”. European Coal and Steel Community and European Respiratory Society. *Eur Respir J.* 1993; 6(Suppl 16):5–40. [PubMed: 8381090]
19. Studnicka M, Frischer T, Neumann M. Determinants of reproducibility of lung function tests in children aged 7 to 10 years. *Pediatr Pulmonol.* 1998; 25:238–243. [PubMed: 9590484]
20. Miller MR, Crapo R, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, Enright P, Grinten CPM, Gustafsson P, Jensen R, Johnson DC, MacIntyre N, McKay R, Navajas D, Pedersen OF, Pellegrino R, Viegi G, Wanger J. General considerations for lung function testing. *Eur Respir J.* 2005; 26:153–161. [PubMed: 15994402]
21. American Thoracic Society Documents. An Official American Thoracic Society/European Respiratory Society Statement: Pulmonary Function Testing in Preschool Children. *Am J Respir Crit Care Med.* 2007; 175:1304–1345. [PubMed: 17545458]
22. Arets HGM, Brackel HJL, van der Ent CK. Forced expiratory manoeuvres in children: do they meet ATS and ERS criteria for spirometry? *Eur Respir J.* 2001; 18:655–660. [PubMed: 11716170]
23. Jedrychowski W, Whyatt R, Camann D, Bawle U, Peki K, Spengler J, Dumyahn T, Perera F. Effect of prenatal PAH exposure on birth outcomes and neurocognitive development among a cohort of Polish mothers and newborns. Study design and preliminary ambient data. *Int J Occup Med Environ Hlth.* 2003; 16:21–29.
24. STATA software for windows, release 10. StaCorp; Texas: 2007.
25. Kohler, U.; Kreuter, F. Data analysis using STATA. Stata Press Publication; College Station, Texas: 2005.
26. Uhl E, Castleman WI, Sorkness RI, Lemanske R, McAllister PK. Parainfluenza virus-induced persistence of airway inflammation, fibrosis, and dysfunction associated with TGF- β 1 expression in Brown Norway rats. *Am J Respir Crit Care Med.* 1996; 154:1834–1842. [PubMed: 8970378]
27. Dosanjh A, Rednam S, Martin M. Respiratory syncytial virus augments production of fibroblast growth factor basic in vitro: implication for a possible mechanism of prolonged wheezing after infection. *Pediatr Allergy Immunol.* 2003; 14:437–440. [PubMed: 14675469]
28. Sanders SP. Asthma, viruses, and nitric oxide. *Proc Soc Exp Biol Med.* 1999; 220:123–132. [PubMed: 10193439]
29. Erlevyn-Lajeunesse MDS, Hunt LP, Pohunek P, Dobson SJ, Kochnar P, Warner JA, Warner JO. Bronchoalveolar lavage MMP-9 and TIM-1 in preschool wheezers and their relationship to persistent wheeze. *Pediatr Res.* 2008; 64:194–199. [PubMed: 18391843]
30. Holgate, S. WHO monograph: Effects of Air Pollution on Children’ Health and Development. Review of evidence. WHO Regional Office for Europe; 2005. Mechanisms by which air pollution injures the child’s respiratory system; p. 29–43.
31. Guerra S, Sherrill DL, Venker C, Halonen M, Martinez FD. Chronic restrictive lung pattern is associated with increased mortality risk. *Am J Respir Crit Care Med.* 2009; 179:A4325.
32. Delacourt C, Benoist MR, Waernessyckle S, Rufin P, Brouard JJ, deBlic J, Scheinmann P. Relationship between bronchial responsiveness and clinical evolution in infants who wheeze. A four-year prospective study. *Am J Respir Crit Care Med.* 2001; 164:1382–1386. [PubMed: 11704582]
33. Lau S, Illi S, Sommerfeld C, Niggemann B, Volkel K, Madloch C, Gruber C, Nickel R, Forster JU, Wahn U. Transient early wheeze is not associated with impaired lung function in 7-yr-old children. *Eur Resp J.* 2003; 21:834–841.
34. Brussee JE, Smit HA, Koopman LP, Wijga AH, Kerkhof M, Corver K, Vos AP, Gerritsen J, Grobbee DE, Brunekreef B. Interrupter resistance and wheezing phenotypes at 4 years of age.

- AmJ Respir Crit Care Med. 2004; 169:209–213. [Published erratum appears in AmJ Respir CritCare Med 2004;169:1262]. [PubMed: 14597483]
35. Henderson J, Granell R, Heron J, Sherriff A, Simpson A, Woodcock A, DP, Strachan DP, Shaheen SO, Sterne JAC. Associations of wheezing phenotypes in the first 6 years of life with atopy, lung function and airway responsiveness in mid-childhood. *Thorax*. 2008; 63:974–980. [PubMed: 18678704]
36. Hyvarinen MK, Kotaniemi-Syrjanen A, Reijonen TM, Korhonen K, Korppi MO. Lung function and bronchial hyper-responsiveness 11 years after hospitalization for bronchiolitis. *Acta Paediatrica*. 2007; 96:1464– 1469. [PubMed: 17880415]
37. Simpson A, Semic-Jusufagic A, Lowe L, Woodcock A, Murray C, Martinez FD, Custovic A. Heterogenous lung function outcome of hospital admission for lower respiratory illness in early life. *Am J Respir Crit Care Med*. 2009; 179:A6004.
38. Wegienka G, Havstad S, Zoratti EM, Ownsby DR, Johnson ChC. Association of early life wheeze and lung function. *Ann Allergy Asthma Immunol*. 2009; 102:29– 34. [PubMed: 19205282]

Figure 1.

Table 1

Characteristics of the groups of children with acceptable and not acceptable spirometry*

	Acceptable spirometry N=139	Not acceptable spirometry N=119	Spirometry (-) N=124	Whole cohort 4 years n=382
Maternal characteristics				
Mother's age:				
mean	27.87	27.56	27.94	27.80
SD	3.290	3.565	3.478	3.433
Maternal education:				
elementary n (%)	12 (8.6)	12 (10.1)	11 (8.9)	35 (9.2)
medium n (%)	33 (23.7)	27 (22.7)	32 (25.8)	92 (24.1)
higher n (%)	94 (67.6)	80 (67.2)	81 (65.3)	255 (66.8)
Maternal atopy (+) n (%)	27 (19.4)	32 (26.9)	35 (28.2)	94 (24.6)
Infant characteristics				
Gender:				
Boys n (%)	67 (48.2)	66 (55.5)	57 (46.0)	190 (49.7)
Girls n (%)	72 (51.8)	53 (44.5)	67 (54.0)	192 (50.3)
Parity:				
1 n (%)	88 (63.3)	73 (61.3)	80 (64.5)	241 (63.1)
2 n (%)	51 (36.7)	46 (38.7)	44 (35.5)	141 (36.9)
Gestational age (weeks):				
mean	39.42	39.29	39.21	39.31
SD	1.31	1.78	1.66	1.58
Birth weight (g):				
mean	3386.6	3448.3	3362.0	3397.9
SD	452.6	514.4	523.2	495.8
Length at birth (cm):				
mean	54.40	55.08	54.29	54.58
SD	2.59	3.13	3.09	2.94
Breastfeeding:				
6 months n (%)	39 (28.1)	35 (29.4)	43 (34.7)	117 (30.6)
> 6 months n (%)	100 (71.9)	84 (70.6)	81 (65.3)	265 (69.4)
Any wheeze in the follow-up n (%)	54 (38.8)	45 (37.8)	56 (45.2)	155 (40.6)

* No characteristics differed significantly across the groups at the p level 0.05

Table 2

Lung function indices grouped by acceptable and not acceptable exhalation efforts

	Acceptable spirometry N=139	Not acceptable spirometry N=119	Total Spirometry N=258	P level for difference between acceptable and not acceptable spirometry
FVC (ml):				
mean	867.3	947.7	904.4	
SD	168.5	256.1	216.8	0.0028
FEV ₁ (ml):				
mean	838.2	912.1	872.2	
SD	157.9	244.9	205.7	0.0038
FEV ₀₅ (ml):				
mean	686.2	734.0	708.2	
SD	145.2	203.7	175.9	0.0291

Table 3

Frequency of wheezing episodes and the number of wheezing days recorded in the follow-up grouped by wheezing phenotype

Wheezing phenotypes	Number of children	Number of wheezing episodes Mean (SD)	Number of wheezing days Mean (SD)
No wheezing	85	-	-
Episodic wheezing	39	1.15 (1.43)	8.89 (14.01)
Transient wheezing	9	3.22 (1.39)	28.78 (16.13)
Recurrent wheezing	6	6.13 (3.38)	39.17 (21.21)
Total	139	0.81 (1.66)	6.13 (13.92)

Table 4

Estimated effect of wheezing experienced by children over the follow-up on lung function indices adjusted for gender and height (the multivariable linear regression models)

	N	Regression coeff.	95% CI	P value
FVC				
<i>1. Wheezing episodes</i>				
No wheezing	85	Reference category		
1 – 2 episodes	43	3.5	-53.5 60.6	0.904
> 2 episodes	11	-120.5	-218.2 -22.8	0.016
<i>2. Wheezing days</i>				
No wheezing	85	Reference category		
1–10 wheezing days	31	1.81	-62.2 65.8	0.955
>10 wheezing days	23	-87.4	-162.4 -12.3	0.016
FEV₁				
<i>1. Wheezing episodes</i>				
No wheezing	85	Reference category		
1 – 2 episodes	43	13.8	-39.2 65.8	0.607
> 2 episodes	11	-98.3	-189.0 -7.5	0.034
<i>2. Wheezing days</i>				
No wheezing	85	Reference category		
1–10 wheezing days	31	10.4	-49.3 70.1	0.731
>10 wheezing days	23	-65.7	-135.7 4.3	0.066
FEV₀₅				
<i>1. Wheezing episodes</i>				
No wheezing	85	Reference category		
1 – 2 episodes	43	6.7	-42.5 56.0	0.788
> 2 episodes	11	-44.7	-129.0 39.6	0.296
<i>2. Wheezing days</i>				
No wheezing	85	Reference category		
1–10 wheezing days	31	9.3	-45.8 64.4	0.740

	N	Regression coeff.	95% CI		P value
>10 wheezing days	23	-46.1	-110.7	18.5	0.161
FEV₀₅/FVC (%)					
<i>1. Wheezing episodes</i>					
No wheezing	85	Reference category			
1 – 2 episodes	43	0.58	-4.23	5.40	0.810
> 2 episodes	11	4.27	-3.61	12.16	0.286
<i>2. Wheezing days</i>					
No wheezing	85	Reference category			
1–10 wheezing days	43	1.20	-4.01	6.42	0.649
>10 wheezing days	11	1.72	-4.63	8.07	0.593