Published in final edited form as:

Gen Hosp Psychiatry. 2013 ; 35(4): 349–353. doi:10.1016/j.genhosppsy.2013.03.009.

Furthering the reliable and valid measurement of mental health screening, diagnoses, treatment, and outcomes through health information technology

Jessica E. Haberer, Tom Trabin, and Michael Klinkman

I. Why is this theme a priority?

Health information technology (IT) may hold the key to overcoming many of the challenges in measurement of mental health. Current measures for screening, diagnosis, treatment, and clinical outcome assessment rely primarily on traditional models of care delivery involving face to face (synchronous) interactions between provider and patient (for example, clinic visits). Relying on synchronous interactions means that measurement is highly likely to take place on an irregular and infrequent basis, reducing the potential effectiveness of measurement-based care. Effective use of health IT can extend care into asynchronous spaces, broaden the populations reached, expand information exchange between provider and patient, and thereby increase the effectiveness and efficiency of mental health care. Fertile areas for research in this area include the invention of new measurement technologies, as well as more effective implementation of existing technologies to serve diverse settings, populations, and health care infrastructures.

II. What has been done in this area?

Mental health care has historically relied on subjective measurement, typically structured patient interview or self-report linked to a direct clinical encounter. For instance, an individual may report depression symptoms on a self-report scale while waiting to see her physician or report on the mental health of others, as a parent reporting on the behavior of a child. Subjective measurement is appealing in that it is inexpensive, fast, and easy to implement, but it can underestimate socially unacceptable behavior (e.g. alcohol consumption) and overestimate socially desirable behavior (e.g. medication adherence)[1]. It is also inherently limited by its dependence on accurate memory and correct attribution of ambiguous symptoms. Individuals may have difficulty recalling events that took place more than a few days ago, especially if cognitive impairment is a factor, but most self-report measures specify longer recall periods on patterns of behavior [2].

Objective measurement has had a less prominent role in behavioral health measurement, but passive methods (biosensors to record sympathetic nervous system activity) and active methods (blood samples to measure serum drug levels) are now available. Compared to subjective measures, objective methods have higher correlation with biological and clinical markers [3, 4]. However, they tend to be expensive and resource intensive, and measurable

© 2013 Elsevier Inc. All rights reserved.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

behaviors or outcomes may not directly correlate with the mental health issue being investigated [5–7].

The reliability and validity of measures can be established through psychometric testing of subjective scales or measures, or formal biologic assay development for serum drug levels, but to achieve widespread use measurement must also be affordable, feasible, and acceptable to clinicians and patients. This is a difficult balance. For example, medication adherence can be assessed through self-report or through any of several objective measures (pill counts, pharmacy refill data, or electronic monitors that record when a medication cap has been removed). While self-report may overestimate adherence [8, 9], the objective measures also present problems: manual pill counts require significant time and cooperation between patient and clinician; pharmacy data can only record whether pills have been obtained, not taken; and an electronic medication bottle may be only be opened once in a week, when the patient fills a weekly pill box.

Mental health screening

Screening or case-finding for mental health problems has traditionally involved questionnaire and scale administration in formal settings, such as a primary care doctor's office or an educational institution. Several reliable and valid self-report questionnaires have been effectively employed in these settings; current examples include the GAD-7 [10, 11], the CAGE alcohol screen [12], the PHQ-2 and PHQ-9 [13, 14], the QIDS-SR [15], the Edinburg Perinatal Depression Scale [16], the Geriatric Depression Scale [17] and many others. However, this approach only reaches individuals who are actively engaged in health care.

Some innovative technological approaches have been developed to cast a wider net in mental health screening. Internet web sites now offer mental health screening for adults [18] and youth [19], and outreach can be enhanced through mobile phone applications [20] and online communities [21]. These approaches may be useful for specific groups who frequent the Internet, such as men who have sex with men [22] or adolescents and young adults [23, 24]. Clinicians or programs may direct individuals to complete screening questionnaires online, or the web sites may be available to all users (for example, www.mentalhealthscreening.org and <http://doyouneedtherapy.com>). Similar mass screening can be conducted off line through computerized forms [25] and may involve novel technologies, such as interactive voice response [26–28]. This approach to screening can be advantageous in that the questionnaires are completed privately or anonymously, thus potentially decreasing social desirability bias [29]; however, individuals must be adept at using technology [30].

Expanded screening has the potential to bring many more individuals into care; however, the reliability and validity of technologically based screening remains unclear and we do not yet know if on-line screening leads to increased rates of treatment. The US Preventive Services Task Force underscores the importance of linking mental health screening to systems of care in recommending screening for depression only when depression care supports are in place [31].

Mental health diagnosis

The standard for mental health diagnosis has long required in-person clinical assessment. In recent years this scenario has been expanded through new technologies including telemedicine [32–35] and computerized questionnaires, some implementing computerized adaptive testing methodologies [36–40]. In this latter approach assessments are tailored to the individual through the use interactive response technology or branching logic based on

specific responses to questions or characteristics (e.g. age, gender), thus leading to diagnostic efficiency. These technologies offer a promising option for reaching rural populations where mental health care resources are extremely limited [32, 34]. Preliminary evidence suggests that the reliability and validity of diagnoses made through the assistance of technology is acceptable when compared to in-person assessment [33, 38, 40, 41].

Mental health treatment- quantity, quality, and adherence

Current approaches to measuring the quantity or quality of treatment vary by type of treatment. Provision of counseling or behavioral therapy can be measured by number of sessions attended or modules completed, and care quality can be assessed through direct observation and critique, by measured adherence to a protocol, or by general measures of provider “quality”. To date, measurement criteria for non-pharmacologic treatment have been largely subjective and based on self-report or clinician judgment [42, 43], quality assessment of providers has been rare outside the training setting, and attention to decision-making ability has often been lacking [44]. Technology-based approaches are now beginning to overcome these limitations. Improved functionality and more widespread implementation of electronic health records has enabled tracking of engagement in care [45], and virtual human agents have been used to assess provider skills [46–48]. Further research on reliability and validity of these new approaches to measurement is needed.

Medication adherence is perhaps the best-studied and implemented category of mental health treatment measurement. As described above, both subjective and objective measures are available but all are subject to imprecision and bias. Recent efforts have focused on technologic enhancements in assessing individual patient medication adherence, such as wireless medication devices [49], frequent self-report through mobile phones [50–52], and Web-based portals [53], with varying levels of success. One novel approach employing mobile phone-based context-sensing to monitor and enhance self-management for depression has just been tested [54].

Mental health outcomes

Incremental changes in behavior or symptom severity over time are routinely assessed through subjective clinical opinion, as well as through scores on validated self-report questionnaires and scales. While an individual’s progress during treatment can be tracked using pen and paper methods, technology-based approaches to outcome monitoring can greatly enhance the integration of outcome measurement into clinical care. Outcome measurement can be in turn linked to clinical decision support tools at the point of care to improve routine practice; this general approach has been given the term *measurement-based care* [55–57]. The literature contains calls for technology to enable rapid development and deployment of various outcome reports, decision support tools, and measurement feedback systems [44, 58, 59]; this work is just now beginning. Outcome measurement instruments have been integrated into electronic health records (EHRs) to routinely appear for individual patients who have the diagnosis of a specific mental health problem [60, 61]; some work has been done to integrate simple “one-click” decision support protocols or prompts that appear for patients with active diagnosis of depression or bipolar disorder [60, 62]. Additionally, mobile devices could allow assessment of actual activity or behavior patterns (e.g. physical activity, duration, continuity of sleep) or direct assessment of behavioral indicators (e.g. voice tone, speed of response) [63][<http://www.cogitocorp.com>]. These tools could help move mental health assessment away from self-report to more objective measures. While these innovations are promising, there is little evidence to date to support their effectiveness.

Technology has also been applied to enhance population-based measurement. Several commercial EHRs and population management software systems now have the capacity to

aggregate outcome measurement across all patients with a specific diagnosis to facilitate population management. Another recent initiative is the development of comprehensive behavioral health reporting systems to facilitate clinical and administrative use of evidence-based practices. In support of this initiative, the National Cancer Institute in collaboration with content area experts has begun work to identify or develop consensus standards for assessing and reporting on key behavioral domains [64, 65].

The design processes underlying system mapping of decision points and distillation of performance information at the individual, caseload, and organizational levels can be implemented to support clinical practice in a wide variety of settings [66, 67]. At the individual level, rapid feedback of outcome measure scores to clinicians can inform midcourse corrections in treatment strategies. At the administrative level, outcome data can be aggregated for all patients of individual clinicians and used as one indicator of clinicians' treatment effectiveness. At the population level, time-series measurements can be compared against expectancy curves of improvement based upon aggregated data for predefined target populations. At this stage of development, validity and reliability remain important concerns, as EHRs have been shown to potentially underreport practice performance [68, 69].

Finally, standardized outcome measurement can support incentive-based reimbursement models. For instance, payment can be provided to the individual or the provider for successful cessation of substance abuse or improved depression remission rate. These arrangements have been found to have substantial and replicable benefits in adult mental health treatment and are currently being tested in youth mental health practice settings [70, 71].

III. What work is needed?

The future of IT for measurement of mental health lies in three key areas: *development and adaptation of reliable and valid measures to provide useful just-in-time data; management of high volume data; and determining the acceptability, feasibility, and sustainability of measurement technology for target populations.*

Development and adaptation of reliable and valid measures to provide useful just-in-time data

Just-in-time is a concept in which information is made available to those who need it at the moment when they need it, and is greatly facilitated by technology [72]. Data can be transmitted in real-time, minimizing recall bias and enabling proactive approaches to disease management. Consider this example.

Mr. Jones uses his smart phone (or his laptop computer, or another personal digital assistant) to interact with his primary care practice. The practice identifies him as a "high-risk" patient for onset of depression due to his family history of depression and comorbid medical problems of diabetes and coronary artery disease, and on May 10 sends out a PHQ-9 scale to all high-risk patients asking them to respond and post results back to the practice inbox. Mr. Jones does so. His score is recorded at 14, representing moderate depression, and is flagged for follow-up. A staff member calls, confirms his symptoms, then schedules an office visit to review his symptoms to see whether treatment is needed. At his appointment 2 weeks later, he completes another PHQ-9 along with screening items for comorbid mental health problems while waiting to be seen. Data is entered directly into the practice EHR, scored, and posted on the front screen of his patient file to be reviewed when his physician opens his record. At the end of the visit, after the diagnosis is confirmed, Mr Jones agrees to start treatment with a selective serotonin reuptake inhibitor. One week after the visit, the practice

EHR pushes an outcome assessment scale including PHQ-9, items assessing side-effects and functional status to his smart phone. He completes the items and sends the scale back to the practice inbox. The EHR compares the PHQ score to previous scores and sends a longitudinal graph back to the smartphone and to his physician's inbox. His physician decides when to schedule the next electronic follow up and/or next office follow up based upon these results.

After several weeks, Mr Jones is doing well, with PHQ score indicating remission of depression. His physician initiates follow up electronic monitoring, sets the interval for his EHR to push the outcome assessment to Mr Jones' smart phone at every 2 months, and sets the threshold for an immediate physician alert to a score of 8. Mr Jones is also shown how to access the outcome assessment at a time of his own choosing if he desires more frequent self-monitoring or feels that his symptoms are recurring. Over the next several months, he completes several measures and remains in remission.

Although not specific to mental health, the New York City Department of Health has implemented such an application of PDAs in primary care clinics [73]. Most of the components needed to implement this approach for depression care have already been introduced (see above), but full integration into practice workflow has yet to be accomplished. The potential response burden for patients with comorbid mental health problems (each with their own measurement protocols) and medical problems can be addressed with use of item response theory-derived personalized measurement, as in the PROMIS system [36, 74]. Considerable investment will be necessary to develop a robust, reliable, and efficient interface to support this level of care.

EHRs can also provide just-in-time support to facilitate treatment measurement, such as electronic prompts for obtaining serum drug levels at recommended intervals or when most recent level is out of a target range. This is becoming standard practice for some medical problems such as anticoagulation management for patients taking warfarin [75, 76]; adapting these protocols to support mental health medication management should be relatively easy to accomplish.

Management of high volume data

Technology readily enables the collection of vast amounts of data. For example, automated text messages can be used to prompt the collection of symptoms of depression on a daily, hourly, or random basis from hundreds - or thousands - of individuals. Similarly, wireless adherence devices can automatically determine if individuals open their medication bottles at recommended times. The number data points quickly add up and can overwhelm a provider, clinic, or service agency. Development of data management systems that can accurately sort out meaningful signal (missed openings of a medication bottle) from background noise (on-time openings of the bottle) will be extremely important for the next generation of mental health information technology.

Determining the acceptability, feasibility, and sustainability of measurement technology in selected populations

The advances in health IT described above have the potential to transform mental health care. To fulfill that potential, the next wave of research will need to carefully address the acceptability, feasibility, and ease of implementation of proposed new technologies very early in the development process. The majority of primary care practices in the US have now implemented an EHR [77], and many will be effectively "locked in" to the specific technology embedded in that EHR for years to come [78]. Mental health measurement technologies that cannot be easily grafted on to existing EHRs, or that do not fit into the

workflow of the developing Patient-Centered Medical Home [79], may prove very difficult to disseminate to or sustain in community practice settings. In specialty mental health practices, traditionally more resistant to adoption of EHRs [80], technology-enabled measurement may be difficult to implement in any form.

In the next wave, it will be important to match technology to the population and setting. Interactive smart phone technology can enhance measurement in practices with highly optimized EHRs and a large proportion of patients who can make use of that technology, but rural practices serving a high proportion of uninsured and indigent patients would not likely benefit from investing in that technology. The frequency and intensity of data collection must be carefully matched to the needs of patients and practices, given the chronic and comorbid nature of many mental health problems. User interfaces must be tailored to the specific needs of patients, providers, and administrators to enable effective use of collected data. Research partnerships that bring together innovative technology developers and investigators in the emerging field of implementation science will be critical to the success of this next wave of research.

IV. Challenges

Along with the promises of health IT come significant challenges: •

- *Privacy and data transmission-* Privacy is a significant concern, given the stigma associated with mental health conditions, and individuals must give informed consent prior to participating in studies or programs. All transmission of mental health IT data, which often involves multiple parties, must be secure and compliant with the Health Insurance Portability and Accountability Act. Security can be achieved through techniques currently in use, such as data encryption, numeric subject identifiers, and personal identification numbers. Alternative methods include voice or voice pattern recognition and biometrics, such as fingerprints and iris scans. Data must also be transmitted and verified in the timely manner associated with the technology.
- *Efficiency of implementation.* Data collection with rapid feedback of results for quality improvement can be accomplished most efficiently with an electronic health record infrastructure. The widespread use of outcome measures to improve treatment quality, support quality assessment, or enhance reimbursement will in the end be somewhat dependent on the (slow-paced) adoption of electronic health record systems.
- *Cost and scalability.* New technology developments, whether hardware, applications, or human oversight, can initially impose significant costs. This barrier can be ameliorated through adaptation of existing technologies and open-source code, as well as collaborations to allow for economies of scale, such as mass production. In resource-limited regions, cost-savings, not just cost-effectiveness, may be required for implementation.
- *Expansion of mental health IT to populations without previous exposure to or understanding of technology, such as the elderly or illiterate.* Electronic measurement and web-based interactions may be impractical or unavailable to these groups. Persons living in remote or rural areas may have more limited access to the Internet or remote measurement technologies. Elderly persons may be less inclined to interact with online mental health interventions or assessment protocols. Persons with more severe or complex emotional disorders may have particular problems in adhering to or following through with on-line protocols. This is an area in which more research is clearly needed.

V. Conclusion

Health IT is a fast-growing field that is infused with an inspiring sense of enthusiasm. New technologies for measurement of mental health are being developed rapidly with many potential applications and clinical benefit; however, scientific rigor is critical for ensuring they have the reliability and validity necessary to effectively support and ideally integrate screening, diagnosis, treatment, and outcomes. Key factors for mental health research include development and adaptation of measures to provide useful just-in-time data; management of high volume data; and determining the acceptability, feasibility, and sustainability of measurement technology for relevant populations. Researchers and implementers must also consider the challenges associated with developing this technology: privacy and data transmission, cost and scalability, and expansion to populations without previous exposure to or understanding of technology.

REFERENCES

1. Sajatovic M, et al. Measurement of psychiatric treatment adherence. *Journal of Psychosomatic Research*. 2010; 69(6):591–599. [PubMed: 21109048]
2. Wilson IB. Improving the self-report of HIV antiretroviral medication adherence: is the glass half full or half empty? *Current HIV/AIDS reports*. 2009; 6(4):177–186. [PubMed: 19849960]
3. Simoni JM. Self-report measures of antiretroviral therapy adherence: A review with recommendations for HIV research and clinical management. *AIDS and behavior*. 2006; 10(3):227–245. [PubMed: 16783535]
4. Nieuwkerk PT. Self-reported adherence to antiretroviral therapy for HIV-1 infection and virologic treatment response: a meta-analysis. *Journal of acquired immune deficiency syndromes (1999)*. 2005; 38(4):445–448. [PubMed: 15764962]
5. Peterson BT, et al. Comparison of actigraphy and polysomnography to assess effects of zolpidem in a clinical research unit. *Sleep Medicine*. 13(4):419–424. [PubMed: 22317945]
6. Razavi N, et al. Measuring motor activity in major depression: The association between the Hamilton Depression Rating Scale and actigraphy. *Psychiatry Research*. 2011; 190(2–3):212–216. [PubMed: 21663976]
7. Fletcher RR, et al. Wearable sensor platform and mobile application for use in cognitive behavioral therapy for drug addiction and PTSD. *Engineering in Medicine and Biology Society, EMBC, 2011 Annual International Conference of the IEEE*. 2011
8. Garber MC, et al. The Concordance of Self-Report with Other Measures of Medication Adherence: A Summary of the Literature. *Medical Care*. 2004; 42(7):649–652. [PubMed: 15213489]
9. Shi L, et al. Concordance of Adherence Measurement Using Self-Reported Adherence Questionnaires and Medication Monitoring Devices. *PharmacoEconomics*. 2010; 28(12):1097–1107. [PubMed: 21080735]
10. Ruiz MA, et al. Validity of the GAD-7 scale as an outcome measure of disability in patients with generalized anxiety disorders in primary care. *Journal of Affective Disorders*. 128(3):277–286. [PubMed: 20692043]
11. Spitzer R, et al. A brief measure for assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine*. 2006; 166(10):1092–1097. [PubMed: 16717171]
12. Ewing JA. Detecting alcoholism. The CAGE questionnaire. *JAMA : the journal of the American Medical Association*. 1984; 252(14):1905–1907. [PubMed: 6471323]
13. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*. 2001; 16(9):606–613. [PubMed: 11556941]
14. Kroenke K, et al. The Patient Health Questionnaire Somatic, Anxiety, and Depressive Symptom Scales: a systematic review. *General Hospital Psychiatry*. 32(4):345–359. [PubMed: 20633738]
15. Rush AJ, et al. The 16-Item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): a psychometric evaluation in patients with chronic major depression. *Biol Psychiatry*. 2003; 54(5):573–583. [PubMed: 12946886]

16. Schaper AM. Use of the Edinburgh Postnatal Depression Scale to identify postpartum depression in a clinical setting. *Journal of reproductive medicine*. 1994; 39(8):620–624. [PubMed: 7996526]
17. Hoyl MT, et al. Development and testing of a five-item version of the Geriatric Depression Scale. *J Am Geriatr Soc*. 1999; 47(7):873–878. [PubMed: 10404935]
18. Bolge SC. Characteristics and symptomatology of patients diagnosed with unipolar depression at risk for undiagnosed bipolar disorder: a bipolar survey. *CNS spectrums*. 2008; 13(3):216–224. [PubMed: 18323755]
19. Diamond G, et al. Development, Validation, and Utility of Internet-Based, Behavioral Health Screen for Adolescents. *Pediatrics*. 2010; 126(1):e163–e170. [PubMed: 20566613]
20. Reid S, et al. A mobile phone application for the assessment and management of youth mental health problems in primary care: a randomised controlled trial. *BMC Family Practice*. 2011; 12(1): 131. [PubMed: 22123031]
21. Giles DC, Newbold J. Self- and Other-Diagnosis in User-Led Mental Health Online Communities. *Qualitative Health Research*. 2011; 21(3):419–428. [PubMed: 20739589]
22. Hirshfield S. Screening for depressive symptoms in an online sample of men who have sex with men. *AIDS care*. 2008; 20(8):904–910. [PubMed: 18720088]
23. Kauer SD. Self-monitoring using mobile phones in the early stages of adolescent depression: randomized controlled trial. *Journal of medical Internet research*. 2012; 14(3):e67. [PubMed: 22732135]
24. Whittaker R, et al. MEMO--a mobile phone depression prevention intervention for adolescents: development process and postprogram findings on acceptability from a randomized controlled trial. *Journal of medical Internet research*. 2012; 14(1):e13. [PubMed: 22278284]
25. Corrigan M. A computerized, self-administered questionnaire to evaluate posttraumatic stress among firefighters after the World Trade Center collapse. *American journal of public health* (1971). 2009; 99(Suppl 3)(s3):S702–S709.
26. Moore HK. A pilot study of an electronic, adolescent version of the quick inventory of depressive symptomatology. *The journal of clinical psychiatry*. 2007; 68(9):1436–1440. [PubMed: 17915985]
27. Moore HK. An examination of 26,168 Hamilton Depression Rating Scale scores administered via interactive voice response across 17 randomized clinical trials. *Journal of clinical psychopharmacology*. 2006; 26(3):321–324. [PubMed: 16702899]
28. Mundt JC, et al. Feasibility and validation of a computer-automated Columbia- Suicide severity rating scale using interactive voice response technology. *Journal of Psychiatric Research*. 2010; 44(16):1224–1228. [PubMed: 20553851]
29. Morrison-Beedy D. Accuracy of audio computer-assisted self-interviewing (ACASI) and self-administered questionnaires for the assessment of sexual behavior. *AIDS and behavior*. 2006; 10(5):541–552. [PubMed: 16721506]
30. Johnson TP. An investigation of the effects of social desirability on the validity of self-reports of cancer screening behaviors. *Medical Care*. 2005; 43(6):565–573. [PubMed: 15908851]
31. Screening for depression in adults: U.S. preventive services task force recommendation statement. *Annals of internal medicine*. 2009; 151(11):784–792. [PubMed: 19949144]
32. Pignatiello A. Child and youth telepsychiatry in rural and remote primary care. *Child and adolescent psychiatric clinics of North America*. 2011; 20(1):13–28. [PubMed: 21092909]
33. Singh SP. Accuracy of telepsychiatric assessment of new routine outpatient referrals. *BMC Psychiatry*. 2007; 7(1):55. [PubMed: 17919329]
34. Saeed SA. Use of telepsychiatry to improve care for people with mental illness in rural North Carolina. *North Carolina Medical Journal* (Durham, N.C.). 2011; 72(3):219–222.
35. Paing WW. Telemedicine in children and adolescents. *Current Psychiatry Reports*. 2009; 11(2): 114–119. [PubMed: 19302764]
36. Cella D. The Patient-Reported Outcomes Measurement Information System (PROMIS): progress of an NIH Roadmap cooperative group during its first two years. *Medical Care*. 2007; 45 suppl 1(5):S3–S11. [PubMed: 17443116]
37. Riley WT. Application of the National Institutes of Health Patient-reported Outcome Measurement Information System (PROMIS) to mental health research. *The journal of mental health policy and economics*. 2011; 14(4):201–208. [PubMed: 22345362]

38. Lin C-C. Web-based tools can be used reliably to detect patients with major depressive disorder and subsyndromal depressive symptoms. *BMC Psychiatry*. 2007; 7(1):12. [PubMed: 17425774]
39. Gibbons RD. Using computerized adaptive testing to reduce the burden of mental health assessment. *Psychiatric Services (Washington, D.C.)*. 2008; 59(4):361–368.
40. Becker J. Functioning and validity of a Computerized Adaptive Test to measure anxiety (A-CAT). *Depression and Anxiety*. 2008; 25(12):E182–E194. [PubMed: 18979458]
41. Garcia-Lizana F. What about telepsychiatry? A systematic review. *Primary Care Companion to the Journal of Clinical Psychiatry*. 2010; 12(2)
42. Poston JM, Hanson WE. Meta-analysis of psychological assessment as a therapeutic intervention. *Psychological Assessment*. 2010; 22(2):203–212. [PubMed: 20528048]
43. Lilienfeld S, Garb H, Wood S. Unresolved questions concerning the effectiveness of psychological assessment as a therapeutic intervention: comment on Poston and Hanson (2010). *Psychological Assessment*. 2011; 23(4):1047–1055. discussion 1056–62. [PubMed: 22122676]
44. Kelley S, Bickman L. Beyond outcomes monitoring: measurement feedback systems in child and adolescent clinical practice. *Current Opinion in Psychiatry*. 2009; 22(4):363–368. [PubMed: 19417669]
45. Robertson L, et al. Using the Internet to Enhance the Treatment of Depression. *Australasian Psychiatry*. 2006; 14(4):413–417. [PubMed: 17116083]
46. Riva G. Virtual reality in psychotherapy: review. *Cyberpsychology & Behavior*. 2005; 8(3):220–230. discussion 231–40. [PubMed: 15971972]
47. Parsons T, et al. A virtual human agent for assessing bias in novice therapists. *Studies in Health Technology and Informatics*. 2009; 142:253–258. [PubMed: 19377161]
48. Fleming M, et al. Virtual Reality Skills Training for Health Care Professionals in Alcohol Screening and Brief Intervention. *The Journal of the American Board of Family Medicine*. 2009; 22(4):387–398.
49. Haberer J, et al. Real-time adherence monitoring for HIV antiretroviral therapy. *AIDS and Behavior*. 2010; 14(6):1340–1346. [PubMed: 20809380]
50. Haberer J, et al. Challenges in Using Mobile Phones for Collection of Antiretroviral Therapy Adherence Data in a Resource-Limited Setting. *AIDS and behavior*. 2010; 14(6):1294–1301. [PubMed: 20532605]
51. Siedner M, et al. High acceptability for cell phone text messages to improve communication of laboratory results with HIV-infected patients in rural Uganda: a crosssectional survey study. *BMC Medical Informatics and Decision Making*. 2012; 12(1):56. [PubMed: 22720901]
52. Granholm E, et al. Mobile Assessment and Treatment for Schizophrenia (MATS): A Pilot Trial of An Interactive Text-Messaging Intervention for Medication Adherence, Socialization, and Auditory Hallucinations. *Schizophrenia Bulletin*. 2012; 38(3):414–425. [PubMed: 22080492]
53. Farrell SP, Mahone IH, Guilbaud P. Web technology for persons with serious mental illness. *Archives of Psychiatric Nursing*. 2004; 18(4):121–125. [PubMed: 15305277]
54. Burns M, et al. Harnessing context sensing to develop a mobile intervention for depression. *Journal of Medical Internet Research*. 2011; 13(3):e55. [PubMed: 21840837]
55. Trivedi MH, et al. Maximizing the Adequacy of Medication Treatment in Controlled Trials and Clinical Practice: STAR*D Measurement-Based Care. *Neuropsychopharmacology*. 2007; 32(12):2479–2489. [PubMed: 17406651]
56. Gaynes BN, et al. Primary versus specialty care outcomes for depressed outpatients managed with measurement-based care: results from STAR*D. *J Gen Intern Med*. 2008; 23(5):551–560. [PubMed: 18247097]
57. Yeung AS, et al. Clinical outcomes in measurement-based treatment (COMET): A trial of depression monitoring and feedback to primary care clinicians. *Depression and Anxiety*. 2012 n/a-n/a.
58. Brown G, et al. Pushing the quality envelope: a new outcomes management system. *Psychiatric Services (Washington, D.C.)*. 2001; 52(7):925–934.
59. Seidman E, et al. A Framework for Measurement Feedback to Improve Decision-Making in Mental Health. *Administration and Policy in Mental Health and Mental Health Services Research*. 2010; 37(1):128–131. [PubMed: 20041342]

60. Gill J, et al. Electronic clinical decision support for management of depression in primary care: a prospective cohort study. *Primary Care Companion to CNS Disorders*. 2012; 14(1)
61. Valuck RJ, et al. Enhancing Electronic Health Record Measurement of Depression Severity and Suicide Ideation: A Distributed Ambulatory Research in Therapeutics Network (DARTNet) Study. *The Journal of the American Board of Family Medicine*. 25(5):582–593.
62. Gill JM, et al. Using Electronic Health Record-Based Tools To Screen for Bipolar Disorder in Primary Care Patients With Depression. *The Journal of the American Board of Family Medicine*. 2012; 25(3):283–290.
63. Berke EM, et al. Objective Measurement of Sociability and Activity: Mobile Sensing in the Community. *The Annals of Family Medicine*. 2011; 9(4):344–350.
64. Glasgow RE, et al. Patient-Reported Measures Of Psychosocial Issues And Health Behavior Should Be Added To Electronic Health Records. *Health Affairs*. 2012; 31(3):497–504. [PubMed: 22392660]
65. Estabrooks PA, et al. Harmonized patient-reported data elements in the electronic health record: supporting meaningful use by primary care action on health behaviors and key psychosocial factors. *Journal of the American Medical Informatics Association*. 2012; 19(4):575–582. [PubMed: 22511015]
66. Chorpita B, et al. Driving with Roadmaps and Dashboards: Using Information Resources to Structure the Decision Models in Service Organizations. *Administration and Policy in Mental Health and Mental Health Services Research*. 2008; 35(1):114–123. [PubMed: 17987376]
67. Seidman E, et al. A framework for measurement feedback to improve decisionmaking in mental health. *Administration and Policy in Mental Health and Mental Health Services Research*. 2011; 37(1–2):128–131. [PubMed: 20041342]
68. Parsons A, et al. Validity of electronic health record-derived quality measurement for performance monitoring. *Journal of the American Medical Informatics Association*. 2012; 19(4):604–609. [PubMed: 22249967]
69. Persell S, et al. Assessing the validity of national quality measures for coronary artery disease using an electronic health record. *Archives of Internal Medicine*. 2006; 166(20):2272–2277. [PubMed: 17101947]
70. Unutzer J, et al. Quality Improvement With Pay-for-Performance Incentives in Integrated Behavioral Health Care. *American Journal of Public Health*. 2012; 102(6):e41–e45. [PubMed: 22515849]
71. Jensen P, Foster M. Closing the Research to Practice Gap in Children's Mental Health: Structures, Solutions, and Strategies. *Administration and Policy in Mental Health and Mental Health Services Research*. 2010; 37(1):111–119. [PubMed: 20217208]
72. Intille SS. Ubiquitous computing technology for just-in-time motivation of behavior change. *Studies in Health Technology and Informatics*. 2004; 107:1434–1437. [PubMed: 15361052]
73. Adusumilli SR, et al. The New York City eClinician Project: using Personal Digital Assistants and wireless internet access to support emergency preparedness and enhance clinical care in community health centers. *AMIA ... Annual Symposium proceedings*. 2006:839. [PubMed: 17238459]
74. Carle AC. Advancing PROMIS's methodology: results of the Third Patient- Reported Outcomes Measurement Information System (PROMIS) Psychometric Summit. *Expert Review of Pharmacoeconomics & Outcomes Research*. 2011; 11(6):677–684. [PubMed: 22098283]
75. Fitzmaurice D, et al. Oral anticoagulation management in primary care with the use of computerized decision support and near-patient testing: A randomized, controlled trial. *Archives of Internal Medicine*. 2000; 160(15):2343–2348. [PubMed: 10927732]
76. Rasmussen R, et al. Effects of computer-assisted oral anticoagulant therapy. *Thrombosis Journal*. 2012; 10(1):17. [PubMed: 22935243]
77. Bazemore A, et al. Establishing a Baseline: Health Information Technology Adoption among Family Medicine Diplomates. *The Journal of the American Board of Family Medicine*. 2011; 24(2):132.
78. Mandl KD, Kohane IS. Escaping the EHR Trap – The Future of Health IT. *New England Journal of Medicine*. 2012; 366(24):2240–2242. [PubMed: 22693995]

79. Nutting PA, et al. Initial lessons from the first national demonstration project on practice transformation to a patient-centered medical home. *Annals of Family Medicine*. 2009; 7(3):254–260. [PubMed: 19433844]
80. Salomon RM, et al. Openness of patients' reporting with use of electronic records: psychiatric clinicians' views. *Journal of the American Medical Informatics Association*. 2010; 17(1):54–60. [PubMed: 20064802]