Published in final edited form as:

J Mol Biol. 2013 October 9; 425(19): 3601–3608. doi:10.1016/j.jmb.2013.01.023.

MicroRNAs and the genetic network in aging

Sachi Inukai and Frank Slack¹

Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT, 06511, USA

Abstract

MicroRNAs (miRNAs) comprise a class of small RNAs important for the post-transcriptional regulation of numerous biological processes. Their combinatorial mode of function, in which an individual miRNA can target many genes and multiple miRNAs share targets, makes them especially suited for regulating processes and pathways at the “network” level. In particular, miRNAs have recently been implicated in aging which is a complex process known to involve multiple pathways. Findings from genome-wide miRNA expression profiling studies highlight three themes in miRNA function during aging: many miRNAs are differentially expressed, many such miRNAs target known aging-associated pathways, and there are global trends in miRNA expression change over time. In addition, several miRNAs have emerged as potentially coordinating multiple pathways during aging. Elucidating the underlying network structure of genes and miRNAs involved in aging processes promises to advance our understanding of not only aging and associated pathogenesis but also of how miRNAs can connect disparate pathways.

Keywords

lifespan; insulin signaling; small non-coding RNAs; *Caenorhabditis elegans*

Introduction

MicroRNAs (miRNAs) are endogenous small RNAs that regulate gene expression largely at the post-transcriptional level and have gained increasing attention in the last decade. Since their discovery in the nematode *Caenorhabditis elegans*, miRNAs have been identified in a variety of animals and plants, and their reported functions span many pathways and processes including timing of developmental decisions, cell differentiation, oncogenesis, and organismal aging.^{1, 2} The human genome codes for over a thousand miRNAs (miRBase release 19),^{3–6} and computational predictions suggest that miRNAs may regulate nearly one-third of all human genes,⁷ making miRNAs one of the largest “families” of regulatory molecules. Understanding how miRNAs fit in to and interact with the overall genetic landscape is a major goal in miRNA research.

The role of miRNAs in aging has only recently begun to be investigated. Aging was long considered a result of random, accumulating molecular deterioration, but research in the last three decades since the isolation of the first genetic lifespan mutants in *C. elegans*^{8–10} has shown a large genetic component influencing organismal aging rates. We now know that

© 2012 Elsevier Ltd. All rights reserved.

¹Corresponding author: Telephone: +1-203-432-3292; Fax: +1-203-432-6161; frank.slack@yale.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

aging is shaped by a network consisting of multiple partially overlapping genetic pathways, many of which center on insulin signaling (see refs. 11 and 12 for review). The involvement of miRNAs in aging therefore is particularly interesting in light of two key features of miRNA-mediated regulation: a single miRNA can target many genes, and multiple miRNAs can target the same mRNA.¹³ This mode of function suggests that miRNAs act as “network-level” regulators modulating complex phenotypes by integrating multiple inputs and outputs. Thus, in the context of aging research, miRNAs may help us understand how diverse genetic pathways are connected and regulated.

This review will present evidence for several conserved roles of miRNAs in determining organismal aging rates. We will highlight some themes learned from genome-wide miRNA expression studies during aging. Other reviews have discussed specific miRNAs and how they have been associated with particular aging-related biological processes.^{14, 15} Here, we will instead focus on how miRNAs connect these processes as potential upstream regulatory factors.

Genetic pathways in aging and *lin-4*

The genes and pathways that are involved in aging rate and lifespan determination have been studied extensively. Basic research conducted in model organisms, particularly *C. elegans*, has identified many single-gene mutants demonstrating lifespan phenotypes, and genome-wide expression studies have revealed distinct expression profiles associated with aging.^{16–18} We know from these studies that specific biological processes are affected with advancing age and that a network of multiple, yet specific, genetic pathways are involved (for reviews, see refs. 11 and 12).

The best characterized aging-associated pathway is insulin/insulin-like growth factor 1 (IGF-1) signaling (IIS). The core components of this pathway and their effects on lifespan are highly conserved: from nematodes to mice, reduced insulin signaling results in extended lifespan.¹⁹ For example, in *C. elegans*, a mutation in the *daf-2* insulin receptor ortholog more than doubled the two-week lifespan of wildtype worms.¹⁰ Inactivation of the IGF-1 receptor gene as well as the insulin receptor substrate gene *IRS1*, which acts downstream of the IGF-1 receptor, in female mice similarly extended lifespan significantly.^{20, 21} Furthermore, longevity in humans has been associated with variations in IIS pathway genes, particularly FOXO3A transcription factor.^{22–26}

IIS plays a central role in orchestrating many cellular processes involved in aging. (We refer readers to ref. 19 for a review on IIS.) At the core of IIS is a phosphorylation cascade, triggered by the binding of insulin or an insulin-like molecule (e.g. IGF-1) to the insulin receptor, culminating in the phosphorylation of forkhead (FOXO) transcription factor which prevents it from localizing to the nucleus. Multiple internal and external signals which influence aging rate converge on the IIS pathway, often on FOXO.²⁷ Some examples of lifespan-determining signals that require DAF-16/FOXO activity include reproductive status²⁸ and environmental stress via *sir-2.1* and Jun kinase signaling pathway.^{29, 30} FOXO in turn regulates the transcription of many downstream effector genes that affect various cellular processes including metabolism, apoptosis, and stress resistance.^{31–33} It is important to note that IIS-independent pathways that affect lifespan have been reported; these include DNA damage checkpoint signaling, developmental and behavioral timing pathway, and mitochondrial genes.^{34–36} We do not know whether or how these pathways are connected to influence aging.

One of the earliest reports of miRNAs associated with lifespan determination also invoked insulin signaling. Our laboratory reported that a mutation in *lin-4*, a heterochronic miRNA studied extensively in the regulation of developmental timing in *C. elegans*,³⁷ shortened

adult lifespan.² This phenotype was dependent on the expression of a *lin-4* target, *lin-14* transcription factor. In turn, the effect of LIN-14 on lifespan was mediated by the insulin signaling pathway, requiring the functions of DAF-2 and DAF-16 FOXO homolog, and by the heat-shock transcription factor HSF-1.² HSF-1 acts downstream of IIS and couples signaling with stress response; *hsf-1* mutants demonstrated shortened lifespan.^{38, 39} This study not only demonstrated that a miRNA can influence lifespan but does so through pathways with known importance in aging. Further, these results suggest that *lin-4* links upstream IIS with downstream heat stress response, possibly to better coordinate the outputs of both pathways.

Genome-wide studies of miRNAs in aging

In an effort to identify other miRNAs affected by aging, genome-wide miRNA expression has been profiled extensively in *C. elegans* (whole animal).^{40–42} Profiling has also been conducted in a variety of tissues and cell types in additional systems. These include the fruit fly *Drosophila melanogaster* brain;⁴³ mouse brain,^{44–46} liver,^{47, 48} and cardiac tissue;⁴⁹ and human skeletal muscle⁵⁰ and blood cells.^{51–53} These studies found that many miRNAs are differentially expressed between old and young subjects. We henceforth refer to these differentially expressed miRNAs as “aging-associated” miRNAs, summarized in Table 1.

One caveat of studies in specific tissue/cell types is the extent to which they represent aging of the whole organism. It is known that tissues age at different rates.⁵⁴ While it can be difficult to extrapolate findings from specific tissues to aging of the whole organism, there are, nonetheless, valuable lessons to gather about both tissue-specific and organismal aging.

Several themes arise from these genome-wide studies. First is that multiple studies have identified common, and often evolutionarily conserved, aging-associated miRNAs. These miRNAs may represent particularly important regulators during aging. Second, aging-associated miRNAs often target pathways and processes that influence aging (Figure 1). Indeed, some miRNAs target multiple aging-related pathways suggesting that miRNAs may have a role in coordinating the outputs of those pathways. Finally, there are clear global trends in miRNA expression change during aging. This suggests that a common process, such as miRNA transcription or biogenesis, may be altered with aging, changing large numbers of miRNA levels simultaneously.

Identification of common aging-associated miRNAs

Common aging-associated miRNAs have been found between published reports. Three separate studies of miRNA expression change with age have been performed in *C. elegans*.^{40–42} While not all reported aging-associated miRNAs overlap, all three studies saw changes in the expression levels of *let-7*, miR-34, miR-35, miR-38, miR-43, miR-70, miR-71, and miR-239. It is plausible that these miRNAs function cooperatively to regulate processes during aging. However, it is important to point out that though alterations were consistent, the directions of change, surprisingly, were not. Notably, some of these miRNAs are conserved in human (*let-7*, miR-34, and miR-43)⁵⁵ and thus may be important for human aging and lifespan determination.

In mouse, Wang and colleagues found commonly upregulated miRNAs in two different aging tissues — the liver and the brain.^{44, 48} They suggested that these miRNAs (miR-30d, miR-34a, miR-468, miR-669b, and miR-709) may contribute to the general aging process, whereas miRNAs found only in one tissue may be implicated in tissue-specific aging.⁴⁴ However, of these core aging miRNAs, only one (miR-669b) was found to be upregulated in old versus young brains of the long-living Ames dwarf mice.⁴⁶ None were found in a study of aging heart tissue⁴⁹ or a different study on the aging brain.⁴⁵ The different experimental

methods and statistical thresholds applied in each study may partly explain why they did not identify overlapping sets of aging-associated miRNAs.

One highly conserved miRNA was identified in multiple aging studies across various organisms and tissues. miR-34 belongs to one of only 19 miRNA families conserved at a sequence level from nematode to mammals.⁵⁶ Moreover, the functions of miR-34 in DNA damage response, senescence, and cell death are highly conserved,^{57–59} and as DNA damage signaling is a known pathway affecting lifespan,³⁴ this suggests that conserved regulation by miR-34 may also exist for aging-associated processes. Its reported expression changes and effects on lifespan, however, have not been consistent. Increased expression with age was reported in *D. melanogaster* brain, *C. elegans* whole animal, rat liver, and several mouse tissues,^{40–42, 60, 61} while decreased expression was observed in the mouse brain.^{45, 62} In *C. elegans*, *mir-34* mutants demonstrated extended lifespan in one study⁶³ (but not in another)⁴¹, but in *D. melanogaster*, overexpression of miR-34 extended lifespan and protected flies from neurodegenerative disease⁴³. Nonetheless, studies point to the importance of miR-34 in lifespan determination across phyla, and further research may unveil conserved regulation mediated by miR-34 that is crucial to normal aging.

Aging-associated pathways are targeted by aging-associated miRNAs

Many aging-associated miRNAs seem to interact with genes and pathways that are relevant to aging. It is worth noting here that the functions of most miRNAs are not known due to the fact that the vast majority of these were identified through cloning and computational prediction alone.^{56, 64–66} The functions of some age-associated miRNAs have been inferred from classic genetic characterizations of individual miRNA mutants while others have been predicted solely from mRNA target predictions coupled with bioinformatics analyses of those targets. Experimentally validating miRNA functions and their putative associations with specific pathways remains to be a challenge.

Many *C. elegans* strains with mutations for aging-associated miRNAs have been characterized in aging,^{41, 67} made possible by a large collection of miRNA loss-of-function mutants.⁶⁸ When characterized during development, most of these strains did not demonstrate gross abnormalities;⁶⁸ nevertheless, mutations for some aging-associated miRNAs have resulted in lifespan phenotypes^{41, 67}: *mir-71* and *mir-246* mutants demonstrated shortened lifespan, and *mir-239* mutants showed extended lifespan. Overexpressing these miRNAs resulted in the respective opposite phenotypes, suggesting specific involvement in aging. Further, the lifespan phenotypes of *mir-71* and *mir-239* mutants were dependent on IIS⁴¹ and FOXO.⁶⁷ miR-71 interacts with at least two other aging-associated pathways: its activity was necessary for the long lifespan of both the DNA damage response gene *cdc-25.1* knockdown animals and of germline-ablated animals.^{41, 67} These findings suggest that miR-71 may act as a link between IIS, the DNA damage checkpoint pathway, and germline signaling. Particularly, the connection with the DNA damage checkpoint pathway is interesting because it functions independently of IIS in lifespan determination.³⁴

The miRNA *let-7* may also link multiple aging-associated pathways. *let-7* is differentially expressed in many aging studies.^{40–42, 45, 48, 50, 53, 69, 70} Nuclear hormone receptor *daf-12* is a known *let-7* target,⁷¹ and mutations in *daf-12* affect longevity.⁷² Further, DAF-12 activity is necessary for the germline signal that determines lifespan.²⁸ *let-7* has also been associated with DNA damage checkpoint genes and mitochondrial respiration genes;^{42, 50} though direct genetic interactions have not been validated, *let-7* may directly or indirectly coordinate these pathways. Furthermore, *let-7* and *daf-12* have been implicated in feedback signaling that is important for the specification of developmental fate in response to environmental cues in *C.*

elegans.^{73, 74} While there currently is no evidence for this interaction during aging, if it exists, it may act to reinforce a specific lifespan phenotype. Unfortunately, most *C. elegans* *let-7* alleles cause developmental defects making aging studies difficult to interpret.

A recent study described a role for *let-7* in the age-related decline of the *Drosophila* testis stem cell niche. Jones and colleagues found that in old *Drosophila* testis, *let-7* targets *Imp* (IGF-II messenger RNA binding protein) which has a role in germline stem cell maintenance by stabilizing the self-renewal factor, *upd*, from being degraded by a short interfering RNA.⁷⁵ Other miRNAs that function to link multiple pathways may also have dual functions as genetic switches during aging.

Most effects of individual miRNAs on target expression levels are thought to be subtle.⁷⁶ Thus, in addition to examining individual miRNAs and their putative targets, it is also informative to investigate the processes that aging-associated miRNAs target as a collective. In a study in the aging mouse brain, Wang and colleagues found that 27 out of 70 upregulated miRNAs target mitochondrial genes.⁴⁴ They showed by mass spectrometry that these genes are indeed affected during aging and correlated with the expression changes of the miRNAs that putatively target them. Driscoll and colleagues found that 31 out of 50 age-associated miRNAs target gerontogenes which they identified from the literature; furthermore, many aging-associated miRNAs were predicted to target genes in the IIS pathway.⁴⁰ Our lab has shown that nearly one-sixth of genes predicted to be targeted by one or more age-associated miRNAs are necessary for normal lifespan in *C. elegans*.⁴²

Systems approaches such as pathway and Gene Ontology term analyses have also identified aging-associated processes as putative targets of aging-associated miRNAs. These include cell cycle control, cell growth, cell proliferation, mitochondrial respiration, and IIS.^{42, 45, 48, 50, 52, 70} Particularly, our lab has shown that each of the core genes in the IIS pathway may potentially be targeted by multiple aging-associated miRNAs in the mouse brain,⁴⁵ again suggesting the centrality of IIS in aging.

Global changes in miRNA expression during aging

Global trends in miRNA expression with increased age have been noted in multiple studies, with many specifically reporting decreases in miRNA abundance.^{40, 41, 43, 44, 47, 48, 51, 53, 70} There were no tissue-specific differences in trends. This observation suggests that in addition to miRNA-specific regulation, there may be more general mechanisms changing the levels of many or most miRNAs over time, such as age-related alterations in miRNA transcription and steps in miRNA biogenesis.

A few miRNA biogenesis genes have been investigated in the context of aging. Bonini and colleagues characterized the lifespan phenotype of hypomorphic *loquacious* (*loqs*) mutants in *D. melanogaster*.⁴³ *Loqs* is a homolog of the human TAR RNA-binding protein (TRBP) and associates with RNase-III enzyme Dicer to facilitate efficient pre-miRNA processing.^{77–79} The *loqs* hypomorph resulted in a significant decrease in *D. melanogaster* lifespan,⁴³ suggesting that miRNAs are required for normal lifespan. How *loqs* expression is affected during natural aging has not been investigated. If its expression is indeed diminished with age, then this may account for the decreased expression of miRNAs during aging in the fly brain.⁴³

Two studies investigated the requirement for Argonaute proteins in aging. Argonautes are key components of the effector complex for miRNA-mediated gene regulation. Our lab studied the effects of adult-specific RNAi-mediated knockdown of *alg-1* (Argonaute-like gene 1, the miRNA-specific Argonaute in *C. elegans*), which resulted in a significant reduction in lifespan compared to control, providing evidence that miRNAs are required for

normal adult lifespan.⁴² Zhang and colleagues examined expression changes of Argonaute (Ago) 1 and 2 in mouse cardiac ventricular tissue with increased age.⁴⁹ The authors found that *Ago1* and *Ago2* mRNA expression increased through mid-adulthood, at which point it peaked, and then gradually decreased with advancing age. As Argonautes can stabilize mature miRNAs and thereby contribute to increased miRNA abundance,⁸⁰ their diminished expression in late adulthood may result in a global reduction in miRNA expression, thereby affecting functions of aging-associated pathways and influencing lifespan.

We know little about how other miRNA biogenesis pathway components are affected during aging. Although many studies have identified specific miRNAs important for aging, perturbations in miRNA biogenesis may too impact aging-associated pathogenesis and decline. Indeed, miRNA biogenesis is frequently deregulated in cancer.^{81–83} Further research is needed to elucidate how aging impacts the upstream regulators of miRNA expression.

Conclusions and future perspectives

Aging is a complex process involving multiple genetic pathways. An outstanding goal of the field is to understand how these pathways interact and result in the final output of aging. The IIS pathway certainly has a central role in coordinating different external and internal signals.¹⁹ However, there are also pathways that affect aging independently of IIS. The implication of miRNAs in regulating processes involved in aging sheds light on a possible mechanism linking these pathways.

Current research suggests that a large number of miRNAs are differentially expressed during aging. Genetic interactions and computational target predictions suggest that many miRNA may influence aging and lifespan by targeting aging-associated pathways. Furthermore, some miRNAs may regulate multiple pathways, suggesting a role in coordinating the functions of disparate pathways during aging. We have only begun to investigate the depth and breadth of miRNA-mediated regulation during aging. There are still many questions to be addressed such as how miRNAs themselves are regulated during aging and how miRNAs are connected to each other.

Investigating the network structure of pathways in aging will allow us to understand how perturbations in a specific part of the network can affect the overall output (i.e. aging), and it promises to advance our knowledge of aging-related pathogenesis. Some miRNAs have been implicated in genetic feedback loops in the context of development⁸⁴ and many more have been predicted.⁸⁵ Further research is needed to identify the exact wiring of the aging network to better understand the systems-level role of miRNAs.

Acknowledgments

The authors would like to thank Zachary Pincus and Chantip Metheetrairut for critical reading of this manuscript. This work was supported by NIH grant (AG033921).

Abbreviation

IIS   Insulin/insulin-like growth factor 1 signaling

References

1. Stefani G, Slack FJ. Small non-coding RNAs in animal development. *Nat Rev Mol Cell Biol.* 2008; 9:219–230.10.1038/nrm2347 [PubMed: 18270516]

2. Boehm M, Slack F. A developmental timing microRNA and its target regulate life span in *C. elegans*. *Science*. 2005; 310:1954–1957.10.1126/science.1115596 [PubMed: 16373574]
3. Kozomara A, Griffiths-Jones S. miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Res*. 2011; 39:D152–7.10.1093/nar/gkq1027 [PubMed: 21037258]
4. Griffiths-Jones S. The microRNA Registry. *Nucleic Acids Res*. 2004; 32:D109–11.10.1093/nar/gkh023 [PubMed: 14681370]
5. Griffiths-Jones S, Grocock RJ, van Dongen S, Bateman A, Enright AJ. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res*. 2006; 34:D140–4.10.1093/nar/gkj112 [PubMed: 16381832]
6. Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ. miRBase: tools for microRNA genomics. *Nucleic Acids Res*. 2008; 36:D154–8.10.1093/nar/gkm952 [PubMed: 17991681]
7. Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell*. 2005; 120:15–20.10.1016/j.cell.2004.12.035 [PubMed: 15652477]
8. Klass MR. A method for the isolation of longevity mutants in the nematode *Caenorhabditis elegans* and initial results. *Mech Ageing Dev*. 1983; 22:279–286. [PubMed: 6632998]
9. Friedman DB, Johnson TE. A mutation in the *age-1* gene in *Caenorhabditis elegans* lengthens life and reduces hermaphrodite fertility. *Genetics*. 1988; 118:75–86. [PubMed: 8608934]
10. Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A *C. elegans* mutant that lives twice as long as wild type. *Nature*. 1993; 366:461–464.10.1038/366461a0 [PubMed: 8247153]
11. Kenyon C. The plasticity of aging: insights from long-lived mutants. *Cell*. 2005; 120:449–460.10.1016/j.cell.2005.02.002 [PubMed: 15734678]
12. Antebi A. Genetics of aging in *Caenorhabditis elegans*. *PLoS Genet*. 2007; 3:1565–1571.10.1371/journal.pgen.0030129 [PubMed: 17907808]
13. Krek A, Grun D, Poy MN, Wolf R, Rosenberg L, Epstein EJ, MacMenamin P, da Piedade I, Gunsalus KC, Stoffel M, Rajewsky N. Combinatorial microRNA target predictions. *Nat Genet*. 2005; 37:495–500.10.1038/ng1536 [PubMed: 15806104]
14. Smith-Vikos T, Slack FJ. MicroRNAs and their roles in aging. *J Cell Sci*. 2012; 125:7–17.10.1242/jcs.099200;10.1242/jcs.099200 [PubMed: 22294612]
15. Kato M, Slack FJ. Ageing and the small, non-coding RNA world. *Ageing Res Rev*. 2012;10.1016/j.arr.2012.03.012
16. Lund J, Tedesco P, Duke K, Wang J, Kim SK, Johnson TE. Transcriptional profile of aging in *C. elegans*. *Curr Biol*. 2002; 12:1566–1573. [PubMed: 12372248]
17. McCarroll SA, Murphy CT, Zou S, Pletcher SD, Chin CS, Jan YN, Kenyon C, Bargmann CI, Li H. Comparing genomic expression patterns across species identifies shared transcriptional profile in aging. *Nat Genet*. 2004; 36:197–204.10.1038/ng1291 [PubMed: 14730301]
18. Zahn JM, Sonu R, Vogel H, Crane E, Mazan-Mamczarz K, Rabkin R, Davis RW, Becker KG, Owen AB, Kim SK. Transcriptional profiling of aging in human muscle reveals a common aging signature. *PLoS Genet*. 2006; 2:e115.10.1371/journal.pgen.0020115.eor [PubMed: 16789832]
19. Tatar M, Bartke A, Antebi A. The endocrine regulation of aging by insulin-like signals. *Science*. 2003; 299:1346–1351.10.1126/science.1081447 [PubMed: 12610294]
20. Holzenberger M, Dupont J, Ducos B, Leneuve P, Geloën A, Even PC, Cervera P, Le Bouc Y. IGF-1 receptor regulates lifespan and resistance to oxidative stress in mice. *Nature*. 2003; 421:182–187.10.1038/nature01298 [PubMed: 12483226]
21. Selman C, Lingard S, Choudhury AI, Batterham RL, Claret M, Clements M, Ramadani F, Okkenhaug K, Schuster E, Blanc E, Piper MD, Al-Qassab H, Speakman JR, Carmignac D, Robinson IC, Thornton JM, Gems D, Partridge L, Withers DJ. Evidence for lifespan extension and delayed age-related biomarkers in insulin receptor substrate 1 null mice. *FASEB J*. 2008; 22:807–818.10.1096/fj.07-9261com [PubMed: 17928362]
22. Pawlikowska L, Hu D, Huntsman S, Sung A, Chu C, Chen J, Joyner AH, Schork NJ, Hsueh WC, Reiner AP, Psaty BM, Atzmon G, Barzilai N, Cummings SR, Browner WS, Kwok PY, Ziv E. Study of Osteoporotic Fractures. Association of common genetic variation in the insulin/IGF1 signaling pathway with human longevity. *Aging Cell*. 2009; 8:460–472.10.1111/j.1474-9726.2009.00493.x [PubMed: 19489743]

23. Willcox BJ, Donlon TA, He Q, Chen R, Grove JS, Yano K, Masaki KH, Willcox DC, Rodriguez B, Curb JD. FOXO3A genotype is strongly associated with human longevity. *Proc Natl Acad Sci U S A*. 2008; 105:13987–13992.10.1073/pnas.0801030105 [PubMed: 18765803]
24. Flachsbarth F, Caliebe A, Kleindorp R, Blanche H, von Eller-Eberstein H, Nikolaus S, Schreiber S, Nebel A. Association of FOXO3A variation with human longevity confirmed in German centenarians. *Proc Natl Acad Sci U S A*. 2009; 106:2700–2705.10.1073/pnas.0809594106 [PubMed: 19196970]
25. Anselmi CV, Malovini A, Roncarati R, Novelli V, Villa F, Condorelli G, Bellazzi R, Puca AA. Association of the FOXO3A locus with extreme longevity in a southern Italian centenarian study. *Rejuvenation Res*. 2009; 12:95–104.10.1089/rej.2008.0827 [PubMed: 19415983]
26. Li Y, Wang WJ, Cao H, Lu J, Wu C, Hu FY, Guo J, Zhao L, Yang F, Zhang YX, Li W, Zheng GY, Cui H, Chen X, Zhu Z, He H, Dong B, Mo X, Zeng Y, Tian XL. Genetic association of FOXO1A and FOXO3A with longevity trait in Han Chinese populations. *Hum Mol Genet*. 2009; 18:4897–4904.10.1093/hmg/ddp459 [PubMed: 19793722]
27. Henderson ST, Johnson TE. *daf-16* integrates developmental and environmental inputs to mediate aging in the nematode *Caenorhabditis elegans*. *Curr Biol*. 2001; 11:1975–1980. [PubMed: 11747825]
28. Hsin H, Kenyon C. Signals from the reproductive system regulate the lifespan of *C. elegans*. *Nature*. 1999; 399:362–366.10.1038/20694 [PubMed: 10360574]
29. Oh SW, Mukhopadhyay A, Svrzikapa N, Jiang F, Davis RJ, Tissenbaum HA. JNK regulates lifespan in *Caenorhabditis elegans* by modulating nuclear translocation of forkhead transcription factor/DAF-16. *Proc Natl Acad Sci U S A*. 2005; 102:4494–4499.10.1073/pnas.0500749102 [PubMed: 15767565]
30. Berdichevsky A, Viswanathan M, Horvitz HR, Guarente L. *C. elegans* SIR-2.1 interacts with 14-3-3 proteins to activate DAF-16 and extend life span. *Cell*. 2006; 125:1165–1177.10.1016/j.cell.2006.04.036 [PubMed: 16777605]
31. Murphy CT, McCarroll SA, Bargmann CI, Fraser A, Kamath RS, Ahringer J, Li H, Kenyon C. Genes that act downstream of DAF-16 to influence the lifespan of *Caenorhabditis elegans*. *Nature*. 2003; 424:277–283.10.1038/nature01789 [PubMed: 12845331]
32. Lee SS, Kennedy S, Tolonen AC, Ruvkun G. DAF-16 target genes that control *C. elegans* life-span and metabolism. *Science*. 2003; 300:644–647.10.1126/science.1083614 [PubMed: 12690206]
33. Oh SW, Mukhopadhyay A, Dixit BL, Raha T, Green MR, Tissenbaum HA. Identification of direct DAF-16 targets controlling longevity, metabolism and diapause by chromatin immunoprecipitation. *Nat Genet*. 2006; 38:251–257.10.1038/ng1723 [PubMed: 16380712]
34. Olsen A, Vantipalli MC, Lithgow GJ. Checkpoint proteins control survival of the postmitotic cells in *Caenorhabditis elegans*. *Science*. 2006; 312:1381–1385.10.1126/science.1124981 [PubMed: 16741121]
35. Lakowski B, Hekimi S. Determination of life-span in *Caenorhabditis elegans* by four clock genes. *Science*. 1996; 272:1010–1013. [PubMed: 8638122]
36. Dillin A, Hsu AL, Arantes-Oliveira N, Lehrer-Graiwer J, Hsin H, Fraser AG, Kamath RS, Ahringer J, Kenyon C. Rates of behavior and aging specified by mitochondrial function during development. *Science*. 2002; 298:2398–2401.10.1126/science.1077780 [PubMed: 12471266]
37. Ambros V. MicroRNAs and developmental timing. *Curr Opin Genet Dev*. 2011; 21:511–517.10.1016/j.gde.2011.04.003 [PubMed: 21530229]
38. Hsu AL, Murphy CT, Kenyon C. Regulation of aging and age-related disease by DAF-16 and heat-shock factor. *Science*. 2003; 300:1142–1145.10.1126/science.1083701 [PubMed: 12750521]
39. Morley JF, Morimoto RI. Regulation of longevity in *Caenorhabditis elegans* by heat shock factor and molecular chaperones. *Mol Biol Cell*. 2004; 15:657–664.10.1091/mbc.E03-07-0532 [PubMed: 14668486]
40. Ibanez-Ventoso C, Yang M, Guo S, Robins H, Padgett RW, Driscoll M. Modulated microRNA expression during adult lifespan in *Caenorhabditis elegans*. *Aging Cell*. 2006; 5:235–246.10.1111/j.1474-9726.2006.00210.x [PubMed: 16842496]

41. de Lencastre A, Pincus Z, Zhou K, Kato M, Lee SS, Slack FJ. MicroRNAs both promote and antagonize longevity in *C. elegans*. *Curr Biol*. 2010; 20:2159–2168.10.1016/j.cub.2010.11.015 [PubMed: 21129974]
42. Kato M, Chen X, Inukai S, Zhao H, Slack FJ. Age-associated changes in expression of small, noncoding RNAs, including microRNAs, in *C. elegans*. *RNA*. 2011; 17:1804–1820.10.1261/rna.2714411 [PubMed: 21810936]
43. Liu N, Landreh M, Cao K, Abe M, Hendriks GJ, Kennerdell JR, Zhu Y, Wang LS, Bonini NM. The microRNA miR-34 modulates ageing and neurodegeneration in *Drosophila*. *Nature*. 2012; 482:519–523.10.1038/nature10810;10.1038/nature10810 [PubMed: 22343898]
44. Li N, Bates DJ, An J, Terry DA, Wang E. Up-regulation of key microRNAs, and inverse down-regulation of their predicted oxidative phosphorylation target genes, during aging in mouse brain. *Neurobiol Aging*. 2011; 32:944–955.10.1016/j.neurobiolaging.2009.04.020 [PubMed: 19487051]
45. Inukai S, de Lencastre A, Turner M, Slack F. Novel microRNAs differentially expressed during aging in the mouse brain. *PLoS One*. 2012; 7:e40028.10.1371/journal.pone.0040028 [PubMed: 22844398]
46. Liang R, Khanna A, Muthusamy S, Li N, Sarojini H, Kopchick JJ, Masternak MM, Bartke A, Wang E. Post-transcriptional regulation of IGF1R by key microRNAs in long-lived mutant mice. *Aging Cell*. 2011; 10:1080–1088.10.1111/j.1474-9726.2011.00751.x;10.1111/j.1474-9726.2011.00751.x [PubMed: 21967153]
47. Bates DJ, Li N, Liang R, Sarojini H, An J, Masternak MM, Bartke A, Wang E. MicroRNA regulation in Ames dwarf mouse liver may contribute to delayed aging. *Aging Cell*. 2010; 9:1–18.10.1111/j.1474-9726.2009.00529.x [PubMed: 19878148]
48. Maes OC, An J, Sarojini H, Wang E. Murine microRNAs implicated in liver functions and aging process. *Mech Ageing Dev*. 2008; 129:534–541.10.1016/j.mad.2008.05.004 [PubMed: 18561983]
49. Zhang X, Azhar G, Wei JY. The expression of microRNA and microRNA clusters in the aging heart. *PLoS One*. 2012; 7:e34688.10.1371/journal.pone.0034688 [PubMed: 22529925]
50. Drummond MJ, McCarthy JJ, Sinha M, Spratt HM, Volpi E, Esser KA, Rasmussen BB. Aging and microRNA expression in human skeletal muscle: a microarray and bioinformatics analysis. *Physiol Genomics*. 2011; 43:595–603.10.1152/physiolgenomics.00148.2010 [PubMed: 20876843]
51. Noren Hooten N, Abdelmohsen K, Gorospe M, Ejiogu N, Zonderman AB, Evans MK. microRNA expression patterns reveal differential expression of target genes with age. *PLoS One*. 2010; 5:e10724.10.1371/journal.pone.0010724 [PubMed: 20505758]
52. Gombor S, Jung HJ, Dong F, Calder B, Atzmon G, Barzilai N, Tian XL, Pothof J, Hoeijmakers JH, Campisi J, Vijg J, Suh Y. Comprehensive microRNA profiling in B-cells of human centenarians by massively parallel sequencing. *BMC Genomics*. 2012; 13:353.10.1186/1471-2164-13-353 [PubMed: 22846614]
53. ElSharawy A, Keller A, Flachsbarth F, Wendschlag A, Jacobs G, Kefer N, Brefort T, Leidinger P, Backes C, Meese E, Schreiber S, Rosenstiel P, Franke A, Nebel A. Genome-wide miRNA signatures of human longevity. *Aging Cell*. 2012; 11:607–616.10.1111/j.1474-9726.2012.00824.x;10.1111/j.1474-9726.2012.00824.x [PubMed: 22533606]
54. Herndon LA, Schmeissner PJ, Dudaronek JM, Brown PA, Listner KM, Sakano Y, Paupard MC, Hall DH, Driscoll M. Stochastic and genetic factors influence tissue-specific decline in ageing *C. elegans*. *Nature*. 2002; 419:808–814.10.1038/nature01135 [PubMed: 12397350]
55. Ibanez-Ventoso C, Vora M, Driscoll M. Sequence relationships among *C. elegans*, *D. melanogaster* and human microRNAs highlight the extensive conservation of microRNAs in biology. *PLoS One*. 2008; 3:e2818.10.1371/journal.pone.0002818 [PubMed: 18665242]
56. Ruby JG, Jan C, Player C, Axtell MJ, Lee W, Nusbaum C, Ge H, Bartel DP. Large-scale sequencing reveals 21U-RNAs and additional microRNAs and endogenous siRNAs in *C. elegans*. *Cell*. 2006; 127:1193–1207.10.1016/j.cell.2006.10.040 [PubMed: 17174894]
57. He L, He X, Lim LP, de Stanchina E, Xuan Z, Liang Y, Xue W, Zender L, Magnus J, Ridzon D, Jackson AL, Linsley PS, Chen C, Lowe SW, Cleary MA, Hannon GJ. A microRNA component of the p53 tumour suppressor network. *Nature*. 2007; 447:1130–1134.10.1038/nature05939 [PubMed: 17554337]

58. Kato M, Paranjape T, Muller RU, Nallur S, Gillespie E, Keane K, Esquela-Kerscher A, Weidhaas JB, Slack FJ. The *mir-34* microRNA is required for the DNA damage response in vivo in *C. elegans* and in vitro in human breast cancer cells. *Oncogene*. 2009; 28:2419–2424.10.1038/onc.2009.106 [PubMed: 19421141]
59. Tazawa H, Tsuchiya N, Izumiya M, Nakagama H. Tumor-suppressive miR-34a induces senescence-like growth arrest through modulation of the E2F pathway in human colon cancer cells. *Proc Natl Acad Sci U S A*. 2007; 104:15472–15477.10.1073/pnas.0707351104 [PubMed: 17875987]
60. Li N, Muthusamy S, Liang R, Sarojini H, Wang E. Increased expression of miR-34a and miR-93 in rat liver during aging, and their impact on the expression of Mgst1 and Sirt1. *Mech Ageing Dev*. 2011; 132:75–85.10.1016/j.mad.2010.12.004 [PubMed: 21216258]
61. Li X, Khanna A, Li N, Wang E. Circulatory miR34a as an RNA-based, noninvasive biomarker for brain aging. *Aging (Albany NY)*. 2011; 3:985–1002. [PubMed: 22064828]
62. Khanna A, Muthusamy S, Liang R, Sarojini H, Wang E. Gain of survival signaling by down-regulation of three key miRNAs in brain of calorie-restricted mice. *Aging (Albany NY)*. 2011; 3:223–236. [PubMed: 21415464]
63. Yang J, Chen D, He Y, Melendez A, Feng Z, Hong Q, Bai X, Li Q, Cai G, Wang J, Chen X. MiR-34 modulates *Caenorhabditis elegans* lifespan via repressing the autophagy gene *atg9*. *Age (Dordr)*. 2011; 10.1007/s11357-011-9324-3
64. Lau NC, Lim LP, Weinstein EG, Bartel DP. An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science*. 2001; 294:858–862.10.1126/science.1065062 [PubMed: 11679671]
65. Lee RC, Ambros V. An extensive class of small RNAs in *Caenorhabditis elegans*. *Science*. 2001; 294:862–864.10.1126/science.1065329 [PubMed: 11679672]
66. Lim LP, Lau NC, Weinstein EG, Abdelhakim A, Yekta S, Rhoades MW, Burge CB, Bartel DP. The microRNAs of *Caenorhabditis elegans*. *Genes Dev*. 2003; 17:991–1008.10.1101/gad.1074403 [PubMed: 12672692]
67. Boulias K, Horvitz HR. The *C. elegans* microRNA *mir-71* acts in neurons to promote germline-mediated longevity through regulation of DAF-16/FOXO. *Cell Metab*. 2012; 15:439–450.10.1016/j.cmet.2012.02.014 [PubMed: 22482727]
68. Miska EA, Alvarez-Saavedra E, Abbott AL, Lau NC, Hellman AB, McGonagle SM, Bartel DP, Ambros VR, Horvitz HR. Most *Caenorhabditis elegans* microRNAs are individually not essential for development or viability. *PLoS Genet*. 2007; 3:e215.10.1371/journal.pgen.0030215 [PubMed: 18085825]
69. Hackl M, Brunner S, Fortschegger K, Schreiner C, Micutkova L, Muck C, Laschober GT, Lepperdinger G, Sampson N, Berger P, Herndler-Brandstetter D, Wieser M, Kuhnel H, Strasser A, Rinnerthaler M, Breitenbach M, Mildner M, Eckhart L, Tschachler E, Trost A, Bauer JW, Papak C, Trajanoski Z, Scheideler M, Grillari-Voglauer R, Grubeck-Loebenstien B, Jansen-Durr P, Grillari J. miR-17, miR-19b, miR-20a, and miR-106a are down-regulated in human aging. *Aging Cell*. 2010; 9:291–296.10.1111/j.1474-9726.2010.00549.x [PubMed: 20089119]
70. Pandey AC, Semon JA, Kaushal D, O'Sullivan RP, Glowacki J, Gimble JM, Bunnell BA. MicroRNA profiling reveals age-dependent differential expression of nuclear factor kappaB and mitogen-activated protein kinase in adipose and bone marrow-derived human mesenchymal stem cells. *Stem Cell Res Ther*. 2011; 2:49.10.1186/scrt90;10.1186/scrt90 [PubMed: 22169120]
71. Grosshans H, Johnson T, Reinert KL, Gerstein M, Slack FJ. The temporal patterning microRNA *let-7* regulates several transcription factors at the larval to adult transition in *C. elegans*. *Dev Cell*. 2005; 8:321–330.10.1016/j.devcel.2004.12.019 [PubMed: 15737928]
72. Larsen PL, Albert PS, Riddle DL. Genes that regulate both development and longevity in *Caenorhabditis elegans*. *Genetics*. 1995; 139:1567–1583. [PubMed: 7789761]
73. Bethke A, Fielenbach N, Wang Z, Mangelsdorf DJ, Antebi A. Nuclear hormone receptor regulation of microRNAs controls developmental progression. *Science*. 2009; 324:95–98.10.1126/science.1164899 [PubMed: 19342589]

74. Hammell CM, Karp X, Ambros V. A feedback circuit involving *let-7*-family miRNAs and DAF-12 integrates environmental signals and developmental timing in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*. 2009; 106:18668–18673.10.1073/pnas.0908131106 [PubMed: 19828440]
75. Toledano H, D'Alterio C, Czech B, Levine E, Jones DL. The *let-7*-Imp axis regulates ageing of the *Drosophila* testis stem-cell niche. *Nature*. 2012; 485:605–610.10.1038/nature11061;10.1038/nature11061 [PubMed: 22660319]
76. Baek D, Villen J, Shin C, Camargo FD, Gygi SP, Bartel DP. The impact of microRNAs on protein output. *Nature*. 2008; 455:64–71.10.1038/nature07242 [PubMed: 18668037]
77. Forstemann K, Tomari Y, Du T, Vagin VV, Denli AM, Bratu DP, Klattenhoff C, Theurkauf WE, Zamore PD. Normal microRNA maturation and germ-line stem cell maintenance requires Loquacious, a double-stranded RNA-binding domain protein. *PLoS Biol*. 2005; 3:e236.10.1371/journal.pbio.0030236 [PubMed: 15918770]
78. Saito K, Ishizuka A, Siomi H, Siomi MC. Processing of pre-microRNAs by the Dicer-1-Loquacious complex in *Drosophila* cells. *PLoS Biol*. 2005; 3:e235.10.1371/journal.pbio.0030235 [PubMed: 15918769]
79. Jiang F, Ye X, Liu X, Fincher L, McKearin D, Liu Q. Dicer-1 and R3D1-L catalyze microRNA maturation in *Drosophila*. *Genes Dev*. 2005; 19:1674–1679.10.1101/gad.1334005 [PubMed: 15985611]
80. Winter J, Diederichs S. Argonaute proteins regulate microRNA stability: Increased microRNA abundance by Argonaute proteins is due to microRNA stabilization. *RNA Biol*. 2011; 8:1149–1157.10.4161/rna.8.6.17665 [PubMed: 21941127]
81. Merritt WM, Lin YG, Han LY, Kamat AA, Spannuth WA, Schmandt R, Urbauer D, Pennacchio LA, Cheng JF, Nick AM, Deavers MT, Mourad-Zeidan A, Wang H, Mueller P, Lenburg ME, Gray JW, Mok S, Birrer MJ, Lopez-Berestein G, Coleman RL, Bar-Eli M, Sood AK. Dicer, Drosha, and outcomes in patients with ovarian cancer. *N Engl J Med*. 2008; 359:2641–2650.10.1056/NEJMoa0803785 [PubMed: 19092150]
82. Thomson JM, Newman M, Parker JS, Morin-Kensicki EM, Wright T, Hammond SM. Extensive post-transcriptional regulation of microRNAs and its implications for cancer. *Genes Dev*. 2006; 20:2202–2207.10.1101/gad.1444406 [PubMed: 16882971]
83. Esquela-Kerscher A, Slack FJ. Oncomirs - microRNAs with a role in cancer. *Nat Rev Cancer*. 2006; 6:259–269.10.1038/nrc1840 [PubMed: 16557279]
84. Herranz H, Cohen SM. MicroRNAs and gene regulatory networks: managing the impact of noise in biological systems. *Genes Dev*. 2010; 24:1339–1344.10.1101/gad.1937010 [PubMed: 20595229]
85. Martinez NJ, Ow MC, Barrasa MI, Hammell M, Sequerra R, Doucette-Stamm L, Roth FP, Ambros VR, Walhout AJ. A *C. elegans* genome-scale microRNA network contains composite feedback motifs with high flux capacity. *Genes Dev*. 2008; 22:2535–2549.10.1101/gad.1678608 [PubMed: 18794350]

Highlights

- miRNAs have been implicated in aging and may function as network-level regulators.
- Differentially expressed miRNAs during aging may target age-associated pathways.
- Specific miRNAs may link and coordinate the functions of aging-associated pathways.

Figure 1. MicroRNAs are part of a complex genetic network implicated in aging

Individual and groups of miRNAs are associated with aging and aging-related pathways that mediate signals from the external and internal environments and center on the insulin signaling pathway. In particular, *let-7*, miR-71, and miR-34 are connected to multiple pathways.

Table 1

Summary of aging-associated microRNAs and their associated species and functions.

miRNAs	species	aging-associated pathways	references
miR-34	<i>C. elegans</i> , <i>D. melanogaster</i> , <i>M. musculus</i> , <i>R. norvegicus</i>	DNA damage response, senescence, cell death	40–42, 43, 45, 60–63
<i>let-7</i>	<i>C. elegans</i> , <i>D. melanogaster</i> , <i>M. musculus</i> , <i>H. sapiens</i>	DNA damage response, mitochondrial respiration, germline maintenance	40–42, 45, 48, 50, 53, 69, 70, 75
miR-71	<i>C. elegans</i>	DNA damage response, IIS, germline signaling	40–42, 67
miR-239	<i>C. elegans</i>	IIS	40–42
miR-35	<i>C. elegans</i>	?	40–42
miR-38	<i>C. elegans</i>	?	40–42
<i>miR-43</i>	<i>C. elegans</i>	?	40–42
miR-70	<i>C. elegans</i>	?	40–42
miR-30d	<i>M. musculus</i>	?	44, 47
miR-468	<i>M. musculus</i>	?	44, 47
miR-669b	<i>M. musculus</i>	?	44, 47
miR-709	<i>M. musculus</i>	?	44, 47
miR-246	<i>C. elegans</i>	?	41