Published in final edited form as:

Dev Psychopathol. 2011 May ; 23(2): 617–627. doi:10.1017/S0954579411000046.

Perceived discrimination, serotonin transporter linked polymorphic region status, and the development of conduct problems

Gene H. Brody,

Center for Family Research, Institute for Behavioral Research, University of Georgia, Department of Behavioral Science and Health Education, Rollins School of Public Health, Emory University

Steven R. H. Beach,

Center for Family Research, Institute for Behavioral Research, University of Georgia

Yi-fu Chen,

Center for Family Research, Institute for Behavioral Research, University of Georgia

Ezemenari Obasi,

Department of Counseling and Human Development Services, College of Education, University of Georgia

Robert A. Philibert,

Department of Psychiatry, University of Iowa

Steven M. Kogan, and

Center for Family Research, Institute for Behavioral Research, University of Georgia

Ronald L. Simons

Department of Sociology, University of Georgia

Abstract

This study examined the prospective relations of adolescents' perceptions of discrimination and their genetic status with increases in conduct problems. Participants were 461 African American youths residing in rural Georgia (*M* age, wave 1 = 15.5 years) who provided three waves of data and a saliva sample from which a polymorphism in the *SCL6A4* (*5-HTT*) gene promoter known as the 5-HTTLPR was genotyped. Data analyses using growth curve modeling indicated that perceived discrimination was significantly related to the slope of conduct problems. As hypothesized, interactions between perceived discrimination and genetic status emerged for male but not female youths. Compared with those carrying two copies of the long allele variant of 5-HTTLPR, male youths carrying one or two copies of its short allele variant evinced higher rates of conduct problems over time when they perceived high levels of racial discrimination. These findings are consistent with resilience and differential susceptibility propositions stating that genes can both foster sensitivity to adverse events and confer protection from those events.

Keywords

African American; conduct problems; discrimination; gender; genetics

For some young people, some forms of externalizing behaviors or conduct problems increase significantly during preadolescence and adolescence (see Moffitt, 1993). Conduct problems encompass a variety of antisocial behaviors, including aggression, destruction of property, deceit, and theft. Sustained or increased conduct problems forecast later aggressive and antisocial behaviors, adult criminality, early mortality, poor physical health, comorbid disorders, academic underachievement, and underemployment during young adulthood (Teichner & Golden, 2000; Wicks-Nelson & Israel, 2003). The processes that contribute to the development and maintenance of conduct problems during this developmental period should be identified, not only to advance knowledge of their etiology and the trajectories they foster but also to inform the design of preventive interventions (Howe, Beach, & Brody, in press). Using a three-wave prospective research design that spanned 3 years, we advanced and evaluated predictions about a salient vulnerability factor for the development of conduct problems among African American youths, perceived racial discrimination. Racial discrimination is a prevalent stressor for African American youths; children and adolescents who perceive high levels of racial discrimination display, over time, increasing rates of conduct problems, and the direction of causality in this association has been demonstrated to flow from discrimination to conduct problems rather than the reverse (Brody, Chen, et al., 2006). It is therefore important to understand better the factors that increase the impact of this contextual stressor.

The primary purpose of this study was to test predictions regarding the genetic moderation of the association between perceived discrimination and development of conduct problems. We did not expect genetic variation to have a direct linear association to the development of conduct problems; instead, we expected genetic status to predict variation in youths' responses to perceived racial discrimination. This perspective is consistent with resilience and differential susceptibility theories, in which genetic variations are hypothesized to render individuals more or less susceptible to environmental risks (Belsky, Bakermans-Kranenburg, & van IJzendoorn, 2007; Caspi, Hariri, Holmes, Uher, & Moffitt, 2010; Cicchetti & Blender, 2006; Kim-Cohen & Gold, 2009). One such genetic factor is variability at *5-HTT*. We tested a hypothesis involving a putative genetic vulnerability factor, the presence of a short allele of a variable nucleotide repeat polymorphism in the promoter region of *5-HTT* known as the 5-HTTLPR (*5-HTT* linked polymorphic region). We proposed that youths who carry one or two copies of this allele, compared with those carrying two copies of the long allele, would display higher rates of conduct problems over time when they perceive high levels of racial discrimination. We also expected rates of conduct problems among youths carrying two copies of the long allele to evince little variation across levels of perceived discrimination. In the following sections, we discuss the hypothesized roles of racial discrimination, the 5-HTTLPR genotype, and their interaction in the development of conduct problems.

Racial discrimination presents daily challenges in the lives of African American youths and their families. Landrine and Klonoff (1996) found that 98% of the African American adults in their sample had experienced a racist event during the past year. All participants reported that, at some point in their lives, they had been treated badly or their intentions were misinterpreted because of their race. Likewise, Sellers, Copeland-Linder, Martin, and Lewis (2006) found that 70% or more of African American adolescents reported racialized insults in which others perceived them as threatening or incompetent. These experiences are demeaning and degrading, inducing stress, frustration, depression, and anxiety. Thoits (1991) maintained that those stressors that threaten the central parts of an individual's identity are particularly pernicious and likely to undermine mental health. Accordingly, African American youths' experiences with discrimination have the potential to provoke problem behaviors arising from anger, a sense of injustice, and other negative emotions.

Survey research with adolescents corroborates the hypothesized association between perceived discrimination and the development of conduct problems; links between these variables have been demonstrated and replicated using both contemporaneous and longitudinal research designs (Brody, Chen, et al., 2006; DuBois, Burk-Braxton, Swenson, Tevendale, & Hardesty, 2002; Simons, Chen, Stewart, & Brody, 2003; Simons et al., 2002; Simons et al., 2006; Szalacha et al., 2003). Even worrying about the prospect of future racial discrimination creates a heightened state of vigilance, placing adolescents' psychological well-being at risk (Fordham & Ogbu, 1986; D. M. Taylor, Wright, & Porter, 1993). These findings resonate with the hypothesis (Franklin, 1993) that ethnic minority individuals' inner vigilance for racial slights can create a state of constant watchfulness that leads to tension and stress.

The first purpose of this study was to replicate findings published in earlier reports (Brody, Chen, et al., 2006; Simons et al., 2003; Simons et al., 2006) in which we found that perceptions of racial discrimination forecast increases in conduct problem behaviors across 5 years for male, but not female, youths. In those studies, we proposed that male youths are more likely than are female youths to become emotionally dysregulated and lose inhibitory controls when experiencing life stresses (Rutter, 1990), leading to an increase in conduct problems in response to perceived discrimination. In the present study, the longitudinal links between perceived racial discrimination and increases in conduct problems were re-examined using a different sample of African American youths who were 15 years of age at the first wave of data collection and 17 years of age at the last wave. We hypothesized that perceptions of high levels of discrimination would forecast higher rates of conduct problems for male youths.

Ecological (Bronfenbrenner & Ceci, 1994), systems (Lerner, 1991), and developmental psychopathology (Cicchetti & Tucker, 1994) approaches to life-span development share the tenet that biological predispositions transact with contextual processes to create variations in phenotypes over time. Such transactions are termed gene $\times$ environment (G $\times$ E) interactions and are evident when genetic variation alters an organism's sensitivity to specific environmental events or when environmental features exert differential control over genetic effects (Kendler & Eaves, 1986). In this study, we focused on the role of genetic status as a moderator of environmental influence. We hypothesized that a functional polymorphism in the promoter region of the serotonin transporter gene, *5-HTT*, would moderate the association of perceived discrimination with longitudinal increases in youths' conduct problems. The *5-HTT* gene is a key regulator of serotonergic neurotransmission, localized to 17p13 and consisting of 14 exons and a single promoter. The common polymorphism in the promoter region results in two variants, a short and a long allele, with the short allele resulting in lower serotonin transporter availability. The short (*s*) variant contains 14, and the long (*l*) variant 16, copies of the repeat element. Youths with one or two copies of the *s* allele are hypothesized to display greater increases in conduct problems when they perceive racial discrimination than are youths who carry two copies of the *l* allele.

Two bodies of research support the proposition that 5-HTTLPR status will moderate the link between perceived discrimination and rates of conduct problems across time. The first was generated by neuroscientists who sought to determine whether 5-HTTLPR status affected the processing of social cues. The second was sponsored by researchers testing G $\times$ E hypotheses involving 5-HTTLPR status and exposure to stressful life events. Research on the neuroscience of information processing indicates that the 5-HTTLPR genotype affects the processing of socially important cues, particularly those related to threats and punishment. Specifically, the *s* allele is associated with relatively greater activity in the amygdala (Heinz et al., 2005), a brain region involved in the processing of verbal and nonverbal threats (Isenberg et al., 1999), and with enhanced amygdala reactivity to

punishment cues in the environment (Battaglia et al., 2005; Hariri et al., 2005; Hariri, Drabant, & Weinberger, 2006; Hariri, Mattay, Tessitore, Fera, & Weinberger, 2003; Hariri et al., 2002). Recent research has extended these findings by demonstrating that *ss* and *sl* allele carriers also display amygdala hyperactivity to nonemotional and neutral cues (Heinz et al., 2007; Munafò, Brown, & Hariri, 2008), direct preferential attention toward threat-related stimuli, and have difficulty disengaging from such stimuli (Beevers, Wells, Ellis, & McGeary, 2009; Osinsky et al., 2008). This suggests that the *s* allele may engender a predisposition toward hypervigilance and high reactivity to environmental events. Presumably, this would have implications for reactions to perceived discrimination.

Recent G×E research involving the *5-HTT* genotype provides evidence that this genotype sponsors differential reactivity to environmental influence (Caspi et al., 2010). Many of these studies focused on determining whether 5-HTTLPR status conditions the associations between stressful life events and depressive symptomatology. These studies described how genetic variability at *5-HTT* alters individuals' reactivity to environmental stressors, resulting in both higher levels of depressive symptoms depending on participants' 5-HTTLR status and the level of environmental stress they encountered. For example, adolescents (Eley et al., 2004), young adults (Wilhelm et al., 2006), and adults (Brummett et al., 2008; Caspi et al., 2003; Lazary et al., 2008; S. E. Taylor et al., 2006; Zalsman et al., 2006) carrying the *ss* or *sl* genotype reported more depressive symptoms when exposed to high levels of stressful life events compared with individuals carrying the *ll* genotype. A study by Surtees et al. (2006), however, did not replicate this pattern (see also Risch et al., 2009).

Other investigators have focused on anxiety and ADHD outcomes to discover whether 5-HTTLPR status contributed to variation in emotional responses to negative environmental events. Gunthert et al. (2007) used a 1-year longitudinal design with a college sample to study moderation effects of daily negative events sampled across 30 days on symptoms of anxiety. They found that, compared with individuals with the *ll* genotype, individuals homozygous for the *s* allele experienced the greatest anxiety in the presence of negative events. Stein, Schork, and Gelernter (2008) examined, with a college sample, the link between retrospectively reported emotional abuse during childhood and anxiety sensitivity. Again, a G×E interaction emerged showing differential outcomes according to the number of *s* alleles participants carried; those with the *ss* genotype scored highest in anxiety sensitivity when reporting high levels of emotional abuse relative to participants with *ll* alleles. Finally, research by Retz, et al. (2008) tested longitudinal moderation effects in a sample of male adults with a history of delinquency. Retrospective assessments of adverse environments and ADHD during childhood and a clinical interview assessment of adult ADHD were obtained to examine the persistence of ADHD symptoms from childhood to adulthood. Results revealed an interaction between 5-HTTLPR genotypes and adverse childhood environments such that, compared with *ll* participants, the association between environmental adversity and symptom persistence was greatest for participants with *ss* or *sl* alleles who experienced childhood adversity. Taken together, the overall findings from this research are consistent with the view that individuals differ in their reactivity, with some more affected than others by stressful experiences.

Thus far, we have reviewed support for the propositions that perceived discrimination forecasts conduct problems, particularly among male youths; that carrying one copy of the *s* allele is associated with heightened reactivity to threatening environmental cues and events; and that studies focusing on 5-HTTLPR within the G×E literature suggest that certain individuals may be more sensitive and reactive than others to negative environmental experiences. The supporting literatures all speak to the primary purpose of this study, which is to determine whether some youths will be affected more than others by perceived

discrimination, resulting in different rates of change in conduct problems across adolescence.

Summary of the Present Research

This study was conducted with rural African American youths using procedures that have been shown in developmental psychopathology research to yield reliable data. These procedures include computer-based interviewing, ethnic matching of interviewers and participants, and extensive reassurances concerning confidentiality of the data (Brody et al., 2004; Patrick et al., 1994). Genetic data were obtained from youths using procedures developed in partnership with rural African American community members (Brody, Beach, & Philibert, in press). We predicted that high levels of perceived discrimination would be associated with longitudinal increases in conduct problems among African American male, but not female, youths, and that this association would be amplified for youths carrying at least one copy of the *s* variant of 5-HTTLPR and ameliorated for those carrying two *l* alleles. We tested the study hypotheses using latent growth curve modeling (Singer & Willett, 2003) with relevant demographic characteristics controlled.

Method

Data were collected as part of a family-based preventive intervention study (see Brody et al., 2004). Random assignment to the prevention or control group did not interact with any of the variables that were addressed in this study. Nevertheless, participation in the prevention or control group was controlled in all of the data analyses reported. Three waves of data on conduct problems were obtained yearly when the youths were 15 through 17 years old. Perceived discrimination was assessed during the first data collection and the genetic data were obtained when the youths were 16.

Participants

The sample for this study included 454 African American youths residing in rural Georgia. Of a sample of families who provided data at Wave 1 ($n = 573$), 91% ($n = 524$) provided it at Wave 3. Youths in 87% ($n = 454$) of these families agreed to provide DNA. Two equivalence analyses were executed to determine whether any differences existed on demographic characteristics (monthly per capita income, maternal education, youth gender, maternal marital status, and program participation) or the study variables (perceived discrimination and conduct problems) between (a) the families with Wave 1 data ($n = 573$) and those with Wave 3 data ($n = 524$) and (b) the target youths who did ($n = 454$) and did not ($n = 70$) agree to provide DNA. To conserve space, the means and standard deviations for these tests are not presented. For these equivalence comparisons, *t* tests with alpha set at $p < .05$ were used for all comparisons except gender, maternal marital status, and program participation, for which chi-square tests with alpha set at $p < .05$ were used. No differences emerged on the demographic or study variables between the youths and families with Wave 1 or Wave 3 data. Some differences emerged between youths who did or did not agree to provide DNA. At Waves 1 and 2, levels of conduct problems were higher for youths who provided DNA than for those who did not provide it; no differences emerged for comparisons at Wave 3.

At the first data collection session, youths' mean age was 15.5 years ($SD = 0.50$). In terms of family structure, 38.7% of the youths were living in single-mother households, 35.3% were living in married-parent households, 18.8% were living with extended family members, 5.1% were living in households with unmarried cohabiting adults, and 2.1% were living in households in which married parents were separated. Mean household gross monthly income was \$2,577 ($SD = \$2,608$) and mean monthly per capita gross income was \$670 (SD

= \$745). Although 72.6% of the mothers were employed outside the home and worked an average of 39.7 hours per week, 46.4% of the families lived below federal poverty standards and another 20.5% lived within 150% of the poverty threshold; they could be described as working poor (Boatright, 2005).

Schools in four rural Georgia counties provided lists of 11-year-old students, from which youth participants were selected randomly (see Brody et al., 2004). Families were contacted and enrolled in the study by community liaisons who resided in the counties where the participants lived. The community liaisons were African American community members, selected on the basis of their social contacts and standing in the community, who worked with the researchers on participant recruitment and retention. The liaisons sent letters to the families and followed up on the letters with phone calls to the mothers. During the phone conversations, the community liaisons answered any questions that the mothers asked. Families who were willing to participate in the pretest were told that a research staff member would contact them to schedule the administration of the assessment in the families' homes. Parents gave written consent to their own and the youths' participation, and youths gave written assent to their own participation. Each family was paid \$100 after each of the six assessments.

Preparation for the Collection of Genetic Data

Several steps were taken to prepare for DNA data collection. The researchers met regularly over a 2-year period to review the genetic and family process literature. These researchers included a psychiatrist who specialized in human genetics; a biostatistician who specialized in the analysis of genetic data; a developmental psychologist; and a clinical psychologist. The meetings they held resulted in the formulation of the hypotheses tested in this report. In addition, two focus groups of rural African Americans were formed, one for parents and one for adolescents. Each group included 10 persons who met for 2 hours. These groups were formed to help the investigators understand any concerns that might arise about the collection of DNA and to develop procedures for dealing with these concerns. Several concerns arose that involved procedural clarity, detection, and potential benefits. Many focus group members wanted a clear explanation of the procedures for obtaining the DNA and they wanted to know how DNA collection would advance knowledge about the development of African American youths. This feedback was incorporated into the development of a brochure, available from the first author, that included answers to frequently asked questions. The information addressed in a straightforward manner the issues the focus groups raised. A copy of the brochure was given to each participating family to provide them with written information that they could consult in addition to the verbal description of the protocol given during DNA collection.

A pilot study was conducted to assess the viability of DNA collection from saliva versus whole blood (Philibert, Zadorozhnyaya, Beach, & Brody, 2008). As predicted, concentrations of DNA were higher in blood than in saliva; the saliva samples, however, contained adequate amounts of DNA to permit genotyping. We concluded that the ease and economy of DNA collection from saliva made it appropriate for the research questions we planned to address.

Procedure

Trained African American field researchers conducted computer-based interviews in participants' homes. Youths were interviewed individually and privately; they were told that their answers were strictly confidential and would not be disclosed to anyone within or outside the family. Mothers provided informed consent for the youths' provision of

interview data and the collection of DNA from the youths; youths assented to their own participation.

Measures

The measures were selected for their relevance to the lives of rural African American youths. They were derived from previous research, which included focus group meetings and pilot testing followed by construct validation of the instruments (Brody, Murry, et al., 2006).

Demographics—Youth age and gender, maternal age, employment status, and monthly income were recorded. Each mother also reported the number of children and adults living in her home, her marital status, and her relationship status.

Perceived discrimination—At the first wave of data collection, the target youths completed 13 items from a revised version of the Schedule of Racist Events (SRE; Landrine & Klonoff, 1996). The SRE was designed for adult respondents; we revised it for youths in late childhood through adolescence. We have used the revised scale successfully in our previous research (Brody, Chen, et al., 2006; Gibbons, Gerrard, Cleveland, Wills, & Brody, 2004; Simons et al., 2003; Simons et al., 2006). The revisions included simplifying the language and replacing items dealing with discrimination in the workplace with items about discriminatory behaviors in the community. The revision process involved presentation of the revised scale to focus groups of African American primary caregivers and youths of the same age as those in the study population; the second step included psychometric and validity analyses. The validity of the revised scale was demonstrated through its association with variables that the literature suggests would be associated with perceived discriminatory experiences, such as anger, a hostile world view, depression, and conduct problems (Clark, Anderson, Clark, & Williams, 1999; Landrine & Klonoff, 1996), in contemporaneous (Simons et al., 2003), 2-year longitudinal (Gibbons et al., 2004), and 5-year longitudinal (Brody, Chen, et al., 2006; Simons et al., 2006) assessments. These associations remained robust when family income, financial stress, negative life events, and parental education were controlled.

The items in the revised SRE assessed the frequency during the past year, ranging from 1 (*never*) to 4 (*several times*), with which the respondent perceived specific discriminatory behaviors. These events include racially based slurs and insults, disrespectful treatment from community members, physical threats, and false accusations from business employees or law enforcement officials. Sample items include, “someone said something insulting to you because you are African American,” “a store owner or sales person working at a business treated you in a disrespectful way because you are African American,” “someone yelled a racial insult at you because you are African American,” and “you encountered Whites who didn’t expect you to do well because you are African American.” Coefficient alpha for the scale was .85.

Conduct problems—Using 14 questions from the National Youth Survey (Elliott, Ageton, & Huizinga, 1985), youths indicated how often, during the past year, they engaged in problem behaviors involving theft, truancy, suspension from school, and fighting. Elliott et al. (1985) illustrated the scoring of this instrument; the number of behaviors that a youth endorses were summed and this sum constituted the conduct problems score. Because this instrument assesses count data, internal consistency analyses were not executed.

Genotyping—Youths’ DNA was obtained using Oragene™ DNA kits (Genetek, Calgary, AB, Canada). Youths rinsed their mouths with tap water, then deposited 4 ml of saliva in the

OraGene sample vial. The vial was sealed, inverted, and shipped via courier to a central laboratory in Iowa City, IA where samples were prepared according to the manufacturer's specifications. Genotype at 5-HTTLPR was determined for each youth as described by Bradley, Dodelzon, Sandhu, and Philibert (2005), using the primers F-GGCGTTGCCGCTCTGAATGC and R-GAGGGACTGAGCTGGACAACCAC, standard *Taq* polymerase and buffer, standard dNTPs with the addition of 100 μ M 7-deaza GTP, and 10% DMSO. The resulting PCR products were electrophoresed on a 6% non-denaturing polyacrylamide gel and products visualized using silver staining. Genotype was then called by two individuals blind to the study hypotheses and other information about the participants. Of the sample, 6.4% were homozygous for the short allele (*ss*), 35.2% were heterozygous (*sl*), and 58.4% were homozygous for the long allele (*ll*). None of the alleles deviated from Hardy–Weinberg equilibrium ($p = .77$, ns). Consistent with prior research (Hariri et al., 2005), genotyping results were used to form two groups of participants: those homozygous for the long allele and those with either one or two copies of the short allele.

Results

Descriptive Statistics

Prevalence of conduct problems increased across time, particularly among male youths. When male youths were 15 years of age, their mean conduct problem score was 4.44 ($SD = 13.58$); at age 17, it was 4.62 ($SD = 11.25$). The corresponding means for female youths were 2.07 ($SD = 7.57$) and 1.93 ($SD = 7.26$), respectively. Means for depressive symptoms at Wave 3 were 10.42 ($SD = 7.77$) for male youths and 11.42 ($SD = 8.34$) for female youths. The gender differences that emerged in this sample are consistent with those found in other longitudinal studies with African American youths residing in urban contexts (Brody, Chen, et al., 2006; Simons et al., 2006).

Plan of Analysis for the Study Hypotheses

Growth curve modeling (GCM) using Hierarchical Linear Modeling (Bryk & Raudenbush, 1992) was used to test the study hypotheses. GCM estimates the mean and variance for the slope and intercept of conduct problems, treating the conduct problem intercept and slope as a random effect. The slope corresponding to change in conduct problems varies depending on the mathematical form being tested (e.g., linear, quadratic, and so forth). In this study, we used a linear model. Testing the hypotheses involved several steps. First, an unconditional growth curve analysis was executed to test the measurement model for conduct problems. Two GCM models were then evaluated with the whole sample. The first model was designed to examine associations among increases in youths' conduct problems, perceived discrimination, and genetic status to determine whether they were statistically significant and to ensure that the hypothesized model would be consistent with the data. The second model was executed to estimate the main effects and interaction of perceived discrimination and genetic status in predicting conduct problem slopes. To test the interaction effect, the interaction term was inserted into the GCM model and its strength was assessed. For the hypothesized interaction between perceived discrimination and genetic status, a significant interaction term indicates that the perceived discrimination-conduct problem relationship differs across genetic status, yielding a G×E interaction. These analyses were repeated separately for male and female youths to test the proposed gendered hypothesis. For all of the models described here, the predictor variables were mean centered and the demographic variables of maternal education, monthly per capita income, youth gender (in analyses of the entire sample), marital status, and program participation were systematically controlled in every equation when predicting the slopes for conduct problems.

Discrimination and Youth Conduct Problems

The theoretical rationale for this study is based on the premise that perceptions of racial discrimination lead to conduct problems rather than the reverse. Structural equation modeling (SEM) was conducted in which the cross-lags (i.e., Wave 1 perceived discrimination to Wave 2 conduct problems and Wave 1 conduct problems to Wave 2 perceived discrimination) were first constrained to be equal; the constrained model was then compared with one in which the two paths were free to vary. The non-constrained model fit the data well, $\chi^2(0) = 0$. A significant reduction in chi-square from the constrained to the non-constrained model indicates that one cross-lag is stronger than the other. In this analysis, only one cross-lag was significant, the path from perceived discrimination to conduct problems ($\beta = .12, p < .01$). The path from conduct problems to perceived discrimination was not significant ($\beta = .05, ns$). Chi-square was reduced significantly from 7.35 ($df = 1$) to 0 ($df = 0$), a change of 7.35 ($df = 1; p < .01$).

Testing the Study Hypotheses Using LGC Modeling

We began by testing the measurement, or unconditional, model for the conduct problem latent growth construct by fitting a univariate growth curve for the construct. The model was based on reported levels of conduct problems at ages 15, 16, and 17. The slope for the three time points of conduct problems reflected the number of years after the first assessment when each subsequent assessment was obtained. The slope variance for the unconditional model was 26.18 ($p < .01$). The significant variance for the slope indicates significant variability around the mean slope for conduct problems.

In the models that tested the study hypotheses, 5-HTTLPR status was dummy coded. Participants with *ss* or *s/* alleles were combined into one genetic status group and assigned a code of 1, participants with *ll* alleles were assigned a code of 0, and the genetic status predictor was regressed on the conduct problems slope. The first model involving the whole sample is presented in Model 1 in Table 1. The results were consistent with the first hypothesis: Perceived discrimination was associated with a significantly higher rate of change in conduct problems across adolescence (perceived discrimination, $\beta = 0.61, p < .01$).

The second model executed on the whole sample (Model 2 in Table 1) was designed to test the proposed G×E hypothesis. When the interaction term was added to the model, the association between perceived discrimination and the slope of youth conduct problems approached significance ($p < .10$). The interaction between perceived discrimination and 5-HTTLPR status yielded significant predictions beyond the previous variables in the model, $\beta = 0.76, p < .01$. To interpret this interaction, we plotted the slopes of conduct problems in Figure 1 for levels of perceived discrimination that range from -3 to $+3$ *SD* from the sample mean, separately for youths with *ss* or *s/* genotypes and youths with *ll* genotypes. A perusal of Figure 1 reveals several notable findings. Consistent with the study hypothesis, compared with youths who had *ll* genotypes, youths with *ss* or *s/* genotypes evinced the highest rates of change in conduct problems when they perceived high levels of racial discrimination. Of particular note were the two findings, illustrated in Figure 2, that the *ll* genotype apparently protected youths exposed to high levels of discrimination from increases in conduct problems and that, under low levels of perceived racial discrimination, *s* carriers evinced lower levels of conduct problems than did *ll* carriers. The latter finding is consistent with Belsky and colleagues' (2007) differential susceptibility hypothesis. We also examined the possibility that a gene-environment correlation could exist between 5-HTTLPR and perceived discrimination. Consistent with our expectations that the *s* allele would be associated with greater vigilance for environmental threat, a significant correlation emerged between genetic status and perceived discrimination ($r = 0.22, p = .05$). To examine the

possibility that this rGE accounted for the association between perceived discrimination and conduct problems, we estimated a mediational model in which we assessed the significance of the indirect effect from genetic status on the conduct problems slope via perceived discrimination. The estimated indirect effect was $\beta = 0.08$, $p = .30$; therefore, the rGE association did not account for the impact of perceived discrimination on outcome.

Gendered models—The model presented in Table 1 was re-run separately for male and female youths to test for hypothesized gender differences. The separate models are presented in Table 1. For male youths, the main effect model presented in Model 3 of Table 1 indicated that perceived discrimination was associated with increases in conduct problems, $\beta = 1.24$, $p < .05$. The second model for male youths, presented in Model 4 in Table 1, tested the $G \times E$ hypothesis that perceived discrimination would interact with genetic status to forecast the slope of conduct problems. The interaction between perceived discrimination and genetic status was significant, $\beta = 1.22$, $p < .01$. Again, to understand how the interaction affected the trajectories of conduct problems for male youths, we plotted the slopes of conduct problems in Figure 2. The forms of the slopes were nearly identical to those depicted in Figure 1. Male youths with the *ss* or *s/* genotype, compared with those who had the *//* genotype, evinced the highest rates of change in conduct problems when they perceived high levels of discrimination. Again, the slope of conduct problems for male youths with the *//* genotype appeared relatively unaffected by perceived racial discrimination, whereas carriers of an *s* allele evinced lower levels of conduct problems than did *//* carriers.

The analyses for female youths (see Models 5 and 6 in Table 1) reveal a different pattern. A main effect for perceived discrimination approached significance ($\beta = 0.19$, $p < .10$), but no $G \times E$ interaction emerged.

Discussion

Sponsored by both differential susceptibility and resilience theories (Belsky et al., 2007; Caspi et al., 2010; Cicchetti & Blender, 2006; Kim-Cohen & Gold, 2009), we predicted that perceived discrimination would be linked to increases in conduct problems when male African American youths carried one or two copies of the *s* allele of the 5-HTTLPR. Consistent with this hypothesis, a clear and pronounced link emerged between perceived discrimination and growth in conduct problems across 3 years among youths with these genotypes. This finding is consistent with the proposition, informed by neuroimaging and prior $G \times E$ research involving the 5-HTTLPR, that male youths who carry the *s* allele, because of their genetic makeup, are more likely to be affected by perceived discrimination than are those with the *//* genotype. The finding that carrying two copies of the */* allele buffered male youths from developing conduct problems when they perceived high levels of discrimination is pertinent to research on youth resilience. This literature has addressed the reasons why some youths who experience many discrete and chronic stressors do not succumb to their negative effects (Luthar, 2006). Typically the locus of these resilience effects is identified through contextual processes at different levels of analysis (family, peer, school, or neighborhood) that alter several types of pathways, including reduction of the effects of risk factors. The present results reinforce recent suggestions that genetic status can also contribute to resilience (Kim-Cohen & Gold, 2009; Moffitt, Caspi, & Rutter, 2006; Rutter & Silberg, 2002). The observed buffering effects of carrying two */* alleles suggest an emotional self-regulatory mechanism in which genotype contributes to youths' downregulation of emotions such as anger, victimization, and feelings of injustice that discrimination occasions (Simons et al., 2003; Simons et al., 2006). Further research is needed to test this hypothesis.

Caution should be used, however, in interpretation and generalization of the protective effects the *II* genotype to all African American youths. Some who seem unfazed by high levels of perceived discrimination may not be as adaptable as they appear; still others may be resilient in some areas but experience distress in others, such as health or aspects of social relationships that do not involve conduct problems. Research with children who have been maltreated (Farber & Egeland, 1987), whose mothers have depression (Hammen, 2003), and whose parents have alcoholism (Zucker, Wong, Puttler, & Fitzgerald, 2003) support this caveat. Even youths who are well adjusted behaviorally can have their resilient trajectories disrupted by the introduction of other risk factors that diminish the protective capacities that the *II* genotype may confer.

The results also supported Belsky and colleagues' (2007) differential susceptibility hypothesis, in which specific gene variants, including the 5-HTTLPR, are proposed to render individuals more susceptible to the surrounding environment whether it is characterized by high positivity or high risk. The finding that, under conditions of low perceived discrimination, *s* carriers evinced lower levels of conduct problems than did *II* carriers does support differential susceptibility predictions.

We also investigated gender differences in the G×E hypothesis that we examined. Although gender has been identified as an important source of individual variability in stress and coping among members of ethnic minority groups (Clark et al., 1999; Slavin, Rainer, McCreary, & Gowda, 1991), few studies other than our own have examined gender differences in youths' responses to discrimination. As expected, the hypothesized G×E effects were apparent for male youths but not for female youths. This finding may reflect two realities. First, the rates of conduct problems for male youths were three times those of female youths across the time frame of the present study. This finding is not specific to the sample in this investigation; other studies report male-to-female ratios of 3:1 or 4:1 for conduct problems (Davison, Neale, Blankstein, & Flett, 2005). The greater prevalence of conduct problems among male youths and the resulting increased variability in change trajectories would render interactions easier to detect. It is possible that a much larger sample than the one included in the present study would allow the identification of a subsample of female youths who display high rates of conduct problems; this would permit an adequately powered test of the G×E hypotheses addressed in this study. Second, gender differences in conduct problem rates may reflect beliefs that aggression and other antisocial behaviors are more acceptable and effective coping tactics for male than female youths. It is also consistent with other studies in which male youths were found to be more likely than female youths to respond to life stress by losing inhibitory controls and expressing anger and frustration through their behaviors (Brody, Chen, et al., 2006; Hetherington, 1989; Rutter, 1990). The results of the present study extend these findings by demonstrating that 5-HTTLPR status may promote a stronger manifestation of conduct problems in adolescence among male than female youths when they experience negative emotions occasioned by high levels of discrimination.

The findings raise questions about the mechanisms through which G×E interactions connect perceive discrimination with the development of conduct problems. The neuroimaging research mentioned in the introduction offers one answer. This research has demonstrated that carriers of one *s* allele evince more amygdala-mediated arousal and increased cortical reactivity in response to threatening stimuli (Battaglia et al., 2005; Hensch et al., 2006). The *s* allele also has been linked with preferential attention toward and difficulty disengaging from threat-related stimuli (Beevers et al., 2009; Osinsky et al., 2008). Together, this literature suggests that male youths who perceive high levels of discrimination and carry one or two copies of the *s* allele develop heightened anticipation of and reactivity to both subtle and overt discrimination. This heightened reactivity in response to perceived discrimination

may, for some youths, engender high levels of negative affectivity that contribute to engagement in and justification of antisocial behavior. This process is consistent with Franklin's (1993) observation that African Americans' inner vigilance toward racial slights can create a state of constant watchfulness that leads to chronic stress and its accompanying emotional costs. The research in this report suggests that this process is most likely to occur among male African Americans who carry an *s* allele. These conjectures await empirical analysis.

Some limitations of the research design should be noted. Only one genetic polymorphism was examined, which does not represent all the variation that could place youths at risk for conduct problems. Many genetic variants may alter risk, the expression of which may only emerge under particular contextual conditions. A corollary of this limitation is the perception that genetic variation confers only risk. Genetic effects may also be protective, and that which is conceptualized as a risk-promoting genetic effect may actually be the absence of protective genes. In this study, the *ll* genotype for 5-HTTLPR protected youths who perceived high levels of discrimination from developing conduct problems. This is an important finding and one that should be explored in future research. Research also is needed to examine the processes that interact with genetic status that have been found to protect youths from the psychological costs of discrimination, such as parenting practices, affiliations with prosocial peers, school efficacy, and racial identity (Brody, Chen, et al., 2006; Simons et al., 2006), to refine understanding of the processes and conditions through which discrimination compromises children's and adolescents' development. Future research with larger samples will be able to address this issue. These limitations notwithstanding, the present study demonstrates the ways in which perceived exposure to racial discrimination and 5-HTTLPR status combine to create different trajectories of conduct problems among male African American youths during adolescence.

Acknowledgments

This research was supported by Award Number 1P30DA027827 from the National Institute on Drug Abuse and Awards Numbers 2R01AA012768 and 3R01AA012768-09S1 from the National Institute on Alcohol Abuse and Alcoholism. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute on Drug Abuse, the National Institute on Alcohol Abuse and Alcoholism, or the National Institutes of Health.

References

- Battaglia M, Ogliaari A, Zanoni A, Citterio A, Pozzoli U, Giorda R, et al. Influence of the serotonin transporter promoter gene and shyness on children's cerebral responses to facial expressions. *Archives of General Psychiatry*. 2005; 62:85–94.10.1001/archpsyc.62.1.85 [PubMed: 15630076]
- Beevers CG, Wells TT, Ellis AJ, McGeary JE. Association of the serotonin transporter gene promoter region (5-HTTLPR) polymorphism with biased attention for emotional stimuli. *Journal of Abnormal Psychology*. 2009; 118:670–681.10.1037/a0016198 [PubMed: 19685963]
- Belsky J, Bakermans-Kranenburg MJ, van IJzendoorn MH. For better *and* for worse: Differential susceptibility to environmental influences. *Current Directions in Psychological Science*. 2007; 16:300–304.10.1111/j.1467-8721.2007.00525.x
- Boatright, SR. The Georgia county guide. 24. Athens, GA: Center for Agribusiness and Economic Development; 2005.
- Bradley SL, Dodelzon K, Sandhu HK, Philibert RA. Relationship of serotonin transporter gene polymorphisms and haplotypes to mRNA transcription. *American Journal of Medical Genetics: Part B. Neuropsychiatric Genetics*. 2005; 136:58–61.10.1002/ajmg.b.30185
- Brody, GH.; Beach, SRH.; Philibert, RA. A contextual-genetics approach to adolescent drug use and sexual risk behavior. In: Thomas, YF.; Price, LN., editors. *Drug use trajectories among minority youth*. New York, NY: Springer; (in press)

- Brody GH, Chen Y-f, Murry VM, Ge X, Simons RL, Gibbons FX, et al. Perceived discrimination and the adjustment of African American youths: A five-year longitudinal analysis with contextual moderation effects. *Child Development*. 2006; 77:1170–1189.10.1111/j.1467-8624.2006.00927.x [PubMed: 16999791]
- Brody GH, Murry VM, Gerrard M, Gibbons FX, Molgaard V, McNair LD, et al. The Strong African American Families program: Translating research into prevention programming. *Child Development*. 2004; 75:900–917.10.1111/j.1467-8624.2004.00713.x [PubMed: 15144493]
- Brody GH, Murry VM, Kogan SM, Gerrard M, Gibbons FX, Molgaard V, et al. The Strong African American Families program: A cluster-randomized prevention trial of long-term effects and a mediational model. *Journal of Consulting and Clinical Psychology*. 2006; 74:356–366.10.1037/0022-006x.74.2.356 [PubMed: 16649880]
- Bronfenbrenner U, Ceci SJ. Nature-nurture reconceptualized in developmental perspective: A bioecological model. *Psychological Review*. 1994; 101:568–586.10.1037/0033-295X.101.4.568 [PubMed: 7984707]
- Brummett BH, Boyle SH, Siegler IC, Kuhn CM, Ashley-Koch A, Jonassaint CR, et al. Effects of environmental stress and gender on associations among symptoms of depression and the serotonin transporter gene linked polymorphic region (5-HTTLPR). *Behavior Genetics*. 2008; 38:34–43.10.1007/s10519-007-9172-1 [PubMed: 17955359]
- Bryk, AS.; Raudenbush, SW. Hierarchical linear models: Applications and data analysis methods. Newbury Park, CA: Sage; 1992.
- Caspi A, Hariri AR, Holmes A, Uher R, Moffitt TE. Genetic sensitivity to the environment: The case of the serotonin transporter gene and its implications for studying complex diseases and traits. *American Journal of Psychiatry*. 2010; 167:509–527.10.1176/appi.ajp.2010.09101452 [PubMed: 20231323]
- Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, et al. Influence of life stress on depression: Moderation by a polymorphism in the 5-HTT gene. *Science*. 2003 Jul 18;301:386–389.10.1126/science.1083968 [PubMed: 12869766]
- Cicchetti D, Blender JA. A multiple-levels-of-analysis perspective on resilience: Implications for the developing brain, neural plasticity, and preventive interventions. *Annals of the New York Academy of Sciences*. 2006; 1094:248–258.10.1196/annals.1376.029 [PubMed: 17347356]
- Cicchetti D, Tucker D. Development and self-regulatory structures of the mind. *Development and Psychopathology*. 1994; 6:533–549.10.1017/S0954579400004673
- Clark R, Anderson NB, Clark VR, Williams DR. Racism as a stressor for African Americans: A biopsychosocial model. *American Psychologist*. 1999; 54:805–816.10.1037/0003-066X.54.10.805 [PubMed: 10540593]
- Davison, GC.; Neale, JM.; Blankstein, KR.; Flett, GL. Abnormal psychology. 2. Ontario, Canada: Wiley Canada; 2005.
- DuBois DL, Burk-Braxton C, Swenson LP, Tevendale HD, Hardesty JL. Race and gender influences on adjustment in early adolescence: Investigation of an integrative model. *Child Development*. 2002; 73:1573–1592.10.1111/1467-8624.00491 [PubMed: 12361320]
- Eley TC, Sugden K, Corsico A, Gregory AM, Sham P, McGuffin P, et al. Gene-environment interaction analysis of serotonin system markers with adolescent depression. *Molecular Psychiatry*. 2004; 9:908–915.10.1038/sj.mp.4001546 [PubMed: 15241435]
- Elliott, DS.; Ageton, SS.; Huizinga, D. Explaining delinquency and drug use. Beverly Hills, CA: Siegel; 1985.
- Farber, EA.; Egeland, B. Invulnerability among abused and neglected children. In: Anthony, EJ.; Cohler, BJ., editors. *The invulnerable child*. New York: Guilford Press; 1987. p. 253-288.
- Fordham S, Ogbu JU. Black students' school success: Coping with the burden of "acting White. *Urban Review*. 1986; 18:176–206.10.1007/BF01112192
- Franklin, AJ. The invisibility syndrome. *Family Therapy Networker*; 1993 Jul-Aug. p. 32-39.
- Gibbons FX, Gerrard M, Cleveland MJ, Wills TA, Brody GH. Perceived discrimination and substance use in African American parents and their children: A panel study. *Journal of Personality and Social Psychology*. 2004; 86:517–529.10.1037/0022-3514.86.4.517 [PubMed: 15053703]

- Gunthert KC, Conner TS, Armeli S, Tennen H, Covault J, Kranzler HR. Serotonin transporter gene polymorphism (5-HTTLPR) and anxiety reactivity in daily life: A daily process approach to gene-environment interaction. *Psychosomatic Medicine*. 2007; 69:762–768.10.1097/PSY.0b013e318157ad42 [PubMed: 17942837]
- Hammen, C. Risk and protective factors for children of depressed parents. In: Luthar, SS., editor. *Resilience and vulnerability: Adaptation in the context of childhood adversities*. New York: Cambridge University Press; 2003.
- Hariri AR, Drabant EM, Munoz KE, Kolachana BS, Mattay VS, Egan MF, et al. A susceptibility gene for affective disorders and the response of the human amygdala. *Archives of General Psychiatry*. 2005; 62:146–152. Retrieved from <http://archpsyc.ama-assn.org/cgi/content/abstract/62/2/146>. [PubMed: 15699291]
- Hariri AR, Drabant EM, Weinberger DR. Imaging genetics: Perspectives from studies of genetically driven variation in serotonin function and corticolimbic affective processing. *Biological Psychiatry*. 2006; 59:888–897.10.1016/j.biopsych.2005.11.005 [PubMed: 16442081]
- Hariri AR, Mattay VS, Tessitore A, Fera F, Weinberger DR. Neocortical modulation of the amygdala response to fearful stimuli. *Biological Psychiatry*. 2003; 53:494–501.10.1016/S0006-3223(02)01786-9 [PubMed: 12644354]
- Hariri AR, Mattay VS, Tessitore A, Kolachana BS, Fera F, Goldman D, et al. Serotonin transporter genetic variation and the response of the human amygdala. *Science*. 2002 Jul 19;297:400–403.10.1126/science.1071829 [PubMed: 12130784]
- Heinz A, Braus DF, Smolka MN, Wrase J, Puls I, Hermann D, et al. Amygdala-prefrontal coupling depends on a genetic variation of the serotonin transporter. *Nature Neuroscience*. 2005; 8:20–21.10.1038/nn1366
- Heinz A, Smolka MN, Braus DF, Wrase J, Beck A, Flor H, et al. Serotonin transporter genotype (5-HTTLPR): Effects of neutral and undefined conditions on amygdala activation. *Biological Psychiatry*. 2007; 61:1011–1014.10.1016/j.biopsych.2006.08.019 [PubMed: 17157270]
- Hensch T, Wargelius HL, Herold U, Lesch KP, Orelund L, Brocke B. Further evidence for an association of 5-HTTLPR with intensity dependence of auditory-evoked potentials. *Neuropsychopharmacology*. 2006; 31:2047–2054.10.1038/sj.npp.1301020 [PubMed: 16421513]
- Hetherington EM. Coping with family transitions: Winners, losers, and survivors. *Child Development*. 1989; 60:1–14.10.1111/1467-8624.ep7250664 [PubMed: 2649320]
- Howe GW, Beach SRH, Brody GH. Microtrial methods for translating gene-environment dynamics into preventive interventions. *Prevention Science*. (in press).
- Isenberg N, Silbersweig D, Engelien A, Emmerich S, Malavade K, Beattie B, et al. Linguistic threat activates the human amygdala. *Proceedings of the National Academy of Sciences of the USA*. 1999; 96:10456–10459. Retrieved from <http://www.pnas.org/content/96/18/10456.abstract>. [PubMed: 10468630]
- Kendler KS, Eaves LJ. Models for the joint effect of genotype and environment on liability to psychiatric illness. *American Journal of Psychiatry*. 1986; 143:279–289. Retrieved from <http://ajp.psychiatryonline.org/cgi/content/abstract/143/3/279>. [PubMed: 3953861]
- Kim-Cohen J, Gold AL. Measured gene-environment interactions and mechanisms promoting resilient development. *Current Directions in Psychological Science*. 2009; 18:138–142.10.1111/j.1467-8721.2009.01624.x
- Landrine H, Klonoff EA. The Schedule of Racist Events: A measure of racial discrimination and a study of its negative physical and mental health consequences. *Journal of Black Psychology*. 1996; 22:144–168.10.1177/00957984960222002
- Lazary J, Lazary A, Gonda X, Benko A, Molnar E, Juhasz G, et al. New evidence for the association of the serotonin transporter gene (SCL6A4) haplotypes, threatening life events, and depressive phenotype. *Biological Psychiatry*. 2008; 64:498–504.10.1016/j.biopsych.2008.03.030 [PubMed: 18486105]
- Lerner RM. Changing organism-context relations as the basic process of development: A developmental context and perspective. *Developmental Psychology*. 1991; 27:27–32.10.1037/0012-1649.27.1.27

- Luthar, SS. Resilience in development: A synthesis of research across five decades. In: Cicchetti, D.; Cohen, DJ., editors. *Developmental psychopathology: Vol. 3. Risk, disorder, and adaptation*. 2. Hoboken, NJ: Wiley; 2006. p. 739-795.
- Moffitt TE. Adolescence-limited and life-course persistent antisocial behavior: A developmental taxonomy. *Psychological Review*. 1993; 100:674–701.10.1037/0033-295X.100.4.674 [PubMed: 8255953]
- Moffitt TE, Caspi A, Rutter ML. Measured gene-environment interactions in psychopathology: Concepts, research strategies, and implications for research, intervention, and public understanding of genetics. *Perspectives on Psychological Science*. 2006; 1:5–27.10.1111/j.1745-6916.2006.00002.x
- Munafò MR, Brown SM, Hariri AR. Serotonin transporter (5-HTTLPR) genotype and amygdala activation: A meta-analysis. *Biological Psychiatry*. 2008; 63:852–857.10.1016/j.biopsych.2007.08.016 [PubMed: 17949693]
- Osinsky R, Reuter M, Küpper Y, Schmitz A, Kozyra E, Alexander N, et al. Variation in the serotonin transporter gene modulates selective attention to threat. *Emotion*. 2008; 8(4):584–588.10.1037/a0012826 [PubMed: 18729589]
- Patrick DL, Cheadle A, Thompson DC, Diehr P, Koepsell T, Kinne S. The validity of self-reported smoking: A review and meta-analysis. *American Journal of Public Health*. 1994; 84:1086–1093.10.2105/AJPH.84.7.1086 [PubMed: 8017530]
- Philibert RA, Zadorozhnyaya O, Beach SRH, Brody GH. Comparison of the genotyping results using DNA obtained from blood and saliva. *Psychiatric Genetics*. 2008; 18:275–281.10.1097/YPG.0b013e3283060f81 [PubMed: 19018232]
- Retz W, Freitag CM, Retz-Junginger P, Wenzler D, Schneider M, Kissling C, et al. A functional serotonin transporter promoter gene polymorphism increases ADHD symptoms in delinquents: Interaction with adverse childhood environment. *Psychiatry Research*. 2008; 158:123–131.10.1016/j.psychres.2007.05.004 [PubMed: 18155777]
- Risch N, Herrell R, Lehner T, Liang KY, Eaves LJ, Hoh J, et al. Interaction between the serotonin transporter gene (5-HTTLPR), stressful life events, and risk of depression: A meta-analysis. *Journal of the American Medical Association*. 2009; 301:2462–2471.10.1001/jama.2009.878 [PubMed: 19531786]
- Rutter, ML. Psychosocial resilience and protective mechanisms. In: Rolf, J.; Masten, AS.; Cicchetti, D.; Nuechterlein, K.; Weintraub, S., editors. *Risk and protective factors in the development of psychopathology*. New York: Cambridge University Press; 1990. p. 181-215.
- Rutter ML, Silberg J. Gene-environment interplay in relation to emotional and behavioral disturbance. *Annual Review of Psychology*. 2002; 53:463–490.10.1146/annurev.psych.53.100901.135223
- Sellers RM, Copeland-Linder N, Martin PP, Lewis RLH. Racial identity matters: The relationship between racial discrimination and psychological functioning in African American adolescents. *Journal of Research on Adolescence*. 2006; 16:187–216.10.1111/j.1532-7795.2006.00128.x
- Simons RL, Chen Y-f, Stewart EA, Brody GH. Incidents of discrimination and risk for delinquency: A longitudinal test of strain theory with an African American sample. *Justice Quarterly*. 2003; 20:827–854.10.1080/07418820300095711
- Simons RL, Murry VM, McLoyd VC, Lin KH, Cutrona CE, Conger RD. Discrimination, crime, ethnic identity, and parenting as correlates of depressive symptoms among African American children: A multilevel analysis. *Development and Psychopathology*. 2002; 14:371–393.10.1017/S0954579402002109 [PubMed: 12030697]
- Simons RL, Simons LG, Burt CH, Drummund H, Stewart EA, Brody GH, et al. Supportive parenting moderates the effect of discrimination upon anger, hostile view of relationships, and violence among African American boys. *Journal of Health and Social Behavior*. 2006; 47:373–389.10.1177/002214650604700405 [PubMed: 17240926]
- Singer, JD.; Willett, JB. *Applied longitudinal data analysis: Modeling change and event occurrence*. New York: Oxford University Press; 2003.
- Slavin LA, Rainer KL, McCreary ML, Gowda KK. Toward a multicultural model of the stress process. *Journal of Counseling and Development*. 1991; 70:156–163. Retrieved from <http://aca.metapress.com/openurl.asp?genre=journal&eissn=1556-6676>.

- Stein MB, Schork NJ, Gelernter J. Gene-by-environment (serotonin transporter and childhood maltreatment) interaction for anxiety sensitivity, an intermediate phenotype for anxiety disorders. *Neuropsychopharmacology*. 2008; 33:312–319.10.1038/sj.npp.1301422 [PubMed: 17460615]
- Surtees PG, Wainwright NWJ, Willis-Owen SAG, Luben R, Day NE, Flint J. Social adversity, the serotonin transporter (5-HTTLPR) polymorphism and major depressive disorder. *Biological Psychiatry*. 2006; 59:224–229.10.1016/j.biopsych.2005.07.014 [PubMed: 16154545]
- Szalacha, LA.; Erkut, S.; García Coll, CT.; Fields, JP.; Alarcón, O.; Ceder, I. Perceived discrimination and resilience. In: Luthar, SS., editor. *Resilience and vulnerability: Adaptation in the context of childhood adversities*. New York: Cambridge University Press; 2003. p. 414-435.
- Taylor, DM.; Wright, SC.; Porter, LE. Dimensions of perceived discrimination: The personal/group discrimination discrepancy. In: Zanna, MP.; Olson, JM., editors. *The psychology of prejudice: The Ontario Symposium*. Vol. 7. Hillsdale, NJ: Erlbaum; 1993. p. 233-255.
- Taylor SE, Way BM, Welch WT, Hilmert CJ, Lehman BJ, Eisenberger NI. Early family environment, current adversity, the serotonin transporter promoter polymorphism, and depressive symptomatology. *Biological Psychiatry*. 2006; 60:671–676.10.1016/j.biopsych.2006.04.019 [PubMed: 16934775]
- Teichner G, Golden CJ. The relationship of neuropsychological impairment to conduct disorder in adolescence: A conceptual review. *Aggression and Violent Behavior*. 2000; 5:509–528.10.1016/S1359-1789(98)00035-4
- Thoits PA. On merging identity theory and stress research. *Social Psychology Quarterly*. 1991; 54:101–112. Retrieved from <http://www.jstor.org/stable/2786929>.
- Wicks-Nelson, R.; Israel, AC. *Behavior disorders of childhood*. 5. Upper Saddle River, NJ: Pearson Education; 2003.
- Wilhelm K, Mitchell PB, Niven H, Finch AW, Wedgwood L, Scimone A, et al. Life events, first depression onset and the serotonin transporter gene. *British Journal of Psychiatry*. 2006; 188:210–215.10.1192/bjp.bp.105.009522 [PubMed: 16507960]
- Zalsman G, Huang Y-y, Oquendo MA, Burke AK, Hu X-z, Brent DA, et al. Association of a triallelic serotonin transporter gene promoter region (5-HTTLPR) polymorphism with stressful life events and severity of depression. *American Journal of Psychiatry*. 2006; 163:1588–1593.10.1176/appi.ajp.163.9.1588 [PubMed: 16946185]
- Zucker, RA.; Wong, MW.; Puttler, LI.; Fitzgerald, HE. Resilience and vulnerability among sons of alcoholics: Relationship to developmental outcomes between early childhood and adolescence. In: Luthar, SS., editor. *Resilience and vulnerability: Adaptation in the context of childhood adversities*. New York: Cambridge University Press; 2003. p. 76-103.

Figure 1. Results of the analysis for the total sample. Slopes of conduct problems for levels of perceived discrimination, ranging from -3 to $+3$ standard deviations from the sample mean, plotted separately for youths with the *ll* genotype and youths with the *ss* or *sl* genotype.

Figure 2.

Results of the analysis for male youths only. Slopes of conduct problems for levels of perceived discrimination, ranging from -3 to $+3$ standard deviations from the sample mean, plotted separately for male youths with the *ll* genotype and male youths with the *ss* or *sl* genotype.

Table 1

Unstandardized Regression Coefficients, Slopes for Conduct Problems

Variables	Full Sample		Male Youths		Female Youths	
	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
Control variables						
Monthly per capita income	0.00	0.00	0.00	0.00	0.00	0.00
Parental education	-0.06	-0.05	-0.13	-0.16	0.00	0.00
Youth gender	-0.75**	-0.68**				
Independent variables						
SRE	0.63*	0.12	1.20**	0.14	0.13 [†]	0.07
5-HTTLPR status	0.40*	0.38*	0.68 [†]	-0.68 [†]	-0.01	-0.01
Two-way interaction term, SRE × 5-HTTLPR status		1.05**		1.79**		0.16
Constant	1.41*	1.27*	0.59	0.07	0.09	0.07
Random effect for slope	1.72**	1.45**	2.84**	2.14**	0.19**	0.18**

Note. SRE = Schedule of Racist Events.

[†] $p < .10$.* $p < .05$.** $p < .01$. All tests two-tailed.