Published in final edited form as:

Org Lett. 2013 February 15; 15(4): 840–843. doi:10.1021/ol303531m.

Rhodium-catalyzed Chemo- and Regioselective Cross-dimerization of Two Terminal Alkynes

Hua-Dong Xu^{*,†,‡}, Ren-Wei Zhang[†], Xiaoxun Li^{†,‡}, Suyu Huang^{†,‡}, Weiping Tang^{†,‡}, and Wen-Hao Hu[†]

[†]Shanghai Engineering Research Centre of Molecular Therapeutics and New Drug Development, East China Normal University, Shanghai, 200062, China

[‡]School of Pharmaceutical Engineering and Life Science, Changzhou University, Changzhou, Jiangsu Province, 213164, China

^{††}School of Pharmacy, University of Wisconsin, Madison, Wisconsin 53705, United States

Abstract

Cross-dimerization of terminal arylalkynes and terminal propargylic alcohols/amides has been achieved in the effect of a rhodium catalyst. This method features high chemo- and regioselectivities rendering convenient and atom economical access to functionalized enynes.

Conjugated 1,3-enynes are an important class of building blocks with diverse applications in organic chemistry¹ and material science.² They are generally prepared from transition metal catalyzed cross-coupling of terminal alkyne and pre-activated alkenes,³ Wittig olefination of conjugated alkynals,⁴ or dehydration of propargylic alcohols.⁵ On the other hand, the more atom economical method⁶ for the preparation of conjugated enynes, namely direct hydroalkynylation across C-C triple bonds or dimerization of alkynes, has found fewer applications due to the challenging issues of chemo-, regio-, and stereoselectivities as illustrated in Scheme 1 for two terminal alkynes.⁷ One way to overcome these obstacles is to take advantage of the steric and/or electronic differences between the two alkynes to render chemo- and regioselectivity. Important progress has been made along this direction.⁸ However, cross-dimerization of two different terminal alkynes with high chemo-, regio-, and stereoselectivity awaits further disclosure. We herein report an efficient and selective cross-dimerization of two different terminal alkynes in a head-to-tail fashion for the preparation of enynes **3a/3a'** (Scheme 1).

Trost's group has extensively studied the addition of terminal alkynes to various electron-deficient internal alkynes using a palladium catalyst.^{8m,n} In the absence of the electron-deficient internal alkyne, the terminal alkyne underwent homodimerization in a head-to-tail

*huadongxu@gmail.com.

Supporting Information Available. Experimental procedures along with characterizing data and copies of NMR spectra is available free of charge via the Internet at <http://pubs.acs.org>.

fashion to form enyne **5a/5a'**. Interestingly, selective cross-dimerization of two terminal alkynes was found to be possible when propargylic alcohols and excess 1,3-enynes were employed.^{8l} The head-to-tail products **3a/3a'** were obtained in synthetically useful yields. Recently, Li's group disclosed a rhodium-catalyzed homodimerization of propargyl tosylamides to give the head-to-tail product (**5a/5a'**) in high selectivity.^{8b} Miura group reported that head-to-head cross-dimerization product (**4a/4a'**) was selectively prepared from two sterically different terminal alkynes.^{8f}

During our development of rhodium-catalyzed intermolecular [5+2] cycloaddition of 3-acyloxy-1,4-enynes and alkynes,⁹ we found homodimerization of propargylic alcohols occurred in the absence of 3-acyloxy-1,4-enynes. We then explored the possibility of heterodimerizing phenylacetylene **1a** and propargylic alcohol **2a** employing a rhodium catalyst. We first examined the ligand effect using [Rh(COD)Cl]₂ as the rhodium precursor (Table 1). Simple Ph₃P ligand afforded the enyne products in 65% yield with a good selectivity for **3a** (entry 1). Replacement of Ph₃P by (*p*-CF₃C₆H₄)₃P or (3,5-CF₃C₆H₄)₃P resulted in lower yields and selectivity (entry 4). For phosphites or more electron-deficient phosphine ligands, the catalyst showed no activity (entries 6-8). When the electron-rich cyclohexylphosphine ligand was employed, the reaction proceeded in lower yield and selectivity (entry 9). With the exception of dppf (entries 10 and 11), most other bidentate ligands appeared to be ineffective (entries 12-16). Ligand dppf actually delivered the highest regioselectivity. Yields of the desired product **3aa**, however, are moderate under these conditions. Without any ligand, the reaction didn't proceed at all (entry 17). Wilkinson catalyst yielded similar results as [Rh(COD)Cl]₂/Ph₃P combination (entries 18 and 19).

Other reaction parameters were explored using the combination of Ph₃P and [Rh(COD)Cl]₂ (Table 2). Since the homo dimerization product **5aa** is much more polar than heterodimers **3aa** and **4aa**, it can be easily removed by column chromatography. Only the ratio of **3aa** and **4aa** was examined. Higher temperature could accelerate the reaction and a slightly higher yield with lower selectivity was observed (entry 1 vs 2). Increasing the ratio of **1a/2a** from 1:2 to 2:3 was beneficial for the yield (entry 4 vs 5). When the reaction was performed in THF, the highest regioselectivity was observed. The yield, however, dropped to 55% (entry 7). While the halogenated solvents behave similarly, DCM was the best in terms of yield and selectivity (entries 8 and 9). Poor yields were obtained with lower catalyst loadings (entries 10, 11). The optimal metal/ligand ratio appeared to be 1:2 (entry 13).

With the optimized conditions in hand, the scope of terminal alkynes was then explored (Figure 1). Various substituents on the phenyl ring of the donor alkyne could be accommodated. Arylacetylenes with para, meta and ortho substituents gave comparable results. Aldehyde, bromide, amide, and even free amino group could be tolerated. The 4-nitro group impaired the ratio of regioisomers **3ca/4ca** to 4:1 in favor of the head-to-tail dimer **3ca**. Interestingly, 3-indole acetylene furnished enyne **3ja** in 42% yield without the observation of the head-to-head dimer. The free hydroxyl group is not required as substrate **2c** also participate in the cross-dimerization. On the other hand, the steric effect on the regioselectivity is pronounced as only head-to-head cross-dimerization product **4ab** was detected by NMR, while secondary propargylic alcohols gave a mixture of homo- and heterodimers with low selectivity. It should be noted that a complex mixture was observed when propargylic alcohol **2a** was reacted with terminal alkyl substituted acetylenes such as 1-heptyne. Similarly, when phenylacetylene **1a** was reacted with homo propargylic alcohols, no significant selectivity was observed, implying that the chelating effect might not play an important role in this reaction and the inductive effect may operate (see the SI Scheme 2 for details). On the other hand, we were pleased to find that propargylic sulfonamide **2d** afforded head-to-tail heterodimer **3ad-3jd** with generally higher yields and higher selectivity

than their corresponding propargylic alcohols. A moderate yield of product **3ae** was obtained from propargylic carbamate **2e**.

The mechanism of this chemoselective cross-dimerization of terminal alkynes involves oxidative addition of rhodium to the arylacetylene C-H bond followed by hydroalkynylation of propargylic alcohols, ethers, or amides.¹⁰ Further study is required to elucidate the details of this reaction, especially the chemo- and regioselectivities.

In summary we have discovered a facile method for the cross-dimerization of arylacetylenes and propargylic alcohols, ethers, or amides via the atom economical direct hydroalkynylation reaction promoted by a rhodium catalyst. We found that the chemo- and regioselectivities were dependent on the steric and electronic nature of both terminal alkynes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

H.-D. Xu thanks the Natural Science Foundation of China (21002032 and 21272077), Shanghai Pujiang Program (11PJ1403100), Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD), and W. T. thanks NIH (R01 GM088285) and University of Wisconsin for financial support.

References

- (1) (a). Werness JB, Tang W. *Org. Lett.* 2011; 13:3664. [PubMed: 21688862] (b) Zhang W, Zheng S, Liu N, Werness JB, Guzei IA, Tang W. *J. Am. Chem. Soc.* 2010; 132:3664. [PubMed: 20192183] (c) Zhang W, Xu H, Xu H, Tang W. *J. Am. Chem. Soc.* 2009; 131:3832. [PubMed: 19245198] (d) Zhang W, Werness JB, Tang W. *Org. Lett.* 2008; 10:2023. [PubMed: 18435544] (e) Wessig P, Müller G. *Chem. Rev.* 2008; 108:2051. [PubMed: 18479169] (f) Kim H, Lee H, Lee D, Kim S, Kim D. *J. Am. Chem. Soc.* 2007; 129:2269. [PubMed: 17279755] (g) Kang B, Kim D.-h. Do Y, Chang S. *Org. Lett.* 2003; 5:3041. [PubMed: 12916976] (h) Schaus SE, Cavalieri D, Myers AG. *Proc. Natl. Acad. Sci. U. S. A.* 2001; 98:11075. [PubMed: 11562456] (i) Stang, PJD. *Francocs Modern Acetylene Chemistry*. VCH; New York, New York:
- (2) (a). Liu Y, Nishiura M, Wang Y, Hou Z. *J. Am. Chem. Soc.* 2006; 128:5592. [PubMed: 16637599] (b) Moonen NNP, Boudon C, Gisselbrecht J-P, Seiler P, Gross M, Diederich F. *Angew. Chem. Int. Ed.* 2002; 41:3044. (c) Campbell K, Kuehl CJ, Ferguson MJ, Stang PJ, Tykwinski RR. *J. Am. Chem. Soc.* 2002; 124:7266. [PubMed: 12071724]
- (3) (a). Ranu BC, Adak L, Chattopadhyay K. *J. Org. Chem.* 2008; 73:5609. [PubMed: 18558765] (b) Thongsornkleeb C, Danheiser RL. *J. Org. Chem.* 2005; 70:2364. [PubMed: 15760233] (c) Negishi E-I, Anastasia L. *Chem. Rev.* 2003; 103:1979. [PubMed: 12744698] (d) Takeuchi R, Tanabe K, Tanaka S. *J. Org. Chem.* 2000; 65:1558. [PubMed: 10814124] (e) Sonogashira K, Tohda Y, Hagihara N. *Tetrahedron Lett.* 1975; 16:4467.
- (4) (a). Deussen H-J, Jeppesen L, Schärer N, Junager F, Bentzen B, Weber B, Weil V, Mozer SJ, Sauerberg P. *Org. Process Res. Dev.* 2004; 8:363. (b) Sauerberg P, Bury PS, Mogensen JP, Deussen H-J, Pettersson I, Fleckner J, Nehlin J, Frederiksen KS, Albrektsen T, Din N, Svensson LA, Ynddal L, Wulff EM, Jeppesen L. *J. Med. Chem.* 2003; 46:4883. [PubMed: 14584939]
- (5). Yan W, Ye X, Akhmedov NG, Petersen JL, Shi X. *Org. Lett.* 2012; 14:2358. [PubMed: 22524511]
- (6). Trost BM. *Science.* 1991; 254:1471. [PubMed: 1962206]
- (7) (a). Jahier C, Zatulochnaya OV, Zvyagintsev NV, Ananikov VP, Gevorgyan V. *Org. Lett.* 2012; 14:2846. [PubMed: 22591008] (b) Sun SK, J. Luo Y, Zhang L. *Synlett.* 2012; 23:54. [PubMed: 22904604] (c) Dash AK, Eisen MS. *Org. Lett.* 2000; 2:737. [PubMed: 10814423] (d) Yi CS, Liu N. *Organometallics.* 1998; 17:3158. (e) Akita MY, H. Nakamura A. *Bull. Chem. Soc. Jpn.* 1984; 57:480. (f) Carlton L, Read G. *J. Chem. Soc., Perkin Trans.* 1978; 1:1631. 0.

- (8) (a). Trost BM, Taft BR, Masters JT, Lumb J-P. J. Am. Chem. Soc. 2011; 133:8502. [PubMed: 21557627] (b) Peng HM, Zhao J, Li X. Adv. Synth. Catal. 2009; 351:1371. (c) Matsuyama N, Hirano K, Satoh T, Miura M. J. Org. Chem. 2009; 74:3576. [PubMed: 19354238] (d) Ogata K, Murayama H, Sugawara J, Suzuki N, Fukuzawa S-I. J. Am. Chem. Soc. 2009; 131:3176. [PubMed: 19226123] (e) Matsuyama N, Tsurugi H, Satoh T, Miura M. Adv. Synth. Catal. 2008; 350:2274. (f) Katagiri T, Tsurugi H, Satoh T, Miura M. Chem. Commun. 2008:3405. (g) Ogata K, Oka O, Toyota A, Suzuki N, Fukuzawa S-i. Synlett. 2008:2663. (h) Tsukada N, Ninomiya S, Aoyama Y, Inoue Y. Pure Appl. Chem. 2008; 80:1161. (i) Tsukada N, Ninomiya S, Aoyama Y, Inoue Y. Org. Lett. 2007; 9:2919. [PubMed: 17602565] (j) Katayama H, Yari H, Tanaka M, Ozawa F. Chem. Commun. 2005; 0:4336. (k) Trost BM, Gunzner JL, Yasukata T. Tetrahedron Lett. 2001; 42:3775. (l) Trost BM, McIntosh MC. Tetrahedron Lett. 1997; 38:3207. (m) Trost BM, Sorum MT, Chan C, Ruehter G. J. Am. Chem. Soc. 1997; 119:698. (n) Trost BM, Chan C, Ruehter G. J. Am. Chem. Soc. 1987; 109:3486. (o) Trost BM, Hachiya I, McIntosh MC. Tetrahedron Lett. 1998; 39:6445.
- (9). Shu XZ, Li X, Shu D, Huang S, Schienebeck CM, Zhou X, Robichaux PJ, Tang W. J. Am. Chem. Soc. 2012; 134:5211. [PubMed: 22364320]
- (10) (a). For oxidative addition of various transition metals including Rh(I) to the acetylene C-H bonds and followed by hydroalkynylation please see Villarino L, García-Fandiño R, López F, Mascareñas JL. Org. Lett. 2012; 14:2996. [PubMed: 22667688] ; Sawano T, Ashouri A, Nishimura T, Hayashi T. J. Am. Chem. Soc. 2012; 134:18936. [PubMed: 23131028] and references cited therein.

Figure 1.

Rhodium-Catalyzed Cross-dimerization of Arylacetylenes and Propargylic Alcohols, Ethers, or Amides^a

^aReaction conditions: arylacetylene 1 (0.5 mmol), propargylic alcohol/ether/amide 2 (0.75 mmol), [Rh(COD)Cl]₂ (5 mol %), Ph₃P (20 mol %), DCM (1 mL), 40 °C, overnight. The ratio of two heterodimers was determined by ¹H NMR of the purified mixtures of isomers after column chromatography and the yields were the isolated yield of two isomers.

Scheme 1.
Possible Products from Dimerization of Two Terminal Alkynes

Table 1

Effect of Ligands^a

entry	ligand	yield (%) ^f , ratio ^h
1	PPh ₃	65% (3aa : 4aa : 5aa = 9:1:2) 62% ^g
2 ^e	PPh ₃	54% (3aa : 4aa : 5aa = 11:1:2.6)
3 ^b	PPh ₃	62% (3aa : 4aa : 5aa = 10:1:1.5) 57% ^g
4	(p-CF ₃ C ₆ H ₄) ₃ P	53% (3aa : 4aa : 5aa = 7:1:2)
5	(3,5-CF ₃ C ₆ H ₄) ₃ P	50% (3aa : 4aa : 5aa = 6:1:1)
6	(CF ₃ CH ₂ O) ₃ P	0%
7	(hfip) ₃ P	0%
8	(C ₆ F ₅) ₃ P	0%
9	Cy ₃ P	40% (3aa : 4aa = 5:1)
10	dppf	52% (3aa : 4aa = 20:1)
11 ^e	dppf	32% (3aa : 4aa = 20:1)
12	dppm	Trace
13	dppe	Trace
14	dppp	Trace
15	dppb	Trace
16	BINAP	33% (3aa : 4aa = 8.5:1)
17 ^c	-	0%
18 ^{d,e}	-	65% (3aa : 4aa : 5aa = 9:1:3)
19d	-	55% (3aa : 4aa : 5aa = 4:1:2)

^aGeneral conditions: **1a** (0.2 mmol), **2a** (0.4 mmol), [Rh(COD)Cl]₂ (5 mol%) and ligand (20 mol %), DCM (1 mL), 40 °C overnight.

^b30 mol% [Rh(COD)Cl]₂ was used

^cNo ligand

^dWilkinson catalyst was used

^eroom temperature

^fNMR yield

^gIsolated yield of **3aa** and **4aa**

^hDetermined by ¹H NMR analysis. hfip = hexafluoroisopropanol

Table 2Optimization of Reaction Conditions^a

entry	1a:2a	x mol % [Rh(COD)Cl] ₂ / y mol % PPh ₃	solvent/T (°C)	yield ^b , (3aa:4aa) ^c
1	1:2	5/20 mol%	DCM/RT	54% (11:1)
2	1:2	5/20 mol%	DCM/40	62% (9:1)
3	1:2	5/30 mol%	DCM/40	62% (10:1)
4	1:2	5/30 mol%	DCM/RT	57% (10:1)
5	2:3	5/30 mol%	DCM/RT	65% (10:1)
6	2:3	5/30 mol%	DCM/40	70% (10:1)
7	2:3	5/30 mol%	THF/40	55% (15:1)
8	2:3	5/30 mol%	DCE/40	62% (10:1)
9	2:3	5/30 mol%	CHCl ₃ /40	65% (5:1)
10	2:3	1/6 mol%	DCM/40	30% (–)
11	2:3	2/12 mol%	DCM/40	39% (10:1)
12	2:3	5/10 mol%	DCM/40	25% (–)
13	2:3	5/20 mol%	DCM/40	73% (10:1)
14	2:3	5/60 mol%	DCM/40	61% (5:1)

^aReaction conditions are essentially the same as those for Table 1, and changes are made according to each entry^bIsolated yield.^cDetermined by ¹H NMR of a partially purified mixture of **3aa** and **4aa** after a short column.