CORRESPONDENCE

Postoperative Care and Follow-Up After Coronary Stenting

by Prof. Dr. med. Tienush Rassaf, PD Dr. med. Stephan Steiner,
Prof. Dr. med. Malte Kelm in volume 5/13

Reality not Reflected

The article provides a well intentioned description, where wishful thinking is mistaken for reality—or rather, “the desired state of affairs” is mistaken for “the real state of affairs”—and does not reflect clinical reality. Members of the German Cardiac Society would confirm this. The indication for invasive primary diagnostic evaluation and stent implantation, as well as for follow-up after stenting, can be assumed to be directly proportional to the number of cardiac catheter laboratories, as the steadily increasing numbers show. Most specialists will be clear that in a scenario of 880 000 catheterizations in 2010—the world record per 1000 population in an industrialized country—competition and financial considerations are the main motivation, rather than guidelines or even patients. And, to remind readers, cardiac catheterization is never trivial but a serious intervention. Possible complications—even if these are rare—include arterial hemorrhage, aortic dissection, or pulmonary embolism.

DOI: 10.3238/arztebl.2013.0403a

REFERENCES

1. Rassaf T, Steiner S, Kelm M: Postoperative care and follow up after coronary stenting. *Dtsch Arztebl Int* 2013; 110(5): 72–82.

Prof. Dr. med. Ulrich Hake, Mainz
u-hake@online.de

Rehabilitation Measures as Therapeutic Cornerstones

The background and results sections of the abstract point out that most patients can return to their everyday routine a few days after a successful percutaneous coronary intervention (PCI).

The section of learning objectives also creates the impression that coronary heart disease is an accident that heals completely after successful therapy using PCI and appropriate, close, medical, and diagnostic aftercare as long as all the rules of the art of medicine are adhered to and much time and effort are expended in order to ensure compliance with the treatment and follow-up examinations.

The section of recommended clinical patient follow-up by primary care physicians and cardiologists suggests conducting clinical, apparatus-based diagnostic evaluation as well as motivating patients to change their lifestyles.

What is not mentioned even in a single word is professional support, which is crucial to help patients learn to cope with their illness (1); this includes influencing lifestyles and modifying risk factor profiles in the context of an indicated follow-up cardiac rehabilitation therapy that is covered by the health insurance companies.

In most hospitals and doctors' minds it takes a contracted injury to the heart (myocardial infarction, the larger the better) or bypass surgery in order to prompt thoughts of cardiac rehabilitation measures.

In this respect, our health insurers are more advanced than those acting at the coalface of acute care. For years, successful coronary intervention has been an acceptable indication for health insurers to cover the cost of subsequent rehabilitation therapy (2), with corresponding successes in the long-term prognosis and avoidance of repeated morbidity (3).

This fact has not become anchored in the minds of specialists in acute medicine; this article yet again shows that no mention is made of the fact that—in addition to medication, cardiac catheterization, coronary computed tomography scanning, magnetic resonance imaging, and stress echocardiography—cardiac rehabilitation should be one of the cornerstones of the therapy provided to patients after PCI.

DOI: 10.3238/arztebl.2013.0403b

REFERENCES

1. Buss U, Herrmann-Lingen C: Der Prozess der Krankheitsbewältigung. In: Rauch B, et al. (eds): *Kardiologische Rehabilitation, Standards für die Praxis nach den Leitlinien der Deutschen Gesellschaft für Prävention und Rehabilitation von Herz-Kreislaufkrankungen-DGPR*. Stuttgart Thieme Verlag 2007; 122–34.
2. Bundesarbeitsgemeinschaft für Rehabilitation – Schriftenreihe der Bundesarbeitsgemeinschaft für Rehabilitation: *Arbeitshilfe zur Rehabilitation Koronarkranker*, last accessed 17 June 1997, p. 61 ff. www.bar-frankfurt.de/fileadmin/dateiliste/publikationen/arbeitshilfen/download/Arbeitshilfe_Koronar.pdf. Last accessed on 14 May 2013.
3. Korsukéwitz C, Falk J, Lindow B: *Kardiologische Rehabilitation in Deutschland*. *Herz* 2012; 37: 12–21.
4. Rassaf T, Steiner S, Kelm M: Postoperative care and follow up after coronary stenting. *Dtsch Arztebl Int* 2013; 110(5): 72–82.

Dr. med. Markus Zirbes
Dr. med. Brigitte Buran-Kilian
Dr. med. Thomas Aupperle
Schwerpunktpraxis, Ambulante kardiologische Rehabilitation, Bonn
markus.zirbes@spkar.de

Cycle Ergometry Is not a Reliable Parameter

We work in an institution that provides follow-up examinations for patients with coronary heart disease and wish to share the following observations.

The authors rightly say that routine CT angiography is not indicated after coronary stent implantation. This

is not only because the coronary stenoses diagnosed on CT are often not hemodynamically relevant. The more important reason is that monitoring stents is affected by their structure and size, among others. Further, in patients with known coronary heart disease the coronary vessels are often subject to severe calcification, which can substantially limit diagnostic accuracy. The consensus recommendations from the working groups of the German Cardiac Society/German Society of Pediatric Cardiology do not regard the indication as definite in asymptomatic and symptomatic patients after coronary intervention or bypass surgery, unless the question is only one of bypass graft patency (1).

Regarding ergometry testing, our critical comment would be that the reported criterion of maximum exertion (85% of age-adjusted heart frequency) does not constitute a reliable parameter. The maximum individual heart frequency is subject to large variability and is furthermore affected by cardiac medication (2, 3). Furthermore, especially untrained elderly patients cannot achieve (cardiac) maximum exertion on the exercise bike because of deficits in the peripheral muscles. Exercise on a treadmill can be helpful in this setting as it typically reflects everyday activity more closely, in order to reach as high a degree of maximum exertion as possible and to improve the sensitivity of the exercise ECG, which is limited anyway. Furthermore, it is possible by using combinations with lactate testing and/or spiro-ergometry testing to establish more objective criteria for maximum exertion, but also to finesse the control of the recommended physical training.

DOI: 10.3238/arztebl.2013.0403c

REFERENCES

1. Achenbach S, Barkhausen J, Beer M, et al.: Konsensusempfehlungen der DRG/DGK/DGPK zum Einsatz der Herzbildgebung mit Computertomografie und Magnetresonanztomografie. *Röfo* 2012; 184: 345–68.
2. Roecker K, Niess A, Horstmann T, Striegel H, Mayer F, Dickhuth HH: Heart rate prescriptions from performance and anthropometrical characters. *Med Sci Sports Exerc* 2002; 34: 881–7.
3. Such M, Meyer T: Die maximale Herzfrequenz. *Dtsch Z Sportmed* 2010; 61: 310–1.
4. Rassaf T, Steiner S, Kelm M: Postoperative care and follow up after coronary stenting. *Dtsch Arztebl Int* 2013; 110(5): 72–82.

Prof. Dr. med. Christof Burgstahler

Prof. Dr. med. Andreas Nieß

Universitätsklinik Tübingen, Abteilung Sportmedizin, Tübingen
christof.burgstahler@med.uni-tuebingen.de

In Reply:

Several studies have investigated the difficulties of systematically implementing guidelines in routine clinical practice. This is the very reason why *Deutsches*

Arzteblatt decided on commissioning this type of article. Our CME article aimed to point out a possible pathway to readers.

Still, our article should be seen merely as a recommendation—just like guidelines themselves. The treating physician needs to consider the risks and benefits of diagnostic cardiac catheterization individually for each patient.

The suggestion made by our colleagues from the department of sports medicine at Tübingen University to use treadmill testing rather than the exercise bike is certainly a good alternative. The same can be said for the combination with lactate diagnosis and/or spiro-ergometry. However, it needs to be borne in mind that these methods are not available everywhere.

We agree that professional support to help patients learn to cope with their illness, including influencing lifestyles and modifying risk profiles, is important.

The German Disease Management Guidelines (1) recommend that multidisciplinary rehabilitation be undertaken after acute ST elevation myocardial infarction and non-ST-elevation myocardial infarction, after coronary bypass surgery, and, in selected cases, after elective percutaneous coronary artery intervention (PCI), for example, in patients with a pronounced risk profile, where the requirement for training is high, or where compliance is a problem. However, the effect of multidisciplinary rehabilitation in patients after non-ST-elevation myocardial infarction and after PCI has not been studied to a satisfactory degree (1).

After PCI, patients are most likely to benefit with regard to lifestyle adjustments, physical performance, and professional rehabilitation. However, they should be technically and professionally monitored during this process.

DOI: 10.3238/arztebl.2013.0404

REFERENCES

1. Bundesärztekammer (BÄK), Kassenärztliche Bundesvereinigung (KBV) Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften (AWMF): Nationale VersorgungsLeitlinie Chronische KHK – Langfassung. Version 1.X. 2006. Available from: www.khk.-versorgungsleitlinien.de; DOI: 10.6101/AZQ/000041 Last accessed on 15. May 2013.
2. Rassaf T, Steiner S, Kelm M: Postoperative care and follow up after coronary stenting. *Dtsch Arztebl Int* 2013; 110(5): 72–82.

Prof Dr. med. Tienush Rassaf

Prof Dr. med. Malte Kelm

Universitätsklinikum Düsseldorf,
Klinik für Kardiologie, Pneumologie, Angiologie
Düsseldorf
Tienush.Rassaf@med.uni-duesseldorf.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists