Published in final edited form as:

J Biomol NMR. 2013 June ; 56(2): 139–148. doi:10.1007/s10858-013-9730-1.

Determining the Depth of Insertion of Dynamically Invisible Membrane Peptides by Gel-Phase ^1H Spin Diffusion Heteronuclear Correlation NMR

T. Wang, H. Yao, and M. Hong*

Department of Chemistry, Iowa State University, Ames, IA 50011

Abstract

Solid-state NMR determination of the depth of insertion of membrane peptides and proteins has so far utilized ^1H spin diffusion and paramagnetic relaxation enhancement experiments, which are typically conducted in the liquid-crystalline phase of the lipid membrane. For membrane proteins or peptide assemblies that undergo intermediate-timescale motion in the liquid-crystalline membrane, these approaches are no longer applicable because the protein signals are broadened beyond detection. Here we show that the rigid-solid HETCOR experiment, with an additional spin diffusion period, can be used to determine the depth of proteins in gel-phase lipid membranes, where the proteins are immobilized to give high-intensity solid-state NMR spectra. Demonstration on two membrane peptides with known insertion depths shows that well-inserted peptides give rise to high lipid cross peak intensities and low water cross peaks within a modest spin diffusion mixing time, while surface-bound peptides have higher water than lipid cross peaks. Furthermore, well-inserted membrane peptides have nearly identical ^1H cross sections as the lipid chains, indicating equilibration of the peptide and lipid magnetization. Using this approach, we measured the membrane topology of the α -helical fusion peptide of the paramyxovirus, PIV5, in the anionic POPC/POPG membrane, in which the fusion peptide undergoes intermediate-timescale motion at physiological temperature. The gel-phase HETCOR spectra indicate that the α -helical fusion peptide is well inserted into the POPC/POPG bilayer, spanning both leaflets. This insertion motif gives insight into the functional role of the α -helical PIV5 fusion peptide in virus-cell membrane fusion.

Keywords

intermediate motion; HETCOR; magic-angle spinning; fusion peptide; membrane topology; solid-state NMR

Introduction

The depth of insertion of membrane peptides and proteins is an important aspect of membrane protein structure determination. A wide range of biophysical techniques such as fluorescence spectroscopy (Chattopadhyay and London, 1987; Kleinschmidt and Tamm, 1996; Voglino et al., 1999; Zoonens et al., 2008), neutron diffraction (Bradshaw et al., 1998; Chenal et al., 2009), hydrogen-deuterium exchange combined with infrared spectroscopy (Hohlweg et al., 2012), have been used to determine membrane protein depths. Compared to these techniques, NMR spectroscopy has the advantages of providing site-specific depth information without introducing bulky probes that may perturb the structure of the

* Corresponding author: Mei Hong Tel: 515-294-3521, Fax: 515-294-0105, mhong@iastate.edu.

membrane proteins or lipids. One NMR approach is to use paramagnetic relaxation enhancement (PRE) (Solomon, 1955) of nuclear spins induced by paramagnetic ions bound to the membrane surface (Buffy et al., 2003; Grobner et al., 1999; Hong and Su, 2011; Hong et al., 2012), dissolved oxygen in the membrane (Al-Abdul-Wahid et al., 2011; Prosser et al., 2000), or spin labels incorporated into lipid molecules (Esposito et al., 1992; Hilty et al., 2004; Jacob et al., 1999). Depending on the distances of the nuclei from the paramagnetic centers, the T_1 or T_2 relaxation rates of the nuclei are enhanced to varying degrees.

The second NMR approach for depth determination utilizes ^1H spin diffusion from lipid and water to the protein (Huster et al., 2002; Kumashiro et al., 1998). In the liquid-crystalline (LC) phase of the membrane where the lipid chains and water are highly dynamic while the protein is immobile, the lipid and water ^1H magnetization can be readily selected and allowed to diffuse to the rigid protein (Fig. 1a). The result of this distance-dependent intermolecular spin diffusion is detected as correlation signals between the protein ^{13}C (or ^{15}N) and lipid/water protons in a 2D heteronuclear correlation (HETCOR) experiment (Huster et al., 2002). This LC-phase ^1H spin diffusion technique has been applied to a number of membrane peptides and proteins such as bacterial toxins (Gallagher et al., 2004; Huster et al., 2002), antimicrobial peptides (Mani et al., 2006; Su et al., 2011; Tang et al., 2009), and cell-penetrating peptides (Su et al., 2008).

However, the LC-phase ^1H spin diffusion technique cannot be applied when the membrane protein of interest undergoes intermediate-timescale motion in the lipid bilayer and causes line broadening and signal loss. This intermediate-exchange broadening results from the intrinsic rotational diffusion of membrane proteins, whose rate depends on the radius of the membrane protein, the viscosity of the lipid bilayer, and the temperature (Saffman and Delbruck, 1975). When the rotational diffusion occurs on the timescale of microseconds in the LC membrane, it interferes with ^1H decoupling and polarization transfer processes in the solid-state NMR experiments, thus causing line broadening. This exchange broadening has been reported for many membrane peptides such as the influenza M2 transmembrane peptide (Cady et al., 2007) and the fusion peptide (FPK4) of the paramyxovirus, parainfluenza virus 5 (PIV5) (Yao and Hong, 2013). In the absence of detectable protein signals, the depth of insertion cannot be determined using the LC-phase ^1H spin diffusion technique.

Fig. 2 shows an example of intermediate-exchange broadening of the ^{13}C magic-angle-spinning (MAS) spectra of the PIV5 fusion peptide bound to the POPC/POPG membrane. The ^{13}C signals of nine labeled residues, distributed throughout the peptide, were undetectable at 293 K, but became strong and well resolved at 243 K, in the gel-phase of the lipid membrane. The $\text{C}\alpha$ and $\text{C}\beta$ chemical shifts of all labeled residues indicate an α -helical conformation (Yao and Hong, 2013). Because of this exchange broadening, it was not possible to determine the membrane topology of the α -helical fusion peptide using the LC-phase ^1H spin diffusion experiment. In contrast, in the neutral POPC membrane, FPK4 adopts a β -strand conformation (Yao and Hong, 2013) and is immobilized at ambient temperature, giving high spectral intensities. Thus, the depth of insertion of the β -strand fusion peptide could be determined using the ^1H spin diffusion technique. The peptide was found to lie on the surface of the POPC membrane, dehydrating the lipid headgroups (Yao and Hong, 2013).

Viral fusion proteins such as the PIV5 F protein have long been important structural biology targets because of their essential role in mediating the entry of enveloped viruses into cells (Harrison, 2008; Lamb and Jardetzky, 2007). Class I viral fusion proteins such as the influenza virus hemagglutinin (HA), the human immunodeficiency virus (HIV) env, and the paramyxovirus F protein, undergo a series of large conformational changes to drive the

merger of the target cell membrane and the virus envelope. During these unfolding and refolding events, an N-terminal fusion peptide, originally enclosed in the interior of the pre-fusion protein globule, becomes exposed and inserts into the target cell membrane. The extended protein intermediate, doubly anchored in the virus envelope and the cell membrane, subsequently folds into a hairpin, in doing so fusing the cell and viral membranes (Yin et al., 2005; Yin et al., 2006). Determining the orientation and depth of insertion of the fusion peptide in lipid membranes is therefore important for elucidating the detailed molecular events during virus-cell fusion.

In this study, we demonstrate a low-temperature gel-phase ^1H -homonuclear-decoupled HETCOR experiment for determining the depth of insertion of membrane proteins that undergo intermediate-timescale motion in the LC membrane. We describe the spectral patterns of surface-bound peptides versus well-inserted peptides obtained from this gel-phase spin diffusion HETCOR experiment using structurally known membrane peptides. We then apply this low-temperature spin diffusion experiment to the α -helical PIV5 fusion peptide in the anionic POPC/POPG membrane. Our results not only reveal the qualitative insertion motif of the peptide, but also show that site-specific depth resolution can be obtained from this gel-phase spin diffusion experiment.

Experimental Methods

Membrane peptides

The PIV5 fusion peptide was synthesized and purified using Fmoc chemistry by PrimmBiotech (Cambridge, MA). The peptide corresponds to residues 103-129 of the PIV5 F protein, with an amino acid sequence of FAGVVIGLAALGVATAAQVTAVALVK. A DIOXA-KKKK tag was appended to the C-terminus of this sequence to increase the solubility of the peptide (Yao and Hong, 2013). Two peptides with different ^{13}C , ^{15}N -labeled residues were used in the current study. The first sample contains ^{13}C , ^{15}N -labeled residues at G114, V115, A126, and L127 (GVAL-FPK4), while the second peptide contains labeled I108, G109, A112, L113, and V125 (IGALV-FPK4). The fusion peptide was reconstituted into POPC and POPC/POPG (4 : 1 mole ratio) membranes using an organic solution mixing protocol. The peptide was dissolved in trifluoroethanol and mixed with lipids in chloroform. The solvents were removed under a stream of nitrogen gas and the mixture was lyophilized. The dry powder was then suspended in 10 mM phosphate buffer (pH = 7.5) with 1 mM EDTA and 1 mM NaN_3 , dialyzed for one day, then subjected to ultracentrifugation at 55,000 rpm for 4 h at 4 °C. The resulting membrane pellets were transferred to 4-mm MAS rotors. The peptide/lipid molar ratio was 1 : 20 for these fusion peptide samples.

As a control, we measured the low-temperature HETCOR spectra of the antimicrobial peptide, protegrin-1 (PG-1), bound to POPE/POPG (3 : 1) membranes at a peptide/lipid molar ratio of 1 : 12.5. This peptide had been extensively studied (Hong and Su, 2011; Tang and Hong, 2009) and is known from the LC-phase ^1H spin diffusion experiment to insert into the hydrophobic part of the POPE/POPG membrane (Mani et al., 2006). The peptide is ^{13}C -labeled at L5 C α and V16 ^{13}CO (Tang et al., 2007).

Solid-State NMR spectroscopy

The 1D ^{13}C CP-MAS spectra of FPK4 in POPC/POPG membranes were measured at 293 K and 243 K on a Bruker DSX-400 MHz (9.4 Tesla) spectrometer. All other NMR experiments were conducted on a Bruker Avance II 600 MHz spectrometer at a field strength of 14.1 Tesla using 4 mm MAS probes. Typical radiofrequency field strengths were 50-70 kHz for ^1H and 50 kHz for ^{13}C . ^{13}C chemical shift were externally referenced to

the ^{13}C signal of α -Gly at 176.465 ppm on the TMS scale, while the ^1H chemical shift was calibrated using N-formyl-Met-Leu-Phe-OH, whose ^1H chemical shifts have been reported (Li et al., 2010).

The gel-phase ^1H spin diffusion HETCOR spectra were measured under 7 kHz MAS using the pulse sequence in Fig. 1b. During the t_1 evolution period, the FSLG sequence (Bielecki et al., 1989) was used to decouple the ^1H - ^1H dipolar interaction. A mixing time after t_1 allowed spin diffusion to occur, followed by Lee-Goldburg cross polarization (CP) (Lee and Goldburg, 1965) to transfer the ^1H magnetization to ^{13}C . The pulse sequence is essentially the rigid-solid HETCOR experiment with an additional spin diffusion period. Applied to LC lipid membranes, this experiment has been used to investigate peptide-water and peptide-lipid intermolecular interactions (Li et al., 2010; Wang et al., 2012).

The temperatures of the gel-phase HETCOR experiments were chosen to be the highest temperature at which the membrane peptides were immobilized. To compare the HETCOR spectra of different samples, we monitored the ^1H linewidths of the lipid chain signals, which reflect the spin diffusion coefficient. For FPK4-containing POPC and POPC/POPG membranes, the HETCOR experiments were carried out at 258 K, which is 13 K below the phase transition temperature of these lipid membranes in the absence of peptides. The PG-1-containing POPE/POPG sample was measured at 253 K, at which the lipid ^1H linewidths were similar to those of the FPK4 samples (see Fig. 7).

The ^1H dimension of the 2D HETCOR spectra was processed using moderate line broadening parameters of $\text{LB} = -5$ and $\text{GB} = 0.1$, which corresponds to a Gaussian full width at half maximum of 0.15 ppm. This line broadening is smaller than the apparent ^1H linewidths of these membrane samples under the conditions of the experiments.

Results and Discussion

The main difference between the LC-phase and gel-phase ^1H spin diffusion HETCOR experiments is that the LC-phase experiment has no ^1H homonuclear decoupling during the t_1 evolution period but has a ^1H T_2 filter before t_1 , while the gel-phase experiment requires ^1H homonuclear decoupling during t_1 but no T_2 filter (Fig. 1). These differences match the dynamic properties of the lipid and peptides in the fluid and gel phases of the lipid membranes. At high temperature, the ^1H T_2 filter and the absence of homonuclear decoupling combine to suppress the ^1H magnetization of the rigid peptide while retaining the magnetization of the mobile lipid and water, which act as the magnetization source. The fast rotational diffusion of lipids in the LC membrane ensures that moderate MAS frequencies alone are sufficient to give high-resolution ^1H spectra (Doherty and Hong, 2009; Oldfield et al., 1987). In contrast, at low temperature where both the peptides and lipids are immobilized, homonuclear decoupling is necessary to obtain high-resolution ^1H spectra of the peptide and lipids. Under this condition, the ^1H dimension of the 2D spectra needs to be assigned to distinguish the peptide and lipid signals and to distinguish intramolecular from intermolecular cross peaks. We show that this assignment is possible based on the characteristic chemical shifts of functional groups as well as the mixing time dependence of spectral intensities.

To investigate how low-temperature HETCOR spectra differ between well-inserted and surface-bound membrane peptides, we first demonstrate the experiment on two peptides whose depths of insertion were previously determined using the LC-phase ^1H spin diffusion technique. The antimicrobial peptide, PG-1, has been extensively studied and is known to insert into the bacteria-mimetic anionic membrane, POPE/POPG (Mani et al., 2006; Tang and Hong, 2009). In comparison, the PIV5 fusion peptide in the POPC membrane is an

example of a surface-bound membrane peptide (Yao and Hong, 2013). Fig. 3 shows the 253 K ^1H - ^{13}C HETCOR spectra of POPE/POPG-bound PG-1. For all samples in this study, we used ^1H spin diffusion mixing times of 0, 0.2 ms, 4 ms and 25 ms. In the absence of spin diffusion, only one-bond cross peaks were detected. Specifically, in the L5 Ca cross section, an H α peak at 4.8 ppm dominates the spectrum, followed by weak H $^{\text{N}}$ and sidechain H β /H γ cross peaks at 8.0 ppm and 0.8 ppm. With a 0.2 ms mixing time, the 0.8-ppm peak becomes the strongest, due to the presence of more sidechain protons than backbone protons. At longer mixing times of 4 ms and 25 ms, the ^1H spectral pattern changed significantly, with a strong ^1H peak at 1.1 ppm and a weak signal at 4.9 ppm, and the H $^{\text{N}}$ signal has disappeared. The 1.1-ppm peak can be assigned to lipid CH $_2$ protons, based on its slightly different chemical shift and its narrower linewidth (1.2 ppm) compared to the L5 H β /H γ linewidths of 1.6 ppm. The assignment of the 4.9-ppm peak to water is based on the similar intensity distribution of the peptide and lipid cross sections at 25 ms mixing and the disappearance of the peptide H $^{\text{N}}$ peak. These observations indicate that spin diffusion has equilibrated the proton reservoir of the peptide and its neighboring lipid and water molecules by 25 ms, and the L5 region of PG-1 is well inserted into the hydrophobic part of the lipid membrane, with shorter distances to the lipid chains than to water. This conclusion is consistent with the LC-phase ^1H spin diffusion results, which found PG-1 to be inserted into the hydrophobic center of the POPE/POPG membrane (Mani et al., 2006). Therefore, the signature of an inserted peptide in the gel-phase HETCOR experiment is the similar intensity distribution between the lipid and peptide cross sections within a modest mixing time.

Fig. 4 shows the gel-phase HETCOR spectra of the β -strand FPK4 bound to the POPC membrane. While the zero-mixing spectrum is similar to the PG-1 spectrum, at 25 ms, the peptide ^1H cross sections are completely different: the peptide cross section has much higher water intensity than the lipid cross section by 25 ms, indicating that the β -strand FPK4 resides on the surface of the POPC bilayer, consistent with the high-temperature ^1H spin diffusion results (Yao and Hong, 2013). In comparison, the lipid cross section at 25 ms is similar to the POPE/POPG cross section in Fig. 3, with the CH $_2$ self-correlation peak dominating the spectrum. The 5.2-ppm peak in the peptide cross sections at 4 ms and 25 ms can be assigned to water rather than H α because its linewidth (0.7 ppm) is much narrower than the H α linewidth (1.8 ppm) in the zero-mixing spectrum. Taken together, Fig. 3 and 4 indicate that membrane-inserted and surface-bound peptides have very different ^1H intensity distributions: the former has strong lipid CH $_2$ peaks and weak water cross peaks, while the latter displays the opposite relative intensities.

Using this gel-phase spin diffusion HETCOR experiment, we investigated the depth of insertion of the PIV5 fusion peptide in the anionic POPC/POPG membrane. Fig. 5 shows representative 2D HETCOR spectra of GVAL-FPK4 and IGALV-FPK4 at 258 K. The sum of the peptide Ca cross sections shows similar ^1H intensity distributions as the lipid CH $_2$ cross section at 25 ms, indicating that the α -helical FPK4 is well inserted into the POPC/POPG membrane. Moreover, the relative intensity of the water and lipid cross peaks differs among different residues. ^{13}C cross sections at the water and lipid CH $_2$ positions indicate that residues near the two ends of the peptide, such as I108, G109 and V125, have higher water/lipid intensity ratios than residues in the middle of the peptide such as A112 and L113 (Fig. 6). The difference is particularly clear at short mixing times but is also detectable at the longest mixing time of 25 ms. Therefore, the N- and C-termini of FPK4 have closer contacts with water than the middle of the peptide, strongly suggesting that FPK4 spans both leaflets of the bilayer (Fig. 6c). The alternative scenario that the peptide might be highly curved such that the N- and C-termini contact water molecules on the same side of the bilayer can be reasonably ruled out, since the ^{13}C chemical shifts of this peptide indicate a relatively straight α -helix, without significant kinks or bend (Yao and Hong, 2013). These data indicate that the gel-phase spin diffusion technique has residue-specific depth resolution, in

contrast to the LC-phase experiment. The reason for this difference is that at low temperature, lipids and peptides have similar spin diffusion efficiencies, thus the rate of magnetization transfer depends on the actual distances of nuclear spins from the magnetization source, rather than the minimum distance of the peptide from the source spins.

In addition to the water/lipid intensity ratio, the ^1H linewidth of the peptide-lipid cross peaks also provides information about the insertion state of the peptide. For the well inserted POPC/POPG-bound FPK4, the peptide Ca to lipid CH_2 cross peak has a ^1H linewidth of 0.9 ppm, which is the same as the ^1H linewidth of the lipid self-correlation peak (Fig. 5b), supporting complete magnetization exchange between the peptide and lipid chains. The same was observed for PG-1 in the POPE/POPE membrane. In contrast, for the POPC-bound FPK4 (Fig. 4), the peptide cross peak with CH_2 groups resonates at a ^1H chemical shift of 1.2 ppm with a linewidth of 1.1 ppm, while the lipid self-correlation peak resonates at 1.4 ppm with a linewidth of 0.8 ppm. These subtle but reproducible differences indicate that peptide sidechain $\text{H}\beta/\text{H}\gamma$ protons make a significant contribution to the 1.2-ppm peak, and the magnetization of the surface-bound FPK4 and the lipid chains are not equilibrated by 25 ms.

The inserted topology of the α -helical PIV5 fusion peptide, together with its intermediate-timescale motion at ambient temperature, gives insight into the structure of FPK4 in the POPC/POPG membrane. If the peptide is monomeric and not significantly tilted, then its dynamics should be much faster than the NMR interactions and fast-averaged spectra with narrow ^{13}C linewidths should be observed. Thus, the fact that the FPK4 spectra are broadened beyond detection between 263 K and 313 K indicates that the α -helical FPK4 is well oligomerized and significantly tilted. Since the water-soluble ectodomain of the F protein is trimeric, the fusion peptide domain may also be trimeric in the lipid membrane. But trimerization alone is unlikely to be sufficient for causing intermediate-timescale motion, since even membrane proteins with seven transmembrane helices have been shown to undergo fast motion in fluid membranes (Lewis et al., 1985). Instead, the fusion peptide assembly is likely significantly tilted in the POPC/POPG membrane in order for intermediate dynamics to persist over a wide temperature range (Fig. 6c).

The membrane-spanning topology of the α -helical PIV5 fusion peptide in the POPC/POPG membrane differs dramatically from the surface-bound topology of the β -strand FPK4 in neutral POPC membranes (Yao and Hong, 2013). Our recent data correlating membrane curvature and hydration with FPK4 conformation suggested the β -strand conformation to be more relevant for membrane intermediates of the fusion pathway, while the α -helical conformation is associated with well-hydrated lamellar membranes that may be relevant for the pre-fusion and post-fusion states. Additional experiments will be necessary to test these hypotheses further. Interestingly, solid-state NMR measurements showed that the HIV fusion peptide forms an oligomeric β -sheet in virus-mimetic lipid membranes and spans one leaflet of the lipid bilayer (Qiang et al., 2009). This partially inserted β -sheet peptide also incurs membrane curvature (Gabrys et al., 2010). Information about α -helical viral fusion peptides so far mainly came from solution NMR studies of micelle-bound influenza HA, which found the peptide to form a bent helix that is shallowly inserted into the detergent micelle (Han et al., 2001; Lorieau et al., 2010).

The gel-phase ^1H spin diffusion HETCOR experiment is best applied at the highest temperature where the membrane peptides are immobilized. At moderate low temperature, the water molecules remain partly mobile, which would give a distinct and narrow ^1H signal that can be readily resolved from the peptide $\text{H}\alpha$ signals. At moderate low temperature, the mixing times required to distinguish well inserted and surface-bound membrane peptides are

longer than at extremely low temperatures, thus facilitating the distinction between topologically different membrane peptides. Finally, the use of moderate low temperature avoids the potential hazard of changing the peptide-lipid interactions compared to the physiological situation. To compare the depths of the membrane peptides in different lipid bilayers, the HETCOR experiments should be conducted at similar reduced temperatures from the membrane phase transition temperature so that the lipid chain dynamics are comparable. Fig. 7 shows the undecoupled ^1H spectra of POPC/POPG, POPC, and POPE/POPG membranes at 253 – 258 K, where the 2D HETCOR spectra were measured. The CH_2^1 linewidths are similar, thus allowing the comparison of the relative depths of the PIV5 fusion peptide and PG-1. ^2H NMR of chain-deuterated lipids can also be used to verify membrane dynamics at the desired temperatures.

In conclusion, the depth of insertion of membrane peptides and proteins that exhibit intermediate-timescale motion in the LC phase of lipid membranes can now be determined using the gel-phase spin diffusion HETCOR experiment. This gel-phase experiment not only overcomes the exchange-broadening problem of many membrane peptides, but also provides site-specific depth resolution, which is absent from the LC-phase ^1H spin diffusion experiment. This gel-phase HETCOR experiment completes the NMR toolbox for determining the membrane topology of peptides and proteins. Further improvement of this technique can be envisioned, for example using advanced ^1H decoupling and isotopic dilution methods to enhance the resolution of the ^1H dimension to better distinguish lipid and protein signals.

Acknowledgments

This work is funded by NIH grant GM066976.

References

- Al-Abdul-Wahid MS, Verardi R, Veglia G, Prosser RS. Topology and immersion depth of an integral membrane protein by paramagnetic rates from dissolved oxygen. *J. Biomol. NMR*. 2011; 51:173–183. [PubMed: 21947925]
- Bielecki A, Kolbert AC, Levitt MH. Frequency-switched pulse sequences: homonuclear decoupling and dilute spin NMR in solids. *Chem. Phys. Lett.* 1989; 155:341–346.
- Bradshaw JP, Davies SM, Hauss T. Interaction of substance P with phospholipid bilayers: A neutron diffraction study. *Biophys. J.* 1998; 75:889–895. [PubMed: 9675189]
- Buffy JJ, Hong T, Yamaguchi S, Waring A, Lehrer RI, Hong M. Solid-State NMR Investigation of the Depth of Insertion of Protegin-1 in Lipid Bilayers Using Paramagnetic Mn^{2+} . *Biophys. J.* 2003; 85:2363–2373. [PubMed: 14507700]
- Cady SD, Goodman C, Tatko C, DeGrado WF, Hong M. Determining the orientation of uniaxially rotating membrane proteins using unoriented samples: a ^2H , ^{13}C , and ^{15}N solid-state NMR investigation of the dynamics and orientation of a transmembrane helical bundle. *J. Am. Chem. Soc.* 2007; 129:5719–5729. [PubMed: 17417850]
- Chattopadhyay A, London E. Parallax method for direct measurement of membrane penetration depth utilizing fluorescence quenching by spin-labeled phospholipids. *Biochemistry*. 1987; 26:39–45. [PubMed: 3030403]
- Chenal A, Prongidi-Fix L, Perier A, Aisenbrey C, Vernier G, Lambotte S, Haertlein M, Dauvergne MT, Fragneto G, Bechinger B, Gillet D, Forge V, Ferrand M. Deciphering membrane insertion of the diphtheria toxin T domain by specular neutron reflectometry and solid-state NMR spectroscopy. *J. Mol. Biol.* 2009; 391:872–883. [PubMed: 19576225]
- Doherty T, Hong M. High-resolution solid-state NMR of anisotropically mobile molecules under very low-power (^1H) decoupling and moderate magic-angle spinning. *J. Magn. Reson.* 2009; 199:225–232. [PubMed: 19501003]

- Esposito G, Lesk AM, Molinari H, Motta A, Niccolai N, Pastore A. Probing protein structure by solvent perturbation of nuclear magnetic resonance spectra. Nuclear magnetic resonance spectral editing and topological mapping in proteins by paramagnetic relaxation filtering. *J. Mol. Biol.* 1992; 224:659–670. [PubMed: 1314901]
- Gabrys CM, Yang R, Wasniewski CM, Yang J, Canlas CG, Qiang W, Sun Y, Weliky DP. Nuclear magnetic resonance evidence for retention of a lamellar membrane phase with curvature in the presence of large quantities of the HIV fusion peptide. *Biochim. Biophys. Acta.* 2010; 1798:194–201. [PubMed: 19616505]
- Gallagher GJ, Hong M, Thompson LK. Solid-state NMR spin diffusion for measurement of membrane-bound peptide structure: gramicidin A. *Biochemistry.* 2004; 43:7899–7906. [PubMed: 15196034]
- Grobner G, Glaubitz C, Watts A. Probing membrane surfaces and the location of membrane-embedded peptides by (^{13}C) MAS NMR using lanthanide ions. *J. Magn. Reson.* 1999; 141:335–339. [PubMed: 10579957]
- Han X, Bushweller JH, Cafiso DS, Tamm LK. Membrane structure and fusion-triggering conformational change of the fusion domain from influenza hemagglutinin. *Nat. Struct. Biol.* 2001; 8:715–720. [PubMed: 11473264]
- Harrison SC. Viral membrane fusion. *Nature Struc. Mol. Biol.* 2008; 15:690–698.
- Hilty C, Wider G, Fernández C, Wüthrich K. Membrane protein-lipid interactions in mixed micelles studied by NMR spectroscopy with the use of paramagnetic reagents. *Chembiochem.* 2004; 5:467–473. [PubMed: 15185370]
- Hohlweg W, Kosol S, Zangger K. Determining the orientation and localization of membrane-bound peptides. *Curr. Protein Pept. Sci.* 2012; 13:267–279. [PubMed: 22044140]
- Hong M, Su Y. Structure and dynamics of cationic membrane peptides and proteins: Insights from solid-state NMR. *Protein Sci.* 2011; 20:641–655. [PubMed: 21344534]
- Hong M, Zhang Y, Hu F. Membrane Protein Structure and Dynamics from NMR Spectroscopy. *Annu. Rev. Phys. Chem.* 2012; 63:1–24. [PubMed: 22136620]
- Hunter D, Yao XL, Hong M. Membrane Protein Topology Probed by ^1H Spin Diffusion from Lipids Using Solid-State NMR Spectroscopy. *J. Am. Chem. Soc.* 2002; 124:874–883. [PubMed: 11817963]
- Jacob J, Baker B, Bryant RG, Cafiso DS. Distance estimates from paramagnetic enhancements of nuclear relaxation in linear and flexible model peptides. *Biophys J.* 1999; 77:1086–1092. [PubMed: 10423452]
- Kleinschmidt JH, Tamm LK. Folding intermediates of a beta-barrel membrane protein. Kinetic evidence for a multi-step membrane insertion mechanism. *Biochemistry.* 1996; 35:12993–13000. [PubMed: 8855933]
- Kumashiro KK, Schmidt-Rohr K, Murphy OJ, Ouellette KL, Cramer WA, Thompson LK. A novel tool for probing membrane protein structure: solid-state NMR with proton spin diffusion and X-nucleus detection. *J. Am. Chem. Soc.* 1998; 120:5043–5051.
- Lamb RA, Jardetzky TS. Structural basis of viral invasion: lessons from paramyxovirus F. *Curr. Opin. Struct. Biol.* 2007; 17:427–436. [PubMed: 17870467]
- Lee M, Goldburg WI. Nuclear-magnetic-resonance line narrowing by a rotating rf field. *Phys. Rev.* 1965; 140:A1261–A1271.
- Lewis BA, Harbison GS, Herzfeld J, Griffin RG. NMR structural analysis of a membrane protein: bacteriorhodopsin peptide backbone orientation and motion. *Biochemistry.* 1985; 24:4671–4679. [PubMed: 4063350]
- Li S, Su Y, Luo W, Hong M. Water-protein Interactions of an Arginine-Rich Membrane Peptide in Lipid Bilayers Investigated by Solid-State Nuclear Magnetic Resonance Spectroscopy. *J. Phys. Chem. B.* 2010; 114:4063–4069. [PubMed: 20199036]
- Lorieau JL, Louis JM, Bax A. The complete influenza hemagglutinin fusion domain adopts a tight helical hairpin arrangement at the lipid:water interface. *Proc Natl Acad Sci U S A.* 2010; 107:11341–11346. [PubMed: 20534508]

- Mani R, Cady SD, Tang M, Waring AJ, Lehrer RI, Hong M. Membrane-dependent oligomeric structure and pore formation of a beta-hairpin antimicrobial peptide in lipid bilayers from solid-state NMR. *Proc. Natl. Acad. Sci. USA*. 2006; 103:16242–16247. [PubMed: 17060626]
- Oldfield E, Bowers JL, Forbes J. High-resolution proton and carbon-13 NMR of membranes: why sonicate? *Biochemistry*. 1987; 26:6919–6923. [PubMed: 3427053]
- Prosser RS, Luchette PA, Westerman PW. Using O2 to probe membrane immersion depth by 19F NMR. *Proc. Natl. Acad. Sci. U.S.A.* 2000; 97:9967–9971. [PubMed: 10954744]
- Qiang W, Sun Y, Weliky DP. A strong correlation between fusogenicity and membrane insertion depth of the HIV fusion peptide. *Proc. Natl. Acad. Sci. U. S. A.* 2009; 106:15314–15319. [PubMed: 19706388]
- Saffman PG, Delbruck M. Brownian motion in biological membranes. *Proc. Natl. Acad. Sci. USA*. 1975; 72:3111–3113. [PubMed: 1059096]
- Solomon I. Relaxation processes in a system of two spins. *Phys. Rev.* 1955; 99:559–565.
- Su Y, Mani R, Hong M. Asymmetric Insertion of Membrane Proteins in Lipid Bilayers by Solid-State NMR Paramagnetic Relaxation Enhancement: a Cell-Penetrating Peptide Example. *J. Am. Chem. Soc.* 2008; 130:8856–8864. [PubMed: 18597439]
- Su Y, Waring AJ, Ruchala P, Hong M. Structures of β -Hairpin Antimicrobial Protegrin Peptides in Lipopolysaccharide Membranes: Mechanism of Gram Selectivity Obtained from Solid-State Nuclear Magnetic Resonance. *Biochemistry*. 2011; 50:2072–2083. [PubMed: 21302955]
- Tang M, Hong M. Structure and mechanism of beta-hairpin antimicrobial peptides in lipid bilayers from solid-state NMR spectroscopy. *Mol. Biosyst.* 2009; 5:317–322. [PubMed: 19396367]
- Tang M, Waring AJ, Hong M. Phosphate-Mediated Arginine Insertion Into Lipid Membranes and Pore Formation by a Cationic Membrane Peptide from Solid-State NMR. *J. Am. Chem. Soc.* 2007; 129:11438–11446. [PubMed: 17705480]
- Tang M, Waring AJ, Hong M. Effects of arginine density on the membrane-bound structure of a cationic antimicrobial peptide from solid-state NMR. *Biochim. Biophys. Acta*. 2009; 1788:514–521. [PubMed: 19059201]
- Voglino L, Simon SA, McIntosh TJ. Orientation of LamB signal peptides in bilayers: influence of lipid probes on peptide binding and interpretation of fluorescence quenching data. *Biochemistry*. 1999; 38:7509–7516. [PubMed: 10360948]
- Wang T, Cady SD, Hong M. NMR Determination of Protein Partitioning into Membrane Domains with Different Curvatures and Application to the Influenza M2 Peptide. *Biophys. J.* 2012; 102:787–794. [PubMed: 22385849]
- Yao H, Hong M. Membrane-Dependent Conformation, Dynamics, and Lipid Interactions of the Fusion Peptide of the Paramyxovirus PIV5 from Solid-State NMR. *J. Mol. Biol.* 2013; 425:563–576. [PubMed: 23183373]
- Yin HS, Paterson RG, Wen X, Lamb RA, Jardetzky TS. Structure of the uncleaved ectodomain of the paramyxovirus (hPIV3) fusion protein. *Proc. Natl. Acad. Sci. U. S. A.* 2005; 102:9288–9293. [PubMed: 15964978]
- Yin HS, Wen X, Paterson RG, Lamb RA, Jardetzky TS. Structure of the parainfluenza virus 5 F protein in its metastable, prefusion conformation. *Nature*. 2006; 439:38–44. [PubMed: 16397490]
- Zoonens M, Reshetnyak YK, Engelman DM. Bilayer interactions of pHLIP, a peptide that can deliver drugs and target tumors. *Biophys. J.* 2008; 95:225–235. [PubMed: 18359793]

Figure 1.

^1H spin diffusion HETCOR pulse sequences. (a) The LC-phase experiment. A ^1H T_2 filter is applied before the t_1 period to suppress the rigid peptide ^1H magnetization. Only the lipid and water ^1H signals are detected in the t_1 dimension. (b) The gel-phase experiment. The ^1H signals of both peptide and lipids are detected in the t_1 dimension.

Figure 2.

1D ^{13}C CP-MAS spectra of the PIV5 fusion peptide in POPC/POPG membranes at 293 K and 243 K. (a) GVAL-FPK4. (b) IGALV-FPK4. The peptide signals disappear at ambient temperature due to intermediate-timescale motion. In the gel phase of the membrane, the peptide signals reappear, and the chemical shifts indicate an α -helical conformation.

Figure 3.

Depth of insertion of PG-1 in POPE/POPG membranes by the gel-phase spin diffusion HETCOR experiment. (a) 2D spectra with 0, 0.2 ms and 25 ms mixing times. (b) ^1H cross sections of the peptide L5 $\text{C}\alpha$ peak at 51 ppm and the lipid CH_2 peak at 33 ppm at varying mixing times. The peptide and lipid cross sections have very similar intensity patterns by 25 ms, indicating that L5 is well inserted into the middle of the POPE/POPG membrane. The spectra were measured at 253 K under 7 kHz MAS. The -1.5 ppm peak is a zero-frequency artifact.

Figure 4.

Depth of insertion of GVAL-FPK4 in the POPC membrane by gel-phase spin diffusion HETCOR. (a) 2D spectra with 0, 0.2 ms and 25 ms mixing. (b) ^1H cross sections of the peptide (red) and lipid CH_2 (black) peaks at various mixing times. The ^1H cross sections of peptide come from the sum of all $\text{C}\alpha$ peaks. The peptide cross section shows much higher water cross peak than the lipid cross section by 25 ms, indicating that FPK4 lies on the surface of the POPC bilayer. The spectra were measured at 258 K under 7 kHz MAS.

Figure 5.

Depth of insertion of the PIV5 fusion peptide in POPC/POPG membranes by gel-phase ^1H spin diffusion HETCOR. (a) 2D spectra of GVAL-FPK4 at 0 and 25 ms mixing. (b) 1D ^1H cross sections of the peptide ^{13}C (red) and lipid CH_2 (black) peaks at various mixing times. The peptide ^1H cross sections are the sum of all $\text{C}\alpha$ peaks and the L127 $\text{C}\beta$ peak. (c) 2D spectra of IGALV-FPK4 at 0 and 25 ms mixing. (d) 1D ^1H cross sections of the peptide (red) and lipid CH_2 (black) signals as a function of mixing time. The similarity of the peptide and lipid cross sections by 25 ms indicate that the α -helical FPK4 is well inserted into the POPC/POPG membrane. All spectra were measured at 258 K under 7 kHz MAS. Figure

Figure 6.

Depth resolution of the gel-phase HETCOR experiment, shown by the spectra of FPK4 bound to the POPC/POPG membrane. (a) IGALV-FPK4 ¹³C cross sections at the lipid CH₂ (black) and water (blue) frequencies from the 4 ms 2D spectrum. The water/lipid intensity ratios are higher for the terminal residues (I108, G109 and V125) than for residues in the middle of the peptide (A112 and L113). (b) GVAL-FPK4 ¹³C cross sections at the lipid CH₂ and water frequencies from the 25 ms 2D spectrum. The water/lipid intensity ratios are higher for the terminal residues A126 and L127 than for the interior residues G114 and V115. (c) Illustration of the full insertion of FPK4 into the POPC/POPG membrane.

Figure 7.

^1H single-pulse spectra of various lipid membranes. (a) FPK4-containing POPC/POPG membrane at 293 K. (b) FPK4-containing POPC/POPG membrane in the gel phase at 258 K. (c) FPK4-containing POPC membrane at 258 K. (d) PG-1 containing POPE/POPG membrane at 253 K. The gel-phase spectra in (b-d) show similar linewidths for the lipid CH_2 peak, indicating comparable spin diffusion coefficients.