Published in final edited form as:

J Clin Psychiatry. 2012 June ; 73(6): 829–836. doi:10.4088/JCP.11m06912.

Prospective, population-based study of the transition from major depressive disorder to bipolar disorder

Stephen E. Gilman, Jamie M. Dupuy, and Roy H. Perlis

Department of Society, Human Development and Health, and Department of Epidemiology, Harvard School of Public Health, Boston, MA (Dr Gilman); Department of Psychiatry, Massachusetts General Hospital, Boston, MA, and the Bipolar Clinic and Research Program, Department of Psychiatry, Massachusetts General Hospital and Harvard Medical School, Boston, MA (Drs Dupuy and Perlis).

Abstract

Objective—It is currently not possible to determine which individuals with unipolar depression are at highest risk for a manic episode. This study investigates clinical and psychosocial risk factors for mania among individuals with major depressive disorder (MDD), indicating diagnostic conversion from MDD to bipolar I disorder.

Methods—We fitted logistic regression models to predict the first onset of a manic episode among 6,214 cases of lifetime MDD according to DSM-IV criteria in the National Epidemiologic Survey on Alcohol and Related Conditions.

Results—Approximately 1 in 20 individuals with MDD transitioned to bipolar disorder during the study's 3-year follow-up period. Demographic risk factors for the transition from MDD to bipolar disorder included younger age, Black race/ethnicity, and less than high school education. Clinical characteristics of depression (e.g., age at first onset, presence of atypical features) were not associated with diagnostic conversion. However, prior psychopathology was associated with the transition to bipolar disorder: history of social phobia (Odds Ratio=2.20; 95% Confidence Interval=1.47, 3.30) and generalized anxiety disorder (OR=1.58; CI=1.06, 2.35). Lastly, we identified environmental stressors over the life course that predicted the transition to bipolar disorder: these include a history of child abuse (OR=1.26; CI=1.12, 1.42) and past-year problems with one's social support group (OR=1.79; CI=1.19, 2.68). The overall predictive power of these risk factors based on a receiver operating curve analysis is modest.

Conclusions—A wide range of demographic, clinical, and environmental risk factors were identified that indicate a heightened risk for the transition to bipolar disorder. Additional work is needed to further enhance the prediction of bipolar disorder among cases of MDD, and to determine whether interventions targeting these factors could reduce the risk of bipolar disorder.

Keywords

unipolar; bipolar; depression; mania; epidemiology

Correspondence to: Stephen E. Gilman, Sc.D., Harvard School of Public Health, 677 Huntington Avenue, Boston, MA 02115. Phone 617-432-6383; Fax 617-432-3755; sgilman@hsph.harvard.edu..

Disclosure: Dr. Perlis received royalties and consulting fees, and held equity in, Concordant Rater Systems.

INTRODUCTION

A majority of individuals with bipolar disorder experience one or more major depressive episodes prior to an initial manic episode.¹ The ability to identify those individuals with unipolar depression who are at greatest risk for a manic episode, and therefore subsequent transition to bipolar depression, could allow for more targeted interventions, symptom surveillance, and potentially for studies aimed at prevention.

What distinguishes individuals with major depressive episodes who will go on to develop mania from those who will not? This question is difficult to answer, because, as Mitchell et al. concluded, “there are no pathognomonic characteristics of bipolar depression compared to unipolar depression (major depressive disorder).”², p.149 There have been many attempts to compare unipolar depressed to bipolar depressed patients cross-sectionally, and based on these comparisons, develop criteria that could be used to indicate probable bipolar disorder.² Features commonly associated with greater risk for bipolar disorder in such studies include more depressive episodes and earlier onset of illness,²⁻⁶ as well as greater depressive severity,⁴ atypical depressive symptoms,^{7,8} psychotic symptoms,^{5,9} and family history of bipolar disorder.^{5,6} While such cross-sectional studies are informative, this approach does not necessarily distinguish features which are present *prior* to the first manic episode, for example in terms of psychiatric comorbidity.³ Moreover, symptoms of bipolar depression are not necessarily stable across episodes.¹⁰ We therefore investigated the clinical and social determinants of the transition from Major Depressive Disorder (MDD) to bipolar disorder in a large two-wave national sample of individuals meeting DSM-IV criteria for MDD. We use data from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC), and attempt to identify risk factors for a manic episode occurring during the study's three-year follow-up period.

METHOD

Study sample

The NESARC is a nationally representative household survey conducted by the National Institute on Alcohol Abuse and Alcoholism.^{11,12} The Wave 1 response rate was 81.2%, resulting in a sample size of 43,093 participants ages 18. The Wave 2 survey, conducted approximately 3 years later, included 34,653 of the Wave 1 participants, representing 86.7% of the 39,959 subjects who were eligible for reinterview. The combined response rate for both waves was 70.2%.¹³ The analytic sample for the current study included all participants with a diagnosis of MDD at Wave 1. This is operationalized in the NESARC as a lifetime history of major depressive episode at Wave 1, without a lifetime history of a manic or hypomanic episode (the DSM-IV's exclusions in major depressive disorder for schizophrenia or psychosis are not implemented in the NESARC).

Measures

The assessment instrument used in the NESARC was the Alcohol Use Disorder and Associated Disabilities Interview Schedule-IV (AUDADIS).¹⁴ The AUDADIS is a fully structured diagnostic interview administered by trained non-clinician interviewers that assessed disorders according to DSM-IV criteria.¹⁵⁻¹⁸ The test-retest reliability of depression diagnoses in the NESARC sample was good (kappa=0.65 for a lifetime diagnosis of major depression, and 0.59 for a past-year diagnosis).¹⁶ The following clinical characteristics of participants' depression were also assessed: age at onset, the presence with atypical features (weight gain, hyperphagia or hypersomnia),¹⁹ number of lifetime depressive episodes, and a high family history loading of depression, defined as a high proportion (top decile in the entire NESARC sample) of first and second-degree relatives

with depression, weighted by their degree of genetic relatedness to the respondent (0.5 for first-degree and 0.25 for second-degree).²⁰

In the sample of individuals with MDD, we investigated the first onset of a manic episode during the NESARC's three-year follow up period, and therefore the transition to a DSM-IV diagnosis of bipolar I disorder. This was assessed at the Wave 2 interview through questions probing about symptoms of mania that occurred since the time of the Wave 1 interview. Grant et al. reported that the test-retest reliability of bipolar diagnoses in the NESARC was good ($\kappa=0.59$).²¹ Though there have not been any clinical reappraisal studies of bipolar diagnoses based on the AUDADIS, the validity of bipolar diagnoses obtained from the Composite International Diagnostic Interview (CIDI), a structured diagnostic interview similar to the AUDADIS, was excellent. Kessler et al. reported almost perfect sensitivity and specificity of CIDI bipolar diagnoses compared against the Structured Clinical Interview for DSM-IV.²²

We included in our analyses participant demographic characteristics (age, sex, race/ethnicity, and educational attainment), lifetime psychiatric disorders assessed at Wave 1, and lifetime stressors. Psychiatric disorders included DSM-IV diagnoses of panic disorder, social phobia, generalized anxiety disorder, alcohol dependence, nicotine dependence, cannabis abuse or dependence, and other substance dependence.²³⁻²⁶ Lifetime stressors included a history of childhood adversities (retrospectively reported at the Wave 2 interview), and past-year stressors (reported at the Wave 1 interview). The childhood adversities included childhood abuse (neglect, verbal abuse, and physical abuse), sexual maltreatment, and economic deprivation. Four types of past-year stressors included social support group problems (loss or injury of a close friend or family member, or separation or divorce), social environmental problems (residential changes or serious problems with a neighbor, friend, or relative), occupational problems (fired or laid off, unemployed, trouble with a boss or coworker, or job changes), and economic problems (household below the United States federal poverty threshold or received welfare). The test-retest reliabilities of these childhood and past-year stressors in the NESARC are excellent.²⁷

Statistical analyses

The study design involved fitting prediction models for a manic episode occurring between the Wave 1 and Wave 2 interviews; restricted to cases of major depression at Wave 1, these models identify risks for the transition from MDD to bipolar disorder. Logistic regression was used, in which the dependent variable was the presence or absence of a manic episode. We conducted two sets of analyses: 1) analyses of clinical characteristics of the participants' depression and lifetime histories of psychiatric disorders; and 2) analyses of childhood and past-year adversities. The primary analytic sample included participants with a lifetime history of MDD, as described above; however, in supplemental analyses, we repeated the sequence of logistic regression models among participants with a *past-year* diagnosis of MDD at Wave 1, and again among participants with lifetime MDD who sought treatment for their depression. These are groups that could arguably resemble patients enrolled into clinical samples more closely than cases identified in the NESARC overall—in terms of having active disorders in the first instance, and in terms of having sought treatment in the second instance. We sought to determine from these analyses whether there were different patterns of clinical and psychosocial predictors of the transition to bipolar I disorder among individuals with presumably more severe depression.

In order to determine the extent to which the covariates studied at baseline predict the transition from MDD to bipolar disorder during the 3-year follow-up period, we conducted a receiver operating characteristic (ROC) analysis.²⁸ We did this by fitting a logistic regression model in the primary analytic sample (i.e., lifetime cases of MDD) that included

all of the participant demographic factors, clinical characteristics of depression, lifetime psychiatric disorders, and childhood and past-year adversities. The predicted probabilities from this model were then used to plot the ROC curve, and the area under the curve (AUC) was computed. An AUC value of 50% indicates the model is no better than chance at predicting transition to bipolar, whereas an AUC of 100% indicates perfect prediction.

We conducted the analyses in SUDAAN,²⁹ which adjusts variances and point estimates for the multi-stage sampling design and differential selection probabilities used to ascertain the NESARC sample. Missing data on Wave 1 covariates (approximately 1% of the sample), and on psychiatric disorders and childhood adversities among Wave 2 non-participants (approximately 20% of the sample), were imputed using the method of multiple imputation as implemented in IVEware,³⁰ with adjustments made for the complex sampling design of the NESARC.³¹ Results were averaged across 20 multiply-imputed datasets.

RESULTS

The lifetime prevalence of MDD in the NESARC is 14.5% (n=6,214). Of these MDD cases, 4.0% (n=245) transitioned to bipolar disorder during the 3-year follow-up period. Among past-year cases of MDD at Wave 1 (40.6% of lifetime cases), 5.6% (n=142) transitioned to bipolar depression during the follow-up period. Further, among lifetime cases of MDD who sought treatment (61.8% of lifetime cases), 4.1% (n=163) transitioned to bipolar disorder.

The demographic and clinical characteristics of participants in the NESARC included in the analyses of risk for transition to bipolar depression are presented in Table 1. These are shown separately for participants with lifetime MDD (first column), participants with past-year MDD (second column), and participants with lifetime MDD who sought treatment (third column). The samples were majority female (approximately two-thirds) and White (approximately four-fifths). Most participants (approximately two-thirds) had attained some college education. Thus, there were no notable differences between the three samples (all MDD cases, past-year cases, and lifetime cases who sought treatment) in the demographic, clinical, and psychosocial factors examined.

We first investigated the phenomenological aspects of MDD that are associated with the transition to bipolar disorder, as well as the influence of lifetime psychiatric disorders on the development of bipolar disorder. Two logistic regression models were fitted, one for the clinical characteristics of depression, and a second that added lifetime disorders (Table 2). Odds ratios greater than 1 indicate factors associated with an increased risk for the transition to bipolar depression. In terms of participant demographic factors, younger age at interview and Black race/ethnicity (compared to White) were associated with higher risks for transition to bipolar depression. In addition, failure to complete high school was associated with a two-fold increased risk for subsequent mania (odds ratio (OR)=1.95; 95% confidence interval (CI)=1.18, 3.22). None of the clinical characteristics of participants' MDD were associated with bipolar transition (e.g., younger age at onset, presence of atypical features); however, a history of other psychiatric disorders was—notably anxiety disorders. Panic disorder (OR=1.53), social phobia (OR=2.20), and generalized anxiety disorder (OR=1.58) were independently associated with the transition to bipolar disorder among participants with a lifetime history of MDD.

The next set of analyses concerned the role of lifetime and past-year stressors in the transition to bipolar disorder. Again, results from two logistic regression models are shown: one for childhood adversities, and a second model adding in past-year stressors. The first finding from these analyses that is notable is that the demographic correlates of transitioning to bipolar disorder observed in Table 2 (higher risks associated with Black race/ethnicity and

lower educational attainment) are substantially reduced in models that include childhood and past-year stressors. Second, a history of childhood abuse, and to a lesser degree sexual maltreatment, were associated with increased risks for transitioning to bipolar depression (OR for childhood abuse=1.26; CI=1.12, 1.42). Third, past-year stressors were significantly associated with bipolar disorder (social support group problems, OR=1.79; economic problems, OR=1.45).

We then conducted an ROC analysis to determine how accurately the group of covariates studied, in the aggregate, predicts the transition to bipolar disorder. The ROC curve is presented in Figure 1—this plots the sensitivity (Y axis) against (1 minus) the specificity (X axis) of cut-points arranged along the distribution of the predicted probabilities of bipolar disorder. The optimal cut-point (indicated by the triangle in Figure 1) has a sensitivity of 65.8% and a specificity of 67.9% for predicting bipolar disorder, and correctly classifies the transition status of 67.9% of the sample. The AUC is 72.4% (CI=69.1%, 75.8%), indicating better-than-chance prediction, but not sufficiently high for use as a clinical screening tool for the future development of bipolar disorder among lifetime cases of MDD.

We conducted two sets of supplementary analyses, in which we repeated the sequence of logistic regression models described above in two subsets of the primary analysis sample: participants with past-year MDD (n=2,585), and participants with lifetime MDD that sought treatment (n=3,798). Similar to the results in Table 2, clinical features of participants' depression were not associated with the risk of bipolar disorder in the supplementary analyses. The pattern of associations between lifetime and recent stressors and subsequent bipolar disorder was also the same in the supplementary analyses. However, subtle differences from the primary sample emerged in the analyses of past-year cases of MDD. Panic disorder was no longer significantly associated with risk for transition to bipolar disorder, whereas cannabis abuse or dependence was associated with it (OR=2.12; CI=1.10, 4.08).

DISCUSSION

Using data from a large national study with 3 years of follow-up, we found that approximately 1 in 20 (4.0%) individuals with a lifetime diagnosis of MDD transitioned to bipolar disorder; the transition rate was similar for individuals with a past-year diagnosis of MDD (5.6%), and for individuals who sought treatment for their depression (4.1%). These transition rates are comparable to rates observed previously in follow-up studies of clinical samples, which have ranged from 4% to 9% across varying time periods,² and the 1% annual conversion rate reported by Angst et al.³²

This study also provides evidence of sociodemographic, clinical, and psychosocial factors that predict the transition from MDD to bipolar disorder. Younger age is one of the strongest demographic correlates of the transition to bipolar depression, which is consistent with a broad literature indicating that the risk for manic episodes peaks between ages 15-20 and declines thereafter.³³ In addition, Black race/ethnicity was associated with a higher risk for bipolar depression, which to our knowledge has not been previously reported, and can be distinguished from the absence of race/ethnicity differences in the risk of Bipolar I disorder in the NESARC sample overall.²¹ Notably, the magnitude of Black-White difference in the transition to bipolar disorder was reduced once past-year stressors were accounted for.

Lower education was also associated with an increased risk of transitioning to bipolar disorder. There is strong evidence that educational attainment is protective against MDD, and in this context, education may also be protective against the transition to bipolar disorder among individuals with MDD.³⁴ Glahn et al. reported that patients with bipolar

disorder had lower levels of educational attainment than demographically matched controls, despite having similar IQ scores; in part their findings were due to the lower likelihood of bipolar patients to complete college.³⁵ This finding illustrates a set of non-causal explanations for our finding that individuals with less than high school educations were twice as likely to transition from MDD to bipolar disorder than individuals with some college education. Consistent with Glahn et al.'s study, our finding may reflect the presence of prodromal symptoms of mania that interfere with the completion of education. This explanation would also be consistent with studies suggesting 'trait' cognitive deficits in bipolar disorder,³⁶ as well as evidence that up to one-third of bipolar individuals recall symptoms at or before age 13.³³ Therefore, while our definition of MDD excluded diagnoses of hypomania, this does not fully eliminate the confounding effects of prodromal or subthreshold symptoms of mania. Additionally, education differences in the risk for mania may reflect the consequences on educational attainment of parental bipolar disorder. Individuals with bipolar disorder are more likely to have parents with bipolar disorder (or other serious mental illness),^{37,38} which itself may interfere with completion of education even in the absence of symptoms in the individual. On the other hand, we note that the association with mania persists even after adjustment for measures of childhood adversity, and that a family history of depression was not predictive of mania.

Among comorbidities, the most robust associations were identified for panic, social phobia and generalized anxiety. Importantly, these symptoms preceded a first manic episode (though not necessarily depressive symptoms), suggesting their potential utility as risk marker for a bipolar course. Previous studies have also linked anxiety to bipolar liability, most notably the findings by Simon et al, which demonstrated a higher prevalence of comorbid panic disorder and generalized anxiety disorder among patients with bipolar disorder.^{4,39} In supplementary analyses (those of past-year cases of MDD), cannabis use disorders were also associated with elevated risk; this finding in particular bears consideration in individuals presenting with a major depressive episode.^{40,41} It is consistent with a previous report associating use of cannabis with risk for subsequent manic symptoms,⁴⁰ and other work associating it with earlier onset age in bipolar or other psychotic disorders.^{42,43}

Our examination of lifetime stressors indicates that childhood adversity was significantly associated with the risk of mania, with an approximately 30% increase in the odds of mania with increasing levels of childhood abuse. These results are consistent with and extend numerous investigations of cohorts of individuals with bipolar disorder that report elevated rates of childhood trauma.⁴⁴⁻⁴⁷ Socioeconomic conditions during childhood and other types of childhood adversities have also been linked with the long-term risk of bipolar disorder.⁴⁸⁻⁵¹

Among stressors in the year prior to initial screen, the most robust risk was associated with social support group problems. This is consistent with a prior study in which the suicide of a family member, in addition to other life events, were associated with risk for first manic hospitalization.⁵² As with our education findings, these may be indicators of more severe illness in general, or emergence of subthreshold manic/hypomanic symptoms, which contribute to conflict with social supports. Alternatively, the social supports may protect against transition to mania, perhaps by buffering the life stresses shown to be associated with mania.⁵²⁻⁵⁴ This model would provide further support for interventions which seek to improve the social functioning of individuals with mood disorder.⁵⁵

The risk factors discussed above could serve as potential risk markers for the transition to bipolar disorder among individuals with MDD. However, in the aggregate, their predictive ability is modest at best, and likely overstates the level of prediction that would be expected

in a replication sample. In the NESARC, a clinical prediction rule based on all of the covariates studied would fail to detect 35% of individuals who transition to bipolar disorder, and falsely identify 32% of individuals as future bipolar cases.

This study focused on the diagnostic conversion from MDD to bipolar I disorder. By not considering hypomania (i.e., bipolar II disorder), our results may not reflect the total impact of the predictors studied on the conversion to bipolar (I and II) disorder. This is notable because evidence suggests that symptoms of mania are common in individuals with MDD, and does not suggest the existence of marked discontinuities between diagnoses of MDD, bipolar II, and bipolar I.^{56,57} This highlights important limitations in the DSM-IV nosology of mood disorders, which might have substantial clinical impact if symptoms of mania go undetected among individuals in treatment for MDD.^{56,58}

As a population-based cohort that relies on structured diagnostic interview administered by non-clinician interviewers, the present study has the following limitations. The assessments of mania in the NESARC have modest psychometric properties, and have not been validated against diagnoses made by clinicians. The lack of assessments of psychosis in the NESARC interview, and small number of individuals with such symptoms, preclude analysis of one of the most frequently identified predictors of bipolar transition. In addition, the NESARC did not assess participants' family history of mania, which is higher among individuals with bipolar disorder than among individuals with MDD.^{4,57} Finally, our analyses of lifetime disorders rely on lifetime reports of disorders, the use of which require additional assumptions be made in mixed-aged samples.^{59,60}

The current study is one of the first population-based, longitudinal studies to investigate the transition from MDD to bipolar disorder, and thus extends the existing evidence which is based on hospitalized or otherwise selectively ascertained populations. In the aggregate, we find some potential predictors of risk for transition to mania which extend those previously reported. None of these is sufficient to identify with confidence those individuals at greatest risk. A key question for future investigation, however, is whether some of these risks might be modifiable: for example, whether strategies to improve resilience or minimize life stresses might reduce the hazard of, or at least delay, onset of mania.

Acknowledgments

We sincerely appreciate the contributions of Ms. Kathleen McGaffigan at the Harvard School of Public Health for data management and statistical programming. A description of the National Epidemiologic Survey on Alcohol and Related Conditions is provided at <http://aspe.hhs.gov/hsp/06/catalog-ai-an-na/nescarc.htm>.

Sources of support: We acknowledge research support from the National Institutes of Health, grants R01MH87544 (PI: Gilman) and R01MH086026 (PI: Perlis).

REFERENCES

1. Perlis RH, Delbello MP, Miyahara S, Wisniewski SR, Sachs GS, Nierenberg AA. Revisiting depressive-prone bipolar disorder: polarity of initial mood episode and disease course among bipolar I systematic treatment enhancement program for bipolar disorder participants. *Biol Psychiatry*. Oct 1; 2005 58(7):549–553. [PubMed: 16197928]
2. Mitchell PB, Goodwin GM, Johnson GF, Hirschfeld RM. Diagnostic guidelines for bipolar depression: a probabilistic approach. *Bipolar Disord*. 2008; 10(1p2):144–152. [PubMed: 18199233]
3. Schaffer A, Cairney J, Veldhuizen S, Kurdyak P, Cheung A, Levitt A. A population-based analysis of distinguishers of bipolar disorder from major depressive disorder. *J Affect Disord*. Mar 9.2010
4. Perlis RH, Brown E, Baker RW, Nierenberg AA. Clinical features of bipolar depression versus major depressive disorder in large multicenter trials. *Am J Psychiatry*. Feb. 2006; 163(2):225–231. [PubMed: 16449475]

5. Andreasen NC, Grove WM, Coryell WH, Endicott J, Clayton PJ. Bipolar versus unipolar and primary versus secondary affective disorder: which diagnosis takes precedence? *J Affect Disord.* 1988; 15(1):69–80. [PubMed: 2970495]
6. Solomon DA, Leon AC, Maser JD, et al. Distinguishing bipolar major depression from unipolar major depression with the screening assessment of depression-polarity (SAD-P). *J Clin Psychiatry.* Mar; 2006 67(3):434–442. [PubMed: 16649831]
7. Serretti A, Mandelli L, Lattuada E, Cusin C, Smeraldi E. Clinical and demographic features of mood disorder subtypes. *Psychiatry Res.* Nov 15; 2002 112(3):195–210. [PubMed: 12450629]
8. Benazzi F. Clinical differences between bipolar II depression and unipolar major depressive disorder: lack of an effect of age. *J Affect Disord.* Jul; 2003 75(2):191–195. [PubMed: 12798259]
9. Guze SB, Woodruff RA Jr, Clayton PJ. The significance of psychotic affective disorders. *Arch Gen Psychiatry.* Sep; 1975 32(9):1147–1150. [PubMed: 1180665]
10. Perlis RH, Ostacher MJ, Uher R, et al. Stability of symptoms across major depressive episodes in bipolar disorder. *Bipolar Disord.* Dec; 2009 11(8):867–875. [PubMed: 19922555]
11. Grant, BF.; Kaplan, K.; Shepard, J.; Moore, T. Source and accuracy statement for Wave 1 of the 2001-2002 National Epidemiology Survey on Alcohol and Related Conditions. National Institute on Alcohol Abuse and Alcoholism; Bethesda, MD: 2003.
12. Grant, BF.; Kaplan, K.; Moore, T.; Kimball, J. Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions: Source and Accuracy Statement. National Institute on Alcohol Abuse and Alcoholism; Bethesda, MD: 2007.
13. Dawson DA, Goldstein RB, Grant BF. Rates and correlates of relapse among individuals in remission from DSM-IV alcohol dependence: a 3-year follow-up. *Alcohol Clin Exp Res.* Dec; 2007 31(12):2036–2045. [PubMed: 18034696]
14. Grant, BF.; Dawson, D. The Alcohol Use Disorder and Associated Disabilities Interview Schedule-IV (AUDADIS-IV). National Institute on Alcohol Abuse and Alcoholism; Rockville, MD: 2001.
15. Grant BG, Towle LH. Standardized diagnostic interviews for alcohol research. *Alcohol Health & Research World.* 1990; 14(4):340–348.
16. Grant BF, Dawson DA, Stinson FS, Chou PS, Kay W, Pickering R. The Alcohol Use Disorder and Associated Disabilities Interview Schedule-IV (AUDADIS-IV): reliability of alcohol consumption, tobacco use, family history of depression and psychiatric diagnostic modules in a general population sample. *Drug Alcohol Depend.* Jul 20; 2003 71(1):7–16. [PubMed: 12821201]
17. Grant BF, Stinson FS, Dawson DA, et al. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry.* Aug; 2004 61(8):807–816. [PubMed: 15289279]
18. American Psychiatric Association. Diagnostic and statistical manual of mental disorders: DSM-IV. 4th ed.. American Psychiatric Association; Washington, DC: 1994.
19. Angst J, Gamma A, Sellaro R, Zhang H, Merikangas K. Toward validation of atypical depression in the community: results of the Zurich cohort study. *J Affect Disord.* Nov; 2002 72(2):125–138. [PubMed: 12200203]
20. Thomas, DC. Statistical methods in genetic epidemiology. Oxford University Press; Oxford ; New York: 2004.
21. Grant BF, Stinson FS, Hasin DS, et al. Prevalence, Correlates, and Comorbidity of Bipolar I Disorder and Axis I and II Disorders: Results From the National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry.* Oct; 2005 66(10):1205–1215. [PubMed: 16259532]
22. Kessler RC, Akiskal HS, Angst J, et al. Validity of the assessment of bipolar spectrum disorders in the WHO CIDI 3.0. *J Affect Disord.* Dec; 2006 96(3):259–269. [PubMed: 16997383]
23. Compton WM, Grant BF, Colliver JD, Glantz MD, Stinson FS. Prevalence of marijuana use disorders in the United States: 1991-1992 and 2001-2002. *JAMA.* May 5; 2004 291(17):2114–2121. [PubMed: 15126440]
24. Grant BF, Dawson DA, Stinson FS, Chou SP, Dufour MC, Pickering RP. The 12-month prevalence and trends in DSM-IV alcohol abuse and dependence: United States, 1991-1992 and 2001-2002. *Drug Alcohol Depend.* Jun 11; 2004 74(3):223–234. [PubMed: 15194200]

25. Grant BF, Hasin DS, Chou SP, Stinson FS, Dawson DA. Nicotine dependence and psychiatric disorders in the United States: results from the national epidemiologic survey on alcohol and related conditions. *Arch Gen Psychiatry*. Nov; 2004 61(11):1107–1115. [PubMed: 15520358]
26. Grant BF, Hasin DS, Stinson FS, et al. Prevalence, correlates, co-morbidity, and comparative disability of DSM-IV generalized anxiety disorder in the USA: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Psychol Med*. Dec; 2005 35(12):1747–1759. [PubMed: 16202187]
27. Ruan WJ, Goldstein RB, Chou SP, et al. The alcohol use disorder and associated disabilities interview schedule-IV (AUDADIS-IV): reliability of new psychiatric diagnostic modules and risk factors in a general population sample. *Drug Alcohol Depend*. Jan 1; 2008 92(1-3):27–36. [PubMed: 17706375]
28. Murphy JM, Berwick DM, Weinstein MC, Borus JF, Budman SH, Klerman GL. Performance of screening and diagnostic tests. Application of receiver operating characteristic analysis. *Arch Gen Psychiatry*. 1987; 44(6):550–555. [PubMed: 3579501]
29. Research Triangle Institute. SUDAAN Language Manual, Release 9.0. Research Triangle. Research Triangle Institute; Park, NC: 2004.
30. Raghunathan TE, Lepkowski JM, Van Hoewyk J, Solenberger P. A multivariate technique for multiply imputing missing values using a sequence of regression models. *Survey Methodology*. 2001; 27(1):85–95.
31. Reiter JP, Raghunathan TE, Kinney SK. The importance of modeling the sampling design in multiple imputation for missing data. *Survey Methodology*. 2006; 32(2):143–149.
32. Angst J, Sellaro R, Stassen HH, Gamma A. Diagnostic conversion from depression to bipolar disorders: results of a long-term prospective study of hospital admissions. *J Affect Disord*. Feb; 2005 84(2-3):149–157. [PubMed: 15708412]
33. Perlis RH, Dennehy EB, Miklowitz DJ, et al. Retrospective age at onset of bipolar disorder and outcome during two-year follow-up: results from the STEP-BD study. *Bipolar Disord*. Jun; 2009 11(4):391–400. [PubMed: 19500092]
34. Lorant V, Deliege D, Eaton W, Robert A, Philippot P, Ansseau M. Socioeconomic inequalities in depression: a meta-analysis. *Am J Epidemiol*. Jan 15; 2003 157(2):98–112. [PubMed: 12522017]
35. Glahn DC, Bearden CE, Bowden CL, Soares JC. Reduced educational attainment in bipolar disorder. *J Affect Disord*. Jun; 2006 92(2-3):309–312. [PubMed: 16524631]
36. Osuji JJ, Cullum CM. Cognition in bipolar disorder. *Psychiatr Clin North Am*. Jun; 2005 28(2):427–441. [PubMed: 15826741]
37. Rende R, Birmaher B, Axelson D, et al. Childhood-onset bipolar disorder: Evidence for increased familial loading of psychiatric illness. *J Am Acad Child Adolesc Psychiatry*. Feb; 2007 46(2):197–204. [PubMed: 17242623]
38. Smoller JW, Finn CT. Family, twin, and adoption studies of bipolar disorder. *Am J Med Genet C Semin Med Genet*. Nov 15; 2003 123(1):48–58. [PubMed: 14601036]
39. Simon NM, Smoller JW, Fava M, et al. Comparing anxiety disorders and anxiety-related traits in bipolar disorder and unipolar depression. *J Psychiatr Res*. May-Jun; 2003 37(3):187–192. [PubMed: 12650739]
40. Henquet C, Krabbendam L, de Graaf R, Ten Have M, van Os J. Cannabis use and expression of mania in the general population. *J Affect Disord*. Oct; 2006 95(1-3):103–110. [PubMed: 16793142]
41. van Laar M, van Dorsselaer S, Monshouwer K, de Graaf R. Does cannabis use predict the first incidence of mood and anxiety disorders in the adult population? *Addiction*. Aug; 2007 102(8):1251–1260. [PubMed: 17624975]
42. Ongur D, Lin L, Cohen BM. Clinical characteristics influencing age at onset in psychotic disorders. *Compr Psychiatry*. Jan-Feb; 2009 50(1):13–19. [PubMed: 19059508]
43. De Hert M, Wampers M, Jendricko T, et al. Effects of cannabis use on age at onset in schizophrenia and bipolar disorder. *Schizophr Res*. Jul 30; 2010
44. Romero S, Birmaher B, Axelson D, et al. Prevalence and correlates of physical and sexual abuse in children and adolescents with bipolar disorder. *J Affect Disord*. Jan; 2009 112(1-3):144–150. [PubMed: 18538857]

45. Leverich GS, Post RM. Course of bipolar illness after history of childhood trauma. *Lancet*. Apr 1; 2006 367(9516):1040–1042. [PubMed: 16581389]
46. Garino JL, Goldberg JF, Ramirez PM, Ritzler BA. Impact of childhood abuse on the clinical course of bipolar disorder. *Br J Psychiatry*. Feb.2005 186:121–125. [PubMed: 15684234]
47. Abram KM, Choe JY, Washburn JJ, Teplin LA, King DC, Dulcan MK. Suicidal ideation and behaviors among youths in juvenile detention. *J Am Acad Child Adolesc Psychiatry*. Mar; 2008 47(3):291–300. [PubMed: 18216737]
48. Tsuchiya KJ, Agerbo E, Byrne M, Mortensen PB. Higher socio-economic status of parents may increase risk for bipolar disorder in the offspring. *Psychol Med*. 2004; 34:1–7.
49. Hyun M, Friedman SD, Dunner DL. Relationship of childhood physical and sexual abuse to adult bipolar disorder. *Bipolar Disord*. Jun; 2000 2(2):131–135. [PubMed: 11252652]
50. Levitan RD, Parikh SV, Lesage AD, et al. Major depression in individuals with a history of childhood physical or sexual abuse: relationship to neurovegetative features, mania, and gender. *Am J Psychiatry*. Dec; 1998 155(12):1746–1752. [PubMed: 9842786]
51. Scott J, Varghese D, McGrath J. As the twig is bent, the tree inclines: adult mental health consequences of childhood adversity. *Arch Gen Psychiatry*. Feb; 2010 67(2):111–112. [PubMed: 20124110]
52. Kessing LV, Agerbo E, Mortensen PB. Major stressful life events and other risk factors for first admission with mania. *Bipolar Disord*. Apr; 2004 6(2):122–129. [PubMed: 15005750]
53. Kennedy S, Thompson R, Stancer HC, Roy A, Persad E. Life events precipitating mania. *Br J Psychiatry*. Apr.1983 142:398–403. [PubMed: 6850179]
54. Ellicott A, Hammen C, Gitlin M, Brown G, Jamison K. Life events and the course of bipolar disorder. *Am J Psychiatry*. Sep; 1990 147(9):1194–1198. [PubMed: 1974746]
55. Miklowitz DJ, Otto MW, Frank E, et al. Intensive psychosocial intervention enhances functioning in patients with bipolar depression: results from a 9-month randomized controlled trial. *Am J Psychiatry*. Sep; 2007 164(9):1340–1347. [PubMed: 17728418]
56. Angst J, Cui L, Swendsen J, et al. Major depressive disorder with subthreshold bipolarity in the National Comorbidity Survey Replication. *Am J Psychiatry*. Oct; 2010 167(10):1194–1201. [PubMed: 20713498]
57. Zimmermann P, Bruckl T, Nocon A, et al. Heterogeneity of DSM-IV major depressive disorder as a consequence of subthreshold bipolarity. *Arch Gen Psychiatry*. Dec; 2009 66(12):1341–1352. [PubMed: 19996039]
58. Keck PE Jr, Kessler RC, Ross R. Clinical and economic effects of unrecognized or inadequately treated bipolar disorder. *J Psychiatr Pract*. May; 2008 14(Suppl 2):31–38. [PubMed: 18677197]
59. Kramer M, von Korff M, Kessler L. The lifetime prevalence of mental disorders: estimation, uses and limitations. *Psychol Med*. Aug; 1980 10(3):429–435. [PubMed: 7443897]
60. Streiner DL, Patten SB, Anthony JC, Cairney J. Has 'lifetime prevalence' reached the end of its life? An examination of the concept. *Int J Methods Psychiatr Res*. Dec; 2009 18(4):221–228. [PubMed: 20052690]

CLINICAL POINTS

Individuals with major depression are at elevated risk for a manic episode, and thus the diagnostic conversion from MDD to bipolar I disorder.

Lower educational attainment, a history of anxiety disorders, and both past-year and childhood psychosocial stressors predicted the development of mania in a population-based sample of individuals with MDD.

The ability of all of these factors combined to predict diagnostic conversion to bipolar I is modest.

Figure 1.

Receiver operating characteristic (ROC) curve for predicting transition to bipolar disorder during the 3 years of follow-up among lifetime cases of MDD in the National Survey on Alcohol and Related Conditions (n=6,214).^a

^aROC curve estimated from a logistic regression model including all of the covariates listed in Tables 2 and 3. The diagonal line indicates chance prediction, corresponding to an area under the curve (AUC) of 0.50. The model AUC is 0.724. The ▲ indicates the optimal cut-point, balancing sensitivity and specificity equally.

Table 1

Descriptive characteristics of participants with lifetime, past-year, and lifetime treated cases of major depressive disorder in the National Epidemiologic Sample on Alcohol and Related Conditions.

	Distribution among participants with lifetime MDD (n=6,214)			Distribution among participants with past-year MDD (n=2,585)			Distribution among participants with lifetime treated MDD (n=3,798)		
	N	%		N	%		N	%	
<i>Demographic factors</i>									
Age									
18-29	1,114	18.9		619	25.0		561	15.7	
30-39	1,339	20.3		563	20.0		800	19.6	
40-49	1,454	25.2		585	24.8		969	27.5	
50-59	1,153	19.5		412	16.5		784	21.2	
60	1,154	16.2		406	13.7		684	16.1	
Sex									
Male	1,827	32.7		751	32.7		946	27.7	
Female	4,387	67.3		1,834	67.3		2,853	72.3	
Race/ethnicity									
White	4,088	78.5		1,582	74.6		2,650	82.2	
Black	856	7.4		412	9.2		418	5.5	
Hispanic	1,004	8.5		463	9.5		584	7.6	
Other	266	5.7		128	6.7		146	4.7	
Education									
Less than high school	950	13.4		478	16.7		473	10.6	
High school or GED	1,683	27.3		733	28.3		975	26.3	
Some college or greater	3,581	59.3		1,374	55.0		2,350	63.1	
<i>Clinical characteristics of depression</i>									
Depression onset 12 years	315	4.9		155	5.7		234	5.9	
First depression onset in the past year	728	11.0		695	26.1		333	7.7	
Presence of atypical features	3,778	60.7		1,653	64.8		2,438	63.4	
>1 Lifetime depressive episode	3,099	50.1		1,649	64.7		2,090	55.5	
High family history loading of depression	1,572	27.2		717	30.1		1,070	30.0	
<i>Lifetime psychiatric disorders</i>									

	Distribution among participants with lifetime MDD (n=6,214)		Distribution among participants with past-year MDD (n=2,585)		Distribution among participants with lifetime treated MDD (n=3,798)	
	N	%	N	%	N	%
Panic disorder	881	14.7	458	18.2	682	18.2
Social phobia	730	12.6	353	15.0	510	14.1
Generalized anxiety disorder	949	16.0	519	20.7	709	19.5
Alcohol dependence	1,223	20.7	541	22.2	739	20.0
Nicotine dependence	1,733	29.8	785	33.0	1,105	30.1
Cannabis abuse or dependence	775	13.2	336	14.2	477	12.7
Other substance dependence	238	3.9	118	4.8	156	4.1
<i>Childhood adversities</i>						
Child abuse (Mean, SE)	0.28	0.0	0.30	0.0	0.3	0.0
Sexual maltreatment (Mean, SE)	4.8	0.0	4.8	0.1	5.0	0.1
Economic deprivation	1,075	15.0	500	16.7	626	14.6
<i>Past-year stressors</i>						
Social support group problems	3,879	62.0	1,739	66.4	2,406	62.7
Social environmental problems	1,786	30.7	907	37.0	1,120	31.2
Occupational problems	3,034	48.9	1,440	55.4	1,936	50.5
Economic problems	1,885	25.6	959	32.1	1,084	23.1

Table 2

Clinical and sociodemographic features associated with the risk for transition from major depressive disorder to bipolar disorder among participants with lifetime MDD in the National Epidemiologic Survey on Alcohol and Related Conditions (n=6,214).

	Model 1: Clinical characteristics of depression		Model 2: Clinical characteristics of depression and lifetime psychiatric disorders	
	OR	CI	OR	CI
<i>Demographic factors</i>				
Age				
18-29	5.58 [*]	2.71, 11.49	5.32 [*]	2.58, 10.99
30-39	5.02 [*]	2.40, 10.49	4.62 [*]	2.20, 9.71
40-49	2.74 [*]	1.31, 5.73	2.33 [*]	1.09, 4.95
50-59	2.85 [*]	1.27, 6.41	2.61 [*]	1.16, 5.88
60	1		1	
Sex				
Male	1.07	0.75, 1.52	1.08	0.74, 1.58
Female	1		1	
Race/ethnicity				
White	1		1	
Black	1.59	0.96, 2.62	1.72 [*]	1.04, 2.84
Hispanic	0.89	0.53, 1.50	0.94	0.55, 1.62
Other	0.72	0.30, 1.73	0.73	0.31, 1.73
Education				
Less than high school	2.00 [*]	1.22, 3.27	1.95 [*]	1.18, 3.22
High school or GED	1.14	0.76, 1.70	1.12	0.75, 1.68
Some college or greater	1		1	
<i>Clinical characteristics of depression</i>				
Depression onset 12 years	1.37	0.79, 2.37	1.26	0.73, 2.16
First depression onset in the past year	0.69	0.39, 1.22	0.69	0.39, 1.22
Presence of atypical features	1.07	0.77, 1.48	1.03	0.74, 1.45
Number of lifetime depressive episodes	1.00 [*]	1.00, 1.01	1.00	0.99, 1.01
High family history loading of depression	1.04	0.74, 1.47	0.93	0.66, 1.31
<i>Lifetime psychiatric disorders</i>				
Panic disorder			1.53	0.99, 2.39
Social phobia			2.20 [*]	1.47, 3.30
Generalized anxiety disorder			1.58 [*]	1.06, 2.35
Alcohol dependence			1.07	0.68, 1.68
Nicotine dependence			1.11	0.75, 1.63
Cannabis abuse or dependence			1.27	0.77, 2.11
Other substance dependence			0.80	0.35, 1.81

*
P<0.05

Table 3

Lifetime and recent stressors associated with the risk for transition from major depressive disorder to bipolar disorder among participants with lifetime MDD in the National Epidemiologic Survey on Alcohol and Related Conditions (n=6,214).

	<u>Model 1: Childhood adversities</u>		<u>Model 2: Childhood adversities and past-year stressors</u>	
	OR	CI	OR	CI
<i>Demographic factors</i>				
Age				
18-29	5.32 *	2.59, 10.91	4.59 *	2.14, 9.81
30-39	4.50 *	2.15, 9.44	4.21 *	1.98, 8.94
40-49	2.54 *	1.21, 5.31	2.40 *	1.14, 5.07
50-59	2.59 *	1.16, 5.77	2.49 *	1.11, 5.60
60	1		1	
Sex				
Male	1.18	0.82, 1.69	1.24	0.87, 1.77
Female	1		1	
Race/ethnicity				
White	1		1	
Black	1.56	0.96, 2.54	1.44	0.88, 2.36
Hispanic	0.87	0.51, 1.47	0.82	0.48, 1.39
Other	0.68	0.28, 1.62	0.65	0.27, 1.56
Education				
Less than high school	1.71 *	1.03, 2.84	1.45	0.90, 2.32
High school or GED	1.06	0.70, 1.59	1.02	0.68, 1.54
Some college or greater	1		1	
<i>Childhood adversities</i>				
Child abuse	1.27 *	1.13, 1.42	1.26 *	1.12, 1.42
Sexual maltreatment	1.05 *	1.00, 1.11	1.05 *	1.00, 1.11
Economic deprivation	1.02	0.70, 1.49	0.98	0.67, 1.43
<i>Past-year stressors</i>				
Social support group problems			1.79 *	1.19, 2.68
Social environmental problems			1.10	0.75, 1.60
Occupational problems			1.18	0.80, 1.72
Economic problems			1.45 *	1.03, 2.06

* P<0.05

Table 4

Clinical and psychosocial predictors of the transition to bipolar disorder among participants with past-year MDD, and with lifetime treated MDD, in the National Epidemiologic Survey on Alcohol and Related Conditions.

	Analysis sample: Past-Year Cases of Depression at Wave 1 (n=2,585)				Analysis sample: Lifetime Cases of Depression who Sought Treatment (n=3,798)			
	Model 1: Clinical characteristics of depression and lifetime psychiatric disorders		Model 2: Childhood adversities and past-year stressors		Model 3: Clinical characteristics of depression and lifetime psychiatric disorders		Model 4: Childhood adversities and past-year stressors	
	OR	CI	OR	CI	OR	CI	OR	CI
<i>Demographic factors</i>								
Age								
18-29	3.14 [*]	1.16, 8.52	2.84 [*]	1.03, 7.78	5.56 [*]	2.11, 14.62	4.75 [*]	1.78, 12.70
30-39	4.19 [*]	1.51, 11.61	4.00 [*]	1.44, 11.12	4.51 [*]	1.74, 11.69	4.41 [*]	1.67, 11.61
40-49	1.50	0.54, 4.21	1.54	0.55, 4.33	1.89	0.72, 4.94	2.06	0.78, 5.46
50-59	2.41	0.84, 6.94	2.05	0.72, 5.86	2.95 [*]	1.04, 8.39	2.86 [*]	1.02, 8.03
60	1		1		1		1	
Sex								
Male	0.78	0.45, 1.36	0.94	0.55, 1.61	0.98	0.61, 1.57	1.23	0.76, 2.00
Female	1		1		1		1	
Race/ethnicity								
White	1		1		1		1	
Black	1.55	0.81, 2.95	1.41	0.74, 2.69	1.39	0.67, 2.87	1.16	0.57, 2.35
Hispanic	0.83	0.40, 1.71	0.63	0.30, 1.33	1.00	0.51, 1.95	0.96	0.51, 1.83
Other	0.49	0.15, 1.63	0.40	0.11, 1.38	1.27	0.54, 2.98	1.06	0.45, 2.49
Education								
Less than high school	1.98 [*]	1.06, 3.68	1.40	0.73, 2.68	1.49	0.79, 2.83	1.14	0.58, 2.24
High school or GED	1.51	0.85, 2.67	1.30	0.73, 2.33	1.11	0.64, 1.93	1.02	0.61, 1.73
Some college or greater	1		1		1		1	
<i>Clinical characteristics of depression</i>								
Depression onset 12 years	1.85	0.95, 3.60			1.33	0.71, 2.49		
First depression onset in the past year	0.57	0.32, 1.03			0.81	0.40, 1.62		
Presence of atypical features	1.36	0.87, 2.14			0.94	0.65, 1.37		

		Analysis sample: Past-Year Cases of Depression at Wave 1 (n=2,585)				Analysis sample: Lifetime Cases of Depression who Sought Treatment (n=3,798)			
		Model 1: Clinical characteristics of depression and lifetime psychiatric disorders		Model 2: Childhood adversities and past-year stressors		Model 3: Clinical characteristics of depression and lifetime psychiatric disorders		Model 4: Childhood adversities and past-year stressors	
		OR	CI	OR	CI	OR	CI	OR	CI
Number of lifetime depressive episodes		1.00	0.98, 1.01			1.00	0.99, 1.01		
High family history loading of depression		0.93	0.55, 1.57			0.91	0.59, 1.39		
<i>Lifetime psychiatric disorders</i>									
Panic disorder		1.04	0.60, 1.78			1.62 [*]	1.04, 2.52		
Social phobia		2.08 [*]	1.19, 3.65			2.12 [*]	1.31, 3.44		
Generalized anxiety disorder		1.35	0.76, 2.41			1.60 [*]	1.00, 2.57		
Alcohol dependence		0.73	0.40, 1.31			1.22	0.73, 2.02		
Nicotine dependence		0.91	0.53, 1.58			1.15	0.72, 1.82		
Cannabis abuse or dependence		2.12 [*]	1.10, 4.08			1.44	0.85, 2.43		
Other substance dependence		1.08	0.42, 2.80			0.75	0.33, 1.70		
<i>Childhood adversities</i>									
Child abuse				1.32 [*]	1.13, 1.54			1.26 [*]	1.06, 1.49
Sexual maltreatment				1.03	0.96, 1.11			1.05	0.98, 1.12
Economic deprivation				0.91	0.54, 1.51			1.00	0.63, 1.57
<i>Past-year stressors</i>									
Social support group problems				1.55	0.92, 2.60			1.76 [*]	1.10, 2.83
Social environmental problems				1.26	0.77, 2.06			1.11	0.71, 1.74
Occupational problems				1.15	0.71, 1.86			1.20	0.76, 1.91
Economic problems				1.62 [*]	1.02, 2.58			1.46	0.96, 2.21

* P<0.05