Published in final edited form as:

J Hosp Med. 2008 May ; 3(3): 212–217. doi:10.1002/jhm.309.

Prevalence and Clinical Outcome of Inpatient Hyperglycemia in a Community Pediatric Hospital

Andres Palacio, M.D., Miguel Ceron, M.D., Robin Klein, M.D., Irene S. Cho, M.D., Dawn Smiley, M.D., Roberto Mejia, and Guillermo E. Umpierrez, M.D.

Department of Medicine, Division of Endocrinology, Emory University School of Medicine, Atlanta, Georgia

Abstract

Background—Inpatient hyperglycemia in adult patients with and without a history of diabetes is a predictor of poor clinical outcome. No previous studies; however, have examined the association of hyperglycemia and clinical outcome in children admitted to a community pediatric hospital.

Methods—Retrospective observational cohort of pediatric patients admitted to a community children's hospital, from January 2004 to August 2004. Medical records of 903 consecutive children admitted to critical and non-critical care areas were reviewed. Of them, 342 patients (38%) had no blood glucose measurements during the hospital stay. In the remaining patients, we determined the prevalence of hyperglycemia and examined the association of hyperglycemia and clinical outcome.

Results—A total of 406 patients (75%) had an admission blood glucose ≥ 120 mg/dl [(mean 98 ± 1 mg/dl ($\pm$ SEM)], 103 children (19%) had an admission blood glucose level between 121–179 mg/dl (mean 143 ± 2 mg/dl), and 32 patients (5.9%) had a blood glucose level ≥ 180 mg/dl (mean 260 ± 18 mg/dl). Seventeen patients (13%) had a known history of diabetes prior to admission. Children with hyperglycemia were more likely to be admitted to the ICU ($p < 0.001$) and had a longer length of ICU stay ($p < 0.001$), but admission hyperglycemia was not associated with longer hospital stay or higher hospital mortality.

Summary—Hyperglycemia is present in one-fourth of children admitted to the hospital, most of them without a history of diabetes prior to admission. Hyperglycemia was associated with a greater need for ICU care and longer ICU stay but not with increased in-hospital mortality.

Introduction

Diabetes is one of the most common diagnoses in hospitalized patients (1; 2). Hyperglycemia is present in 38% of adults admitted to the hospital and one third of these patients had no history of diabetes before the admission (3). The impact of inpatient hyperglycemia on clinical outcome in adult patients has been increasingly appreciated. Extensive evidence from observational studies indicates that hyperglycemia in patients with or without a history of diabetes is an important marker of poor clinical outcome (3–12). Several prospective randomized trials in patients with critical illness have shown that aggressive glycemic control improves short- and long-term mortality, multiorgan failure and systemic infection, and length of hospitalization (13–17). The importance of glucose control also applies to adult patients admitted to general surgical and medical wards (3; 6; 18). In

such patients, we recently reported that the presence of hyperglycemia is associated with prolonged hospital stay, infection, disability after hospital discharge, and death (3; 6; 18). Despite the extensive data in adult patients, there is little information on the impact of inpatient hyperglycemia in pediatric patients. Only a few observational studies in critically ill children admitted to pediatric ICU, with severe brain injury or with extensive burn injuries have shown a positive association between inpatient hyperglycemia and increased length of hospital and ICU stay, and higher risk of complication and mortality rates (19–23). No previous studies; however, have examined the association of hyperglycemia and clinical outcome in children admitted to a general community pediatric hospital. Therefore, in this study we determined the prevalence of inpatient hyperglycemia and examined the impact of hyperglycemia on morbidity and mortality in children admitted to Hughes Spalding Children's Hospital, a large community hospital serving the inner city and indigent pediatric population in Atlanta, Georgia.

Methods

This is a retrospective observational cohort of pediatric patients consecutively admitted to Hughes Spalding Children's Hospital in Atlanta, from January 2004 to August 2004. The Hughes Spalding Children's Hospital is a general community pediatric hospital part of the Grady Health System in Atlanta, a large healthcare organization that operates under the auspices of the Fulton-Dekalb Hospital Authority -- the major counties in metropolitan Atlanta -- to deliver care to their uninsured and underserved populations. Ninety percent of the organization's inpatient cases are either uninsured or dependent on Medicaid. This is a broad-based pediatric hospital without cardiac surgery, burn unit, or dedicated inpatient hematology-oncology unit. Patients are managed by members of the pediatric residency program and supervised by a faculty member from Emory University School of Medicine. The Institutional Review Board of Emory University and Grady Health System Oversight Research Committee approved methods for data collection and analysis used in the study and waived the need for informed consent.

The medical records of 903 consecutive pediatric patients admitted to both critical and noncritical care areas were reviewed. For the analysis, patients were divided according to a known history of diabetes prior to admission, and according to the admission blood glucose concentration. A normoglycemic group included patients with normal plasma glucose and without a history of diabetes. Serum or plasma glucose measured in the laboratory was assumed to be equivalent to blood glucose measured by finger stick at bedside using a glucose meter. Hyperglycemia was defined as an admission or in-hospital blood glucose level greater than 120 mg/dl. High blood glucose was subsequently divided into those with blood glucose between 120 and 179 mg/dl and those with blood glucose $\geq$ 180 mg/dl. Patient information was collected regarding demographic characteristics, blood glucose levels on admission and during hospital stay, concurrent medical diagnoses, medical treatment, and hospital outcome (including mortality and disposition at discharge).

The primary objectives of this study were to determine the prevalence of in-hospital hyperglycemia and to examine the association of hyperglycemia and mortality in children with critical and noncritical illness in a community pediatric hospital. Secondary end points included length of hospital, requirement of intensive care, and treatment of hyperglycemia. In addition to blood glucose levels, prognostic variables included gender, age, body mass index, admission diagnosis, presence of co-morbidities, intensive care unit admission.

Statistical Analysis

To compare demographics and clinical characteristics between groups, independent t-test and ANOVA with Sheffé's method was used for continuous variables. Levine's test for homogeneity of variances and log transformations were used when necessary. For categorical variables, χ^2 analysis was used. $P < .05$ was considered significant. SPSS version 12.0 (SPSS, Inc., Chicago, IL), was the statistical software used for the analysis.

Results

Of the 903 admitted patients, 342 patients (38%) had no blood glucose measurement during the hospital stay and were excluded from the analysis. Three patients with a length of stay greater than 6 months were excluded. In addition, 16 patients admitted with diabetic ketoacidosis (DKA) and 1 subject with hyperglycemic hyperosmolar syndrome (HHS) were also excluded from the analysis. The remaining 542 patients constituted the study population. The majority of these patients, 406 patients (75%) had an admission blood glucose concentration equal to or less than 120 mg/dl [mean 98 ± 1 mg/dl ($\pm$ SEM), median: 93 mg/dl]. A total of 103 children (19%) had an admission blood glucose level between 121–179 mg/dl (mean 143 ± 2 mg/dl, median: 140 mg/dl), and 32 patients (5.9%) had an admission blood glucose level greater than 180 mg/dl (mean 260 ± 18 mg/dl, median: 211 mg/dl) (Figure 1).

The clinical characteristics of study patients are shown in Table 1. Most patients in this study were from minority ethnic groups and included 82% blacks, 12% Hispanics, 2% other minority groups, and 4.2% Caucasians. There were no significant differences in the mean age, gender, racial distribution, or body mass index among the three groups. A total of 409 patients (75.5%) were admitted to general pediatric wards and 133 patients (24.5%) were admitted to the surgical unit. There were no differences in the admission blood glucose between patients admitted to general pediatric wards (112.2 mg/dl) compared to that of children admitted to surgical areas (115.7 mg/dl, $p > 0.05$). The most common diagnoses in the severe hyperglycemia group were trauma/surgery (25%), pulmonary disease (18.8%), metabolic disorders (12.5%), and infection (6.3%). Most children admitted with hyperglycemia had no previous history of diabetes prior to admission. Among the 135 children with admission hyperglycemia (blood glucose > 120 mg/dl), 17 patients (13%) had a known history of diabetes or were receiving therapy prior to admission. The mean admission blood glucose was 162.4 mg/dl (range 121 to 480 mg/dl) in children with new hyperglycemia and was 369.8 mg/dl (range 145 to 678 mg/dl) in those children with a known history of diabetes, ($p < 0.01$). Among children without a history of diabetes, 33 out of 118 children (28%) with admission hyperglycemia had one or more glucose values > 120 mg/dl during the hospitalization course. Twenty five children had blood glucose between 121–179 mg/dl (mean 109 ± 5 mg/dl) and 8 children had a blood glucose ≥ 180 mg/dl (mean 159 ± 13 mg/dl). Most patients with a history of diabetes were admitted with significant hyperglycemia. One patient (1%) between the 121–179 mg/dL category, and 16 (50%) of patients had glucose levels greater than 180 mg/dL.

The presence of hyperglycemia on admission in pediatric patients was not associated with increased mortality or with increased hospital stay. Only one death was reported during the study period and it occurred in a patient with respiratory failure due to bronchiolitis who was admitted with admission blood glucose of 151 mg/dl. The mean length of stay for patients with normoglycemia was 3.83 ± 0.2 days, which increased to 5.36 ± 1.0 days and 5.68 ± 1.8 days for children with blood glucose between 120–179 mg/dl and ≥ 180 mg/dl, respectively ($p > 0.05$).

Children with hyperglycemia were more likely to be admitted to the ICU, and had a longer length of ICU stay. Admission to the ICU was needed in 10% of children with an admission blood glucose < 120 mg/dl, 18% of children with a blood glucose between 120–179 mg/dl and in 40% of subjects with an admission blood glucose ≥ 180 mg/dl ($p < 0.01$). In addition, the ICU length of stay was significantly longer in hyperglycemic children, particularly among those patients with glucose levels ≥ 180 mg/dl ($p < 0.001$). The mean ICU length of stay (ICU) was 0.56 ± 0.1 days for patients with normoglycemia, and 1.1 ± 0.4 days and 3.6 ± 1.9 days for patients with blood glucose between 120–179 mg/dl and ≥ 180 mg/dl, respectively ($p < 0.01$).

Newly diagnosed hyperglycemia was frequently left untreated. Only 3 children without a history of diabetes but with hyperglycemia recorded during the hospital stay received insulin therapy. New hyperglycemia patients received regular insulin per sliding scale as the main insulin regimen in the hospital. In contrast, all patients with previous history of diabetes were treated with insulin during the hospital stay.

Discussion

Diabetes mellitus represents a significant public health burden on the basis of increased morbidity, mortality, and economic cost. Increasing evidence from observational and prospective interventional studies have shown that inpatient is a predictor of poor clinical outcome in adult subjects (3–13; 16; 17). Admission hyperglycemia has been associated with increased morbidity and mortality in patients with critical illnesses, as well as in non-critically ill adult subjects admitted to general surgical and medical wards (3; 6; 18). In this study we also found that hyperglycemia is a common finding in children admitted with critical and non-critical illness, and that most children had no history of diabetes before the admission. Admission hyperglycemia was present in one-fourth of the children admitted to the hospital. Children with hyperglycemia were more likely to be admitted to the ICU and had a longer length of ICU stay; however, inpatient hyperglycemia was not associated with higher hospital mortality or longer hospital stay than children with normoglycemia. Our findings suggest that recognition of inpatient hyperglycemia can be improved because screening for hyperglycemia was not performed in more than one-third of patients (38%) during the hospital stay.

The prevalence of inpatient hyperglycemia in children varies according to the severity of illness and study population. In critically ill children, Ruiz Magro et al (21) reported that 50% of 353 patients without diabetes mellitus had initial glucose values >120 mg/dL. Faustino et al (20) in a study of 942 nondiabetic patients found a prevalence of hyperglycemia within 24 hours of admission to the ICU of 70.4% in patients with a glucose value >120 mg/dL, 44.5% in patients with a glucose value >150 mg/dL, and 22.3% with a value > 200 mg/dL. The prevalence of hyperglycemia for noncritically ill children seen in the emergency department is much lower with a range from 3.8% to 5.0% (based on the initial blood glucose being > 150 mg/dL) (19; 24). In agreement with these studies, we found that inpatient hyperglycemia is a common finding among hospitalized children. Approximately 75% of our patients had a normal blood glucose on admission, 19% had an admission blood glucose between 121–179 mg/dl (mean 143 ± 2 mg/dl), and 5.9% of children had an admission blood glucose ≥ 180 mg/dl (mean 260 ± 18 mg/dl). Only 13 per cent of our patients had a known history of diabetes prior to admission suggesting that the hyperglycemia was the result of stress of the medical or surgical illness. Stress hyperglycemia, defined as a transient increase in blood glucose level during an acute physiological stress, has been reported to occur in 4% of children with an acute non-critical illness, and in > 50% of children in the ICU.

A few studies have reported on the impact of inpatient hyperglycemia in children with acute critical illness [10–15]. Three retrospective studies have demonstrated that admission hyperglycemia is also a predictor of adverse outcomes in the pediatric intensive care unit (20; 22). Srinivasan and colleagues (22) demonstrated that 86% of patients in their pediatric intensive care unit had a glucose value >126 mg/dL at some point during their stay. In addition, they showed that the duration of hyperglycemia and the peak glucose were also associated with mortality. Faustino and Apkon (20) demonstrated that hyperglycemia occurs frequently among critically ill nondiabetic children and is correlated with a greater in-hospital mortality rate and longer length of stay in the ICU. They reported a 2.5-fold increased risk of dying if the maximum glucose obtained within 24 hours of admission to the ICU was >150 mg/dL. More recently, Yates et al (25) reported that hyperglycemia in the postoperative period was associated with increased morbidity and mortality in postoperative pediatric cardiac patient. Other studies in children with traumatic brain or head injury have also shown an association between poor neurologic outcome and elevated admission blood glucose (24; 26–28). Brain trauma patients with permanent neurological deficit and vegetative state were found to have significantly higher admission blood glucose concentration than children with good neurological recovery or minimal deficit. In addition, the development of inpatient hyperglycemia in children with extensive burn injuries greater than 60% of total body surface area was found to increase the risk of bacteremia, fungemia, reduced skin graft take, and increased mortality rate (29). These data show an association with initial glucose, peak glucose, and duration of hyperglycemia with increased incidence of morbidity and mortality in children with acute critical illness. We found no association between initial blood glucose and risk for death. This is in contrast to our previous results in adult subjects in whom inpatient hyperglycemia was found to represent an important marker of increased morbidity and mortality among both critically ill and non-critically ill adult patients (3). It is important to notice that the overall mortality rate reported in children with hyperglycemia relates to severity of illness and is significantly lower than that of adults (30). In most critically ill pediatric series, the hospital mortality range from 2% to 5.3% and is higher in patients with severe trauma and major cardiac surgery (23; 31). The mortality in children without critical illness admitted to general pediatric wards is significantly lower (30).

In agreement with the increasing rate of obesity among children with diabetes (32; 33), especially in minority populations, we found that hospitalized children with a history of diabetes and glucose > 180 mg/dL had higher body mass index (BMI) than those with normoglycemia ($p<0.001$). Obesity in children has been associated with an the presence of several comorbidities and an increased risk of hospital complications (34; 35). There is also increasing evidence among patients admitted to the intensive care unit (ICU) that obesity contributes to increased morbidity and to a prolonged length of stay (35). Because of the higher rate of hyperglycemia, diabetes, and hospital complications, we believe that obese children should be screened for hyperglycemia and diabetes.

We acknowledge the following limitations in this study. The main limitation of this study is its retrospective nature. The method of blood glucose collection and analysis was not standardized, thus it prevented uniformity in the collection of serum glucose values for individual patients. We arbitrarily used three cutoff glucose values in this study (<120 mg/dL, 120–179 mg/dL, and > 180 mg/dL). Although similar values have been used in inpatient diabetes studies (20–22), there is no uniform definition of hyperglycemia in hospitalized patients and the clinical significance of these cutoff values in pediatric population has not been determined. The study was conducted in a single institution in Atlanta with a population and disease spectrum that might be different from the other pediatric institutions. Our study did not address the question of whether treatment of hyperglycemia may improve outcome of length of hospital stay in patient with hyperglycemia. We believe that newly

diagnosed hyperglycemia is usually considered a transient finding in response to the acute illness not requiring medical intervention, as indicated by the fact that more than half of these patients did not receive antidiabetic therapy. Another limitation of our study is that we were not able to determine the percentage of patients with latent or unrecognized diabetes because of the lack of hemoglobin A1C testing and the lack of follow-up after discharge. A prospective, randomized trial of strict glycemic control is certainly needed to address these issues.

In summary, inpatient hyperglycemia is a common finding in children with and without critical illness. Hyperglycemia is present in one-fourth of the children admitted to the hospital, most of them without a history of diabetes prior to admission. Although we found a higher need for ICU admission and a longer length of ICU stay, the presence of hyperglycemia was not associated with higher hospital mortality compared to children with normoglycemia. Several observational studies have reported an association of hyperglycemia and poor clinical outcome in critical ill children; however, no prospective controlled studies have assessed the effect of tight glucose control in pediatric populations. These studies need to be prospective randomized multicenter trials of sufficient magnitude to provide a power analysis to allow for multiple observations and evaluation of subsets of critically and noncritically ill pediatric patients.

Acknowledgments

Dr Umpierrez is supported by research grants from the American Diabetes Association (7-03-CR-35), American Heart Association (0555306B), and National Institutes of Health: R03 DK073190-01 and General Clinical Research Center Grant M01 RR-00039.

REFERENCES

1. Mokdad AH, Ford ES, Bowman BA, Nelson DE, Engelgau MM, Vinicor F, Marks JS. Diabetes trends in the U.S.: 1990–1998. *Diabetes Care*. 2000; 23:1278–1283. [PubMed: 10977060]
2. Levetan CS, Passaro M, Jablonski K, Kass M, Ratner RE. Unrecognized diabetes among hospitalized patients. *Diabetes Care*. 1998; 21:246–249. [PubMed: 9539990]
3. Umpierrez GE, Isaacs SD, Bazargan N, You X, Thaler LM, Kitabchi AE. Hyperglycemia: an independent marker of in-hospital mortality in patients with undiagnosed diabetes. *J Clin Endocrinol Metab*. 2002; 87:978–982. [PubMed: 11889147]
4. Braithwaite SS, Buie MM, Thompson CL, Baldwin DF, Oertel MD, Robertson BA, Mehrotra HP. Hospital hypoglycemia: not only treatment but also prevention. *Endocr Pract*. 2004; 10(Suppl 2): 89–99. [PubMed: 15251646]
5. Capes SE, Hunt D, Malmberg K, Gerstein HC. Stress hyperglycaemia and increased risk of death after myocardial infarction in patients with and without diabetes: a systematic overview. *Lancet*. 2000; 355:773–778. [PubMed: 10711923]
6. Clement S, Braithwaite SS, Magee MF, Ahmann A, Smith EP, Schafer RG, Hirsh IB. Management of diabetes and hyperglycemia in hospitals. *Diabetes Care*. 2004; 27:553–597. [PubMed: 14747243]
7. Estrada CA, Young JA, Nifong LW, Chitwood WR Jr. Outcomes and perioperative hyperglycemia in patients with or without diabetes mellitus undergoing coronary artery bypass grafting. *Ann Thorac Surg*. 2003; 75:1392–1399. [PubMed: 12735552]
8. Mizock BA. Blood glucose management during critical illness. *Rev Endocr Metab Disord*. 2003; 4:187–194. [PubMed: 12766547]
9. Montori VM, Bistrian BR, McMahon MM. Hyperglycemia in acutely ill patients. *Jama*. 2002; 288:2167–2169. [PubMed: 12413377]
10. Umpierrez GE, A.E. K. ICU Care for Patients with Diabetes. *Current Opinions Endocrinol*. 2004; 11:75–81.

11. Norhammar AM, Ryden L, Malmberg K. Admission plasma glucose. Independent risk factor for long-term prognosis after myocardial infarction even in nondiabetic patients. *Diabetes Care*. 1999; 22:1827–1831. [PubMed: 10546015]
12. Finney SJ, Zekveld C, Elia A, Evans TW. Glucose control and mortality in critically ill patients. *Jama*. 2003; 290:2041–2047. [PubMed: 14559958]
13. Krinsley JS. Association between hyperglycemia and increased hospital mortality in a heterogeneous population of critically ill patients. *Mayo Clin Proc*. 2003; 78:1471–1478. [PubMed: 14661676]
14. Kitabchi AE, Umpierrez GE, Murphy MB, Barrett EJ, Kreisberg RA, Malone JI, Wall BM. Management of hyperglycemic crises in patients with diabetes. *Diabetes Care*. 2001; 24:131–153. [PubMed: 11194218]
15. Malmberg K. Prospective randomised study of intensive insulin treatment on long term survival after acute myocardial infarction in patients with diabetes mellitus. DIGAMI (Diabetes Mellitus, Insulin Glucose Infusion in Acute Myocardial Infarction) Study Group. *Bmj*. 1997; 314:1512–1515. [PubMed: 9169397]
16. Van den Berghe G, Wilmer A, Hermans G, Meersseman W, Wouters PJ, Milants I, Van Wijngaerden E, Bobbaers H, Bouillon R. Intensive insulin therapy in the medical ICU. *N Engl J Med*. 2006; 354:449–461. [PubMed: 16452557]
17. van den Berghe G, Wouters P, Weekers F, Verwaest C, Bruyninckx F, Schetz M, Vlasselaers D, Ferdinande P, Lauwers P, Bouillon R. Intensive insulin therapy in the critically ill patients. *N Engl J Med*. 2001; 345:1359–1367. [PubMed: 11794168]
18. Pomposelli JJ, Baxter JK 3rd, Babineau TJ, Pomfret EA, Driscoll DF, Forse RA, Bistrian BR. Early postoperative glucose control predicts nosocomial infection rate in diabetic patients. *JPEN J Parenter Enteral Nutr*. 1998; 22:77–81. [PubMed: 9527963]
19. Bhisitkul DM, Morrow AL, Vinik AI, Shults J, Layland JC, Rohn R. Prevalence of stress hyperglycemia among patients attending a pediatric emergency department. *J Pediatr*. 1994; 124:547–551. [PubMed: 8151468]
20. Faustino EV, Apkon M. Persistent hyperglycemia in critically ill children. *J Pediatr*. 2005; 146:30–34. [PubMed: 15644818]
21. Ruiz Magro P, Aparicio Lopez C, Lopez-Herce Cid J, Martinez Campos M, Sancho Perez L. Metabolic changes in critically ill children. *An Esp Pediatr*. 1999; 51:143–148. [PubMed: 10495500]
22. Srinivasan V, Spinella PC, Drott HR, Roth CL, Helfaer MA, Nadkarni V. Association of timing, duration, and intensity of hyperglycemia with intensive care unit mortality in critically ill children. *Pediatr Crit Care Med*. 2004; 5:329–336. [PubMed: 15215001]
23. Tenner PA, Dibrell H, Taylor RP. Improved survival with hospitalists in a pediatric intensive care unit. *Crit Care Med*. 2003; 31:847–852. [PubMed: 12626995]
24. Valerio G, Franzese A, Carlin E, Pecile P, Perini R, Tenore A. High prevalence of stress hyperglycaemia in children with febrile seizures and traumatic injuries. *Acta Paediatr*. 2001; 90:618–622. [PubMed: 11440092]
25. Yates AR, Dyke PC 2nd, Taeed R, Hoffman TM, Hayes J, Feltes TF, Cua CL. Hyperglycemia is a marker for poor outcome in the postoperative pediatric cardiac patient. *Pediatr Crit Care Med*. 2006; 7:351–355. [PubMed: 16738506]
26. Cochran A, Scaife ER, Hansen KW, Downey EC. Hyperglycemia and outcomes from pediatric traumatic brain injury. *J Trauma*. 2003; 55:1035–1038. [PubMed: 14676647]
27. Chiaretti A, De Benedictis R, Langer A, Di Rocco C, Bizzarri C, Iannelli A, Polidori G. Prognostic implications of hyperglycaemia in paediatric head injury. *Childs Nerv Syst*. 1998; 14:455–459. [PubMed: 9808254]
28. Paret G, Barzilai A, Lahat E, Feldman Z, Ohad G, Vardi A, Ben-Abraham R, Barzilai Z. Gunshot wounds in brains of children: prognostic variables in mortality, course, and outcome. *J Neurotrauma*. 1998; 15:967–972. [PubMed: 9840769]
29. Gore DC, Chinkes D, Hegggers J, Herndon DN, Wolf SE, Desai M. Association of hyperglycemia with increased mortality after severe burn injury. *J Trauma*. 2001; 51:540–544. [PubMed: 11535907]

30. Landrigan CP, Srivastava R, Muret-Wagstaff S, Soumerai SB, Ross-Degnan D, Graef JW, Homer CJ, Goldmann DA. Impact of a health maintenance organization hospitalist system in academic pediatrics. *Pediatrics*. 2002; 110:720–728. [PubMed: 12359785]
31. Chang RK, Klitzner TS. Can regionalization decrease the number of deaths for children who undergo cardiac surgery? A theoretical analysis. *Pediatrics*. 2002; 109:173–181. [PubMed: 11826192]
32. Rosenbloom AL, Joe JR, Young RS, Winter WE. Emerging epidemic of type 2 diabetes in youth. *Diabetes Care*. 1999; 22:345–354. [PubMed: 10333956]
33. Von Karla V, Hewett ML. Type 2 diabetes in children and adolescents: screening, diagnosis, and management. *Jaapa*. 2007; 20:51–54. [PubMed: 17416023]
34. Nafiu OO, Reynolds PI, Bamgbade OA, Tremper KK, Welch K, Kasa-Vubu JZ. Childhood body mass index and perioperative complications. *Paediatr Anaesth*. 2007; 17:426–430. [PubMed: 17474948]
35. Carroll CL, Bhandari A, Zucker AR, Schramm CM. Childhood obesity increases duration of therapy during severe asthma exacerbations. *Pediatr Crit Care Med*. 2006; 7:527–531. [PubMed: 17006390]

Figure 1.

Hyperglycemia: A Common Comorbidity in Medical-Surgical Patients Admitted to a Community Pediatric Hospital.

A total of 24.9 per cent of children admitted to a community children's hospital were found to have high blood glucose on admission. Hyperglycemia was defined as an admission blood glucose level > 120 mg/dl and subsequently divided into those with blood glucose between 120 and 179 mg/dl and those with blood glucose ≥ 180 mg/dl.

Table 1

	BG <120 mg/dl	BG 121–179 mg/dl	BG ≥ 180 mg/dl
No. of patients (%)	406 (75)	103 (19)	32(6)
Mean age (yr)	7.0 ± .4	6.8 ± .6	7.8 ± 1.1
Gender (M/F)	50 / 50	57 / 43	50 / 50
Race			
Caucasian	4	8	9
Black	80	80	84
Hispanic	15	10	6
Other	1	2	0
Weight on admission (kg)	29 ± 2	26 ± 3	32 ± 6
Height on admission (cm)	79 ± 4	94 ± 9	74 ± 19
Body mass index (kg/m ²)	17 ± 5	18 ± 4	37 ± 16 ^a
Mean admission BG	92 ± 1	143 ± 2 ^b	260 ± 18 ^b
Mean inpatient BG	96 ± 3	109 ± 5	159 ± 13 ^b
Mean length hospital stay	3.8 ± 0.2	5.4 ± 1.0	5.7 ± 1.8
Mean length ICU stay	0.6 ± 0.1	1.1 ± .4 ^a	3.6 ± 1.9 ^b
Admission service (%)			
Pediatrics	79.6	58.8	72.4
Surgery	20.4	41.2	27.6

Results are ± SEM

* Diabetes Ketoacidosis (DKA) patients (n = 16) were excluded from analyses.

^aP < 0.05 vs. normoglycemia.

^bP < 0.001 vs. normoglycemia.