Published in final edited form as:

Curr Protoc Nucleic Acid Chem. 2012 June ; 0 1: Unit1.26. doi:10.1002/0471142700.nc0126s49.

Two-Step, One-Pot Synthesis of Inosine, Guanosine, and 2'-Deoxyguanosine O⁶-Ethers via Intermediate O⁶-(Benzotriazol-1-yl) Derivatives

Hari Prasad Kokatla¹ and Mahesh K. Lakshman¹

¹The City College and The City University of New York, New York

Abstract

A simple method for the etherification at the O⁶-position of silyl-protected inosine, guanosine, and 2'-deoxyguanosine is described. Typically, a THF solution of the silylated nucleoside is treated with 1*H*-benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and Cs₂CO₃, under a nitrogen atmosphere. Conversion to the O⁶-(benzotriazol-1-yl) ethers occurs within about 10 minutes for inosine, and within about 60 minutes for guanosine and 2'-deoxyguanosine. Then, for reaction with alcohols, the reaction mixture is evaporated and the O⁶-(benzotriazol-1-yl) ether is treated with Cs₂CO₃ and an appropriate alcohol, at room temperature. On the other hand, for reaction with phenols, Cs₂CO₃ and the appropriate phenol are added to the reaction mixture without evaporation, and the reaction is carried out at 70°C. Subsequently, workup, isolation, and purification lead to the requisite O⁶-alkyl or -aryl ethers in good to excellent yields.

Keywords

inosine; guanosine; 2'-deoxyguanosine; BOP; reactive nucleosides; ethers; benzotriazolyl

INTRODUCTION

The ubiquitous presence of nucleosides in living systems makes them excellent scaffolds for chemical modification, and both the heterocycle as well as the sugar moiety are targets for this purpose. The ensuing modified nucleoside analogues are highly important for a variety of reasons. To name a few, these include treatment of diseases, nucleic acid labeling, study of DNA damage, as probes for base-pair recognition and of other biological processes.

In this unit a two-step, one-pot operational procedure is described for the modification of the purine bases of nucleosides. Specifically, an efficient protocol for etherification of the C-6 amide carbonyl in silyl-protected inosine, guanosine, and 2'-deoxyguanosine is the focus. The silylated nucleoside precursors are readily accessible from commercially available inosine, guanosine, and 2'-deoxyguanosine. Reactions of these with 1*H*-benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and Cs₂CO₃ result in efficient conversion of each to a reactive intermediate, the corresponding O⁶-(benzotriazol-1-yl) derivative. These O⁶-(benzotriazol-1-yl) derivatives then undergo reaction with an alcohol or a phenol, also in the presence of Cs₂CO₃, to produce the purine nucleoside O⁶-ether derivatives. In the broader context, this approach provides a relatively simple procedure for the synthesis of various ether derivatives from purine nucleosides, and Basic Protocol 1 describes the synthesis of alkyl and aryl ethers from three purine nucleosides.

This protocol is related to one previously published (Bae et al., 2009), wherein we explained the synthesis of O^6 -(benzotriazol-1-yl)inosine and 2'-deoxyinosine derivatives as *discrete isolated reactive entities*. In a subsequent step, these isolated O^6 -(benzotriazol-1-yl) nucleoside derivatives were converted to C-6 modified purine nucleosides via reactions with amine, thiol, and phenol nucleophiles. The original work on hypoxanthine nucleosides (Bae and Lakshman, 2007) was then extended to guanine nucleosides (Lakshman and Frank, 2009). Related to these reports, one-pot conversions of inosine and 2'-deoxyinosine to adenosine derivatives has been accomplishable via the use of BOP (Wan et al., 2005). What has been less easy to achieve is a one-pot etherification of purine nucleosides, which eliminates isolation of the O^6 -(benzotriazol-1-yl) intermediates. This is the subject of the present unit.

CAUTION: All operations involving organic solvents and reagents should be conducted in a well-ventilated hood. Use of protective equipment such as gloves, safety glasses and laboratory coats is strongly recommended.

BASIC PROTOCOL 1 SYNTHESIS OF INOSINE, GUANOSINE, AND 2'-DEOXYGUANOSINE O^6 -ETHERS VIA INTERMEDIATE O^6 -(BENZOTRIAZOL-1-YL) DERIVATIVES

Synthesis of silyl-protected inosine, guanosine, and 2'-deoxyguanosine

As required, inosine **S.1**, guanosine **S.5**, or 2'-deoxyguanosine **S.9** is dried by addition and evaporation of anhydrous pyridine. Dry *N,N*-dimethylformamide (DMF) is added, followed by addition of imidazole and *tert*-butyldimethylsilyl chloride (TBDMS-Cl). After allowing the reactions to proceed for an appropriate length of time, the silylated product **S.2**, **S.6**, or **S.10** is isolated by precipitation into water, the solids are redissolved in dichloromethane and dried, and the solvent is evaporated. The crude material is purified by column chromatography on silica gel, using an appropriate eluting solvent.

Synthesis of the O^6 -alkyl and -aryl ethers of inosine, guanosine and 2'-deoxyguanosine derivatives from the silylated precursors **S.2**, **S.6**, and **S.10**

The substrates and nucleophiles used in this protocol, as well as the yields of the O^6 -ethers that were synthesized are shown in Table X.XX.1. (1) Synthesis of O^6 -alkyl ethers (**S.4a–e**, **S.8a**, and **S.12a**): Reaction of silyl-protected nucleosides **S.2** or **S.6** or **S.10** and BOP is conducted at room temperature, in anhydrous THF, with Cs_2CO_3 as base. Typically, formation of the intermediate O^6 -(benzotriazol-1-yl)inosine is complete within 10 min, whereas formation of O^6 -(benzotriazol-1-yl)guanosine and O^6 -(benzotriazol-1-yl)-2'-deoxyguanosine is complete within 1 h. The reaction mixture is evaporated to dryness on a rotary evaporator. Cs_2CO_3 and the appropriate alcohol are added and the reaction mixture is stirred at room temperature for an appropriate length of time (**S.4a** = 10 min, **S.4b** = 20 min, **S.4c** = 10 min, **S.4d** = 7 hr, **S.4e** = 2 hr, **S.8a** = 2 hr, and **S.12a** = 3.5 hr). (2) Synthesis of O^6 -aryl ethers (**S.4f**, **S.8b**, **S.12b**): Reaction of silyl-protected nucleosides **S.2** or **S.6** or **S.10** and BOP was conducted at room temperature, in anhydrous THF, with Cs_2CO_3 as base. Typically, formation of the intermediate O^6 -(benzotriazol-1-yl)inosine is complete within 10 minutes at room temperature, whereas formation of O^6 -(benzotriazol-1-yl)guanosine and O^6 -(benzotriazol-1-yl)-2'-deoxyguanosine is complete within 1 h. The reaction mixture is *not* evaporated, but Cs_2CO_3 and the appropriate phenol are added to the reaction mixture. The mixture is stirred at 70°C for an appropriate length of time (**S.4f** = 4 hr, **S.8b** = 1.5 hr, **S.12b** = 1.5 hr).

For either (1) or (2) above, workup of the reaction mixture, to isolate the crude material, followed by chromatographic purification on silica gel then affords the desired products.

Materials

Inosine (**S.1**), 99% (Acros)
Guanosine (**S.5**), 98% (Lancaster)
2'-Deoxyguanosine (**S.9**), 98% (Transgenomic)
Imidazole, 99% pure (Sigma)
tert-Butyldimethylsilyl chloride (TBDMS-Cl), 98% pure (Acros)
1*H*-Benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP), 98% pure (Chem-Impex)
Cesium carbonate (Cs₂CO₃), 99% pure (Aldrich)
Anhydrous pyridine, distilled from KOH (stored over KOH)
N,N-Dimethylformamide (DMF), anhydrous (Aldrich)
Tetrahydrofuran (THF): distilled over lithium aluminum hydride (LiAlH₄) and then distilled over sodium just prior to use
Ethyl acetate (EtOAc), HPLC grade (Fisher Scientific)
Hexanes, ACS grade (Fisher Scientific)
Dichloromethane (CH₂Cl₂), ACS grade (Fisher Scientific)
Sodium sulfate (Na₂SO₄), anhydrous, 99% pure (Spectrum)
200–300-mesh silica gel (Natland)
Nitrogen gas
Alcohols for reaction with **S.2**, **S.6**, and **S.10**:
 Methanol (for **S.4a**, **S.8a**, and **S.12a**)
 Allyl alcohol (for **S.4b**)
 Propargyl alcohol (for **S.4c**)
 Isopropyl alcohol (for **S.4d**)
 Ethylene glycol (for **S.4e**)

p-Nitrophenol (for **S.4f**)

p-Methoxyphenol (for **S.8b**)
 Phenol (for **S.12b**)
50- and 100-mL round-bottom flasks
Rotary evaporator equipped with a water aspirator
Magnetic stirrer and stir bars
Büchner funnel
Water aspirator
Glass funnel, plugged with cotton

Oil pump for vacuum drying

TLC plates: 200- μ m aluminum foil-backed silica gel plates with fluorescent indicator (for TLC analysis; Analtech)

4-mL clear glass vials with Teflon/rubber-lined, closed-top, screw caps (Wheaton)

Fraction collector

70°C temperature-controlled sand bath

60- and 125-mL separatory funnels

Dual-wavelength UV lamp (254 and 365 nm; for TLC analysis)

Additional reagents and equipment for performing thin-layer chromatography (TLC; *APPENDIX 3D*) and silica gel column chromatography (*APPENDIX 3E*)

NOTE: Except where indicated above, all reagents were obtained from commercial sources and used without further purification.

Silylate hydroxyl groups

Inosine

1. Using a round-bottom flask dry inosine (**S.1**) two times by addition and evaporation of anhydrous pyridine (add an appropriate volume of pyridine to completely wet the nucleoside), using a rotary evaporator. Dissolve the residue in anhydrous DMF, and then add imidazole and TBDMS-Cl with magnetic stirring. Stir the reaction mixture at room temperature as indicated below.

Reagent quantities for **S.2**: 1.0 g (3.72 mmol) inosine **S.1**, 2.03 g (29.76 mmol, 8 eq), imidazole, 2.24 g (14.8 mmol, 4 eq), TBDMS-Cl, and 5 mL DMF. Reaction time: 16 hr.

Volumes for some reagents (e.g., pyridine in this step, water and hexanes in the next step) are not provided for methods such as co-evaporation, precipitation, and washing because they are considered standard operating procedure.

2. Add water to the mixture and stir for 10 min. Filter the white precipitate in a Büchner funnel, using a water aspirator, and then wash the precipitate with water followed by hexanes.
3. Dissolve the resulting white precipitate in CH_2Cl_2 . Dry over anhydrous Na_2SO_4 and filter out the drying reagent using a glass funnel plugged with cotton. Evaporate the filtrate to dryness, using a rotary evaporator, and then dry the product overnight under oil pump vacuum.

Compound **S.2** may need purification to remove partially protected products. Purify by silica gel column chromatography, using 3% (v/v) methanol in CH_2Cl_2 solvent system.

4. Characterize the product by TLC and ^1H NMR.

2',3',5'-Tri-O-(tert-butyldimethylsilyl)inosine (**S.2**). Yield of white powder 2.1 g (94%). R_f : 0.41 (9:1 v/v $\text{CH}_2\text{Cl}_2/\text{MeOH}$). ^1H NMR (500 MHz, CDCl_3): δ 12.74 (br s, 1H, NH), 8.23 (s, 1H, purinyl-H), 8.08 (s, 1H, purinyl-H), 6.01 (d, 1H, H-1', $J = 4.9$ Hz), 4.51 (t, 1H, H-2', $J = 4.4$ Hz), 4.30 (t, 1H, H-3', $J = 3.9$ Hz), 4.13 (m, 1H, H-4' Hz), 3.99 (dd, 1H, H-5', $J = 3.9, 11.2$ Hz), 3.80 (dd, 1H, H-5', $J = 2.6, 11.2$ Hz), 0.96, 0.93,

and 0.81 (3s, 27H, tert-Bu), 0.15, 0.14, 0.10, 0.09, -0.02, and -0.18 (6s, 18H, SiCH₃).

Guanosine

1. Using a round-bottom flask dry guanosine (**S.5**) two times by addition and evaporation of anhydrous pyridine (add an appropriate volume of pyridine to completely wet the nucleoside), using a rotary evaporator. Dissolve the residue in anhydrous DMF, and then add imidazole and TBDMS-Cl with magnetic stirring. Stir the reaction mixture at room temperature as indicated below.

Reagent quantities for **S.6**: 1.0 g (3.53 mmol) guanosine **S.5**, 1.92 g (28.2 mmol, 8 eq), imidazole, 2.12 g (14.1 mmol, 4 eq), TBDMS-Cl, and 5 mL DMF. Reaction time: 16 hr.

Volumes for some reagents (e.g., pyridine in this step, water and hexanes in the next step) are not provided for methods such as co-evaporation, precipitation, and washing because they are considered standard operating procedure.

2. Add water to the mixture and stir for 10 min. Filter the white precipitate in a Büchner funnel, using a water aspirator, and then wash the precipitate with water followed by hexanes.
3. Dissolve the resulting white precipitate in CH₂Cl₂. Dry over anhydrous Na₂SO₄ and filter out the drying reagent using a glass funnel plugged with cotton. Evaporate the filtrate to dryness, using a rotary evaporator, and then dry the product overnight under oil pump vacuum.

Compound **S.6** may need purification to remove partially protected products. Purify by silica gel column chromatography, using 3% (v/v) EtOAc in hexanes solvent system.

4. Characterize the product by TLC and ¹H NMR.

2',3',5'-Tri-O-(tert-butyldimethylsilyl)guanosine (**S.6**). Yield of white powder 2.15 g (97%). R_f: 0.50 (9:1 v/v CH₂Cl₂/MeOH). ¹H NMR (500 MHz, CDCl₃): δ 12.01 (br s, 1H, NH), 7.90 (s, 1H, purinyl-H), 6.30 (br s, 2H, NH₂), 5.81 (d, 1H, H-1', J = 2.9 Hz), 4.42 (m, 1H, H-2' Hz), 4.27 (t, 1H, H-3', J = 3.9 Hz), 4.09 (m, 1H, H-4'), 3.98 (dd, 1H, H-5', J = 2.4, 11.2 Hz), 3.79 (dd, 1H, H-5', J = 2.4, 11.2 Hz), 0.95, 0.91, and 0.86 (3s, 27H, tert-Bu), 0.13, 0.12, 0.09, 0.08, 0.01, and -0.04 (3s, 18H, SiCH₃).

2'-Deoxyguanosine

- 1 Using a round-bottom flask dry 2'-deoxyguanosine (**S.9**) two times by addition and evaporation of anhydrous pyridine (add an appropriate volume of pyridine to completely wet the nucleoside), using a rotary evaporator. Dissolve the residue in anhydrous DMF, and then add imidazole and TBDMS-Cl with magnetic stirring. Stir the reaction mixture at room temperature as indicated below.

Reagent quantities for **S.10**: 1.0 g (3.7 mmol) 2'-deoxyguanosine **S.9**, 1.5 g (22.2 mmol, 6 eq), imidazole, 1.6 g (11.1 mmol, 3 eq), TBDMS-Cl, and 5 mL DMF. Reaction time: 16 hr.

Volumes for some reagents (e.g., pyridine in this step, water and hexanes in the next step) are not provided for methods such as co-

evaporation, precipitation, and washing because they are considered standard operating procedure.

- 2 Add water to the mixture and stir for 10 min. Filter the white precipitate in a Büchner funnel, using a water aspirator, and then wash the precipitate with water followed by hexanes.
- 3 Dissolve the resulting white precipitate in CH_2Cl_2 . Dry over anhydrous Na_2SO_4 and filter out the drying reagent using a glass funnel plugged with cotton. Evaporate the filtrate to dryness, using a rotary evaporator, and then dry the product overnight under oil pump vacuum.

Compound **S.10** may need purification to remove partially protected products. Purify by silica gel column chromatography, using 10% (v/v) MeOH in CH_2Cl_2 solvent system.

- 4 Characterize the product by TLC and ^1H NMR.

2',3'-Di-O-(tert-butyldimethylsilyl)-2'-deoxyguanosine (**S.10**). Yield of white powder 1.75 g (95%). R_f : 0.42 (9:1 v/v $\text{CH}_2\text{Cl}_2/\text{MeOH}$). ^1H NMR (500 MHz, CDCl_3): δ 10.61 (br s, 1H, NH), 7.87 (s, 1H, purinyl-H), 6.47 (br s, 2H, NH_2), 6.10 (t, 1H, H-1', $J = 6.8$ Hz), 4.48 (m, 1H, H-3'), 3.80 (m, 1H, H-4'), 3.69 (dd, 1H, H-5', $J = 6.3, 11.2$ Hz), 3.64 (dd, 1H, H-5', $J = 4.3, 11.2$ Hz), 2.64 (app quint, 1H, H-2', $J_{\text{app}} = 6.0$ Hz), 2.23 (ddd, 1H, H-2', $J = 3.1, 6.0, 13.5$ Hz), 0.88, and 0.86 (2s, 18H, tert-Bu), 0.10, 0.04, 0.03 (3s, 12H, SiCH_3).

Step one of the two-step etherification process: activate **S.2** or **S.6** or **S.10** with BOP and Cs_2CO_3

- 5 To a solution of **S.2** or **S.6** or **S.10** and BOP in anhydrous THF, add Cs_2CO_3 under a nitrogen atmosphere (balloon filled with nitrogen gas). Stir the reaction mixture at room temperature for the period of time indicated below.

Reagent quantities for **S.4a-f**: 100 mg (0.163 mmol) silyl-protected inosine **S.2**, 144 mg (0.326 mmol) BOP, 106 mg (0.326 mmol) Cs_2CO_3 , and 1.5 mL THF. Reaction time: 10 min.

Reagent quantities for **S.8a,b**: 50 mg (0.079 mmol) silyl-protected guanosine **S.5**, 71 mg (0.159 mmol) BOP, 52 mg (0.159 mmol) Cs_2CO_3 , and 1.5 mL THF. Reaction time: 1 hr.

Reagent quantities for **S.12a,b**: 50 mg (0.100 mmol) silyl-protected 2'-deoxyguanosine **S.10**, 88.5 mg (0.200 mmol) BOP, 65 mg (0.200 mmol) Cs_2CO_3 , and 1 mL THF. Reaction time: 1 hr.

- 6 Evaporate the reaction mixture on a rotary evaporator. Skip this step for synthesis of compounds **S.4f**, **S.8b**, and **S.12b**, and go directly to step 8.

Step two of the two-step etherification process: reaction with alcohol or phenol, and Cs_2CO_3

- 7 For **S.4a-e**: add 106 mg (0.326 mmol) Cs_2CO_3 , the appropriate alcohol (quantities given below), and stir the reaction mixture at room temperature (reaction times given below).

S.4a: 132 μL (3.26 mmol) methanol, 10 min

S.4b: 222 μL (3.26 mmol) allyl alcohol, 20 min

S.4c: 188 μL (3.26 mmol) propargyl alcohol, 10 min

S.4d: 251 μL (3.26 mmol) isopropyl alcohol, 7 hr

S.4e: 182 μL (3.26 mmol) ethylene glycol, 2 hr

For **S.8a:** add 52 mg (0.159 mmol) Cs_2CO_3 , 72 μL (1.59 mmol) methanol, and stir the reaction mixture at room temperature for 2 hr

For **S.12a:** add 65 mg (0.200 mmol) Cs_2CO_3 , 81 μL (2 mmol) methanol, and stir the reaction mixture at room temperature for 3.5 hr

- 8** For **S.4f**, **S.8b**, and **S.12b:** add Cs_2CO_3 the appropriate phenol (quantities given below), and stir the reaction mixture at 70°C (reaction times given below)

S.4f: 106 mg (0.326 mmol) Cs_2CO_3 , 45 mg (0.326) *p*-nitrophenol, 4 hr

S.8b: 52 mg (0.159 mmol) Cs_2CO_3 , 20 mg (0.159 mmol) *p*-methoxyphenol, 1.5 hr

S.12b: 65 mg (0.200 mmol) Cs_2CO_3 , 19 mg (0.200 mmol) phenol, 1.5 hr

Workup of the reaction mixture, isolation of the crude products, and their purification

- 9** Transfer the reaction mixture to a 60-mL separatory funnel. Dilute the reaction mixture with water (10 mL) and extract with EtOAc (3×10 mL). Separate and combine the organic layers, dry over Na_2SO_4 , and evaporate to dryness on a rotary evaporator.

- 10** Purify the crude product by silica gel column chromatography, using the appropriate solvent system given below:

For **S.4a:** 20% (v/v) EtOAc in hexanes

For **S.4b:** 20% (v/v) EtOAc in hexanes

For **S.4c:** 20% (v/v) EtOAc in hexanes

For **S.4d:** 15% (v/v) EtOAc in hexanes

For **S.4e:** 70% (v/v) EtOAc in hexanes (15% of a dimer arising from reaction with both hydroxyl groups of ethylene glycol was also isolated as an early eluting material)

For **S.4f:** 20% (v/v) EtOAc in hexanes

For **S.5a:** 20% (v/v) EtOAc in hexanes

For **S.5b:** 20% (v/v) EtOAc in hexanes

For **S.12a:** 20% (v/v) EtOAc in hexanes

For **S.12b:** 20% (v/v) EtOAc in hexanes

- 11** Analyze fractions obtained from the column by TLC, combine fractions containing product, and remove the solvent on a rotary evaporator. Finally dry the product under vacuum.

- 12** Characterize the products by TLC, ^1H NMR, ^{13}C NMR and HRMS.

2',3',5'-Tri-O-(tert-butyldimethylsilyl)-O⁶-methylinosine (**S.4a**).

Yield of a yellowish solid 96 mg (94%). R_f (SiO_2 /20% v/v EtOAc in hexanes) = 0.4. ^1H NMR: δ 8.52 (s, 1H, purinyl-H), 8.32 (s, 1H,

purinyl-H), 6.08 (d, 1H, H-1', J = 4.8 Hz), 4.62 (t, 1H, H-2', J = 4.6 Hz), 4.31 (t, 1H, H-3', J = 4.1 Hz), 4.18 (s, 3H, OCH₃), 4.13 (q, 1H, H-4', J = 3.1 Hz), 4.03 (dd, 1H, H-5', J = 3.9, 11.2 Hz), 3.79 (dd, 1H, H-5', J = 2.6, 11.2 Hz), 0.95, 0.93, and 0.75 (3s, 27H, tert-Bu), 0.14, 0.13, 0.10, 0.09, -0.04, and -0.22 (6s, 18H, SiCH₃). ¹³C NMR: δ 161.2, 152.2, 151.9, 141.3, 122.2, 88.6, 85.6, 76.3, 72.1, 62.7, 54.3, 26.3, 26.0, 25.8, 18.7, 18.3, 18.0, -4.1, -4.4, -4.8, -5.1. HRMS (ESI) calcd for C₂₉H₅₇N₄O₅Si₃ [M + H]⁺ 625.3631, found 625.3636.

O⁶-(Allyl)-2',3',5'-tri-O-(tert-butyldimethylsilyl)inosine (**S.4b**).

Yield of a white solid 96 mg (90%). R_f (SiO₂/30% v/v EtOAc in hexanes) = 0.7. ¹H NMR: δ 8.50 (s, 1H, purinyl-H), 8.30 (s, 1H, purinyl-H), 6.17 (m, 1H, -CH=), 6.08 (d, 1H, H-1', J = 4.8 Hz), 5.45 (d, 1H, =CH_{trans}, J = 17.0 Hz), 5.29 (d, 1H, =CH_{cis}, J = 10.2 Hz), 5.15 (br d, 2H, OCH₂, J = 5.8 Hz), 4.63 (t, 1H, H-2', J = 8.3 Hz), 4.31 (t, 1H, H-3', J = 3.4 Hz), 4.13 (br d, 1H, H-4', J = 2.4 Hz), 4.02 (dd, 1H, H-5', J = 3.4, 11.2 Hz), 3.79 (d, 1H, H-5', J = 11.2 Hz), 0.95, 0.93, and 0.78 (3s, 27H, tert-Bu), 0.14, 0.13, 0.10, 0.09, -0.04, and -0.23 (6s, 18H, SiCH₃). ¹³C NMR: δ 160.5, 152.1, 141.3, 132.6, 122.1, 118.7, 88.5, 85.7, 76.3, 72.1, 67.7, 62.7, 26.3, 26.0, 25.8, 18.7, 18.3, 18.0, -4.1, -4.4, -4.8, -5.1. HRMS (ESI) calcd for C₃₁H₅₉N₄O₅Si₃ [M + H]⁺ 651.3788, found 651.3787.

2',3',5'-Tri-O-(tert-butyldimethylsilyl)-O⁶-propargylinosine (**S.4c**).

Yield of a white solid 91 mg (86%). R_f (SiO₂/30% v/v EtOAc in hexanes) = 0.7. ¹H NMR: δ 8.54 (s, 1H, purinyl-H), 8.36 (s, 1H, purinyl-H), 6.08 (d, 1H, H-1', J = 4.8 Hz), 5.23 (dd, 2H, OCH₂, J = 1.4, 2.4 Hz), 4.59 (t, 1H, H-2', J = 4.4 Hz), 4.32 (t, 1H, H-3', J = 4.1 Hz), 4.13 (q, 1H, H-4', J = 3.4 Hz), 4.00 (dd, 1H, H-5', J = 3.6, 11.4 Hz), 3.79 (dd, 1H, H-5', J = 2.4, 11.7 Hz), 2.49 (t, 1H, ≡C-H, J = 2.4 Hz), 0.95, 0.92, and 0.79 (3s, 27H, tert-Bu), 0.14, 0.13, 0.10, 0.09, -0.04, and -0.21 (6s, 18H, SiCH₃). ¹³C NMR: δ 159.5, 152.3, 151.9, 141.7, 122.1, 88.7, 85.6, 78.3, 76.4, 75.3, 71.9, 62.6, 54.3, 26.3, 26.0, 25.8, 18.7, 18.2, 18.0, -4.1, -4.5, -4.7, -5.1. HRMS (ESI) calcd for C₃₁H₅₇N₄O₅Si₃ [M + H]⁺ 649.3631, found 649.3634.

2',3',5'-Tri-O-(tert-butyldimethylsilyl)-O⁶-isopropylinosine (**S.4d**).

Yield of clear gummy material 80 mg (75%). R_f (SiO₂/30% v/v EtOAc in hexanes) = 0.7. ¹H NMR: δ 8.48 (s, 1H, purinyl-H), 8.26 (s, 1H, purinyl-H), 6.08 (d, 1H, H-1', J = 4.8 Hz), 5.66 (septet, 1H, OCH, J = 6.3 Hz), 4.63 (t, 1H, H-2', J = 4.6 Hz), 4.31 (t, 1H, H-3', J = 3.6 Hz), 4.12 (br d, 1H, H-4', J = 2.4 Hz), 4.01 (dd, 1H, H-5', J = 3.6, 11.4 Hz), 3.79 (dd, 1H, H-5', J = 1.9, 11.2 Hz), 1.47 (d, 6H, (CH₃)₂, J = 6.3 Hz), 0.94, 0.92, and 0.78 (3s, 27H, tert-Bu), 0.13, 0.12, 0.10, 0.09, -0.04, and -0.22 (6s, 18H, SiCH₃). ¹³C NMR: δ 160.7, 152.2, 152.1, 141.0, 122.3, 88.5, 85.6, 76.2, 72.1, 70.4, 62.7, 26.3, 26.0, 25.8, 22.1, 18.7, 18.3, 18.0, -4.1, -4.4, -4.8, -5.1. This compound has been previously reported (Bae and Lakshman, 2007).

2',3',5'-Tri-O-(tert-butyldimethylsilyl)-O⁶-(2-hydroxyethyl)inosine (**S.4e**).

Yield of a white solid 68 mg (63%). R_f (SiO₂/EtOAc) = 0.3. ¹H NMR: δ 8.49 (s, 1H, purinyl-H), 8.35 (s, 1H, purinyl-H), 6.87 (d, 1H, H-1', J = 4.4 Hz), 4.72 (m, 2H, OCH₂), 4.61 (t, 1H, H-2', J = 4.4 Hz), 4.31 (m, 1H, H-3'), 4.13 (br s, 1H, H-4'), 4.02 (m, 3H, H-5' and OCH₂), 3.79 (d, 1H, H-5', J = 11.2 Hz), 3.52 (s, 1H, OH), 0.95, 0.92, and 0.79 (3s, 27H, tert-Bu), 0.14, 0.13, 0.09, 0.08, -0.04, and -0.22 (6s, 18H, SiCH₃). ¹³C NMR: δ 160.8, 152.2, 152.0, 141.6, 122.0, 88.6, 85.7, 76.4, 72.0, 69.8, 62.6, 61.9, 26.3, 26.0, 25.8, 18.7, 18.2, 18.0, -4.1, -4.5, -4.8, -5.1. HRMS (ESI) calcd for C₃₀H₅₉N₄O₆Si₃ [M + H]⁺ 655.3737, found 655.3739.

The reaction with ethylene glycol also gave a product arising from reaction of both hydroxyl groups (data below).

Ethylene glycol bis-O⁶-[9-(2',3',5'-tri-O-tert-butyltrimethylsilyl)- β -D-ribofuranosyl]]purinyl ether.

Minor product obtained as a white solid (32 mg, 15%) in the reaction of 3 with ethylene glycol. R_f (SiO₂/20% v/v EtOAc in hexanes) = 0.4. ¹H NMR: δ 8.49 (s, 2H, purinyl-H), 8.29 (s, 2H, purinyl-H), 6.09 (d, 2H, H-1', J = 4.8 Hz), 5.06 (m, 4H, OCH₂), 4.61 (t, 2H, H-2', J = 4.4 Hz), 4.31 (t, 2H, H-3', J = 2.9 Hz), 4.13 (br d, 2H, H-4', J = 2.4 Hz), 4.02 (dd, 2H, H-5', J = 3.4, 11.2), 3.79 (d, 2H, H-5', J = 11.7 Hz), 0.94, 0.92, and 0.79 (3s, 54H, tert-Bu), 0.13, 0.12, 0.10, 0.09, -0.04, and -0.22 (6s, 36H, SiCH₃). ¹³C NMR: δ 160.5, 152.2, 152.0, 141.3, 122.0, 88.6, 85.7, 76.3, 72.1, 64.9, 62.7, 26.3, 26.0, 25.9, 18.7, 18.3, 18.0, -4.1, -4.5, -4.7, -5.1. HRMS (ESI) calcd for C₅₈H₁₁₁N₈O₁₀Si₆ [M + H]⁺ 1247.7033, found 1247.7032.

2',3',5'-Tri-O-(tert-butyltrimethylsilyl)-O⁶-(4-nitrophenyl)inosine (**S. 4f**).

Yield of a white solid 102 mg (85%). R_f (SiO₂/20% v/v EtOAc in hexanes) = 0.5. ¹H NMR: δ 8.52 (s, 1H, purinyl-H), 8.50 (s, 1H, purinyl-H), 8.33 (d, 2H, Ar-H, J = 8.7 Hz), 7.47 (d, 2H, Ar-H, J = 8.7 Hz), 6.14 (d, 1H, H-1', J = 4.7 Hz), 4.60 (t, 1H, H-2', J = 4.4 Hz), 4.33 (t, 1H, H-3', J = 3.9 Hz), 4.17 (m, 1H, H-4'), 4.04 (dd, 1H, H-5', J = 3.3, 11.4), 3.82 (d, 1H, H-5', J = 11.3 Hz), 0.97, 0.93, and 0.81 (3s, 27H, tert-Bu), 0.16, 0.15, 0.11, 0.10, -0.01, and -0.18 (6s, 18H, SiCH₃). ¹³C NMR: δ 159.0, 157.5, 153.5, 151.9, 145.3, 143.0, 125.6, 122.6, 122.3, 88.8, 85.7, 76.6, 71.9, 62.5, 26.3, 26.0, 25.8, 18.7, 18.2, 18.0, -4.1, -4.4, -4.7, -5.1. This compound has been previously reported (Bae and Lakshman, 2007).

2',3',5'-Tri-O-(tert-butyltrimethylsilyl)-O⁶-methylguanosine (**S.8a**).

Yield of an orange solid 37 mg (73%). R_f (SiO₂/20% v/v EtOAc in hexanes) = 0.3. ¹H NMR: δ 7.97 (s, 1H, purinyl-H), 5.91 (d, 1H, H-1', J = 5.3 Hz), 5.12 (s, 2H, NH₂), 4.46 (t, 1H, H-2', J = 4.9 Hz), 4.26 (t, 1H, H-3', J = 4.3 Hz), 4.08 (m, 1H, H-4'), 4.04 (s, 3H, OCH₃), 3.96 (dd, 1H, H-5', J = 3.9, 11.2 Hz), 3.76 (dd, 1H, H-5', J = 2.4, 11.2 Hz), 0.94, 0.92, and 0.79 (s, 27H, tert-Bu), 0.13, 0.12, 0.11, 0.09, -0.04, and -0.18 (6s, 18H, SiCH₃). ¹³C NMR: δ 161.6, 159.5, 153.8, 137.9, 116.0, 87.7, 85.4, 76.4, 72.2, 62.8, 53.9, 26.2, 26.0, 25.9, 18.7, 18.2, 18.1, -4.1, -4.5, -4.8, -5.2. This compound has been previously reported (Lakshman and Frank, 2009).

2',3',5'-Tri-O-(tert-butyldimethylsilyl)-O⁶-(4-methoxyphenyl)guanosine (**S.8b**).

Yield of a white solid 44 mg (75%). *R_f* (SiO₂/20% v/v EtOAc in hexanes) = 0.4. ¹H NMR: δ 8.08 (s, 1H, purinyl-H), 7.16 (d, 2H, Ar-H, *J* = 8.8 Hz), 6.91 (d, 2H, Ar-H, *J* = 8.8 Hz), 5.92 (d, 1H, H-1', *J* = 4.5 Hz), 4.79 (s, 2H, NH₂), 4.51 (t, 1H, H-2', *J* = 4.4 Hz), 4.31 (t, 1H, H-3', *J* = 4.0 Hz), 4.16 (m, 1H, H-4'), 3.99 (dd, 1H, H-5', *J* = 3.5, 11.4 Hz), 3.82 (s, 3H, OCH₃), 3.79 (dd, 1H, H-5', *J* = 1.9, 11.4 Hz), 0.95, 0.92, and 0.83 (3s, 27H, tert-Bu), 0.14, 0.13, 0.10, 0.09, 0.00, and -0.10 (6s, 18H, SiCH₃). ¹³C NMR: δ 160.9, 159.3, 157.0, 154.7, 146.2, 138.9, 122.9, 116.1, 114.4, 88.2, 85.1, 76.3, 71.9, 62.6, 55.7, 26.3, 26.0, 25.9, 18.7, 18.7, 18.2, 18.1, -4.1, -4.4, -4.6, -5.1. HRMS (ESI) calcd for C₃₅H₆₂N₅O₆Si₃ [M + H]⁺ 732.4002, found 732.4007.

3',5'-Di-O-(tert-butyldimethylsilyl)-O⁶-methyl-2'-deoxyguanosine (**S.12a**).

Yield of a yellowish solid 38 mg (75%). *R_f* (SiO₂/40% v/v EtOAc in hexanes) = 0.3. ¹H NMR: δ 7.70 (s, 1H, purinyl-H), 6.31 (t, 1H, H-1', *J* = 6.3 Hz), 4.89 (s, 2H, NH₂), 4.57 (m, 1H, H-3'), 4.05 (s, 3H, OCH₃), 3.89 (m, 1H, H-4'), 3.80 (dd, 1H, H-5', *J* = 4.4, 11.2 Hz), 3.74 (dd, 1H, H-5', *J* = 2.9, 11.2 Hz), 2.55 (app quint, 1H, H-2', *J_{app}* ~ 6.3 Hz), 2.34 (ddd, 1H, H-2', *J* = 2.2, 6.0, 13.1 Hz), 0.90 (br s, 18H, tert-Bu), 0.08, 0.07, and 0.06 (3s, 12H, SiCH₃). ¹³C NMR: δ 161.7, 159.5, 153.6, 137.7, 116.1, 87.8, 83.7, 72.0, 63.0, 53.9, 41.1, 26.1, 25.9, 18.6, 18.1, -4.4, -4.5, -5.2, -5.3. This compound has been previously reported (Lakshman and Frank, 2009).

3',5'-Di-O-(tert-butyldimethylsilyl)-O⁶-phenyl-2'-deoxyguanosine (**S.12b**).

Yield of a yellowish solid 42 mg (73%). *R_f* (SiO₂/40% v/v EtOAc in hexanes) = 0.3. ¹H NMR: δ 8.00 (s, 1H, purinyl-H), 7.40 (t, 2H, Ar-H, *J* = 7.7 Hz), 7.25 (m, 3H, Ar-H), 6.34 (t, 1H, H-1', *J* = 6.4 Hz), 4.77 (s, 2H, NH₂), 4.60 (m, 1H, H-3'), 3.99 (m, 1H, H-4'), 3.83 (dd, 1H, H-5', *J* = 4.1, 11.1 Hz), 3.77 (dd, 1H, H-5', *J* = 2.6, 11.1 Hz), 2.59 (app quint, 1H, H-2', *J_{app}* ~ 6.4 Hz), 2.37 (ddd, 1H, H-2', *J* = 1.4, 5.8, 12.9 Hz), 0.923 and 0.920 (2s, 18H, tert-Bu), 0.10 and 0.09 (2s, 12H, SiCH₃). ¹³C NMR: δ 160.6, 159.3, 154.7, 152.7, 138.7, 129.4, 125.4, 122.1, 116.2, 87.9, 83.9, 72.1, 63.0, 41.2, 26.1, 25.9, 18.6, 18.2, -4.4, -4.5, -5.1, -5.2. This compound has been previously reported (Lakshman and Frank, 2009).

COMMENTARY

Background Information

In 2000, Lin and Robins reported that activation of the C-6 amide functionality of inosine and 2'-deoxyinosine can be accomplished by treatment of the acetyl-protected nucleosides to PPh₃/I₂/(*iso*-Pr)₂NEt (Lin and Robins, 2000). Addition of morpholine, piperidine, or imidazole to the reaction mixtures led to the formation of C-6 morpholinyl, piperidinyl, and imidazolyl derivatives, respectively (Lin and Robins, 2000). Subsequently, in 2004 a similar procedure was applied to the synthesis of C-6 imidazolyl derivatives from guanosine and 2'-deoxyguanosine (Janeba et al., 2004). A year later, Wan et al. reported a one-pot synthesis of adenosine analogues by reaction of acetyl-protected inosine with 1*H*-benzotriazol-1-

ylxyo-tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and amines, in the presence of (*iso*-Pr)₂NEt in DMF as solvent (Wan et al., 2005). They also demonstrated that sugar-unprotected inosine and 2'-deoxyinosine (1 example) could be utilized for this conversion (Wan et al., 2005). In this work, they report the formation of a C-6 phosphonium ion as a putative intermediate (such an intermediate was reported to be observed by liquid chromatography/mass spectrometry). The plausible mechanisms of the reactions described above are shown in Figure X.XX.1.

We became interested in the mechanism by which BOP reacts at the C-6 amide bond since we believed that two possible mechanisms could operate. This is shown in Figure X.XX.2. In principle, as reported by Wan et al. (Wan et al., 2005), the reaction could proceed via the phosphonium salt as a result of deprotonation at N1 by the base, followed by attack of the amide oxygen atom at the phosphorus atom of BOP (pathway **a** in Figure X.XX.2). The alternative is deprotonation at N1 by base and attack of the amide oxygen at the nitrogen atom in BOP, in an S_N2'-like fashion, with expulsion of HMPA (pathway **b** in Figure X.XX.2).

We therefore investigated the mechanism of this reaction by ³¹P{¹H} NMR and discovered that the reaction does not instantly produce HMPA, and that the nucleoside phosphonium ion does make an appearance (Bae and Lakshman, 2007). This was evidenced by a singlet at $\delta = 35$ ppm in the ³¹P{¹H} NMR in addition to the PF₆⁻ resonance at $\delta = -143$ ppm. Also, the ¹H NMR spectrum of the isolated intermediate indicated the presence of the -P(NMe)₂ with a doublet at 2.92 ppm (*J*_{H-P} = 11.2 Hz). Thus, pathway **a** is definitely operational in these types of transformations, although pathway **b** cannot be excluded.

On the basis of these results we then developed a second generation synthesis of *O*⁶-(benzotriazol-1-yl)inosine as well as the 2'-deoxy analogue (Bae and Lakshman, 2008a), and also a "catch-and-release" approach for high-throughput type applications, using polymer-supported 1-hydroxy-1*H*-benzotriazole (Bae and Lakshman, 2008b). In addition, we evaluated the use of BOP for modifying the C-2 position. For this purpose, reactions of disilyl *O*⁶-benzyl-2'-deoxyxanthosine were studied (Bae and Lakshman, 2008c). Here we discovered that a *O*²-(benzotriazol-1-yl) derivative is not formed, but that the reactions terminate at the stage of the phosphonium ion. The C-2 phosphonium derivative is an excellent reagent for synthesis of 2'-deoxyguanosine derivatives, and its applicability was exemplified via rapid access to an acrolein adduct of 2'-deoxyguanosine (Bae and Lakshman, 2008c).

We then developed a method for the functionalization at the C-6 position of guanine nucleosides, via reaction of silylated guanosine and 2'-deoxyguanosine with BOP, and isolation of the *O*⁶-(benzotriazol-1-yl) derivatives (Lakshman and Frank, 2009). In the cases of the hypoxanthine (inosine and 2'-deoxyinosine) and guanine nucleosides, the *O*⁶-(benzotriazol-1-yl) derivatives showed extremely good reactivity towards amine, alcohol, phenol, and thiol nucleophiles.

Building on these developments, we next discovered a method for the etherification of the C-6 amide linkage in inosine, guanosine, and 2'-deoxyguanosine (Kokatla and Lakshman, 2010). The present protocol describes this methodology and there are some notable differences when compared with the one-pot amination reactions (Wan et al., 2005, and Lakshman and Frank 2009). Importantly, these etherification reactions are successful as a *two-step, one-pot operation*. This is shown in Figure X.XX.3.

S_NAr, or nucleophilic aromatic substitution, is a two-step process where a nucleophile first adds to a *sp*² carbon atom of an aromatic ring that bears a good leaving group. The ensuing

Meisenheimer complex then undergoes expulsion of the leaving group with concomitant rearomatization. Such a process is quite facile with nucleosides and therefore this reaction has found wide application for modifying the heterocyclic moieties of nucleosides.

For the S_NAr reaction, several leaving groups have been used at the C-6 position of purines, and the references below provide some examples:

- a. chloro (Robins and Basom, 1973; Robins and Uzna ski, 1981a; Véliz and Beal, 2000; Francom et al., 2002; Francom and Robins, 2003)
- b. bromo (Nair and Richardson, 1980; Lagisetty et al., 2006; Véliz and Beal, 2001)
- c. iodo (Cosstick and Douglas, 1991; van der Wenden et al., 1995; Liu et al., 2004)
- d. fluoro (Robins and Basom, 1973; Robins and Uzna ski, 1981b)
- e. phenoxy (Gao et al., 1992; Ferentz and Verdine, 1992; Lakshman and Zajc, 1996; Allerson et al., 1997)
- f. aryl sulfonyl (Hayakawa et al., 1993; Allerson et al., 1997; Lakshman et al., 2005)
- g. alkyl sulfonyl (Nagatsugi et al., 1997), pyridyl (Adamiak et al., 1985; Fathi et al., 1990),
- h. sulfone (Lin and Robins, 2000)
- i. imidazolyl (Lin and Robins, 2000; Janeba et al., 2004)
- j. 1,2,4-triazol-4-yl (Samano et al., 1994; Miles et al., 1995).

Many of the above mentioned derivatives have been utilized for the synthesis of C-6 ethers of purine nucleoside. As examples, a C-6 chloride has been displaced with alkoxide (Nair et al., 1988; Aso et al., 2000), a C-6 pentafluorophenyl ether of 2'-deoxyguanosine was converted to the methyl ether (Gao et al., 1992), we (Lakshman et al., 2000) and others (Kaloudis et al., 2009) have converted C-6 arylsulfonyl guanine nucleosides to O^6 -ethers via a quaternary ammonium salt, a C-6 imidazolyl guanosine derivative was converted to the methyl ether (Janeba et al., 2004), as also a C-6 1,2,4-triazol-4-yl analogue (Samano et al., 1994; Miles et al., 1995).

In all of these cases, the methodology typically requires the synthesis and isolation of a discrete reactive nucleoside derivative, prior to the etherification step. Another approach is the generation of a reactive intermediate in situ, which then undergoes reaction with an alcohol. A classic example within the area of purine nucleoside modification is the Mitsunobu reaction. Pfeleiderer et al. developed the Mitsunobu reaction as a method to introduce the base-labile *p*-nitrophenylethyl group (Himmelsbach et al., 1984). This method was then modified and applied to the synthesis of silyl-protected, reactive nucleoside derivatives such as 2-fluoro-2'-deoxyinosine (Zajc et al., 1992), 2-bromo-2'-deoxyinosine (Harwood et al., 1999), 2-bromoinosine (Qian and Glaser, 2005), and 8-bromo-2'-deoxyguanosine (Meier and Gräsl, 2002).

The protocol described herein is a bridge between the actual isolation of the reactive nucleoside derivatives and in situ formation of a reactive species that undergoes further transformation. That is, as described earlier in our original publications (Bae and Lakshman 2007; Lakshman and Frank 2009) and a previous protocol (Bae et al., 2009), isolation of the reactive nucleoside derivatives **S.3**, its 2'-deoxy analogue, **S.7**, and **S.11**, followed by displacement with alcohols as a separate, second step is eminently possible. However, as described herein, isolation is not specifically necessary, and the etherification can be

conducted as a two-step, one-pot operation ensuring that **S.3** or **S.7** or **S.11** are preformed at the end of the first step.

Critical Parameters and Troubleshooting

The silylation reactions of inosine **S.1**, guanosine **S.5**, and 2'-deoxyguanosine **S.9** are reasonably simple and are not typically expected to produce any problems. Methods for the conversion of the silylated nucleosides to the *O*⁶-(benzotriazol-1-yl) derivatives **S.3**, **S.7**, and **S.11** are also relatively simple. In these procedures, it is important to ensure that the equipment and solvents used for the reaction are dry. Some important noteworthy points are as follows. It appears that BOP and an alcohol do not react, since exposure of BOP to methanol *in the absence of a base* led to no discernable reaction over 1 hr (as judged by ³¹P{¹H} NMR, Kokatla and Lakshman, 2010). However, in the presence of Cs₂CO₃, a reaction between BOP and methanol was observed, leading to the formation of hexamethylphosphorus triamide (HMPA) and 1-methoxy-1*H*-1,2,3-benzotriazole. It appears, from preliminary experiments, that 1,8-diazabicycloundec-7-ene (DBU) can also be used in place of Cs₂CO₃, although the reaction may be slower (Kokatla and Lakshman, 2010). It is also possible that reactions of alcohols, like those of phenols, can be conducted without evaporation of the solvent after step one (this has been demonstrated with methanol; Kokatla and Lakshman, 2010). However, evaporation of the reaction mixture after step one ensures good reactivity of alcohols and produces generally faster reactions. In unpublished experiments, attempts at preparing the acetyl-protected *O*⁶-allyl derivative of 2'-deoxyguanosine (acetyl-protected 2'-deoxy analogue of **S.4b**) were unsuccessful. This is possibly due to the reactivity of acetate esters to nucleophilic conditions.

The labile nature of the nucleosides needs to be recognized, and these compounds can be intolerant of certain conditions. Typically, glassware and other equipment that nucleosides are expected to come into contact with should be acid free. Routinely, glassware is washed with 10% aqueous ammonium hydroxide, water, and acetone, followed by drying in an oven. Use of high temperatures for the water bath during rotary evaporation procedures should be avoided.

Generally, the procedures contained in this protocol can be considered simple. Nevertheless, familiarity with standard operating procedures of a chemical synthesis laboratory is essential. These include but are not limited to ability to setup, monitor, and work reactions up, perform extraction procedures, and thin layer as well as column chromatography, familiarity with evaporation and vacuum drying protocols. The ability to use NMR and mass spectrometry instrumentation typically requires training, and the ability to interpret data obtained by spectroscopic means requires adequate knowledge. The authors suggest that the methods outlined in this protocol be adhered to as closely as possible.

Compound Characterization

For monitoring reactions, aluminum foil-backed, silica gel TLC of 200 μm thickness were used. For purifications by column chromatography 200–300-mesh silica gel was used. Please refer to the individual compound headings for eluting solvents used for chromatography, and for *R*_f values. In this protocol, the ¹H NMR spectra reported were recorded at 500 MHz and are referenced to the residual protonated solvent. ¹³C NMR spectra reported here were recorded at 125 MHz and are referenced to the ¹³C resonance of the deuterated solvent. Chemical shifts (δ) are reported in parts per million and coupling constants (*J*) are in hertz (Hz). It is expected that lower field strength instruments (400 MHz for ¹H/100 MHz for ¹³C or 300 MHz for ¹H/75 MHz for ¹³C) will provide adequate resolution for compound characterization. The purine is numbered 1–9 by the accepted convention, and the sugar carbon atoms are numbered 1'–5' starting at the anomeric carbon

atom and proceeding via the carbon chain to the primary carbinol carbon. Multiplicities in the ^1H NMR are reported using standard abbreviations (s = singlet, d = doublet, t = triplet, q = quartet, quint = quintet, m = multiplet, app = apparent), based upon the apparent signal multiplicities observed.

Anticipated Results

Yields for all products are indicated in Table X.XX.1 and in the characterization data for each compound. Following the procedures presented here, it is anticipated that comparable or better results should be obtainable for the synthesis of the various $\mathcal{O}^6$ -alkyl and -aryl ethers of inosine, guanosine, and 2'-deoxyguanosine. The method should also be applicable to silyl-protected 2'-deoxyinosine (not described here). Some of the nucleoside ethers described herein could have additional uses. As examples, the propargyl derivative can be used in Cu-catalyzed azide-alkyne ligation chemistry (Tornøe, et al., 2002 and Rostovtsev, et al., 2002), the ethylene glycol derivative with a free terminal hydroxyl group can be used for additional functionalization or to append other molecules, the *p*-nitrophenoxy, *p*-methoxyphenoxy, and phenoxy derivatives can be used in subsequent $\text{S}_{\text{N}}\text{Ar}$ reactions, and may be amenable to oligonucleotide assembly procedures. Such an approach had been demonstrated previously (Ferentz and Verdine, 1992; Allerson et al., 1997). *p*-Nitrophenoxy derivatives of nucleosides are attractive for processes wherein the released *p*-nitrophenoxide can be assessed by UV-Vis spectrometry (Zemlicka and Endo, 1996).

Time Considerations

Step 1, the conversion of silyl-protected inosine (**S.2**) to the $\mathcal{O}^6$ -(benzotriazol-1-yl) ether **S.3**, takes literally 10–15 minutes to reach completion with silyl-protected inosine. This step takes about 1 hr with silyl-protected guanosine (**S.6** $\rightarrow$ **S.7**) and 2'-deoxyguanosine (**S.10** $\rightarrow$ **S.11**). These reactions can be monitored for completion by TLC, using a UV lamp. Evaporation of the reaction mixtures should take a few minutes, depending upon the volume of the relatively volatile THF (65–67°C). Reactions with alcohols take anywhere from 10 min to 7 hr, at room temperature, for the alcohols reported in this protocol. Reaction times with other alcohols may vary, but completion of reaction can be again monitored by TLC, using a UV lamp. Reactions with phenols reported herein take 1.5–4 hr at 70°C. It should be noted that the reported reactions include an electron-deficient as well as and electron-rich phenol, and thus the reaction times may represent general times for reactions with other phenols. Nevertheless, reactions with other phenols will need to be monitored by TLC for completion. The entire chemical process from start to finish, which includes isolation of the crude products and chromatographic purification should typically take no more than two days. Characterization of the products by NMR and mass spectrometry will take additional time depending upon the experiments being performed. More than one reaction can be performed on a single day, with subsequent purifications on a second day. However, this may require greater skills and time management.

In their pure form, products **S.4a–f**, **S.8a,b**, and **S.12a,b** are all likely to be quite stable to prolonged storage in a freezer at -20°C , although individual stability assays have not been performed for each product.

Acknowledgments

This work was partially supported by NIH Grant S06 GM008168-30 and by a PSC CUNY award to M.K.L. Infrastructural support at CCNY was provided by NIH/NCRR/RCMI Grant G12 RR03060.

Literature Cited

- Adamiak RW, Biała E, Skalski B. New, ionic side-products in oligonucleotide synthesis: Formation and reactivity of fluorescent N-/purin-6-yl/pyridinium salts. *Nucleic Acids Res.* 1985; 13:2989–3003. [PubMed: 4000969]
- Aso M, Norihisa K, Tanaka M, Koga N, Suemune H. Synthesis of a new class of spin-labeled purine ribonucleosides and development of a novel nucleophilic reaction to form 2,6,8-trifunctionalized purine derivatives. *J Chem Soc, Perkin Trans.* 2000; 2:1637–1638.
- Allerson CR, Chen SL, Verdine GL. A chemical method for site-specific modification of RNA: The convertible nucleoside approach. *J Am Chem Soc.* 1997; 119:7423–7433.
- Bae S, Lakshman MK. O^6 -(Benzotriazol-1-yl)inosine derivatives: Easily synthesized, reactive nucleosides. *J Am Chem Soc.* 2007; 129:782–789. [PubMed: 17243813]
- Bae S, Lakshman MK. Unusual deoxygenation and reactivity studies related to O^6 -(benzotriazol-1-yl)inosine derivatives. *J Org Chem.* 2008a; 73:1311–1319. [PubMed: 18205381]
- Bae S, Lakshman MK. A novel polymer-supported approach to nucleoside modification. *J Org Chem.* 2008b; 73:3707–3713. [PubMed: 18429630]
- Bae S, Lakshman MK. Synthetic utility of an isolable nucleoside phosphonium salt. *Org Lett.* 2008c; 10:2203–2206. [PubMed: 18476663]
- Bae S, Chaturvedi S, Lakshman MK. O^6 -(Benzotriazol-1-yl)inosine derivatives for C6 modification of purine nucleosides. *Curr Protoc Nucleic Acid Chem.* 2009; 36:1.22.1–.122.23.
- Cosstick R, Douglas ME. Synthesis of a dinucleoside monophosphate analogue containing 6-*N*-(2-aminoethyl)-2'-deoxyadenosine. A novel approach to sequence specific cross-linking in synthetic oligonucleotides. *J Chem Soc, Perkin Trans.* 1991; 1:1035–1040.
- Fathi R, Goswami B, Kung PP, Gaffney BL, Jones RA. Synthesis of 6-substituted 2'-deoxyguanosine derivatives using trifluoroacetic anhydride in pyridine. *Tetrahedron Lett.* 1990; 31:319–322.
- Ferentz AE, Verdine GL. Aminolysis of 2'-deoxyinosine aryl ethers: nucleoside model studies for the synthesis of functionally tethered oligonucleotides. *Nucleosides Nucleotides.* 1992; 11:1749–1763.
- Francom P, Janeba Z, Shibuya S, Robins MJ. Nucleic acid related compounds. 116 Nonaqueous diazotization of aminopurine nucleosides Mechanistic considerations and efficient procedures with *tert*-butyl nitrite or sodium nitrite. *J Org Chem.* 2002; 67:6788–6796. [PubMed: 12227811]
- Francom P, Robins MJ. Nucleic acid related compounds. 118 Nonaqueous diazotization of aminopurine derivatives Convenient access to 6-halo and 2,6-dihalopurine nucleosides and 2'-deoxynucleosides with acyl or silyl halides. *J Org Chem.* 2003; 68:666–669. [PubMed: 12530909]
- Gao H, Fathi R, Gaffney BL, Goswami B, Kung PP, Rhee Y, Jin R, Jones RA. 6-*O*-(Pentafluorophenyl)-2'-deoxyguanosine: A versatile synthon for nucleoside and oligonucleotide synthesis. *J Org Chem.* 1992; 57:6954–6959.
- Harwood EA, Sigurdsson ST, Edfeldt NBF, Reid BR, Hopkins PB. Chemical synthesis and preliminary structural characterization of a nitrous acid interstrand cross-linked duplex DNA. *J Am Chem Soc.* 1999; 121:5081–5082.
- Hayakawa Y, Hirose M, Noyori R. *O*-Allyl protection of guanine and thymine residues in oligodeoxyribonucleotides. *J Org Chem.* 1993; 58:5551–5555.
- Himmelsbach F, Schulz BS, Trichtinger R, Charubala R, Pfeleiderer W. The *p*-nitrophenylethyl (NPE) group: a versatile new blocking group for phosphate and aglycone protection in nucleosides and nucleotides. *Tetrahedron.* 1984; 40:59–72.
- Janeba Z, Lin X, Robins MJ. Functionalization of guanosine and 2'-deoxyguanosine at C6: A modified Appel process and S_NAr displacement of imidazole. *Nucleosides Nucleotides & Nucleic Acids.* 2004; 23:137–147.
- Kaloudis P, Paris C, Vrantz D, Encinas C, Pérez-Ruiz R, Miranda MA, Gimisis T. Photolabile *N*-hydroxypyrid-2(1*H*)-one derivatives of guanine nucleosides: a new method for independent guanine radical generation. *Org Biomol Chem.* 2009; 7:4965–4972. [PubMed: 19907788]
- Kokatla HP, Lakshman MK. One-pot etherification of purine nucleosides and pyrimidines. *Org Lett.* 2010; 12:4478–4481. [PubMed: 20845979]
- Lagisetty P, Russon LM, Lakshman MK. A general synthesis of C6-azolyl purine nucleosides. *Angew Chem Int Ed.* 2006; 45:3660–3663.

- Lakshman MK, Frank J. A simple method for C-6 modification of guanine nucleosides. *Org Biomol Chem.* 2009; 7:2933–2940. [PubMed: 19582304]
- Lakshman MK, Zajc B. A rapid, high-yield method for 5'-hydroxyl protection in very reactive and amino group modified nucleosides using dimethoxytrityl tetrafluoroborate. *Nucleosides Nucleotides.* 1996; 15:1029–1039.
- Lakshman MK, Gunda P, Pradhan P. Mild and room temperature C–C bond forming reactions of nucleoside C-6 arylsulfonates. *J Org Chem.* 2005; 70:10329–10335. [PubMed: 16323841]
- Lakshman MK, Ngassa FN, Keeler JC, Dinh YQV, Hilmer JH, Russon LM. Facile synthesis of *O*⁶-alkyl, *O*⁶-aryl and diaminopurine nucleosides from 2'-deoxyguanosine. *Org Lett.* 2000; 2:927–930. [PubMed: 10768188]
- Lin X, Robins MJ. Mild and efficient functionalization at C6 of purine 2'-deoxynucleosides and ribonucleosides. *Org Lett.* 2000; 2:3497–3499. [PubMed: 11082018]
- Liu J, Janeba Z, Robins MJ. S_NAr iodination of 6-chloropurine nucleosides: Aromatic Finkelstein reactions at temperatures below –40°C. *Org Lett.* 2004; 6:2917–2919. [PubMed: 15330647]
- Meier C, Gräsl S. Highly efficient synthesis of a phosphoramidite building block of C8-deoxyguanosine adducts of aromatic amines. *Synlett.* 2002:802–804.
- Miles RW, Samano V, Robins MJ. Nucleic acid related compounds. 86 Nucleophilic functionalization of adenine, adenosine, tubercidin, and formycin derivatives via elaboration of the heterocyclic amino group into a readily displaced 1,2,4-triazol-4-yl substituent. *J Am Chem Soc.* 1995; 117:5951–5957.
- Nagatsugi F, Uemura K, Nakashima S, Maeda M, Sasaki S. 2-Aminopurine derivatives with C6-substituted olefin as novel cross-linking agents and the synthesis of the corresponding β-phosphoramidite precursors. *Tetrahedron.* 1997; 53:3035–3044.
- Nair V, Richardson SG. Utility of purinyl radicals in the synthesis of base-modified nucleosides and alkylpurines: 6-amino group replacement by hydrogen, chlorine, bromine, and iodine. *J Org Chem.* 1980; 45:3969–3974.
- Nair V, Turner GA, Buenger GS, Chamberlain SD. New methodologies for the synthesis of C-2 functionalized hypoxanthine nucleosides. *J Org Chem.* 1988; 53:3051–3057.
- Qian M, Glaser R. Demonstration of an alternative mechanism for G-to-G cross-link formation. *J Am Chem Soc.* 2005; 127:880–887. [PubMed: 15656626]
- Robins MJ, Basom GL. Nucleic acid related compounds. 8 Direct conversion of 2'-deoxyinosine to 6-chloropurine 2'-deoxyriboside and selected 6-substituted deoxynucleosides and their evaluation as substrates for adenosine deaminase. *Can J Chem.* 1973; 51:3161–3169.
- Robins MJ, Uzna ski B. Nucleic acid related compounds. 33 Conversions of adenosine and guanosine to 2,6-dichloro, 2-amino-6-chloro, and derived purine nucleosides. *Can J Chem.* 1981a; 59:2601–2607.
- Robins MJ, Uzna ski B. Nucleic acid related compounds. 34 Nonaqueous diazotization with *tert*-butyl nitrite Introduction of fluorine, chlorine, and bromine at the C-2 of purine nucleosides. *Can J Chem.* 1981b; 59:2608–2611.
- Rostovtsev VV, Green LG, Fokin VV, Sharpless KB. A stepwise Huisgen cycloaddition process: copper(I)-catalyzed regioselective “ligation” of azides and terminal alkynes. *Angew Chem Int Ed.* 2002; 41:2596–2599.
- Samano V, Miles RW, Robins MJ. Efficient conversion of 6-aminopurines and nucleosides into 6-substituted analogues via novel 6-(1,2,4-triazol-4-yl)purine derivatives. *J Am Chem Soc.* 1994; 116:9331–9332.
- Tornøe CW, Christensen C, Meldal M. Peptidotriazoles on solid phase: [1,2,3]-triazoles by regiospecific copper(I)-catalyzed 1,3-dipolar cycloadditions of terminal alkynes to azides. *J Org Chem.* 2002; 67:3057–3064. [PubMed: 11975567]
- van der Wenden EM, von Frijtag Drabbe Künzel JK, Mathôt RAA, Danhof M, IJzerman AP, Soudijn W. Ribose-modified adenosine analogs as potential partial agonists for the adenosine receptor. *J Med Chem.* 1995; 38:4000–4006. [PubMed: 7562934]
- Véliz EA, Beal PA. C6 substitution of inosine using hexamethylphosphorous triamide in conjunction with carbon tetrahalide or *N*-halosuccinimide. *Tetrahedron Lett.* 2000; 41:1695–1697.

- Véliz EA, Beal PA. 6-Bromopurine nucleosides as reagents for nucleoside analogue synthesis. *J Org Chem.* 2001; 66:8592–8598. [PubMed: 11735542]
- Wan ZK, Binnun E, Wilson DP, Lee J. A highly facile and efficient one-step synthesis of *N*⁶-adenosine and *N*⁶-2'-deoxyadenosine derivatives. *Org Lett.* 2005; 7:5877–5880. [PubMed: 16354089]
- Zajc B, Lakshman MK, Sayer JM, Jerina DM. Epoxide and diol epoxide adducts of polycyclic aromatic hydrocarbons at the exocyclic amino group of deoxyguanosine. *Tetrahedron Lett.* 1992; 33:3409–3412.
- Zemlicka J, Endo T. O⁶-(4-Nitrophenyl)inosine and -guanosine as chromogenic substrates for adenosine deaminase. *Nucleosides Nucleotides.* 1996; 15:619–629.

Figure X.XX.1.

Plausible mechanism for the activation of the C-6 amide in inosine nucleosides and conversion to adenosine derivatives.

Figure X.XX.2.

Two possible modes by which the C-6 amide could react with BOP leading to O⁶-(benzotriazol-1-yl) inosine derivatives.

Figure X.XX.3.

Synthesis of silylated nucleoside precursors, their conversion to the O^6 -(benzotriazol-1-yl) derivatives (which are not isolated), and conversion to the O^6 -alkyl and -aryl ethers.

Table X.XX.1

Substrates and alcohols used for the two-step, one-pot etherification, product numbers and the corresponding yields.

Nucleoside substrate	R-OH or Ar-OH	Product	R or Ar =	% Yield
<p>S.2</p>	CH ₃ -OH	S.4a	CH ₃ —	94
		S.4b		90
		S.4c		86
		S.4d		75
	HO-CH ₂ -CH ₂ -OH	S.4e		63 (+ 15% dimer)
		S.4f		85
<p>S.6</p>	CH ₃ -OH	S.8a	CH ₃ —	73
		S.8b		75
<p>S.10</p>	CH ₃ -OH	S.12a	CH ₃ —	75
		S.12b		73