Published in final edited form as:

Angew Chem Int Ed Engl. 2013 July 15; 52(29): 7590–7592. doi:10.1002/anie.201303457.

Deprotonation of a Borohydride: Synthesis of a Carbene-Stabilized Boryl Anion**

David A. Ruiz, Gaël Ung, Mohand Melaimi, and Guy Bertrand

UCSD-CNRS Joint Research Chemistry Laboratory (UMI 3555) Department of Chemistry and Biochemistry University of California San Diego La Jolla, CA 92093-0343 (USA)

Guy Bertrand: guybertrand@ucsd.edu

Abstract

An acidic hydride! Thanks to the presence of a π -acceptor cyclic (alkyl)(amino)carbene (CAAC) and of two electron-withdrawing nitrile groups, a borohydride reacts with a base to give a carbene-stabilized boryl anion, which reacts with carbon and metal electrophiles at the boron center.

Keywords

Boron; anion; gold

Due to their electron deficiency, trivalent boron compounds are usually electrophilic. As a result, boranes and borohydrides are regarded as the archetypical Lewis acids and hydride donors, respectively. On the other hand, the synthesis of trivalent nucleophilic boron derivatives has eluded chemists for decades, and only recently has their isolation been achieved.^[1] In 2006, Yamashita, Nozaki, and co-worker have reported the preparation and full characterization of the boryl anion **A**,^[2, 3] and since then a few boron centered nucleophiles have been described (Fig. 1). Braunschweig *et al.* have isolated the N-heterocyclic carbene (NHC) stabilized borole **B**^[4] as well as the metalloborylene **C**,^[5] which feature a π -nucleophilic and an sp-hybridized boron atom, respectively. Curran *et al.* have generated the NHC-stabilized parent boryl anion **D**, and although it could not be isolated, they trapped it with a variety of electrophiles.^[6] Also noteworthy is the synthesis by Bernhardt *et al.*, of the dianion $[\text{B}(\text{CN})_3]^{2-}$ **E**, generated by reduction of $\text{K}[\text{B}(\text{CN})_4]$ with potassium.^[7] Lastly in 2011, our group isolated the neutral tricoordinate nucleophilic boron **F**, which is isoelectronic with amines and phosphines.^[8] Unfortunately, the steric

**The authors gratefully acknowledge financial support from the NIH (R01 GM 68825) and DOE (DE-SC0009376), and Dr. C. Moore for assistance with X-ray crystallography.

Correspondence to: Guy Bertrand, guybertrand@ucsd.edu.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

environment around the boron center of **F** hampered its reactivity from any electrophile larger than a proton.

The synthesis of all of these nucleophilic boron compounds **A–F** has been achieved through the reduction of the boron halide or pseudo-halide precursors. So far, any attempts to deprotonate a borohydride to generate a boryl species have failed. Nozaki and Yamashita reported that the boryl anion **A** cannot be prepared by deprotonation, instead a Lewis acid-base adduct is formed.^[3] Curran also mentioned that strong bases did not lead to desired boryl anion **D**.^[6] Indeed, due to the higher electronegativity of hydrogen (2.20) compared to boron (2.04), hydrogens bonded to boron are regarded as hydrides, and thus deprotonation does not seem feasible.

Herein we report the first deprotonation reaction of a neutral borohydride, the isolation of the corresponding carbene-stabilized boryl anion, and the reactivity of the latter as a boron-centered nucleophile.

Inspired by the work of Braunschweig *et al.*^[4] and Curran *et al.*,^[6, 9] we targeted a carbene-stabilized boryl anion. To favor the deprotonation of a borohydride, the obvious option is to place electron-withdrawing groups at boron. We chose a cyclic (alkyl)(amino)carbene (CAAC)^[10] as the Lewis base since they are slightly more nucleophilic but much more electrophilic than NHCs.^[11] The recently prepared CAAC(BH₃) adduct **1**^[12] provided a suitable and simple starting point for our study (Scheme 1). However, all attempts to deprotonate **1** failed, indicating that additional electron-withdrawing groups at boron were necessary. Reacting **1** with two equivalents of trifluoromethane sulfonic acid (TfOH) led to rapid evolution of hydrogen,^[9b] and after work-up, the bis(triflate) derivative **2** was isolated in 80% yield. The ¹¹B NMR displays a broad signal at –2.2 ppm, significantly upfield from **1** (–30.6 ppm), and a single crystal X-ray diffraction study^[13] reveals that both of the triflate groups are covalently bound to boron. Disappointingly, all attempts to deprotonate **2** failed again. Thus, the more electron-withdrawing cyano groups were installed by simple displacement of the triflate substituents with sodium cyanide. Compound **3**, which was isolated in 87% yield, gives a ¹¹B NMR signal at –34.6 ppm (d, *J*_{BH} = 94 Hz), and was fully characterized, including an X-ray diffraction study (Fig. 2, left).

Reacting dicyanide **3** with KHMDS leads to a yellow solution, which displays a broad singlet at –17.9 ppm in the ¹¹B NMR spectrum. After work up a yellow solid was obtained (95% yield), and yellow single crystals were grown by slow evaporation of a THF solution. An X-ray diffraction study reveals the polymeric nature of the boryl anion **4**.^[13] Subsequent treatment with dibenzo-18-crown-6 allows for the isolation of the monomeric boryl anion **4**, with the potassium cation trapped within the sphere of the crown ether (Fig 2, right). As observed by Bernhardt *et al.*, in the case of dianion **E**,^[7] the potassium cation is bound to a nitrile nitrogen (2.7884(15) Å). The B–C_{nitrile} bond lengths in **4** (B1–C2: 1.553(2), B1–C3: 1.552(2) Å) are slightly shorter than in **3** (B1–C2: 1.605(6), B1–C3: 1.595(6) Å), but longer than in **E** (1.513 Å), indicating a weaker delocalization of the negative charge on the nitrile groups. Concomitantly, the B1–C1 bond distance in **4** (1.473(2) Å) is considerably shortened compared to **3** (1.617(6) Å) indicating a strong delocalization of the electron density to the carbene moiety. This is confirmed by the significant lengthening of the C1–N1 bond (**3**: 1.294(4), **4**: 1.396(2) Å), and the trigonal planar geometry of boron characteristic of an sp² hybridization (sum of the angles: 359.96°). Clearly, the π-accepting properties of CAACs play an important role in the stabilization of **4**, as shown by the borataalkene resonance form.^[14]

Boryl anion **4** is stable in the solid state and in solution for weeks under an inert atmosphere with no signs of decomposition, although it is extremely air and moisture sensitive. It reacts

readily with HCl giving back borohydride **3** (Scheme 2). More interestingly, despite the presence of the potassium cation at nitrogen, the addition of isopropyl iodide gave the isopropylborane **5**, as shown by the high field ^{11}B NMR signal (-21.9 ppm), and by an X-ray diffraction study (Fig. 3, left).^[13] The boryl behaviour of **4** was confirmed by the reaction with (trimethylphosphine)gold chloride, which afforded the rare boryl-gold complex **6**.^[14, 15] The ^{11}B NMR of this compound displayed a doublet at -26.2 ppm, which is due to the boron-phosphorus coupling ($J_{\text{BP}} = 30$ Hz). An X-ray diffraction study of **6** (Fig. 3, right) revealed a B1-Au bond length of $2.210(5)$ Å whereas the B1-C1 bond becomes elongated to 1.544 Å compared to **4**, although being shorter than in **3**.

In summary, this work demonstrates that the use of both an electrophilic carbene and electron-withdrawing substituents makes a neutral carbene-borohydride adduct acidic enough to be deprotonated by a strong base. The resulting boryl anion is perfectly stable in the absence of air and moisture. Work is in progress to examine further the reactivity and ligand properties of **4**. So far, we found that electrophiles react at boron. However, we wish to investigate the potential ambiphilic nature of the BCN fragments in the hope of preparing neutral and even cationic tricoordinated boron nucleophiles by mono- and di-N-alkylation, respectively.

Experimental Section

Synthesis of **3**

Derivative **2** (5.0 g, 7.8 mmol) was dissolved in THF (80 mL) and NaCN (0.81 g, 16.5 mmol) was added under a flow of argon. The mixture was stirred for three days, while monitoring the reaction by ^{19}F and ^{11}B NMR. The solvent was removed under vacuum, and the residue extracted with 80 mL of dichloromethane. After evaporation under vacuum, the resulting solid was washed with hexanes (3 X 20 mL) and dried to afford **3** (2.6 g, 87% yield). Colorless single crystals were obtained by vapor diffusion of hexanes in a chloroform solution. Mp: 282 °C; ^{11}B NMR (160 MHz, CDCl_3): $\delta = -34.6$ (d, $J_{\text{BH}} = 94$ Hz). HRMS: m/z calculated for $\text{C}_{25}\text{H}_{37}\text{BN}_3$ ($\text{M}+\text{H}$) $^+$ 390.3080, found 390.3064.

Synthesis of carbene-boryl **4** and **4'**

Derivative **3** (366 mg, 0.94 mmol) and potassium bis(trimethylsilyl)amide (200 mg, 1.00 mmol) were loaded in a Schlenk tube under argon. THF (15 mL) was added to the solids and the mixture stirred for 45 minutes. The volatiles were evaporated under vacuum and the residue was washed with ether (15 mL), then dried, affording **4'** as a yellow solid (382 mg, 95% yield). Single yellow crystals were obtained by slow evaporation of a THF solution. Mp: 345 °C; ^{11}B NMR (160 MHz, THF-d_8): $\delta = -17.9$. HRMS: m/z calculated for $\text{C}_{25}\text{H}_{35}\text{BN}_3$ (M) $^-$ 388.2930, found 388.2992. An excess of dibenzo-18-crown-6 was added to THF solution of **4'**. Single crystals of **4** were obtained by slow evaporation. The ^{11}B NMR spectrum is identical to that of **4'**.

Synthesis of **5**

4' (100 mg, 0.23 mmol) was loaded in a Schlenk tube under argon and dissolved in THF (5 mL). Isopropyl iodide (80 mg, 0.46 mmol) was added and the mixture stirred overnight. Potassium iodide precipitated out of the reaction. The solvent was evaporated and the residue was extracted with dichloromethane (10 mL). The solvent was partly evaporated to obtain a saturated solution. Hexanes were added and the suspension was stored overnight at -20 °C. After filtration **5** (76 mg, 75% yield) was obtained as a solid. Single colorless crystals were obtained by vapor diffusion of hexanes in a chloroform solution. Mp: 294 °C; ^{11}B NMR (160 MHz, CDCl_3): $\delta = -21.9$. HRMS: m/z calculated for $\text{C}_{28}\text{H}_{42}\text{BN}_3\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 453.3400, found 453.3401.

Synthesis of 6

4' (100 mg, 0.23 mmol) and (trimethylphosphine)gold chloride (73 mg, 0.24 mmol) were loaded in a Schlenk tube and 10 mL of THF was added. The mixture was stirred for 2 hours and a precipitate appeared. The solvent was removed under vacuum and the residue was extracted with dichloromethane (10 mL). The solvent was removed and the solid was washed with ether (15 mL) to afford **6** (107 mg, 69% yield) as an off white solid. Single crystals were grown from a saturated chloroform/ether solution. Mp: 160 °C; ^{11}B NMR (160 MHz, CDCl_3): δ = -26.2 (d, J_{BP} = 30 Hz); $^{31}\text{P}\{^1\text{H}\}$ NMR (202 MHz, CDCl_3): δ = 10.1 (br m). HRMS: m/z calculated for $\text{C}_{28}\text{H}_{45}\text{AuBN}_3\text{P}$ ($\text{M}+\text{H}$) $^+$ 662.3109, found 662.3110.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. For reviews, see: Cid J, Gulyas H, Carbo JJ, Fernandez E. Chem. Soc. Rev. 2012; 41:3558–3570. [PubMed: 22337581] Weber L. Eur. J. Inorg. Chem. 2012:5595–5609. Gulyas H, Bonet A, Pubill-Ulldemolins C, Sole C, Cid J, Fernandez E. Pure Appl. Chem. 2012; 84:2219–2231. Yamashita M. Bull. Chem. Soc. Jpn. 2011; 84:983–999. Yamashita M, Nozaki K. J. Synth. Org. Chem. Jpn. 2010; 68:359–369.
2. Segawa Y, Yamashita M, Nozaki K. Science. 2006; 314:113–115. [PubMed: 17023656]
3. For recent papers, see: Dettenrieder N, Dietrich HM, Schaedle C, Maichle-Moessmer C, Tornroos KW, Anwander R. Angew. Chem. 2012; 124:4537–4541. Angew. Chem. Int. Ed. 2012; 51:4461–4465. Yamashita M. Bull. Chem. Soc. Jpn. 2011; 84:983–999. Okuno Y, Yamashita M, Nozaki K. Eur. J. Org. Chem. 2011:3951–3958. Cheung MS, Marder TB, Lin Z. Organometallics. 2011; 30:3018–3028. Okuno Y, Yamashita M, Nozaki K. Angew. Chem. 2011; 123:950–953. Angew. Chem. Int. Ed. 2011; 50:920–923. Terabayashi T, Kajiwaru T, Yamashita M, Nozaki K. J. Am. Chem. Soc. 2009; 131:14162–14163. [PubMed: 19775080]
4. a) Braunschweig H, Chiu CW, Kupfer T, Radacki K. Inorg. Chem. 2011; 50:4247–4249. [PubMed: 21480647] b) Braunschweig H, Chiu C-W, Radacki K, Kupfer T. Angew. Chem. 2010; 122:2085–2088. Angew. Chem. Int. Ed. 2010; 49:2041–2044.
5. a) Braunschweig H, Damme A, Dewhurst RD, Kramer T, Östreicher S, Radacki K, Vargas A. J. Am. Chem. Soc. 2013; 135:2313–2320. [PubMed: 23331127] b) Braunschweig H, Brenner P, Dewhurst RD, Kaupp M, Müller R, Ostreicher S. Angew. Chem. 2009; 121:9916–9919. Angew. Chem. Int. Ed. 2009; 48:9735–9738. c) Braunschweig H, Burzler M, Dewhurst RD, Radacki K. Angew. Chem. 2008; 120:5732–5735. Angew. Chem. Int. Ed. 2008; 47:5650–5653.
6. Monot J, Solovyev A, Bonin-Dubarle H, Derat E, Curran DP, Robert M, Fensterbank L, Malacria M, Lacôte E. Angew. Chem. 2010; 122:9352–9355. Angew. Chem. Int. Ed. 2010; 49:9166–9169.
7. Bernhardt E, Bernhardt-Pitchougina V, Willner H, Ignatiev N. Angew. Chem. 2011; 123:12291–12294. Angew. Chem. Int. Ed. 2011; 50:12085–12088.
8. Kinjo R, Donnadiou B, Celik MA, Frenking G, Bertrand G. Science. 2011; 333:610–613. [PubMed: 21798945]
9. a) Curran DP, Solovyev A, Makhlof Brahmi M, Fensterbank L, Malacria M, Lacôte E. Angew. Chem. 2011; 123:10476–10500. Angew. Chem. Int. Ed. 2011; 50:10294–10317. b) Solovyev A, Chu Q, Geib SJ, Fensterbank L, Malacria M, Lacôte E, Curran DP. J. Am. Chem. Soc. 2010; 132:15072–15080. [PubMed: 20886852]
10. a) Martin D, Melaimi M, Soleilhavoup M, Bertrand G. Organometallics. 2011; 30:5304–5313. [PubMed: 22707843] b) Melaimi M, Soleilhavoup M, Bertrand G. Angew. Chem. 2010; 122:8992–9032. Angew. Chem. Int. Ed. 2010; 49:8810–8849. c) Jazzar R, Dewhurst RD, Bourg J-B, Donnadiou B, Canac Y, Bertrand G. Angew. Chem. 2007; 119:2957–2960. Angew. Chem. Int. Ed. 2007; 46:2899–2902. d) Lavallo V, Canac Y, Prasang C, Donnadiou B, Bertrand G. Angew. Chem. 2005; 117:5851–5855. Angew. Chem. Int. Ed. 2005; 44:5705–5709.

11. a) Back O, Henry-Ellinger M, Martin CD, Martin D, Bertrand G. *Angew. Chem.* 2013; 125:3011–3015. *Angew. Chem. Int. Ed.* 2013; 52:2939–2943. b) Frey GD, Lavallo V, Donnadiou B, Schoeller WW, Bertrand G. *Science*. 2007; 316:439–441. [PubMed: 17446400] c) Lavallo V, Canac Y, Donnadiou B, Schoeller WW, Bertrand G. *Angew. Chem.* 2006; 118:3568–3571. *Angew. Chem. Int. Ed.* 2006; 45:3488–3491.
12. Frey GD, Masuda JD, Donnadiou B, Bertrand G. *Angew. Chem.* 2010; 122:9634–9637. *Angew. Chem. Int. Ed.* 2010; 49:9444–9447.
13. CCDC 931828 (2), 931829 (3), 931831 (4), 931830 (4[†]), 931832 (5) and 931833 (6) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif
14. a) Olmstead MM, Power PP, Weese KJ. *J. Am. Chem. Soc.* 1987; 109:2541–2542. b) Chiu CW, Gabbai FP. *Angew. Chem.* 2007; 119:1753–1755. *Angew. Chem. Int. Ed.* 2007; 46:1723–1725. c) Chiu CW, Gabbai FP. *Angew. Chem.* 2007; 119:7005–7005. *Angew. Chem. Int. Ed.* 2007; 46:6878–6881.
15. Segawa Y, Yamashita M, Nozaki K. *Angew. Chem.* 2007; 119:6830–6833. *Angew. Chem. Int. Ed.* 2007; 46:6710–6713.
16. Other types of boron-gold complexes have been described: Braunschweig H, Brenner P, Dewhurst RD, Kaupp M, Mueller R, Ostreicher S. *Angew. Chem.* 2009; 121:9916–9919. *Angew. Chem. Int. Ed.* 2009; 48:9735–9738. Bontemps S, Bouhadir G, Gu W, Mercy M, Chen C-H, Foxman BM, Maron L, Ozerov OV, Bourissou Didier. *Angew. Chem.* 2008; 120:1503–1506. *Angew. Chem. Int. Ed.* 2008; 47:1481–1484. Sircoglou M, Bontemps S, Mercy M, Saffon N, Takahashi M, Bouhadir G, Maron L, Bourissou D. *Angew. Chem.* 2007; 119:8737–8740. *Angew. Chem. Int. Ed.* 2007; 46:8583–8586. Bontemps S, Bouhadir G, Miqueu K, Bourissou D. *J. Am. Chem. Soc.* 2006; 128:12056–12057. [PubMed: 16967941]

Figure 1.
Known nucleophilic boron compounds (Dipp: 2,6-diisopropylphenyl, Mes: 2,4,6-trimethylphenyl)

Figure 2.
Structures of **3** and **4** in the solid state. Hydrogen atoms except H1 are omitted for clarity.

Figure 3. Structures of **5** and **6** in the solid state. Hydrogen atoms and co-crystallized solvent molecules are omitted for clarity.

Scheme 1.
Synthesis of the borohydride precursor **3**.

Scheme 2.
Synthesis and reactivity of boryl anion **4**.