Published in final edited form as:

J Comp Neurol. 2012 October 1; 520(14): 3236–3255. doi:10.1002/cne.23095.

Localization of Serotonin in the Nervous System of *Biomphalaria glabrata*, an Intermediate Host for Schistosomiasis

Nadia Delgado, Deborah Vallejo, and Mark W. Miller*

Institute of Neurobiology, and Department of Anatomy & Neurobiology, University of Puerto Rico, Medical Sciences Campus, San Juan, Puerto Rico 00901

Abstract

The digenetic trematode *Schistosoma mansoni* that causes the form of schistosomiasis found in the Western Hemisphere requires the freshwater snail *Biomphalaria glabrata* as its primary intermediate host. It has been proposed that the transition from the free-living *S. mansoni* miracidium to parasitic mother sporocyst depends on uptake of biogenic amines, e.g. serotonin, from the snail host. However, little is known about potential sources of serotonin in *B. glabrata* tissues. This investigation examined the localization of serotonin-like immunoreactivity (5HTli) in the central nervous system (CNS) and peripheral tissues of *B. glabrata*. Emphasis was placed on the cephalic and anterior pedal regions that are commonly the sites of *S. mansoni* miracidium penetration. The anterior foot and body wall were densely innervated by 5HTli fibers but no peripheral immunoreactive neuronal somata were detected. Within the CNS, clusters of 5HTli neurons were observed in the cerebral, pedal, left parietal, and visceral ganglia, suggesting that the peripheral serotonergic fibers originate from the CNS. Double-labeling experiments (biocytin backfill × serotonin immunoreactivity) of the tentacular nerve and the three major pedal nerves (Pd n. 10, Pd n. 11, and Pd n. 12) disclosed central neurons that project to the cephalopedal periphery. Overall, the central distribution of 5HTli neurons suggests that, as in other gastropods, serotonin regulates the locomotion, reproductive, and feeding systems of *Biomphalaria*. The projections to the foot and body wall indicate that serotonin may also participate in defensive, nociceptive, or inflammation responses. These observations identify potential sources of host-derived serotonin in this parasite-host system.

INDEXING TERMS

Schistosoma mansoni; pulmonate mollusk; pond snail; miracidium

The parasitic disease schistosomiasis is estimated to presently affect more than 207 million people worldwide (World Health Organization, 2011) and its economic impact on developing countries is second only to malaria (Hotez, 2008; King, 2010). The digenetic trematode blood flukes that cause schistosomiasis require specific aquatic snail species to serve as intermediate hosts, where a sexual reproduction generates the cercariae that are capable of infecting their vertebrate definitive hosts. Within the definitive host, schistosomes reproduce sexually to produce eggs that release free-living miracidia into aquatic habitats where they in turn infect the intermediate snail host.

The trematode species *Schistosoma mansoni* that causes the form of human schistosomiasis found in the Western Hemisphere employs the planorbid snail *Biomphalaria glabrata* as its major intermediate host (Rollinson and Chappell, 2002; Bayne, 2009; Toledo and Fried, 2010). Early investigations reported the presence of serotonin in *B. glabrata*, and advanced the hypothesis that biogenic amines could stimulate miracidium swimming and attraction to the foot and tentacle regions of the snail (Chiang et al., 1974; Etges et al., 1975). Moreover, transition from the miracidium to the parasitic primary sporocyst form of *S. mansoni* that occurs within the *Biomphalaria* integument is proposed to require uptake of serotonin from the snail host (Boyle et al., 2000, 2003; Yoshino et al., 2001; Boyle and Yoshino, 2005). Finally, serotonergic signaling is considered to represent a potential target for parasite manipulation of *B. glabrata* behavior (Manger et al., 1996; Santhanagopalan and Yoshino, 2000; Boyle and Yoshino, 2002) and snail control strategies (Muschamp and Fong, 2001). To date, however, the sources of host-derived serotonin are not well understood and the neural circuitry that controls *Biomphalaria* behavior remains largely unexplored.

Serotonin is a major neurotransmitter and modulator of central neural circuits in gastropods (Gerschenfeld, 1973; Kupfermann et al., 1979; Walker, 1986; Satterlie and Norekian, 1996). Intensive study supports its participation in producing a defensive arousal state in response to aversive stimuli (Brunelli et al., 1976; Jing and Gillette, 2000; Katz et al., 2001; Marinesco and Carew, 2004a, b). In the marine opisthobranch *Aplysia*, serotonin has also been proposed to act in the periphery as a mediator of nociceptive or inflammation signals (Billy and Walters, 1989; Weregoda and Walters, 2007). This study explored the localization of serotonin-like immunoreactive material in the central nervous system and in peripheral tissues of *B. glabrata*. Emphasis was placed on identification of serotonergic neurons that could 1) participate in the infection process by *S. mansoni* miracidia and their transformation to parasitic sporocysts; 2) serve as potential targets for parasite manipulation of snail behavior; and 3) provide targets for novel approaches to vector control. Preliminary reports of these observations were presented in abstract form (Delgado et al., 2010, 2011).

MATERIALS AND METHODS

Specimens

Experiments were conducted on laboratory-reared *B. glabrata* (6–8 mm shell diameter). These specimens were considered sexually mature, as evidenced by their capacity to lay eggs. Snails were housed in plastic aquaria at room temperature (21–23°C) and fed carrots *ad libitum*. All protocols were approved by the Institutional Animal Care and Use Committee (IACUC) of the University of Puerto Rico Medical Sciences Campus (Protocol #3220110).

Nerve backfills

Tissues were dissected in snail saline with the following composition (in mmol l⁻¹): NaCl 51.3, KCl 1.7, MgCl₂ 1.5, CaCl₂ 4.1, HEPES 5, pH 7.8. Dissected ganglia were positioned with minuten pins near a small petroleum jelly (Vaseline) enclosure (3–5 mm diameter) on the surface of a Sylgard-lined Petri dish. The nerve of interest was cut and its end was drawn into the Vaseline-lined pool. The saline was withdrawn from the pool and replaced with a saturated solution (1.4 mg/50 µl dH₂O) of biocytin (Sigma-Aldrich, St. Louis, MO). The enclosure was sealed with Vaseline and the preparation was incubated overnight at room temperature to allow migration of the biocytin. The nerve was then extracted from the pool and the ganglia were re-pinned and washed 3–5 times with saline. Tissues were immersed in 0.5% protease (Type XIV; Sigma-Aldrich) for 30 to 40 minutes and fixed for 1 hour in cold 4% paraformaldehyde prepared in 80 mM phosphate buffer (PB: 24 mM KH₂PO₄, 56 mM Na₂HPO₄, pH 7.4) containing 24% sucrose. Following fixation, tissues were transferred to

microcentrifuge tubes, washed 5 times (30 minutes each) with PTA solution (0.1 M PB containing 2% Triton X-100 and 0.1% sodium azide) and incubated in Alexa Avidin 488 (Molecular Probes, Eugene, OR) diluted 1:1,000 to 1:2,000 in PTA (24–48 hours, room temperature). The preparations were assessed daily until the quality of the backfill staining was determined to be sufficient for advancing to immunohistochemical processing.

Wholemount immunohistochemistry

The dissection, protease, and fixation protocols were performed as described above. Following preincubation with normal goat serum (0.8%, 3–5 hours, room temperature), tissues were immersed (1–7 days, room temperature) in a rabbit polyclonal antiserum (Sigma-Aldrich; #S5545; JCN Antibody Data Base v. 7, #4516) diluted 1:2,000 in PTA. Following repeated PTA washes (5 times, at least 30 minutes each, room temperature), ganglia were incubated in second antibodies conjugated to fluorescent markers (Alexa 488 goat antirabbit IgG (H + L) conjugate or Alexa 546 goat antirabbit IgG (H + L); Molecular Probes). The second antibody dilutions ranged from 1:1,000 to 1:600 μ l and incubation times ranged from 2–10 days. Preparations were examined on a Nikon Eclipse or on a Zeiss Pascal laser scanning confocal microscope (LSCM). Images were captured with the Nikon ACT-1 (v. 2.10) software of the Eclipse or the Zeiss LSM 5 Image Browser (v. 3.1.0.11) program of the LSCM. In most cases, confocal imaging was performed from both the dorsal and ventral surfaces to a depth of 100–150 μ m (2–10 μ m optical sections), depending on the region of interest. Stacks, z-series, overlays, and calibrations were generated using the ImageJ Program (v. 1.43u, NIH public domain). Images were imported to CorelDraw 10 or Microsoft PowerPoint v. 14.0.0 files for addition of labels, adjustment of brightness/contrast, and organization of panels.

Controls, nomenclature, and measurements

The immunogen, serotonin creatinine complex was conjugated with formaldehyde to bovine serum albumin (BSA). Labeling was blocked following preincubation of the antibody (1:1,000) with serotonin creatinine sulfate coupled to BSA with paraformaldehyde (Immunostar product #20081; 20 μ g/ml; overnight; 4°C). It was not reduced following preabsorption with BSA (0.25%; overnight, 4°C).

When possible, the nomenclature for cell clusters and individual neurons were assigned to correspond with designations in more extensively studied pulmonates, such as *Lymnaea stagnalis* (Slade et al., 1981; Benjamin and Winlow, 1981; Croll and Chiasson, 1989) and *Helisoma trivolvis* (Syed et al., 1993). Cluster labels included the ganglion (abbreviated and italicized: cerebral, *Ce*; pedal, *Pe*; pleural, *Pl*; parietal, *Pa*; visceral, *V*; buccal, *B*) and a designation (S) to signify that the serotonergic neurons represent, in most cases, a subset of the entire cluster (see Newcomb et al., 2006). These labels did not further specify the position of the cluster (e.g., dorsal vs. ventral, medial vs. lateral) within the ganglion. Cells were counted on the Nikon fluorescence microscope directly from mounted ganglia. All results reported in this study were observed in a minimum of three specimens.

RESULTS

The general topography of the central ganglia of *Biomphalaria* is similar to other pulmonates (Slade et al., 1981; Kyriakides et al., 1989; Kiehn et al., 1991; Hernádi and Elekes, 1999). The central nervous system (CNS) consists of five paired ganglia (cerebral, pedal, pleural, parietal, and buccal) and one unpaired visceral ganglion (Lever et al., 1965; Chiang et al., 1972). The nervous system has an epiathroid organization (see Chase, 2002) and the most evident asymmetry is found in the parietal ganglia, where the left ganglion is approximately three times larger than the right. In this respect, the *Biomphalaria* CNS corresponds to the

sinistral pulmonates, such as *Helisoma* and *Melampus* (Kahan and Moffett, 1979; Kiehn et al., 1991), and appears as a mirror image of the dextral pulmonates, e.g., *Lymnaea stagnalis* and *Helix* (Slade et al., 1981; Kyriakides et al., 1989; Chase, 2002).

The paired cerebral and pedal ganglia form the major components of the circumesophageal ring, located dorsal and ventral to the esophagus, respectively. The isolated CNS thus possesses a 3D conformation that precludes access to the dorsal surface of the pedal ganglia and the ventral surface of the cerebral ganglia (Fig. 1A). Two manipulations are implemented to render the pulmonate CNS in a more planar configuration that enables visual and physical access to all ganglion surfaces (see Kemenes et al., 1989; Malyshev and Balaban, 2002). In some experiments (Figs. 4, 7–9, 11), the cerebral commissure was severed, and the cerebral hemiganglia were reflected to expose the dorsal surface of the pedal ganglia (Fig. 1B, D, E). For experiments in which it was required to maintain the cerebral ganglia in their natural conformation (Figs. 2, 3, 12), the pedal commissure was severed and the pedal hemiganglia were rotated laterally (Fig. 1C, F). In both configurations the reflected ganglia were viewed from an oblique angle that was predominantly the reverse of the remaining CNS.

Cerebral ganglia

Serotonin-like immunoreactivity was observed in four groups of neurons in each cerebral hemiganglion (Fig. 2). Three clusters, *CeSB*, *CeSC*, and *CeSF*, were visible from the dorsal surface (Fig. 2A) and an additional group, *CeSD*, was observed when preparations were viewed from the ventral surface (Fig. 2B). Frames from a z-stack were examined to estimate the distance of each cluster from the two surfaces of the ganglion. The *CeSD* cluster was located most superficially on the ventral surface (Fig. 2C). While the *CeSB* and *CeSC* clusters were near the dorsal surface, they were commonly observed in sections that were 30–80 μm below the most superficial level, reflecting both their depth and the convex curvature of the surface (Fig. 2D, E). The *CeSF* cluster was located on the most dorsal surface (Fig. 2E).

No major asymmetries were detected in the distribution of serotonin-like immunoreactive neurons between the two cerebral hemiganglia. The *CeSB* cluster was composed of 4 to 6 neurons, one of which was significantly larger (40–50 μm) than the others (10–20 μm). The large *CeSB* neuron, which was found to project to the buccal ganglion (Fig. 3A, B), was designated C1 due to its proposed homology with identified neurons in other gastropods (Sakharov, 1976; Granzow and Rowell, 1981; Kemenes et al., 1989; see Discussion). The *CeSC* cluster was composed of 4 to 7 small (10–20 μm) cells located slightly anterior to the juncture of the cerebral commissure with each cerebral hemiganglion. The *CeSD* cluster, the only serotonergic cluster on the ventral surface of the ganglion, consisted of 6 to 10 moderately sized (20–30 μm) neurons. The dorsal *CeSF* cluster was composed of five brightly stained neurons of varied sizes (20–40 μm in diameter). The axons of most *CeSF* cluster neurons could be followed as they projected in the anterior direction. These axons then turned sharply, contributing to a prominent fiber bundle projecting toward the two cerebral-pedal connectives (observed in Fig. 2B).

Double-labeling experiments were conducted to explore the projection patterns of cerebral serotonergic neurons. Backfills of the cerebral-buccal connective (c-b c.; Fig. 3A) labeled neurons in the ipsi- and contralateral cerebral ganglion, the ipsilateral pleural and parietal ganglia, and the visceral ganglion. When backfilled preparations were processed for serotonin-like immunoreactivity, double-labeling was only detected in C1 (Fig. 3B). This finding is in agreement with observations in several gastropod species, where the sole central serotonergic innervation of the buccal circuitry originates from a single pair of giant cerebral interneurons (Weiss and Kupfermann, 1976; Croll, 1987). A second series of

double-labeling experiments examined central projections to the tentacular nerve (Fig. 3C–E). These experiments showed that neurons of the *CeSD* cluster project to the ipsilateral tentacular nerve. Double-labeling was not detected in any additional neurons in the *Ce g.* or in other ganglia.

Pedal ganglia

The paired pedal ganglia contained the largest number of serotonergic neurons (Fig. 4). On the dorsal surface, a cluster of 10 to 12 neurons of varying size (10–40 μm) extended across the medial region of each hemiganglion. Based on its position and similarities to a serotonergic neuron cluster in *Lymnaea* (Croll and Chiasson, 1989), this group was designated the *PdSIa* cluster (Fig. 4A). A prominent band of 5HTli fibers passing through the pedal commissure appeared to link the two *PdSIa* clusters. As in *Lymnaea*, the *PdSIa* cluster was contiguous to an asymmetric *PdSIb* group of 25 to 30 neurons that extended in the lateral direction on the side of the reproductive organs (the left side in *Biomphalaria*). It was usually possible to distinguish the boundary between the two groups of cells, as the *PdSIb* cluster was composed of neurons that were somewhat smaller (10–20 μm) and more uniform in size than the *PdSIa* cluster (Fig. 4A). Additional unpaired large serotonergic neurons were observed in the right hemiganglion (Fig. 4A). One such cell was designated RPD1 in view of its proposed homology to a giant serotonergic neuron that has been described in other species (Cottrell et al., 1979; Audesirk et al., 1985; Kyriakides et al., 1989; Syed et al., 1993).

Serotonin-immunoreactive neurons were present in four bilateral clusters on the ventral surface of the pedal ganglia (Fig. 4B). The largest ventral cells (30–50 μm) comprised the *PdSA* cluster lining the posteromedial border of each hemiganglion. An expansive cluster of 25 to 30 neurons extended across the entire anterior margin of each Pd g., from the commissure to the base of the middle pedal nerve (Pd n. 11). As it was not possible to partition this assembly into the three divisions (A, B, and C) described by Slade et al. (1981) in *Lymnaea*, it was collectively denoted *PdSB*. Finally, two clusters were distinguished in the lateral region of each hemiganglion, corresponding to the L region in *Lymnaea* (Slade et al., 1981). The more medial group, denoted *PdSLa*, consisted of 7 to 11 neurons and the lateral *PdSLb* clusters were composed of 6 to 10 cells (Fig. 4B). The *PdSLb* cluster was located on the lateral margin of the Pd g. where previous authors described extensions of the dorsal Ib clusters wrapping around to the ventral surface in *Lymnaea* (Slade et al., 1981; Croll and Chiasson, 1989). However, unlike *PdSIb* cells, several neurons in this group had distinctive elongated perikarya and appeared to project in the medial direction. Double-labeling experiments described below support their association with the ventral surface of the Pd g. and their designation as a division of the L cluster. In addition to the neurons in the A, B, and L clusters, several large serotonin-immunoreactive cells were observed on the ventral surface of the Pd g. (Fig. 4B).

Buccal, pleural, parietal, and visceral ganglia

No serotonin-immunoreactive cell bodies were detected in the buccal ganglion (Fig. 5A), the left and right pleural ganglia (Fig. 5B, C), or in the right parietal ganglion (Fig. 5D). Several large-caliber fibers coursed through these ganglia, giving rise to finer collaterals that produced a rich innervation of substantial portions of the neuropil core. In the left parietal ganglion, immunoreactive cell bodies covered the posterior part of the dorsal surface (Fig. 5E). A cluster of smaller cells (10–20 μm) was located near the parietal-visceral connective (*LPaSD*) and a group of slightly larger neurons (20–30 μm) was located more laterally (*LPaSE*), in the central part of the ganglion.

On the dorsal surface of the visceral ganglion, a group of smaller (10–20 μm) serotonergic neurons (VSC) was located near the left parieto-visceral connective (Fig. 5E). Slightly more laterally, larger cells (20–40 μm) extended toward the right parieto-visceral connective (VSH, see Benjamin and Winlow, 1981; Syed et al., 1993). A separate group (VSA) of large neurons (30–50 μm) was located in the anterior region of the *Vg*. The VSA group could also be observed from the ventral perspective of the ganglion (Fig. 5F).

Innervation of the cephalopodal integument

As infection of *B. glabrata* by *S. mansoni* miracidia is commonly achieved by boring through the cephalopodal or tentacular epithelium (see Maldonado, 1967), it was of interest to examine the serotonergic innervation of these tissues. Wholemout processing for serotonin-like immunoreactivity was performed on the anterior region of the foot following a lengthwise mid-dorsal incision and removal of the buccal mass and CNS (Fig. 6). When viewed from the exterior surface, the epithelium exhibited patches of punctate 5HTli stippling (Fig. 6A). In contrast, the internal surface of the pedal epithelium was uniformly lined with a plexus of varicose serotonergic fibers (Fig. 6B). The entire muscular lining of the anterior foot, as well as the lips and more dorsal portions of the head were also innervated by a more diffuse serotonergic fiber system (Fig. 6C). No peripheral serotonergic cell bodies were detected. The remnants of the pedal nerves that remained following removal of the nervous system contained several serotonergic fibers, however, suggesting that the peripheral innervation originates from the CNS. This hypothesis was supported by the observation of numerous 5HTli fibers in each of the three major pedal nerves at their origin in the pedal ganglion (Fig. 6D; superior pedal n., *Pd n 10*; middle pedal n., *Pd n 11*; and inferior pedal n., *Pd n 12*).

Double-labeling experiments (biocytin backfill/serotonin immunohistochemistry) were performed on the three major pedal nerves to localize central serotonergic neurons that project into the foot (Figs. 7–9). The superior pedal nerve (Pedal nerve 10), which innervates the most anterior portion of the foot (see Fig. 5C), originates from the dorsolateral region of the Pd g. (Fig. 6D). Backfills of Pd n. 10 labeled neurons throughout the CNS (Fig. 7A). After processing for serotonin-like immunoreactivity, double-labeling was only detected in the ipsi- and contralateral *PdSLa* clusters (Fig. 7B–H) on the dorsal surface of the pedal ganglia. Interestingly, the number of double-labeled cells in the contralateral *PdSLa* cluster (Fig. 7F–H) exceeded the number in the ipsilateral Pd. g (Fig. 7C–E).

The middle pedal nerve (Pd n. 11), which innervates more central portions of the foot, originates from the ventrolateral margin of the Pd g. (see Fig. 6D). Backfills of Pd n. 11 labeled neurons throughout the CNS (Fig. 8A). After processing for serotonin-like immunoreactivity (Fig. 8B), double-labeling was observed in the ipsilateral *PdSLb* cluster (Fig. 8C–E) and in one additional cell on the ventral surface of the ipsilateral Pd g. (Fig. 8F–H). No contralateral double-labeled cells were detected. Within the *PdSLb* cluster, 3 to 5 of the larger neurons were consistently double-labeled. Their perikarya had an elongated form, with initial processes that were oriented toward the origin of Pd n. 11 (see Fig. 8E). The single double-labeled cell, which was located in the posteromedial quadrant of the ventral surface, had a more rounded form (Fig. 8H).

The inferior pedal nerve (Pd n. 12), which innervates more posterior regions of the foot, descends from the ventral surface of the Pd g. to reach the integument directly below the CNS (Fig. 6D). In our experiments Pd n. 12 was often reflected in the anterior direction to perform the backfill protocol and to facilitate subsequent visualization of the Pd g. (Fig. 9A). Backfills of Pd n. 12 produced labeling of neurons in all central ganglia (Fig. 9A). Subsequent immunostaining for serotonin (Fig. 9B) disclosed double-labeling in the majority of ipsilateral *PdSLa* neurons (Fig. 9A2, B). Two additional double-labeled cells

were located in the medial quadrant of the ventral surface of the ipsilateral Pd g. (Fig. 9C–E). The position, size, and shape of the larger of these neurons resembled the medial double-labeled cell observed in backfills of Pd n. 11 (Fig. 8F–H), suggesting that this cell could project to both nerves. No contralateral double-labeled cells were observed.

Serotonergic innervation of the reproductive system

A survey of the *Biomphalaria* hermaphroditic reproductive system suggested the involvement of serotonin in both female and male reproductive functions (Fig. 10). The most distal structures of the female reproductive tract, including the gonopore, spermathecum, and uterus, were innervated by a dense plexus of 5HTli fibers (Fig. 10A). In the male reproductive tract, serotonin-immunoreactive fibers were observed on the spermiduct (Fig. 10B), as well as on distal structures such as the penis retractor muscle and penis sheath that are critical determinants for the expression of male reproductive behavior (Fig. 10C). No peripheral 5HTli cell bodies were associated with the reproductive organs that were examined.

The neural control of female reproductive function in gastropods is mediated predominantly via the intestinal nerve of the visceral ganglion (see de Vlieger, 1968; Ferguson et al., 1993). Biocytin backfills of the visceral nerve labeled neurons in each of the circumesophageal ganglia (Fig. 11A). When backfilled preparations were processed for 5HTli (Fig. 11B), double-labeling was observed in two neurons near the posterior pole of the left parietal ganglion and in a cluster of superficial dorsal V g. cells belonging to the VSH cluster (Fig. 11C–E). The remaining ganglia were devoid of double-labeled neurons, with the exception of the R Pd g., where a single large serotonergic cell was located on the posterior margin of the ganglion (Fig. 11F–H). This cell corresponded to the RPdD1 neuron described above. RPdD1 was previously shown to project to the intestinal nerve in the sinistral pond snail *Helisoma trivolvis*. (see Syed et al., 1993).

The central control of male reproductive structures is mediated via the penial nerve, which originates from the dorsal surface of the left cerebral hemiganglion. The majority of neurons labeled with backfills of the penial nerve (p n.) were located in clusters on the left side of the CNS (Fig. 12A). These included the prominent ventral lobe of the cerebral ganglion and a cluster of 15 to 20 cells in the pedal ganglion near the cerebral-pedal connective. When backfilled preparations were processed for 5HTli, double-labeling was only observed in the pedal ganglion (Fig. 12B–E). These cells belong to the PdSIb cluster described above (see Fig. 4A), the major asymmetric serotonergic cluster in the Pd g. Notably, not all of the PdSIb neurons were backfilled and not all of the neurons backfilled in this region were serotonergic (Fig. 12E).

DISCUSSION

This study surveyed serotonin-like immunoreactivity in the central nervous system and peripheral tissues of *B. glabrata* (Fig. 13). Previous observations suggest participation of serotonin in the *B. glabrata*–*S. mansoni* host-parasite system (Etges et al., 1975; Manger et al., 1996; Boyle et al., 2000, 2003; Boyle and Yoshino, 2005). Hypotheses concerning the functional properties of the *Biomphalaria* serotonergic system are also suggested by studies conducted in related gastropods (Jing and Gillette, 2000, 2003; Katz et al., 2001; Weragoda and Walters, 2007).

Serotonergic innervation of the cephalopedal integument

Early investigators used microfluorimetry and histochemical fluorescence microscopy to demonstrate the presence of serotonin in the cephalopedal tissues of *B. glabrata* (Chiang et

al., 1974). Etges et al. (1975) reported that *S. mansoni* miracidia were activated by serotonin and proposed that serotonin could form part of a complex attractant “miraxone” released by *B. glabrata* (see Chernin, 1970). The patches of serotonin-immunoreactive punctae observed on the external surface of the foot (Fig. 6A) may reflect an anatomical substrate for serotonin release to the surrounding medium.

Several studies have demonstrated the involvement of serotonin in transformation from the free-living *S. mansoni* miracidium to the parasitic mother sporocyst stage that occurs following penetration of the skin. Infection is followed by an increased dependence on uptake of exogenous serotonin by primary sporocysts where it produces muscular contractions that promote the emergence of daughter sporocysts (Boyle et al., 2000, 2003; Boyle and Yoshino, 2005). Moreover, increased motility of the mother sporocyst is promoted via surface-exposed serotonin receptors (Boyle et al., 2000). The subcutaneous serotonergic fiber plexus and the axons innervating the muscles of the cephalopodal regions (Fig. 6B, C) could serve as sources of host-derived serotonin required for transformation of the *S. mansoni* miracidium to its parasitic sporocyst form and for its subsequent phase of multiplication.

In several opisthobranch and pulmonate species, a plexus of serotonin-immunoreactive fibers innervates the pedal sole ciliary cells that produce locomotion (Syed et al., 1988; Moroz et al., 1997; McKenzie et al., 1998). The network of subcutaneous 5HT₁ fibers observed after removal of the peripheral muscles (Fig. 6B) is likely to correspond to this ciliary motor plexus. As peripheral serotonergic cell bodies are rarely detected, it is thought that this innervation originates from central neurons (Moroz et al., 1997). Longley (2008) described seven serotonergic neurons in the pedal ganglion of *Lymnaea stagnalis appressa* that give rise to the pedal sole plexus. The serotonergic neurons labeled in this study with backfills of the pedal nerves are likely to include cells that contribute to the subcutaneous pedal plexus of *Biomphalaria*.

Serotonin-immunoreactive fibers have also been observed innervating the body wall musculature of several gastropods (McPherson and Blankenship, 1991, 1994; Satterlie et al., 1995; Moroz et al., 1997; McKenzie et al., 1998). In the swimming opisthobranch *Aplysia brasiliana* and in the pteropod *Clione limacina*, serotonergic neurons project from the pedal ganglion to specialized parapodial structures that produce undulating movements (McPherson and Blankenship, 1991; Satterlie et al., 1995). In nonswimming *Aplysia* species, e.g., *Aplysia californica*, pedal serotonergic neurons potentiate motor neuron-evoked contractions in the foot and body wall, (McPherson and Blankenship, 1992). In all cases, the serotonergic neurons exert modulatory actions, i.e., they enhance the contractions produced by motor neurons but do not themselves cause contractions. It was proposed that these serotonergic modulatory neurons act to enhance escape crawling following noxious stimuli (McPherson and Blankenship, 1992).

Tissue damage or inflammation caused by penetration of the cephalopodal skin of *Biomphalaria* by miracidia could stimulate release of serotonin from the subcutaneous or neuromuscular axons observed in this study. In addition to its proposed motor functions considered above, Walters and coworkers advanced the hypothesis that serotonin can induce hyperexcitability of peripheral sensory axons following noxious stimuli in *Aplysia* (Billy and Walters, 1989; Weregoda and Walters, 2007). In support of this proposal, serotonergic pedal ganglion neurons with projections to the periphery were shown to respond to aversive stimuli with increased firing rates (Marinesco et al., 2004a). Moreover, levels of serotonin in the hemolymph of *Aplysia* are increased following sensitizing stimuli (Levenson et al., 1999).

Higher-order serotonergic neurons

A large distinctive neuron in the anterior region of the cerebral ganglion has been observed in a wide range of gastropods (Cottrell and Macon, 1974; Gerschenfeld and Paupardin-Tritsch, 1974; Berry and Pentreath, 1976; Sudlow et al., 1998). While different names have been assigned to these neurons in various species (*Helix*, *Limax*: metacerebral giant cells [MCG; Kandel and Tauc, 1966; Wieland et al., 1987]; *Helisoma*: serotonergic cerebral cell [SCC; Granzow and Fraser Rowell, 1981]; *Aplysia*: metacerebral cell [MCC; see Weiss and Kupfermann, 1976], *Lymnaea stagnalis*: C1 or cerebral giant cell [CGC; see Alania et al., 2008]; *Tritonia*: C1 [Fickbohm et al., 2001]) strong arguments have been advanced proposing their homology (Sakharov, 1976; Weiss and Kupfermann, 1976; Croll, 1987). As a metacerebrum was not readily observed in the *Biomphalaria* cerebral ganglion, and several additional neurons of comparable dimensions were detected, we selected the nondescriptive nomenclature of C1 for the apparent homolog in *B. glabrata*.

The buccal ganglia lack 5HT-immunoreactive somata in all gastropods that have been examined to date (Ono and McCaman, 1984; Longley and Longley, 1986; Croll and Chiasson, 1989; Kemenes et al., 1989; Satterlie et al., 1995; Sudlow et al., 1998; Fickbohm et al., 2001). In all species, the C1 homologs project to the buccal ganglion where they produce modulatory actions on the feeding system (Gillette and Davis, 1977; Kupfermann and Weiss, 1981; Straub and Benjamin, 2001). As the C1 homologs are not *sensu stricto* participants in the feeding central pattern generator (CPG) circuit, this modulatory architecture is designated “extrinsic” (Katz, 1995, 1998). The ability of single serotonergic neurons to exert such broad and coherent actions on the circuits controlling ingestive behavior could provide precise targets for parasite-induced modification of snail feeding and growth (Williams and Gilbertson, 1983; see also de Jong-Brink et al., 2001).

The large neuron in the posterior region of the right pedal ganglion was designated RPdD1 due to its proposed homology with the RPdD1 neuron of the sinistral pulmonate *Helisoma* (Syed et al., 1993) and the LPdD1 neuron of the dextral *Lymnaea* (Cottrell et al., 1979; Audesirk et al., 1985; Kyriakides et al., 1989). As observed in those species, the *Biomphalaria* RPdD1 axon projects through the ipsilateral pleural and parietal ganglia to the visceral ganglion, giving rise to fibers in the parietal and intestinal nerves (Fig. 11H). While the RPdD1 neuron of *Helisoma* was found to possess some synaptic interactions with interneurons involved in the control of respiratory behavior, its function remains incompletely understood (Syed et al., 1993).

The CeSF cluster of five serotonergic neurons (Fig. 2A) is noteworthy due to its possible role in responding to noxious stimuli, such as penetration of miracidia. Katz et al. (2001) described common features of a conserved cluster of five serotonergic neurons on the dorsal surface of the cerebral ganglion in several opisthobranch groups. In the nudibranch *Tritonia diomedea*, the notaspid *Pleurobranchaea californica*, and the gymnosome *Clione limacina* three of these neurons act as interneuronal modulators of swimming (Getting, 1981; Arshavsky et al., 1992; Katz et al., 1994; Satterlie and Norekian, 1995, 1996; Jing and Gillette, 1999). In each of these species, as well as in nonswimming opisthobranchs, the neurons in this cluster respond to noxious stimuli and appear to contribute to a general arousal state (see also Jing and Gillette, 2000; Xin et al., 2001; Marinesco et al., 2004b; Jing et al., 2009). It was proposed that the rhythmic opisthobranch swim interneurons evolved from a nonrhythmic system of serotonergic cells that was responsive to noxious stimuli and that activated several motor systems (Katz et al., 2001). While functional properties of possible homologs in pulmonates have not been examined (see Croll and Chiasson, 1989), we hypothesize that neurons in the CeSF cluster could similarly contribute to the general arousal state (increased heart rate, locomotion, and feeding) observed in *Biomphalaria*.

following infection by *S. mansoni* miracidia (Lee and Cheng, 1971; Williams and Gilbertson, 1983; Boissier et al. 2003).

Serotonergic neurons and the control of reproduction

The array of neuron clusters labeled with backfills of the *Biomphalaria* penial nerve resembled a mirror image of the pattern observed in dextral pulmonates (Li and Chase, 1995; de Boer et al., 1996; Koene et al., 2000). While the prominent cluster of neurons at the origin of the penial nerve in the cerebral ganglion appears to correspond to the cerebral extension of the sinistral snail *Helisoma trivolvis* (Young et al., 1999), we adopted the more widely employed nomenclature of ventral lobe (see Koene, 2010). Previous investigators showed that serotonergic pharmacological antagonists caused penis eversion in *Biomphalaria* and postulated that serotonin signaling promotes penis retraction (Muschamp and Fong, 2001; Fong et al., 2005). This hypothesis is supported by two findings from the present study: 1) the presence of serotonergic fibers and varicosities on the penis sheath and the penis retractor muscle (PRM; Fig. 10C), and 2) the demonstration of a serotonergic projection to the penial nerve from the *PeSIb* cluster (Fig. 12). In *Helix pomatia*, serotonin was shown to produce relaxation of the PRM (Wabnitz and von Wachtendonk, 1976). Moreover, stimulation of individual neurons in the lateral region of the right pedal ganglion produced either PRM contraction or relaxation (Eberhardt and Wabnitz, 1979). Evidence from several species supports the serotonergic phenotype of the pedal Ib cluster (Croll and Chiasson, 1989; Hernádi et al., 1989; Kemenes et al., 1989).

Infection of *B. glabrata* by *S. mansoni* has complex effects on reproduction, including a reduction of egg laying referred to as “parasitic castration” (Minchella and Loverde, 1981; Crews and Yoshino, 1989; Boyle and Yoshino, 2000). Manger et al. (1996) measured significant decreases in the concentration of serotonin in extracts of the CNS and plasma during the prepatency period of infection. As injection of infected snails with exogenous serotonin was found to restore control levels of egg laying, it was proposed that parasitic castration is attributable, at least in part, to the diminished levels of serotonin caused by larval *S. mansoni* infection (Manger et al., 1996). While ovulation in pulmonates is initiated by neuroendocrine factors (Geraerts et al., 1988), several behavioral components of the egg-laying process are triggered by feedback to the CNS, primarily via the intestinal nerve (Ferguson et al., 1993). The double-labeling of neurons in the left parietal, visceral, and right pedal ganglion observed in our experiments with backfills of the intestinal nerve can be expected to reflect the broadest spectrum of potential serotonergic cells involved in motor and sensory function of the genital tract. As this approach will also result in labeling of neurons projecting to additional peripheral structures innervated by the intestinal nerve (see de Vlieger, 1968), more refined anatomical and electrophysiological approaches will be required to identify individual neurons involved in the control of specific stages of oogenesis, and the feedback control of egg-laying behavior.

CONCLUSION

The serotonergic neural system described in this study could act at the interface between *S. mansoni* and *B. glabrata* at all stages of the infection process, ranging from the initial response to miracidium penetration to later phases when competition for energy and resources can occur (see de Jong Brink et al., 2001; Bayne, 2009). Numerous structural and functional commonalities have been recognized between the serotonergic systems of mollusks and mammals, the two hosts of *S. mansoni* (Jacobs and Gelperin, 1981). In both groups, small assemblies of serotonergic neurons or even individual cells have the capacity to broadly impact neural circuits and behavior (Weiger, 1997; Gillette, 2006). This neural architecture led to the proposal that pivotal aminergic systems could provide effective channels for host responses to infection and for parasite manipulation of host behavior (see

Helluy and Holmes, 1990; Maynard et al., 1996; Adamo, 2002; Lèfevre et al., 2009). Our observations should stimulate further investigation of these hypotheses at the level of individual identified neurons.

Acknowledgments

The authors thank Dr. Fred A. Lewis (Biomedical Research Institute), Dr. Adelfa Serrano, and Dr. José F. Maldonado for their encouragement and support.

Grant sponsor: National Institutes of Health; Grant numbers: RCMI RR-03051, NIGMS MBRS: GM-08224, GM-061838; Grant sponsor: National Science Foundation; Grant number: DBI-0115825.

LITERATURE CITED

- Adamo SA. Modulating the modulators: parasites, neuromodulators and host behavioral change. *Brain Behav Evol.* 2002; 60:370–377. [PubMed: 12563169]
- Alania M, Vorontsov DD, Sakharov DA. Higher-order control of the feeding network in *Lymnaea*. *Acta Biol Hung.* 2008; 59(Suppl):22–28.
- Audesirk G. Amine-containing neurons in the brain of *Lymnaea stagnalis*: distribution and effects of precursors. *Comp Biochem Physiol.* 1985; 81A:359–365.
- Bayne CJ. Successful parasitism of vector snail *Biomphalaria glabrata* by the human blood fluke (trematode) *Schistosoma mansoni*: a 2009 assessment. *Molec Biochem Parasitol.* 2009; 165:8–18. [PubMed: 19393158]
- Benjamin PR, Winlow W. The distribution of three wide-acting synaptic inputs to identified neurons in the isolated brain of *Lymnaea stagnalis* (L). *Comp Biochem Physiol A.* 1981; 70:293–307.
- Berry MS, Pentreath VW. Properties of a symmetric pair of serotonin-containing neurones in the cerebral ganglia of *Planorbis*. *J Exp Biol.* 1976; 65:361–380. [PubMed: 187713]
- Billy AJ, Walters ET. Modulation of mechanosensory threshold in *Aplysia* by serotonin, small cardioactive peptide B (SCPB), FMRFamide, acetylcholine, and dopamine. *Neurosci Lett.* 1989; 105:200–204. [PubMed: 2577223]
- Boissier J, Rivera ER, Moné H. Altered behavior of the snail *Biomphalaria glabrata* as a result of infection with *Schistosoma mansoni*. *J Parasitol.* 2003; 89:429–433. [PubMed: 12880237]
- Boyle JP, Yoshino TP. The effect of water quality on oviposition in *Biomphalaria glabrata* (Say, 1818) (Planorbidae), and a description of the stages of the egg-laying process. *J Molluscan Stud.* 2000; 66:83–93.
- Boyle JP, Yoshino TP. Monoamines in the albumen gland, plasma, and central nervous system of the snail *Biomphalaria glabrata* during egg-laying. *Comp Biochem Physiol A Mol Integr Physiol.* 2002; 132:411–422. [PubMed: 12020657]
- Boyle JP, Yoshino TP. Serotonin-induced muscular activity in *Schistosoma mansoni* larval stages: importance of 5-HT transport and role in daughter sporocyst production. *J Parasitol.* 2005; 91:542–550. [PubMed: 16108544]
- Boyle JP, Zaide JV, Yoshino TP. *Schistosoma mansoni*: effects of serotonin and serotonin receptor antagonists on motility and length of primary sporocysts in vitro. *Exp Parasitol.* 2000; 94:217–226. [PubMed: 10831389]
- Boyle JP, Hillyer JF, Yoshino TP. Pharmacological and autoradiographical characterization of serotonin transporter-like activity in sporocysts of the human blood fluke, *Schistosoma mansoni*. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol.* 2003; 189:631–641. [PubMed: 12861423]
- Brunelli M, Castellucci V, Kandel ER. Synaptic facilitation and behavioral sensitization in *Aplysia*: possible role for serotonin and cyclic AMP. *Science.* 1976; 194:1178–1181. [PubMed: 186870]
- Chernin E. Behavioral responses of miracidia of *Schistosoma mansoni* and other trematodes to substances emitted by snails. *J Parasitol.* 1970; 56:287–296. [PubMed: 5445826]
- Chiang PK, Bourgeois JG, Bueding E. Histochemical distribution of acetylcholinesterase in the nervous system of the snail, *Biomphalaria glabrata*. *Int J Neurosci.* 1972; 3:47–60. [PubMed: 4573018]

- Chiang PK, Bourgeois JG, Bueding E. 5-Hydroxytryptamine and dopamine in *Biomphalaria glabrata*. *J Parasitol*. 1974; 60:264–271. [PubMed: 4821112]
- Cottrell GA, Abernathy KB, Barrand MA. Large amine-containing neurons in the central ganglia of *Lymnaea stagnalis*. *Neuroscience*. 1979; 4:685–689. [PubMed: 450258]
- Crews AE, Yoshino TP. *Schistosoma mansoni*: effect of infection on reproduction and gonadal growth in *Biomphalaria glabrata*. *Exp Parasitol*. 1989; 68:326–334. [PubMed: 2703029]
- Croll RP. Identified neurons and cellular homologies. In: Ali, MA., editor. *Nervous systems in invertebrates*. New York: Plenum; 1987. p. 41–59.
- Croll RP, Chiasson BJ. Postembryonic development of serotoninlike immunoreactivity in the central nervous system of the snail, *Lymnaea stagnalis*. *J Comp Neurol*. 1989; 280:122–142. [PubMed: 2918092]
- de Boer PACM, Jansen RF, ter Maat A. Copulation in the hermaphrodite snail *Lymnaea stagnalis*: a review. *Invert Reprod Dev*. 1996; 30:167–176.
- de Boer PACM, Ter Maat A, Pieneman AW, Croll RP, Kurukowa M, Jansen RF. Functional role of peptidergic anterior lobe neurones in male sexual behavior of the snail *Lymnaea stagnalis*. *J Neurophysiol*. 1997; 78:2823–2833. [PubMed: 9405503]
- de Jong-Brink M, Bergamin-Sassen M, Solis Soto M. Multiple strategies of schistosomes to meet their requirements in the intermediate snail host. *Parasitology*. 2001; 123(Suppl):S129–141. [PubMed: 11769278]
- Delgado, N.; Vallejo, DI.; Miller, MW. Program No 783.1. 2010 Neuroscience Meeting Planner. San Diego, CA: Society for Neuroscience, Online; 2010. Localization of biogenic amines in the pond snail *Biomphalaria glabrata*.
- Delgado, N.; Vallejo, DI.; Miller, MW. Program No 707.05. 2011 Neuroscience Meeting Planner. Washington, DC: Society for Neuroscience, Online; 2011. Localization of serotonin in the central nervous system and periphery of *Biomphalaria glabrata*, an intermediate host for schistosomiasis.
- de Vlieger TA. An experimental study of the tactile system of *Lymnaea stagnalis* (L.). *Neth J Zool*. 1968; 18:105–154.
- Dorsett DA. Giant neurones and axon pathways in the brain of *Tritonia*. *J Exp Biol*. 1967; 46:137–115.
- Eberhardt B, Wabnitz RW. Morphological identification and functional analysis of central neurons innervating the penis retractor muscle of *Helix pomatia*. *Comp Biochem Physiol*. 1979; 63A:599–613.
- Etges FJ, Carter OS, Webbe G. Behavioral and developmental physiology of schistosome larvae as related to their molluscan hosts. *Ann N Y Acad Sci*. 1975; 266:480–496. [PubMed: 801112]
- Ferguson GP, Pieneman AW, Jansen RF, Ter Maat A. Neuronal feedback in egg-laying behavior of the pond snail *Lymnaea stagnalis*. *J Exp Biol*. 1993; 178:251–259.
- Fickbohm DJ, Lynn-Bullock CP, Spitzer N, Caldwell HK, Katz PS. Localization and quantification of 5-hydroxytryptophan and serotonin in the central nervous systems of *Tritonia* and *Aplysia*. *J Comp Neurol*. 2001; 437:91–105. [PubMed: 11477599]
- Fong PP, Olex AL, Farrell JE, Majchrzak RM, Muschamp JW. Induction of preputium eversion by peptides, serotonin receptor antagonists, and selective serotonin reuptake inhibitors in *Biomphalaria glabrata*. *Invert Biol*. 2005; 124:296–302.
- Geraerts, WPM.; Ter Maat, A.; Vreugdenhil, E. The peptidergic neuroendocrine control of egg laying behavior in *Aplysia* and *Lymnaea*. In: Laufer, H.; Downer, R., editors. *Invertebrate endocrinology*, vol. 2. *Endocrinology of selected invertebrate types*. New York: Alan Liss; 1988. p. 144–231.
- Gerschenfeld HM. Chemical transmission in invertebrate central nervous systems and neuromuscular junctions. *Physiol Rev*. 1973; 53:1–119. [PubMed: 4405559]
- Gerschenfeld HM, Paupardin-Tritsch D. On the transmitter function of 5-hydroxytryptamine at excitatory and inhibitory monosynaptic junctions. *J Physiol Lond*. 1974; 243:457–481. [PubMed: 4155768]
- Getting PA. Mechanisms of pattern generation underlying swimming in *Tritonia*. I. Neuronal network formed by monosynaptic connections. *J Neurophysiol*. 1981; 46:65–79. [PubMed: 6267210]
- Gillette R. Evolution and function in serotonergic systems. *Integr Comp Biol*. 2006; 46:838–846. [PubMed: 21672789]

- Gillette R, Davis WJ. The role of the metacerebral giant neuron in the feeding behavior of *Pleurobranchaea*. *J Comp Physiol*. 1977; 116:129–159.
- Granzow B, Fraser Rowell CH. Further observations on the serotonergic cerebral neurones of *Helisoma* (Mollusca, Gastropoda): The case for homology with the metacerebral giant cells. *J Exp Biol*. 1981; 90:283–305.
- Helluy S, Holmes JC. Serotonin, octopamine, and the clinging behavior induced by the parasite *Polymorphus paradoxus* (Acanthocephala) in *Gammarus lacustis* (Crustacea). *Can J Zool*. 1990; 68:1214–1220.
- Hernádi L. Topographic organization of efferent neurons with different neurochemical characters in the cerebral ganglia of the snail *Helix aspersa*. *Microsc Res Tech*. 2000; 49:521–533. [PubMed: 10862109]
- Hernádi L, Elekes K. Topographic organization of serotonergic and dopaminergic neurons in the cerebral ganglia and their peripheral projection patterns in the head areas of the snail *Helix aspersa*. *J Comp Neurol*. 1999; 411:274–287. [PubMed: 10404253]
- Hernádi L, Elekes K, S-Rózsa K. Distribution of serotonin-containing neurons in the central nervous system of the snail *Helix pomatia*. *Cell Tissue Res*. 1989; 257:313–323.
- Hotez, PJ. Forgotten people, forgotten diseases: the neglected tropical diseases and their impact on global health and development. Washington, DC: American Society of Microbiology; 2008.
- Jacobs, BL.; Gelperin, A., editors. Serotonin neurotransmission and behavior. Cambridge, MA: MIT Press; 1981.
- Jing J, Gillette R. Central pattern generator for escape swimming in the notaspid sea slug *Pleurobranchaea californica*. *J Neurophysiol*. 1999; 81:654–667. [PubMed: 10036268]
- Jing J, Gillette R. Escape swim network interneurons have diverse roles in behavioral switching and putative arousal in *Pleurobranchaea*. *J Neurophysiol*. 2000; 83:1346–1355. [PubMed: 10712462]
- Jing J, Gillette R. Directional avoidance turns encoded by single interneurons and sustained by multifunctional serotonergic cells. *J Neurosci*. 2003; 23:3039–3051. [PubMed: 12684491]
- Jing J, Vilim FS, Cropper EC, Weiss KR. Neural analog of arousal: Persistent conditional activation of a feeding modulator by serotonergic initiators of locomotion. *J Neurosci*. 2008; 28:12349–12361. [PubMed: 19020028]
- Jing J, Gillette R, Weiss KR. Evolving concepts of arousal: insights from simple model systems. *Rev Neurosci*. 2009; 20:405–427. [PubMed: 20397622]
- Kahan LB, Moffett S. Cobalt mapping of neurons with processes in pedal nerves in the pulmonate snail *Melampus*. *Zoomorphologie*. 1979; 94:81–91.
- Katz PS. Intrinsic and extrinsic neuromodulation of motor circuits. *Curr Opin Neurobiol*. 1995; 5:99–808.
- Katz PS. Neuromodulation intrinsic to the central pattern generator for escape swimming in *Tritonia*. *Ann N Y Acad Sci*. 1998; 860:181–188. [PubMed: 9928311]
- Katz PS, Getting PA, Frost WN. Dynamic neuromodulation of synaptic strength intrinsic to a central pattern generator circuit. *Nature*. 1994; 367:729–731. [PubMed: 8107867]
- Katz PS, Fickbohm DJ, Lynn-Bullock CP. Evidence that the central pattern generator for swimming in *Tritonia* arose from a non-rhythmic neuromodulatory arousal system: implications for the evolution of specialized behavior. *Am Zool*. 2001; 41:962–975.
- Kemenes K, Elekes K, Hiripi L, Benjamin PR. A comparison of four techniques for mapping the distribution of serotonin and serotonin-containing neurons in fixed and living ganglia of the snail, *Lymnaea*. *J Neurocytol*. 1989; 18:193–208. [PubMed: 2732758]
- Kiehn L, Saleuddin S, Lange A. Dopaminergic neurons in the brain and dopaminergic innervation of the albumen gland in mated and virgin *Helisoma duryi* (Mollusca: Pulmonata). *BMC Physiol*. 2001; 1:9. [PubMed: 11513757]
- King CH. Parasites and poverty: the case of schistosomiasis. *Acta Trop*. 2010; 113:95–104. [PubMed: 19962954]
- Koene JM. Neuro-endocrine control of reproduction in hermaphroditic freshwater snails: mechanisms and evolution. *Front Behav Sci*. 2010; 4:167.

- Koene JM, Jansen RF, Ter Maat A, Chase R. A conserved location for the central nervous system control of mating in gastropod molluscs: evidence from a terrestrial snail. *J Exp Biol.* 2000; 203:1071–1080. [PubMed: 10683166]
- Kupfermann, I.; Weiss, KR. The role of serotonin in arousal and feeding behavior of *Aplysia*. In: Jacobs, BL.; Gelperin, A., editors. Serotonin neurotransmission and behavior. Cambridge, MA: MIT Press; 1981. p. 255–287.
- Kupfermann I, Cohen JL, Mandelbaum DE, Schonberg M, Susswein AJ, Weiss KR. Functional role of serotonergic neuromodulation in *Aplysia*. *Fed Proc.* 1979; 38:2095–2102. [PubMed: 221269]
- Kyriakides M, McCrohan CR, Slade CT, Syed NI, Winlow W. The morphology and electrophysiology of the neurons of the paired pedal ganglia of *Lymnaea stagnalis* (L.). *Comp Biochem Physiol A Comp Physiol.* 1989; 93:861–876. [PubMed: 2570671]
- Lee F, Cheng T. *Schistosoma mansoni* infection in *Biomphalaria glabrata*: alterations in heart rate and thermal tolerance in the host. *J Invert Pathol.* 1971; 18:412–418.
- Lefèvre T, Adamo SA, Biron DG, Missé D, Hughes D, Thomas D. Invasion of the body snatchers: the diversity and evolution of manipulative strategies in host-parasite interactions. *Adv Parasitol.* 2009; 68:45–83. [PubMed: 19289190]
- Levenson J, Byrne JH, Eskin A. Levels of serotonin in the hemolymph of *Aplysia* are modulated by light/dark cycles and sensitization training. *J Neurosci.* 1999; 19:8094–8103. [PubMed: 10479709]
- Lever J, de Vries CM, Jager JC. On the anatomy of the central nervous system and the location of neurosecretory cells in *Australorbis glabratus*. *Malacologia.* 1965; 2:219–230.
- Li G, Chase R. Correlation of axon projections and peptide immunoreactivity in mesocerebral cells of the snail *Helix aspersa*. *J Comp Neurol.* 1995; 353:9–17. [PubMed: 7714252]
- Longley RD. Development of the 5-HT-like immunoreactive pedal plexus in the pond snail *Lymnaea stagnalis appressa*. *Biol Bull.* 2008; 215:280–294. [PubMed: 19098149]
- Longley RD, Longley AJ. Serotonin immunoreactivity of neurons in the gastropod *Aplysia californica*. *J Neurobiol.* 1986; 17:339–358. [PubMed: 3018150]
- Maldonado, JF. Schistosomiasis in America. Barcelona: Editorial Científico-Médica; 1967.
- Manger P, Li J, Christensen BM, Yoshino TP. Biogenic monoamines in the freshwater snail, *Biomphalaria glabrata*: influence of infection by the human blood fluke, *Schistosoma mansoni*. *Comp Biochem Physiol.* 1996; 114A:227–234.
- Marinesco S, Kolkman KE, Carew TJ. Serotonergic modulation in *Aplysia*. I. Distributed serotonergic network persistently activated by sensitizing stimuli. *J Neurophysiol.* 2004a; 92:2468–2486. [PubMed: 15140903]
- Marinesco S, Wickremasinghe N, Kolkman KE, Carew. Serotonergic modulation in *Aplysia*. II. Cellular and behavioral consequences of increased serotonergic tone. *J Neurophysiol.* 2004b; 92:2487–2496. [PubMed: 15140904]
- Maynard BJ, DeMartini L, Wright WG. *Gammarus lacustris* harboring *Polymorphus paradoxus* show altered patterns of serotonin-like immunoreactivity. *J Parasitol.* 1996; 82:663–666. [PubMed: 8691384]
- McCrohan CR, Benjamin PR. Patterns of activity and axonal projections of the cerebral giant cells of the snail, *Lymnaea stagnalis*. *J Exp Biol.* 1980; 85:149–169. [PubMed: 6154759]
- McKenzie JD, Counce M, Hetherington MS, Winlow W. Serotonergic innervation of the foot of the pond snail *Lymnaea stagnalis* (L.). *J Neurocytol.* 1998; 27:459–470. [PubMed: 10192527]
- McPherson DR, Blankenship JE. Neural control of swimming in *Aplysia brasiliana*. III. Serotonergic modulatory neurons. *J Neurophysiol.* 1991; 66:1366–1379. [PubMed: 1662263]
- McPherson DR, Blankenship JE. Neuronal modulation of foot and body wall contractions in *Aplysia californica*. *J Neurophysiol.* 1992; 67:23–28. [PubMed: 1552321]
- Minchella DJ, Loverde PT. A cost of increased early reproductive effort in the pond snail *Biomphalaria glabrata*. *Am Natural.* 1981; 118:876–881.
- Moroz LL, Sudlow LC, Jing J, Gillette R. Serotonin-immunoreactivity in peripheral tissues of the opisthobranch molluscs *Pleurobranchaea californica* and *Tritonia diomedea*. *J Comp Neurol.* 1997; 382:176–188. [PubMed: 9183687]

- Muschamp JW, Fong PP. Effects of the serotonin receptor ligand methiothepin on reproductive behavior of the freshwater snail *Biomphalaria glabrata*: reduction of egg laying and induction of penile erection. *J Exp Zool.* 2001; 289:202–207. [PubMed: 11170017]
- Newcomb JM, Fickbohm DJ, Katz PS. Comparative mapping of serotonin-immunoreactive neurons in the central nervous systems of nudibranch molluscs. *J Comp Neurol.* 2006; 499:485–505. [PubMed: 16998939]
- Ono JK, McCaman RE. Immunocytochemical localization and direct assays of serotonin-containing neurons in *Aplysia*. *Neuroscience.* 1984; 11:549–560. [PubMed: 6717803]
- Pentreath, VW.; Berry, MS.; Osborne, NN. The serotonergic cerebral cells in gastropods. In: Osborne, NN., editor. *Biology of serotonergic transmission*. Chichester, UK: Wiley Press; 1982. p. 457–513.
- Rollinson, D.; Chappell, LH. Flukes and snails revisited. London: Cambridge University Press; 2002.
- Sakharov, DA. Nerve cell homologues in gastropods. In: Salánki, J., editor. *Neurobiology of invertebrates: gastropod brain*. Budapest: Akadémiai Kiadó; 1976. p. 27–40.
- Satterlie RA, Norekian TP. Serotonergic modulation of swimming speed in the pteropod mollusk *Clione limacina*. III. Cerebral neurons. *J Exp Biol.* 1995; 198:917–930. [PubMed: 7730754]
- Satterlie RA, Norekian TP. Modulation of swimming speed in the pteropod mollusk, *Clione limacina*: role of a compartmental serotonergic system. *Invert Neurosci.* 1996; 1996:157–165. [PubMed: 9372161]
- Satterlie RA, Norekian TP, Jordan S, Kazilek CJ. Serotonergic modulation of swimming speed in the pteropod mollusk *Clione limacina*. *J Exp Biol.* 1995; 198:895–904. [PubMed: 7730752]
- Slade CT, Mills J, Winlow W. The neuronal organization of the paired pedal ganglia of *Lymnaea stagnalis* (L.). *Comp Biochem Physiol A.* 1981; 69:789–803.
- Sudlow LC, Jing J, Moroz LL, Gillette R. Serotonin immunoreactivity in the central nervous system of the marine molluscs *Pleurobranchaea californica* and *Tritonia diomedea*. *J Comp Neurol.* 1998; 395:466–480. [PubMed: 9619500]
- Syed NI, Roger I, Ridgway RL, Bause LG, Lukowiak K, Bulloch AGM. Identification, characterization, and in vitro reconstruction of an interneuronal network of the snail, *Helisoma trivolvis*. *J Exp Biol.* 1993; 174:19–44. [PubMed: 8440965]
- Toledo, R.; Fried, B., editors. *Biomphalaria snails and larval trematodes*. New York: Springer Press; 2010.
- Wabnitz RW, von Wachtendonk D. Evidence for serotonin (5-hydroxytryptamine as transmitter in the penis retractor muscle of *Helix pomatia* L. 1976. *Cell Molec Life Sci.* 1976; 32:707–709.
- Walker, RJ. Transmitters and modulators. In: Willows, AOD., editor. *The Mollusca*, vol. 9. *Neurobiology and behavior*, part 2. Orlando, FL: Academic Press; 1986. p. 279–485.
- Weiger WA. Serotonergic modulation of behavior: a phylogenetic overview. *Biol Rev Camb Philos Sci.* 1997; 72:61–95.
- Weiss KR, Kupfermann I. Homology of the giant serotonergic neurons metacerebral cells in *Aplysia* and pulmonate molluscs. *Brain Res.* 1976; 117:33–49. [PubMed: 990936]
- Weregoda RMS, Walters ET. Serotonin induces memory-like, rapamycin-sensitive hyperexcitability in sensory axons of *Aplysia* that contributes to injury responses. *J Neurophysiol.* 2007; 98:1231–1239. [PubMed: 17634332]
- Wieland SJ, Jahn E, Gelperin A. Localization and synthesis of monoamines in regions of *Limax* CNS controlling feeding behavior. *Comp Biochem Physiol C.* 1987; 86:125–130. [PubMed: 2881707]
- Williams CL, Gilbertson DE. Altered feeding responses as a cause for the altered heartbeat rate and locomotor activity of *Schistosoma mansoni*-infected *Biomphalaria glabrata*. *J Parasitol.* 1983; 69:671–676. [PubMed: 6631636]
- World Health Organization. WHO Media Centre. Schistosomiasis. Fact sheet No 115. Geneva: WHO; 2011. www.who.int/mediacentre/factsheets/fs115/en/index.html
- Xin Y, Koester J, Jing J, Weiss KR, Kupfermann I. Cerebral-abdominal interganglionic coordinating neurons in *Aplysia*. *J Neurophysiol.* 2001; 85:174–186. [PubMed: 11152718]
- Yoshino TP, Boyle JP, Humphries JE. Receptor-ligand interactions and cellular signaling at the host-parasite interface. *Parasitology.* 2001; 123:S143–S157. [PubMed: 11769279]

Young KG, Chang JP, Goldberg JJ. Gonadotropin-releasing hormone neuronal system of the freshwater snails *Helisoma trivolvis* and *Lymnaea stagnalis*: possible involvement in reproduction. *J Comp Neurol.* 1999; 404:427–437. [PubMed: 9987988]

Figure 1.

Biomphalaria central nervous system: topography and experimental manipulations. **A:** The circumesophageal ring of Bassomatophoran pulmonates consists of paired cerebral ganglia (*L Cer g.*, *R Cer g.*), pleural ganglia (*L Pl g.*, *R Pl g.*), and parietal ganglia (*L Pa g.*, *R Pa g.*). A single visceral ganglion (*V g.*) is located in the most posterior position of the ring and a pair of buccal ganglia (*B g.*) is located more anteriorly on the pharynx. The left parietal ganglion is markedly larger than the right, reflecting the sinistral orientation of major reproductive and respiratory structures in *Biomphalaria*. The paired cerebral and pedal ganglia are connected by the cerebral commissure (*C c.*) and pedal commissure (*P c.*), respectively. The ventral surface of the cerebral ganglia (dark shading) lies in close apposition to the dorsal surface of the pedal ganglia (lighter shading) limiting much of their accessibility and visibility. **B:** In one of the dissections used in this study (Figs. 4, 7–9, 11), the cerebral commissure was severed prior to fixation and the two cerebral ganglia were reflected laterally to expose the dorsal surface of the pedal ganglia. **C:** In the other configuration employed in this study (Figs. 2, 3, 12), the pedal commissure was severed and two pedal hemiganglia were reflected laterally. **D:** Low-power view of serotonin-like immunoreactivity in a nervous system prepared as illustrated in B. When viewing the dorsal surface of the pedal, parietal, and visceral ganglia, the separated cerebral hemiganglia are observed primarily from their dorsomedial aspect. **E:** When a preparation dissected in the configuration shown in B is observed from the opposite perspective, with the plane of focus on the ventral surfaces of the pedal, parietal, and visceral ganglia (note left–right reversal), the divided cerebral hemiganglia are viewed primarily from their dorsolateral aspect. **F:** Serotonin-like immunoreactivity in a wholemount CNS oriented as illustrated in C. In this example, the cerebral, parietal, and visceral ganglia are observed from their ventral aspect

(note left–right reversal), and the divided pedal ganglia are viewed from a dorsolateral perspective. Scale bar = 200 μm (applies to D–F).

Figure 2.

Serotonin-like immunoreactivity in the cerebral ganglion. **A:** The paired cerebral ganglia viewed from the dorsal surface. Three clusters of neurons, *CeSB*, *CeSC*, and *CeSF*, are located in each hemiganglion. **B:** Same preparation as A viewed from the ventral aspect. An additional cluster, the *CeSD* group, is observed on the ventral surface of each hemiganglion. The large *CI* neuron is located within the *CeSB* cluster in the anterior region of each hemiganglion. **C–F:** Individual frames from the z-stack used to create the projection shown in B. Images captured at distances of C: 30 μm , D: 150 μm , E: 180 μm , and F: 220 μm from the ventral surface of the right cerebral hemiganglion. Clusters correspond to enclosed regions in B: *CeSD*: dotted ellipse; *CeSC*: dotted rectangle; *CeSB*: dashed ellipse; *CeSF*: dashed rectangle, applies to all panels.

Figure 3.

Projections from serotonergic neurons in the cerebral ganglion. **A,B:** Serotonergic projection to the buccal neuromuscular system originates from C1. **A:** Neurobiotin backfill of the cerebral-buccal connective (*C-b c.*) labels several cell bodies in the anterior region of the cerebral ganglion (ventral surface of the left *Ce g.* shown). **B:** After processing for serotonin-like immunoreactivity (same preparation as A), double-labeling was only detected in C1 of the *C&SB* cluster. The intensity of 5HTli observed in the C1 was typically weaker than the other neurons in the *C&SB* cluster, possibly reflecting its comparatively large nucleus or lower somatic serotonin levels. **C–E:** Serotonergic projection to the tentacular nerve originates in the *C&SD* cluster. **C:** Neurobiotin backfill of the tentacular nerve labeled neurons throughout the cerebral ganglia, including a cluster of tightly apposed cells near the ventral surface of the ipsilateral hemiganglion (enclosed by white dashed ellipse; right hemiganglion shown). **D:** After processing for serotonin-like immunoreactivity, neurons of the *C&SD* cluster (white dashed ellipse) appeared to correspond to the cells labeled by the tentacular nerve tracing. **E:** Overlay of C, D images confirmed double-labeling (cells appear yellow) of all neurons in the ipsilateral *C&SD* cluster. Scale bars = 50 μm in B (applies to A); 100 μm in E (applies to C,D).

Figure 4.

Serotonin-like immunoreactivity in the pedal ganglia. **A:** Dorsal surface of the paired pedal ganglia. Serotonergic neurons of varied sizes (10–40 μm) are located in the bilateral *PdSLa* clusters that form a band across the dorsal surface of each hemiganglion. The left *PdSLa* cluster is contiguous with the unilateral *PdSLb* cluster of smaller (10–20 μm) neurons which extend to the lateral edge of the ganglion. A bundle of immunoreactive fibers is present in the pedal commissure (*Pd c.*). The giant *RPdD1* neuron is visible on the posterior edge of the *RPd g.* **B:** Serotonin-like immunoreactivity on the ventral surface of the pedal ganglia (same preparation as A). Large (20–60 μm) neurons are present in the central *PdSA* clusters on each side of the *Pd c.* The bilateral *PdSB* clusters form prominent bands across the anterior edge of each hemiganglion and the *PdSLa* and *PdSLb* clusters of smaller (10–20 μm) neurons are located more laterally. Additional large immunoreactive cells that are not part of these clusters are observed on both surfaces of the *Pe g.* Scale bar = 100 μm .

Figure 5.

Serotonin-like immunoreactivity in the buccal, pleural, parietal, and visceral ganglia. **A–D:** No serotonin-immunoreactive neurons were detected in the buccal ganglion (A), the left (B) and right (C) pleural ganglia (*L Pl g.*, *R Pl g.*), or (D) the right parietal ganglion (*R Pa g.*). Immunoreactive axons coursed through each of these ganglia, giving rise to extensive networks of fine fibers. Immunoreactive fibers were also observed in most peripheral nerves. Abbreviations: *b c.*, buccal commissure; *c-b c.*, cerebral-buccal connective; *e t.*, esophageal trunk; *pb n.*, posterobuccal nerve. **E:** Clusters of 5HTli neurons extended across the posterior regions of the left parietal ganglion (*L Pa g.*) and the visceral ganglion (*V g.*). The *L Pa g.* group was composed of a cluster of smaller cells (*LPaSD*) located near the left parietal-visceral connective and a neighboring cluster (*LPaSE*) of larger cells located more laterally toward the central region of the ganglion. In the visceral ganglion, a cluster of smaller cells (*VSC*) was located near the left parietal-visceral connective. Lateral to the *VSC* cluster a group of larger neurons (*VSH*, see Benjamin and Winlow, 1981; Syed et al., 1993) extended in the anterior direction toward the right parietal-visceral connective. A separate group of larger neurons (*VSA*) located in the anterior region of the *V g.* could also be seen from the ventral perspective (F). **F:** A few 5HTli neurons were observed on the ventral surfaces of the *V g.* and the *L Pa g.* (same preparation as E). The nerves of the *L Pa g.* were rich in immunoreactive fibers. Scale bars = 50 μ m in A, F (applies to E); 100 μ m in D (applies to B, C).

Figure 6.

Serotonin-like immunoreactivity in the cephalopedal region of *Biomphalaria*. **A:** Irregular patches of 5HTli punctae viewed from the external surface of the pedal epithelium. **B:** The entire internal surface of the epithelium was covered by a dense network of varicose immunoreactive fibers. **C:** Low-power image of the anterior foot and lips. The buccal mass was removed exposing the muscles associated with the anterior foot and the lips. Muscles were innervated by a plexus of serotonergic fibers. The insertions of the left and right pedal nerves 10 (*Pd n 10*) were captured in this image. **D:** Right pedal ganglion, ventral surface. Numerous serotonin-immunoreactive fibers were present in each of the three major pedal nerves that innervate the foot (superior pedal n., *Pd n 10*, medial pedal n., *Pd n 11*, and inferior pedal n., *Pd n 12*). Scale bars = 100 μ m in A–C; 50 μ m in D.

Figure 7.

Localization of serotonergic neurons that project to pedal nerve 10. **A:** Biocytin backfill of the left pedal nerve 10 (*LPd n. 10*) labeled neurons in all of the central ganglia. **B:** Serotonin-like immunoreactivity in the same preparation as A. Double-labeling was only detected in the left and right dorsal *PaSIa* clusters (dashed rectangle and dotted rectangle, respectively). **C–E:** Double-labeling in the left *PaSIa* cluster. **C:** Enlarged image of area enclosed by the dashed rectangle in A. Several neurons were labeled in the region of the serotonergic *PaSIa* cluster. Only two cells (arrows) were subsequently shown to contain 5HTli. **D:** Enlarged image of area enclosed by dashed rectangle in B. Arrows indicate two neurons in the *PaSIa* cluster that correspond to backfilled cells in C. **E:** In an overlay of C, D, only the two neurons denoted by arrows in those panels appear white. **F–H:** Double-labeling in the right *PaSIa* cluster. **F:** Enlarged image of area enclosed by the dotted rectangle in A. Several neurons were labeled in the region of the serotonergic *PaSIa* cluster. Seven neurons (marked by asterisks) were subsequently shown to contain 5HTli. **G:** Enlarged image of area enclosed by dotted rectangle in B. Asterisks mark neurons in the *PaSIa* cluster that correspond to the backfilled cells in F. **H:** In an overlay of F, G, the neurons marked by asterisks in those panels appear white. Scale bars = 200 μm in B (applies to A); 50 μm in E, H (applies to C, D, F, G).

Figure 8.

Localization of serotonergic neurons that project to pedal nerve 11. **A:** Biocytin backfill of the right pedal nerve 11 (*R Pd n. 11*) labeled neurons in all of the central ganglia. **B:** Serotonin-like immunoreactivity on the ventral surface of the same preparation as in A. Double-labeling was detected in neurons on the lateral edge in the ipsilateral *PdSLb* cluster and in one additional ventral neuron (dashed oval and dotted rectangle, respectively). **C–E:** Double-labeling in the right *PdSLb* cluster. **C:** Enlarged image of area enclosed by dashed oval in A. Several neurons were labeled on the lateral edge of the ganglion in the *PdSLb* cluster. **D:** Enlarged image of area enclosed by dashed oval in B. Oval surrounds four neurons in the *PdSLb* cluster that correspond to the backfilled cells in C. **E:** In an overlay of C, D, only the four neurons that are marked by both the backfill and immunohistochemistry appear white. **F–H:** Double-labeling of a ventromedial neuron. **F:** Enlarged image of area enclosed by the dotted rectangle in A. A single neuron labeled in the central region of the ganglion (arrow) was subsequently shown to contain 5HTli. **G:** Enlarged image of area enclosed by dotted rectangle in B. Arrow denotes immunoreactive neuron that corresponds to the backfilled cell in F. **H:** In an overlay of F, G, the neuron marked by the arrow in those panels appears white. Scale bars = 150 μ m in B (applies to A); 30 μ m in E, H (applies to C, D, F, G).

Figure 9.

Localization of serotonergic neurons that project to pedal nerve 12. **A:** Biocytin backfill of the right pedal nerve 12 (*R Pd n. 12*) labeled neurons in all of the central ganglia (ventral surface shown). **B:** Serotonin-like immunoreactivity on the ventral surface of the same preparation as in A. Double-labeling was detected in neurons in the region of the ipsilateral *PdSLa* cluster and in two additional ventral neurons (dashed rectangle and dotted rectangle, respectively). **C–E:** Double-labeling in the right *PdSLa* cluster. **C:** Enlarged image of area enclosed by the dashed rectangle in A. Several neurons were labeled in the medial right *PdSLa* cluster (surrounded by dashed line). **D:** Enlarged image of area enclosed by dashed rectangle in B. Dashed line surrounds eight neurons in the right *PdSLa* cluster that correspond to the backfilled cells in C. **E:** In an overlay of C, D, only the eight neurons that are marked by both the backfill and immunohistochemistry appear white. **F–H:** Double-labeling of two additional ventromedial neurons. **F:** Enlarged image of area enclosed by the dotted rectangle in B. Two neurons labeled in the central region of the ganglion (arrow and arrowhead) were subsequently shown to contain 5HTli. **G:** Enlarged image of area enclosed by dotted rectangle in A. Arrow and arrowhead denote immunoreactive neurons that correspond to the backfilled cells in F. **H:** In an overlay of F, G, the two neurons marked by the arrow and arrowhead in those panels appears white. Scale bars = 150 μ m in B (applies to A); 20 μ m in E (applies to C, D); 30 μ m in H (applies to F, G).

Figure 10.

Serotonergic innervation of reproductive structures. **A:** Serotonergic fibers covered the distal structures of the female reproductive tract, including the spermatheca (*Sp*) and the uterus (*Ut*). The fiber network diminished abruptly near the juncture of the *Ut* with the nidamental gland (*Ng*). **B:** Serotonin-like immunoreactivity was observed in more proximal segments of the reproductive tract, including the spermiduct (*Sd*) as it coursed in contiguity with the larger oviduct (*Ov*), which was not innervated. **C:** Serotonergic fibers were also found on distal regions of the male reproductive tract, including the penis retractor muscle (*Prm*) and the preputium (*Pp*). The most intense staining was observed lining the region of insertion of the *Pp* by the *Prm*. Scale bar = 200 μ m (applies to all panels).

Figure 11.

Localization of serotonergic neurons that project to the intestinal nerve. **A:** Biocytin backfill of the intestinal nerve (*Int n.*) labeled neurons in all of the central ganglia (dorsal surface shown). **B:** Serotonin-like immunoreactivity on the dorsal surface of the same preparation as **A**. Double-labeling was detected in two large neurons near the posterior pole of the *L Pa g.* and in several cells associated with the *VSH* cluster in the visceral ganglion (dashed rectangles in **A**, **B**). Double-labeling was also observed in the giant *RPd1* cell in the right pedal ganglion (dotted rectangles in **A** and **B**). **B:** Double-labeling in cells on the dorsal surface of the left parietal ganglion and the visceral ganglion. **C:** Enlarged image of area enclosed by the dashed rectangle in **A**. Backfills of the intestinal nerve labeled several neurons in the *L Pa g.* and the *V g.* **D:** Serotonin-like immunoreactivity, same field as **C**, area enclosed by dashed rectangle in **B**. Clusters of neurons were labeled in the *L Pa g.* and the *V g.* **E:** Overlay of **C**, **D** images disclosing double-labeling (white) in two large neurons

at the posterior pole of the *L Pa g.* and in the VSH cluster of the visceral ganglion. **F–H:** Double-labeling in the RPeD1 neuron. F: Enlarged image of area enclosed by the dotted rectangle in A. Backfills of the intestinal nerve labeled a single giant neuron in the posterior region of the right pedal ganglion. G: Serotonin-like immunoreactivity in the same field as C (enclosed by dotted rectangle in B) marked the giant RPeD1 neuron. H: Overlay of F, G images demonstrates double-labeling in RPeD1. Scale bars = 200 μm in B (applies to A); 100 μm in E (applies to C, D); 50 μm in H (applies to F, G).

Figure 12.

Localization of central serotonergic neurons that project to the penial nerve. **A:** The majority of neurons labeled by backfilling the penial nerve (*p n.*) were located in five clusters on the left side of the CNS: the left ventral lobe (*LVL*) near the origin of the *p n.* in the left cerebral ganglion (*L Ce g.*), single clusters in the left pedal ganglion (*L Pd g.*) and left pleural ganglion (*L Pl g.*), and two groups of cells in the left parietal ganglion (*L Pa g.*). Additional individual neurons were labeled in the visceral ganglion (*V g.*), the right cerebral ganglion (*R Ce g.*), the right pedal ganglion (*R Pd g.*), and the buccal ganglion (not shown). **B:** The serotonergic projection to the penial nerve was located in the *PdSIb* cluster (see Fig. 4A) on the dorsal surface of the left pedal ganglion. Optimal imaging of the serotonergic projection was obtained with preparations oriented as shown in Fig. 1F. In this image the ventral lobe is viewed from the ventral aspect of the left cerebral ganglion (different preparation than A). 5HTli neurons were labeled (magenta) throughout the CNS, with double-labeling (white) limited to cells in the *PdSIb* cluster of the left pedal ganglion. **C:** Enlarged image of the region enclosed by dashed rectangle in B. Only the backfill is shown. **D:** Enlarged image of the region enclosed by dashed rectangle in B. Only the 5HTli is shown. **E:** Overlay of C, D. Double-labeling was only observed in a cluster of small neurons near the origin of the cerebral-pedal connective (*Ce-pd c.*). Scale bars = 100 μ m in A, B; 50 μ m in C (applies to D, E).

Figure 13.

Schematic summary of the results of this study. To facilitate future investigations, all drawings show the CNS in one of the planar configurations shown in Fig. 1. Neurons are not drawn to scale. **A:** Localization of serotonin-like immunoreactive neurons (red fill) in the central ganglia of *Biomphalaria*. Drawings show the CNS with the pedal commissure severed and the pedal ganglia reflected in the lateral direction (see Fig. 1C, F). The labels "Dorsal" and "Ventral" refer to all ganglia except the pedals, which appear in reversed orientation. **B:** Serotonergic neurons labeled with backfills of various nerves. Neuron fill colors correspond to the nerves and nerve labels. Drawings show the CNS with the cerebral commissure severed and the cerebral ganglia reflected in the lateral direction (see Fig. 1B). The labels "Dorsal" and "Ventral" refer to all ganglia except the cerebrals, which appear in inverted orientation.