Survivin Inhibitor YM-155 Sensitizes Tumor Necrosis Factor–Related Apoptosis-Inducing Ligand-Resistant Glioma Cells to Apoptosis through Mcl-1 Downregulation and by Engaging the Mitochondrial Death Pathway

Daniel R. Premkumar, Esther P. Jane, Kimberly A. Foster, and Ian F. Pollack

Department of Neurosurgery (D.R.P., E.P.J., K.A.F., I.F.P.), University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania (D.R.P., E.P.J., I.F.P.), and University of Pittsburgh Cancer Institute Brain Tumor Program (I.F.P.), Pittsburgh, Pennsylvania

Received March 12, 2013; accepted May 28, 2013

ABSTRACT

Induction of apoptosis by the death ligand tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) is a promising antitumor therapy. However, not all tumor cells are sensitive to TRAIL, highlighting the need for strategies to overcome TRAIL resistance. Inhibitor of apoptosis family member survivin is constitutively activated in various cancers and blocks apoptotic signaling. Recently, we demonstrated that YM-155 [3-(2-methoxyethyl)-2-methyl-4,9-dioxo-1-(pyrazin-2-ylmethyl)-4,9-dihydro-3H-naphtho[2,3-d]imidazol-1-ium bromide], a small molecule inhibitor, downregulates not only survivin in gliomas but also myeloid cell leukemia sequence 1 (Mcl-1), and it upregulates proapoptotic Noxa levels. Because Mcl-1 and survivin are critical mediators of resistance to various anticancer therapies, we questioned whether YM-155 could sensitize resistant glioma cells to TRAIL. To address this hypothesis, we combined YM-155 with TRAIL and examined the effects on cell survival and apoptotic signaling. TRAIL or YM-155 individually

induced minimal killing in highly resistant U373 and LN308 cell lines, but combining TRAIL with YM-155 triggered a synergistic proapoptotic response, mediated through mitochondrial dysfunction via activation of caspases-8, -9, -7, -3, poly-ADP-ribose polymerase, and Bid. Apoptosis induced by combination treatments was blocked by caspase-8 and pan-caspase inhibitors. In addition, knockdown of Mcl-1 by RNA interference overcame apoptotic resistance to TRAIL. Conversely, silencing Noxa by RNA interference reduced the combined effects of YM-155 and TRAIL on apoptosis. Mechanistically, these findings indicate that YM-155 plays a role in counteracting glioma cell resistance to TRAIL-induced apoptosis by downregulating Mcl-1 and survivin and amplifying mitochondrial signaling through intrinsic and extrinsic apoptotic pathways. The significantly enhanced antitumor activity of the combination of YM-155 and TRAIL may have applications for therapy of malignant glioma.

Introduction

Human malignant gliomas are aggressive tumors that generally respond poorly to current therapy with surgery, radiation, and conventional chemotherapy (Pollack, 1994; DeAngelis, 2001; Wen and Kesari, 2008). Although molecularly targeted single-agent therapy holds the promise of more effective treatment in many cancers, it has failed to offer long-term survival benefit in malignant gliomas (Pollack et al.,

2007; Prados et al., 2009). Therefore, novel approaches are required, potentially taking into account multiagent combinations. Recently, we performed a small interfering RNA screen to identify the mechanisms underlying glioma cell survival. We determined that nuclear factor κ B (NF- κ B), proteasomal components, and Akt and Bcl2 family members (Thaker et al., 2009; Thaker et al., 2010a,b; Kitchens et al., 2011), when inhibited, enhanced cytotoxicity in glioma cells.

Tumor necrosis factor-related apoptosis inducing ligand (TRAIL) is a promising agent because of its tumor-specific induction of apoptosis. The receptors for TRAIL, DR4, and DR5 contain a cytoplasmic “death domain” capable of engaging the cell suicide apparatus through an adaptor molecule intermediate, such as Fas-associated death domain

This work was supported by National Institutes of Health [Grant P01NS40923] (to I.F.P.); by the Walter L. Copeland fund of the Pittsburgh Foundation (to D.R.P. and K.A.F.); and by a grant from Ian's Friends Foundation (to I.F.P.) in honor of Ian Yagoda.
dx.doi.org/10.1124/jpet.113.204743.

ABBREVIATIONS: AIF, apoptosis-inducing factor; CHAPS, 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonic acid; DiOC6, 3',3'-dihexyloxacarbo-cyanine iodide; DMSO, dimethylsulfoxide; E64d, (2S,3S)-trans-epoxysuccinyl-L-leucylamido-3-methylbutane ethyl ester; IAP, inhibitor of apoptosis protein; Mcl-1, myeloid cell leukemia sequence 1; MTS, 3-[4,5-dimethylthiazol-2-yl]-5-[3-carboxymethoxyphenyl]-2-[4-sulfophenyl]-2H, tetrazolium; NF- κ B, nuclear factor κ B; PAGE, polyacrylamide gel electrophoresis; PARP, poly-ADP-ribose polymerase; PBS, phosphate-buffered saline; PI, propidium iodide; shRNA, small hairpin RNA; TBS, Tris-buffered saline; TRAIL, tumor necrosis factor-related apoptosis-inducing ligand; YM-155, 3-(2-methoxyethyl)-2-methyl-4,9-dioxo-1-(pyrazin-2-ylmethyl)-4,9-dihydro-3H-naphtho[2,3-d]imidazol-1-ium bromide; z-DEVD-fmk, N-benzyloxycarbonyl-Asp-Glu-Val-Asp-fluoromethyl ketone; z-IETD-fmk, N-benzyloxycarbonyl-Ile-Glu-Thr-Asp-fluoromethyl ketone; z-LEHD-fmk, N-benzyloxycarbonyl-Leu-Glu-His-Asp-fluoromethyl ketone; z-VAD-fmk, N-benzyloxycarbonyl-Val-Ala-Asp-fluoromethyl ketone.

protein, promoting death-inducing signaling complex (DISC) assembly, activation of caspase-8, and further activation of caspase-3 (Gong and Almasan, 2000a,b). The TRAIL-mediated apoptotic process can be also activated by the mitochondrial pathway (Srivastava, 2001). In such cases, Bid is cleaved by caspase-8, the apoptosome is assembled, caspase-9 is activated, and it further activates effector caspase-3, leading to accumulation of cytochrome *c* in the cytosol. Furthermore, the active truncated form of Bid (tBid) triggers the mitochondrial activation of caspase-9 by inducing the homooligomerization and allosteric activation of Bak or Bax, two multidomain proapoptotic members of the Bcl-2 family (Wei et al., 2001).

Although most human cancer cell lines express death receptors for Apo2L/TRAIL, many remain resistant to TRAIL-induced death. Various mechanisms of resistance have been reported in different cancers (Wang et al., 2001; Russo et al., 2010; Yang et al., 2010). We, among others, have indicated that dysregulation of the NF- κ B, protein kinase C, Bcl-2, and Akt pathways may be integrally involved in mediating TRAIL resistance in glioma (Okhrimenko et al., 2005; Jane et al., 2011; Opel et al., 2011; Premkumar et al., 2012). In a prior study, we demonstrated that TRAIL sensitivity was not correlated with the relative expression of functional TRAIL receptors or decoy receptors; however, bortezomib (an NF- κ B/proteasomal inhibitor) sensitized TRAIL-resistant glioma cells (Jane et al., 2011).

Survivin is a member of the inhibitor of apoptosis protein (IAP) family (Salvesen and Duckett, 2002), which also may contribute to TRAIL resistance. In recent studies, we and others have shown that survivin is strongly expressed in pediatric and adult gliomas (Chakravarti et al., 2002, 2004; Okada et al., 2008). Survivin overexpression has been associated with cancer progression, multidrug resistance, poor prognosis, and short survival in several tumor types (Tanaka et al., 2000; Karam et al., 2007; Troeger et al., 2007; Lei et al., 2010; Reis et al., 2011). Survivin can inhibit apoptosis by blocking a common step downstream of mitochondrial cytochrome *c* release by inhibiting terminal effectors caspase-3 and caspase-7 and by interfering with caspase-9 activity and processing (Shin et al., 2001; Asanuma et al., 2005; Bilancio et al., 2006; Xia et al., 2006).

YM-155 is a novel survivin suppressant that is currently in clinical trials (Kummar et al., 2009; Satoh et al., 2009). We recently showed that YM-155 suppressed survivin expression, with little effect on the expression levels of other IAP family members and inhibited growth and viability of certain glioma cell lines, in addition to downregulating myeloid cell leukemia sequence 1 (Mcl-1) levels (Jane et al., 2013). Recent studies showed that upregulation of survivin by gene transfer enhanced resistance to TRAIL-induced apoptosis (Kim et al., 2011; Raviv et al., 2011), whereas transfection with survivin antisense enhanced sensitivity to TRAIL-induced apoptosis (Li et al., 2005; Azuhata et al., 2006). Because Mcl-1 is also a critical mediator of cellular resistance to various anticancer therapies, including suppression of TRAIL-induced cell death (Kobayashi et al., 2005; Ricci et al., 2007; Kim et al., 2008; Oh et al., 2012), we questioned whether YM-155 could sensitize resistant glioma cells to TRAIL, either by inhibition of survivin or Mcl-1 or both. In this study, we observed YM-155 sensitized glioma cells to TRAIL by promoting signaling through both the intrinsic and extrinsic

apoptotic pathways. Our results demonstrate that therapeutic agents that downregulate Mcl-1 or survivin may promote the efficacy of TRAIL in the clinical setting.

Materials and Methods

Cell Lines. The established malignant glioma cell lines U87, U373, LN229, A172, and T98G were obtained from the American Type Culture Collection (Manassas, VA). LN18, LN2428, and LN2308 were provided by Dr. Nicolas de Tribolet (Lausanne, Switzerland). Human astrocytes (HAs) and growth media were obtained from ScienCell Research Laboratories (Carlsbad, CA). Cell culture conditions of these cell lines were as previously described (Jane et al., 2011, 2013; Premkumar et al., 2012).

Reagents and Antibodies. Soluble human recombinant Super-KillerTRAIL (referred to as TRAIL in this article) was purchased from Enzo Biochemicals (Enzo Life Sciences, Farmingdale, NY). YM-155 was purchased from Chemie Tek (Indianapolis, IN). Caspase inhibitors (z-VAD-fmk, z-IETD-fmk, z-DEVD-fmk, and z-LEHD-fmk) were purchased from R&D Systems (Minneapolis, MN). The following antibodies were used: Mcl-1 (#4572), Bak (3814), Bax (#2774), Bid (#2002), cytochrome *c* (#4280), cleaved poly-ADP-ribose polymerase (PARP, #9546), cleaved caspase-3 (#9664), cleaved caspase-8 (#9496), cleaved caspase-9 (#9501), and β -actin (#4970) were from Cell Signaling Technology (Beverly, MA). Noxa (sc-26917) and apoptosis-inducing factor (AIF; sc-5586) were from Santa Cruz Biotechnology (Santa Cruz, CA). Monoclonal anti-Bax (#556467) and monoclonal anti-Bak (AM03) were from BD Pharmingen (San Diego, CA) and Calbiochem (San Diego, CA), respectively.

Cell Proliferation and Cytotoxicity Assay. Cells were seeded in 96-well plates (5000 cells/well) in 100 μ l of growth medium and incubated at 37°C for 24 hours before the addition of inhibitors or vehicle for 3 days. Cells were washed, and viable cells were quantitated using a colorimetric assay (CellTiter96 Aqueous Non-Radioactive Cell Proliferation Assay; Promega, Madison, WI) as described previously (Jane et al., 2013). The IC₅₀ value was calculated as the mean drug concentration required to inhibit cell proliferation by 50% compared with vehicle controls.

Annexin V Apoptosis Assay. Apoptosis was evaluated using a fluorescent annexin V/propidium iodide (PI) assay kit (Invitrogen, Carlsbad, CA) as described previously (Jane et al., 2011, 2013; Premkumar et al., 2012). Cells were treated with or without inhibitors for various intervals, harvested, and pelleted by centrifugation (1000 rpm for 5 minutes); washed in ice-cold phosphate-buffered saline (PBS); and resuspended in the annexin V- fluorescein isothiocyanate/PI reagent in the dark for 15 minutes before flow cytometric analysis using a FACSCalibur flow cytometer (BD Biosciences, San Jose, CA).

DiOC6 Labeling and Detection of Mitochondrial Membrane Depolarization. Mitochondrial membrane depolarization was measured as described previously (Jane et al., 2011, 2013; Premkumar et al., 2012). Nonadherent cells were collected, and attached cells were trypsinized and resuspended in phosphate-buffered saline (PBS). Cells were loaded with 50 nM 3',3'-dihexyloxacarbo-cyanine iodide (DiOC6; Invitrogen), which accumulates in intact mitochondria, at 37°C for 15 minutes. Cells were then spun at 3000g, rinsed with PBS, and resuspended. Fluorescence intensity was detected by flow cytometry and analyzed with CellQuest (BD Biosciences, San Jose, CA) and FlowJo (Tree Star, Inc., Ashland, OR) analysis software.

Immunoprecipitation and Western blotting Analysis. Cells were washed in ice-cold PBS and lysed in buffer containing 30 mM HEPES, 10% glycerol, 1% Triton X-100, 100 mM NaCl, 10 mM MgCl₂, 5 mM EDTA, 2 mM Na₃VO₄, 2 mM β -glycerophosphate, 1 mM phenylmethylsulfonyl fluoride, 1 mM 4-(2-aminoethyl) benzenesulfonyl fluoride, 0.8 μ M aprotinin, 50 μ M bestatin, 15 μ M E-64, 20 μ M leupeptin, and 10 μ M pepstatin A for 15 minutes on ice. Samples were centrifuged at 12,000g for 15 minutes, supernatants were isolated, and protein was quantified using protein assay reagent (Pierce

Chemical, Rockford, IL). Equal amounts of protein were separated by SDS-PAGE and electrotransferred onto a nylon membrane (Invitrogen). Nonspecific antibody binding was blocked by incubation of the membranes with 4% bovine serum albumin in Tris-buffered saline (TBS)/Tween 20 (0.1%). The membranes were incubated with primary antibody overnight at 4°C, washed in TBS/Tween 20, and incubated with a 1:2000 dilution of horseradish peroxidase-conjugated secondary antibody in TBS/Tween 20 at room temperature for 1 hour. Proteins were visualized by Western blot chemiluminescence reagent (Cell Signaling). Where indicated, the membranes were reprobed with antibodies against β -actin to ensure equal loading and transfer of proteins.

For immunoprecipitation, cell extracts were prepared by lysing 5×10^6 cells on ice for 30 minutes in CHAPS lysis buffer (10 mM HEPES, pH 7.4, 150 mM NaCl, 1% CHAPS, protease, phosphatase inhibitors). Lysates were clarified by centrifugation, and equal amounts of protein extracts were incubated overnight with primary antibody. Dynabeads Protein G (Invitrogen) was added for 2 hours, followed by magnetic separation of the immunoprecipitated fraction; Western blot analysis was conducted as outlined already herein. Scanning densitometry was performed using acquisition into Adobe Photoshop (Adobe Systems, Inc., San Jose, CA) followed by image analysis (UN-SCAN-IT gel, version 6.1; Silk Scientific, Orem, UT).

Subcellular Fractionation. Cells were treated with or without inhibitors, and cytosolic proteins were fractionated as described previously (Premkumar et al., 2012). Cells were resuspended in lysis buffer containing 0.025% digitonin, sucrose (250 mM), HEPES (20 mM, pH 7.4), magnesium chloride (5 mM), potassium chloride (10 mM), EDTA (1 mM), phenylmethylsulfonyl fluoride (1 mM), 10 μ g/ml aprotinin, and 10 μ g/ml leupeptin. After 10 minutes of incubation at 4°C, cells were centrifuged (2 minutes at 13,000g), and the supernatant (cytosolic fraction) was removed and frozen at -80°C for subsequent use.

Transient Transfection. Optimal 29mer-pRS-small hairpin (sh) RNA constructs were obtained from Origene (Rockville, MD). Sequences specific for human Mcl-1 (ACC TAG AAG GTG GCA TCA GGA ATG TGC TG), Noxa (GGA GGT GCT ACA CAA TGT GGC GTC GGC AC), and control sequences (GCA CTA CCA GAG CTA ACT CAG ATA GTA CT) (nontarget shRNA) were used for this study. Glioma cells were seeded in six-well plates and allowed to reach 70% confluence. Transfection of targeting or control shRNA was performed using FuGENE6 per the manufacturer's recommendations (Roche Applied Science, Indianapolis, IN). One microgram of Mcl-1 or Noxa or nontargeting shRNA in 100 μ l of Opti-MEM medium was mixed with 2 μ l of FuGENE6 and incubated at room temperature for 20 minutes, followed by the addition of complete medium to make the total volume up to 2 ml. After 48 hours, media were changed and cells were incubated with inhibitors or vehicle for 24 hours. Assessment of cell viability (annexin V binding), loss of mitochondrial membrane potential, or Western blot analysis was carried out as already described.

Statistical Analysis. Unless otherwise indicated, data are expressed as mean $\pm$ S.D. The significance of differences between experimental conditions was determined using a two-tailed Student's *t* test. Differences were considered significant at *P* values $\leq$ 0.05.

Results

Differential Apoptotic Responses of Glioma Cell Lines to TRAIL. Our recent study demonstrated a wide range of TRAIL sensitivity to apoptosis induction in glioma cell lines (Jane et al., 2011). As shown in Fig. 1A, annexin V/PI flow cytometry analysis clearly demonstrated that the percentage of apoptotic cells was increased to ~80% when LN18 and T98G (TRAIL-sensitive) cells were treated with TRAIL (with dramatic apoptotic responses to concentrations

as low as 5 ng/ml for 24 hours), whereas U373 and LN308 cells were resistant to TRAIL (~12% cell death at 50 ng/ml TRAIL). Western blot analyses [time-course (Fig. 1B) and dose-response (Fig. 1C) studies] clearly demonstrate activation of the caspase cascade with cleavage of the initiator caspase-8 and the main effector caspase-3 in LN18 and T98G (TRAIL-sensitive) cells. For example, after 6 hours of treatment (TRAIL, 10 ng/ml), the exogenous caspase-8 was activated by proteolytic cleavage as shown by the appearance of the cleaved intermediate (43-kDa) and the active caspase-8 (18-kDa) fragments. As a result of caspase-3 activity, cleavage of its substrate PARP was seen in TRAIL-sensitive cell lines (LN18 and T98G) but not in TRAIL-resistant LN308 cell line. The active form of caspase-9 (37 kDa) and the reduction of procaspase-10 were observed after TRAIL treatment in LN18 and T98G cell lines. The downstream caspases-7 and -3, as well as the proapoptotic Bcl-2 family member protein Bid, were also cleaved (Fig. 1D). However, no Bid cleavage was observed in LN308 and U373 (TRAIL-resistant) cell lines (Fig. 1D). As caspases have a key role in TRAIL-mediated apoptosis, we preincubated LN18 cells with caspase-specific inhibitors (z-DEVD-fmk, caspase-3 inhibitor; z-IETD-fmk, caspase-8 inhibitor, and z-VAD-fmk, pan-caspase inhibitor) and examined the effect by Western blot analysis. Our results indicate that caspase-8 inhibition suppressed TRAIL-mediated caspase-3 and -7 cleavage, suggesting that caspase-8 is necessary to engage the caspase effector cascade activation in response to TRAIL (Fig. 1E). The cleavage of Bid and caspase-9 induced by TRAIL and cell death protection by z-LEHD-fmk (caspase-9 inhibitor) suggests that the mitochondrial apoptotic pathway is also activated (data not shown). To confirm this, the release of cytochrome *c* and AIF was measured after TRAIL treatment. The results showed that the amount of cytochrome *c* and AIF in the cytosolic fraction was increased in LN18 and T98G (TRAIL-sensitive, Fig. 1F) but not in TRAIL-resistant LN308 (Fig. 1F) and U373 (data not shown) cell lines. Together, these results confirm that TRAIL induces the activation of the mitochondrial pathway in TRAIL-sensitive glioma cell lines.

YM-155 Synergizes with TRAIL to Induce Apoptosis in TRAIL-Resistant Glioma Cell Lines. In our recent studies, we demonstrated that the survivin inhibitor YM-155 exhibited significant activity against glioma cells (Jane et al., 2013). Because IAP family proteins block TRAIL-mediated apoptotic signaling through inhibition of caspase activity (Azuhata et al., 2001, 2006; Kim et al., 2004; Nakao et al., 2006; Gaiser et al., 2008), we hypothesized that combining YM-155 and TRAIL would potentiate apoptosis induction in TRAIL-resistant gliomas. To explore the possibility of sensitizing TRAIL-induced cytotoxicity in human glioma cells by YM-155, we treated TRAIL-resistant cells (LN308 and U373) with TRAIL in the presence or absence of YM-155 for 24 hours. Then cell death was assessed by annexin V/PI flow cytometry and Western blot analysis. Although TRAIL or YM-155 alone was minimally effective, cotreatment with YM-155 significantly enhanced the effect of TRAIL-induced apoptosis. As shown in Fig. 2A, the number of viable cells decreased to 92% with YM-155 (25 nM) and to 91% with TRAIL (25 ng/ml) in LN308. Likewise, limited effects were seen in U373. In contrast, when cells were cotreated with YM-155 and TRAIL, the number of viable cells reduced to 30% (Fig. 2A). Western blot analysis demonstrated that the combination of YM-155

and TRAIL potently induced activation of caspase-3, caspase-7, and PARP cleavage in both LNZ308 and U373 cell lines (Fig. 2B). The combination of YM-155 and TRAIL strongly induced caspase-3 processing with 17- and 15-kDa cleavage products and 89-kDa cleavage product of PARP in TRAIL-resistant glioma cells. Bid and caspase-8 were also activated (Fig. 2C). Incubation with caspase-9 inhibitor (z-LEHD-fmk) was able to salvage LNZ308 cells partially from YM-155 +

TRAIL-induced apoptosis (~50% protection from YM-155 + TRAIL-induced apoptosis, Fig. 2D). Chemical inhibitors of caspase-8 (z-IETD), caspase-3 (z-DEVD-fmk), and pan-caspase inhibitor (z-VAD-fmk) also significantly blocked the apoptotic activity of combination therapy with YM-155 and TRAIL (Fig. 2D), suggesting that both the intrinsic and extrinsic pathways play a role in cell death induced by these agents. Overall, these results indicate that YM-155 substantially

Fig. 2. YM-155 synergizes with TRAIL to induce apoptosis in TRAIL-resistant glioma cell lines. (A) TRAIL-resistant (LNZ308 and U373) cells were seeded at 60% confluence, allowed to attach overnight, and treated with indicated concentrations of YM-155 with or without TRAIL (25 ng/ml) for 24 hours. Control cells received an equivalent amount of DMSO (0). Apoptosis was analyzed by flow cytometry. The percentages of cells in each quadrant are indicated. Bar chart represents data from three independent experiments (** $P < 0.005$). (B) LNZ308 and U373 cells were treated with YM-155 (25 nM) or TRAIL (25 ng/ml) or the combination of both for 24 hours. Equal amounts of protein were separated by SDS-PAGE and subjected to Western blotting analysis with the indicated antibodies. (C) LNZ308 cells were pretreated with 25 μ M z-IETD (caspase-8 inhibitor), z-DEVD (caspase-3 inhibitor), or z-VAD-fmk (pan-caspase inhibitor) for 2 hours, followed by YM-155 (25 nM) + TRAIL (25 ng/ml) for 24 hours. Whole-cell extracts were prepared as described in *Materials and Methods*. Equal amounts of protein were separated by SDS-PAGE and subjected to Western blotting analysis with the indicated antibodies. (D) LNZ308 cells were pretreated with caspase inhibitors, as in C, followed by YM-155 (25 nM) + TRAIL (25 ng/ml) [Y + T] for 24 hours. Apoptosis was analyzed by flow cytometry. Data are representative of triplicate studies from three independent experiments (** $P < 0.005$). (E) Non-neoplastic human astrocytes (HAs) were seeded at 60% confluence, allowed to attach overnight, and treated with TRAIL (25 ng/ml) or YM-155 (25 nM) or the combination of both (T + Y) for 24 hours. Control cells received an equivalent amount of DMSO. Apoptosis was analyzed by flow cytometry. The percentages of cells in each quadrant are indicated (upper panel). In parallel, HAs were incubated with varying concentrations of TRAIL with or without YM-155 (25 nM). Control cells were treated with equivalent concentrations of vehicle (DMSO). After a 72-hour treatment, cell proliferation was assessed semiquantitatively by spectrophotometric measurement of MTS bioreduction. Points represent the mean of three measurements $\pm$ S.D. (lower panel).

increases the apoptotic potential of TRAIL-resistant glioma cells. We further addressed whether treatment with YM-155, TRAIL, or both would be toxic to normal human astrocytes. Annexin V/PI (Fig. 2E, upper panel) and cell proliferation (assessed by MTS cell proliferation assay; Fig. 2E, lower panel) assay clearly demonstrate that both YM-155 and TRAIL were not toxic to non-neoplastic astrocytes, suggesting that the combination of YM-155 and TRAIL may selectively kill malignant glioma cells.

YM-155 Enhances TRAIL-Mediated Signaling by Engaging the Mitochondrial Death Pathway. Having shown that YM-155-induced sensitization to TRAIL involved caspase activation, we next investigated whether the amplification of death signaling was a function of mitochondrial

outer membrane permeabilization. To analyze this hypothesis, we measured the disruption of the transmembrane mitochondrial potential ($\Delta\psi_m$) using the fluorescent dye DiOC6 in glioma cells after TRAIL plus YM-155 treatment. No reduction of the $\Delta\psi_m$ was observed with TRAIL-resistant LNZ308 cells treated with TRAIL (Fig. 3A) or YM-155 alone (Fig. 3B), whereas cotreatments with YM-155 substantially increased the number of cells with a reduced $\Delta\psi_m$ (Fig. 3C). A similar result was observed in U373 cell line (data not shown). The disruption of $\Delta\psi_m$ was largely prevented by the caspase-8 inhibitor (z-IETD-fmk), caspase-3 inhibitor (z-DEVD-fmk), and the pan-caspase inhibitor z-VAD-fmk in these cells (Fig. 3D). Western blot analysis of the kinetics of the release of

Fig. 3. YM-155 enhances TRAIL-mediated signaling by engaging the mitochondrial death pathway. LNZ308 cells were treated with indicated concentrations of TRAIL (A) YM-155 (B), or both (C) for 24 hours. Control cells received equivalent amounts of DMSO. The integrity of the mitochondrial membranes was examined by DiOC6 staining as described in *Materials and Methods*. The decrease in fluorescence intensity reflected the loss of $\Delta\psi_m$. Histogram (upper panel) and bar chart (lower panel) represent the mean number of $\Delta\psi_m$ cells acquired from three independent experiments. (** $P < 0.05$, compared with control). (D) LNZ308 cells were pretreated with 25 μ M z-IETD-fmk (caspase-8 inhibitor), z-DEVD-fmk (caspase-3 inhibitor), or z-VAD-fmk (pan-caspase inhibitor) for 2 hours followed by YM-155 (25 nM) + TRAIL (25 ng/ml) for 24 hours. Loss of mitochondrial membrane potential was determined by DiOC6 staining and flow cytometry. Histogram (left panel) and bar chart (right panel) represent the mean number of $\Delta\psi_m$ cells acquired from three independent experiments (** $P < 0.005$, compared with control). (E) TRAIL-resistant LNZ308 cells were incubated with YM-155 (25 nM) or TRAIL (25 ng/ml) or the combination of both for the indicated duration. Control cells received equal amounts of DMSO. Cytosolic extract was prepared as described in *Materials and Methods*. Equal amounts of protein were separated by SDS-PAGE and subjected to Western blotting analysis with the indicated antibodies.

mitochondrial amplification factors revealed that YM-155 + TRAIL treatment resulted in a significant time-dependent increase of cytosolic cytochrome *c* and AIF levels, whereas TRAIL or YM-155 alone had minimal effect on the cytosolic translocation of these proteins (Fig. 3E). Taken together, these results show that the disruption of the $\Delta\psi_m$ induced by TRAIL + YM-155 is highly dependent on caspase activation and loss of mitochondrial membrane potential is a crucial event driving initiator caspase activation and apoptosis.

Cotreatment of YM-155 and TRAIL Induces Bax and Bak Conformational Changes.

Because activation of Bax

or Bak is essential for the loss of mitochondrial membrane potential, given that cells lacking both proteins fail to undergo mitochondrial membrane outer membrane potential loss and apoptosis in response to diverse intrinsic stimuli (Wei et al., 2001), we examined the effect of TRAIL on Bax activation in glioma cells by analyzing activity-related conformational changes by immunoprecipitation followed by Western blot analysis with antibodies recognizing active Bax or Bak (monoclonal anti-bax, 6A7; Sigma-Aldrich, St. Louis, MO; or monoclonal anti-Bak, Ab-1; Calbiochem). LN18 and T98G (TRAIL-sensitive) cells exposed to TRAIL displayed a significant

Fig. 4. Cotreatment of YM-155 and TRAIL induces Bax and Bak conformational changes. (A) TRAIL-sensitive LN18 and T98G cells were treated with TRAIL (10 ng/ml) for indicated duration and lysed with 1% CHAPS buffer. An equal amount of protein (400 μ g) was immunoprecipitated (IP) with monoclonal anti-Bax (6A7; Sigma-Aldrich) antibody and then immunoblotted with polyclonal anti-Bax antibody (Cell Signaling Technology). (B) TRAIL-resistant LNZ308 and U373 cells were incubated with TRAIL (25 ng/ml) or YM-155 (25 nM) and subjected to IP and immunoblotting as in (A). (C) LNZ308 cells were treated with the indicated concentrations of TRAIL with or without YM-155 (25 nM) for 24 hours. Bak immunoprecipitates using monoclonal anti-Bak (Ab-1; Calbiochem) were subjected to Western blot (WB) analysis using polyclonal anti-Bak antibody (Cell Signaling Technology). Samples of whole-cell lysates (10% input) were subjected to Western blot analysis to monitor the total level of Bak protein.

increase in Bax conformational change (Fig. 4A). When TRAIL-resistant cells were exposed to TRAIL or YM-155, a minimal Bax and Bak conformational change was observed. In contrast, cells coexposed to YM-155 and TRAIL displayed a significant increase in Bax (Fig. 4B) and a modest increase in Bak (Fig. 4C) conformational change compared with cells treated with YM-155 or TRAIL alone, suggesting that activation of Bak, and particularly Bax, may be responsible for the marked induction of apoptosis in cells coexposed to YM-155 and TRAIL.

Downregulation of Mcl-1 and Noxa Activation Are Crucial Events in TRAIL-Induced Apoptosis. Recently, several groups have shown that Mcl-1 downregulation (Meng et al., 2007; Ricci et al., 2007; Rosato et al., 2007) or Noxa upregulation (Shankar et al., 2008; Naumann et al., 2011; Opel et al., 2011) by various means dramatically enhances TRAIL lethality in diverse human tumor cells. As we (Jane et al., 2013) observed, YM-155 synergistically enhanced ABT-737-induced apoptosis, at least in part by downregulating Mcl-1 expression; we then examined the functional relevance of Mcl-1 and Noxa on TRAIL-induced apoptosis. As shown in Fig. 5A, Western blot analysis demonstrated that YM-155 induced Noxa protein expression and caused downregulation of Mcl-1 in a dose-dependent manner, resulting in Mcl-1, the antiapoptotic protein that Noxa binds, being decreased in this cell line, and resulting in a net increase in Noxa relative to its

target. YM-155 had little effect on other IAP family member proteins (Fig. 5B). In addition, Bcl-2, Bim, and Bcl-xL expression were relatively unchanged (data not shown). Because Noxa exerts its proapoptotic effects through binding to the prosurvival protein Mcl-1, resulting in dissociation and activation of Bax and Bak, followed by mitochondrial outer membrane permeabilization and induction of apoptosis (Okumura et al., 2008), we examined the effect of YM-155 treatment on the interaction between Noxa and its high-affinity partner Mcl-1 (Chen et al., 2005). Immunoprecipitation followed by Western blot analysis revealed that an increased Mcl-1-Noxa complex was detected after YM-155 treatment (Fig. 5C). We also examined whether upregulation of Noxa can activate Bak by displacing it from Mcl-1. As shown in Fig. 5C, treatment with YM-155 released Bak from its interaction with Mcl-1. These observations suggest that induction of Noxa by YM-155 can sequester Mcl-1 and also release Bak from Mcl-1, which may contribute to the enhanced apoptotic effect of TRAIL in glioma cells.

To validate the hypothesis that Mcl-1 downregulation and Noxa induction are required for TRAIL-induced apoptosis, we performed RNA interference experiments to knock down the expression of these proteins. Cells were transfected with Noxa- or Mcl-1-specific or nontarget shRNA, as described in *Materials and Methods*. Whereas Mcl-1 shRNA mimicked the inhibitory effect on Mcl-1 expression obtained with YM-155, Noxa shRNA counteracted the Noxa induction obtained with this agent (Fig. 5D). To examine the role of Mcl-1 and Noxa on cell viability, transiently transfected cells were treated as indicated and apoptosis was detected by Annexin V/PI staining by flow cytometry. As shown in Fig. 5D, a decrease in Mcl-1 significantly increased TRAIL-induced apoptosis, whereas a decrease in Noxa levels resulted in reduced TRAIL cytotoxicity. These data show that modulation of Mcl-1 and Noxa by YM-155 could play a pivotal role in apoptotic susceptibility to TRAIL in glioma cell lines.

Discussion

TRAIL induces apoptosis of many cancer cell lines in vitro, and its tumoricidal activity has been confirmed in different animal models of human cancer. Although most human cancer cell lines express death receptors for TRAIL (e.g., DR4 and DR5), many remain resistant to TRAIL. Therefore, identifying the signals responsible for protecting tumor cells against TRAIL-induced apoptosis may have a major effect on using TRAIL therapeutically. Considering its role in cell proliferation and apoptosis, survivin is a promising therapeutic target in this regard. Inhibitors targeting this pathway may reduce cellular antiapoptotic activity by downregulating survivin expression, thereby increasing the efficacy of cotherapies. Recent studies showed that upregulation of survivin by gene transfer enhanced resistance to TRAIL-induced apoptosis (Kim et al., 2011; Raviv et al., 2011), whereas transfection with survivin antisense rendered resistant cells susceptible to TRAIL-induced apoptosis (Li et al., 2005; Azuhata et al., 2006). In addition, because it was shown that Mcl-1 downregulation could cooperate with TRAIL (Meng et al., 2007; Kim et al., 2008) and that its inhibition could be required for apoptosis induction and promoted by YM-155 treatment (Jane et al., 2013), we hypothesized that

Fig. 5. Downregulation of Mcl-1 and Noxa activation are crucial events in TRAIL-induced apoptosis. (A and B) Logarithmically growing glioma cells were treated with the indicated concentrations of YM-155 for 24 hours, and Western blot analysis was performed with the indicated antibodies. (C) LNZ308 and U373 cells were treated with indicated concentrations of YM-155 for 24 hours. An equal amount of protein (400 μ g) was immunoprecipitated (IP) with Mcl-1 antibody and subjected to Western blot analysis using the indicated antibodies. (D) LNZ308 cells were transfected with nontarget (NT) or Noxa or Mcl-1 shRNA. Forty-eight hours post-transfection, cells were treated with the indicated concentrations of TRAIL or YM-155 or the combination of both for 24 hours, and viability was assessed by annexin V/PI apoptosis assay (** $P < 0.005$ versus control). In parallel, cell lysates were collected, and protein was subjected to Western blot analysis using anti-Noxa or anti-Mcl-1 antibody. Immunoblots were stripped and reprobed with β -actin. Data are representative of triplicate studies from three independent experiments.

combining YM-155 and TRAIL could achieve synergistic cytotoxicity in TRAIL-resistant glioma cells.

In this study, we showed that YM-155 has such an action, suppressing survivin and Mcl-1 expression, and at clinically achievable concentrations plays a significant role in enhancing apoptosis in TRAIL-resistant glioma cells mediated by the loss of mitochondrial membrane polarization in a caspase-dependent manner. We observed both caspase-8 and caspase-9 activation with YM-155 and TRAIL, suggesting that both intrinsic and extrinsic caspase pathways are involved. Our data suggesting that caspase inhibition could block YM-155 and TRAIL-induced apoptosis are consistent with a process that may involve caspase pathway activation as a fundamental component of the response.

Interestingly, among prosurvival Bcl-2 family proteins, Mcl-1 is the only member that is downregulated by YM-155 (Jane et al., 2013). Mcl-1 plays a prominent role in the inhibition of apoptosis, mediating its effects primarily through interaction with proapoptotic members of the Bcl-2 family at the level of the mitochondria. Mcl-1 is a critical molecule for glioma cell survival (Rieger et al., 1998), making YM-155 a potentially useful therapeutic agent. Previously, we showed that Mcl-1 is a major determinant of ABT-737 toxicity and that YM-155-mediated downregulation of survivin and

Mcl-1 significantly enhanced ABT-737 efficacy (Jane et al., 2013). Regarding the mechanism, Tang et al. (2011) determined that treatment with YM-155 results in the downregulation of Mcl-1 at the transcription level. Prior reports that Bcl-2 and Mcl-1 exert inhibitory functions on TRAIL-induced apoptosis (Kobayashi et al., 2005; Ricci et al., 2007; Kim et al., 2008; Oh et al., 2012; Premkumar et al., 2012; Chanvorachote and Pongrakhananon, 2013) are consistent with our observation that Mcl-1 downregulation by YM-155 or by genetic manipulation (shRNA studies) plays a pivotal role in TRAIL-induced apoptosis. In parallel with the decrease in Mcl-1, we observed an increase in the BH-3-only protein Noxa, an antagonist of Mcl-1, suggesting that Noxa upregulation in response to YM-155 may serve to post-transcriptionally inactivate Mcl-1, causing displacement of proapoptotic proteins bound to the Mcl-1 protein. Because downregulation of Noxa (by genetic interference) inhibited YM-155 and TRAIL-induced apoptosis to some extent, a role of Noxa in YM-155 + TRAIL-mediated apoptosis can be assumed. This assumption is consistent with our previous reports that the antiglioma activity of bortezomib is the result of Noxa upregulation and Mcl-1 cleavage/degradation (Premkumar et al., 2013).

Cross-talk between the extrinsic and intrinsic pathways has been previously reported to be mediated by activated Bid

(tBid), a BH3-only proapoptotic BCL-2 protein, observed to initiate cytochrome *c* release from mitochondria, thereby initiating intrinsic apoptosis. Likewise, proapoptotic tBid activates the death receptor pathway initiator caspase-8, displacing Bax and Bak and allowing them to oligomerize and induce mitochondrial cytochrome *c* release, an important early step in mitochondrial-mediated apoptosis (Tait and Green, 2010). Our study demonstrated that cotreatment with YM-155 and TRAIL induces Bid activation, cytochrome *c* release, and mitochondrial membrane dysfunction. Inhibiting caspase activity with caspase inhibitors blocks mitochondrial depolarization seen with YM-155 and TRAIL and resultant cell death. Furthermore, Mcl-1 downregulation by YM-155 may elicit apoptosis via Bak released from Mcl-1/Bak complexes. We also demonstrated that Bak and Bax, major proapoptotic effectors, undergo conformational changes to mediate mitochondrial outer membrane permeabilization. Bak and Bax conformational change is known to follow caspase-8 activation and is accompanied by pore formation in the outer mitochondrial membrane and the release of cytochrome *c* from mitochondria (Tait and Green, 2010). Together, Bak and Bax conformational change, activation of Bid, and loss of mitochondrial membrane potential may provide a mechanistic explanation for the enhanced cell death induced by YM-155 in TRAIL-resistant glioma cell lines. In summary, combined treatment with YM-155 and TRAIL might be a promising experimental therapy because YM-155 sensitizes glioma cells to TRAIL-induced apoptosis via various mechanisms, thereby minimizing the risk of acquired tumor cell resistance.

Acknowledgments

The authors thank Joe D. DiDomenico, Robert Lacomy, and Alexis Styche for technical assistance.

Authorship Contributions

Participated in research design: Premkumar, Pollack.

Conducted experiments: Jane, Premkumar, Foster.

Performed data analysis: Jane, Premkumar, Foster, Pollack.

Wrote or contributed to the writing of the manuscript: Premkumar, Pollack.

References

- Asanuma H, Torigoe T, Kamiguchi K, Hirohashi Y, Ohmura T, Hirata K, Sato M, and Sato N (2005) Survivin expression is regulated by coexpression of human epidermal growth factor receptor 2 and epidermal growth factor receptor via phosphatidylinositol 3-kinase/AKT signaling pathway in breast cancer cells. *Cancer Res* **65**:11018–11025.
- Azuhata T, Scott D, Griffith TS, Miller M, and Sandler AD (2006) Survivin inhibits apoptosis induced by TRAIL, and the ratio between survivin and TRAIL receptors is predictive of recurrent disease in neuroblastoma. *J Pediatr Surg* **41**:1431–1440.
- Azuhata T, Scott D, Takamizawa S, Wen J, Davidoff A, Fukuzawa M, and Sandler A (2001) The inhibitor of apoptosis protein survivin is associated with high-risk behavior of neuroblastoma. *J Pediatr Surg* **36**:1785–1791.
- Bilancio A, Okkenhaug K, Camps M, Emery JL, Ruckle T, Rommel C, and Vanhaesebroeck B (2006) Key role of the p110delta isoform of PI3K in B-cell antigen and IL-4 receptor signaling: comparative analysis of genetic and pharmacologic interference with p110delta function in B cells. *Blood* **107**:642–650.
- Chakravarti A, Noll E, Black PM, Finkelstein DF, Finkelstein DM, Dyson NJ, and Loeffler JS (2002) Quantitatively determined survivin expression levels are of prognostic value in human gliomas. *J Clin Oncol* **20**:1063–1068.
- Chakravarti A, Zhai GG, Zhang M, Malhotra R, Latham DE, Delaney MA, Robe P, Nestler U, Song Q, and Loeffler J (2004) Survivin enhances radiation resistance in primary human glioblastoma cells via caspase-independent mechanisms. *Oncogene* **23**:7494–7506.
- Chanvorachote P and Pongrakhananon V (2013) Ouabain downregulates Mcl-1 and sensitizes lung cancer cells to TRAIL-induced apoptosis. *Am J Physiol Cell Physiol* **304**:C263–C272.
- Chen L, Willis SN, Wei A, Smith BJ, Fletcher JI, Hinds MG, Colman PM, Day CL, Adams JM, and Huang DC (2005) Differential targeting of prosurvival Bcl-2 proteins by their BH3-only ligands allows complementary apoptotic function. *Mol Cell* **17**:393–403.
- DeAngelis LM (2001) Brain tumors. *N Engl J Med* **344**:114–123.
- Gaiser T, Becker MR, Habel A, Reuss DE, Ehemann V, Rami A, and Siegelin MD (2008) TRAIL-mediated apoptosis in malignant glioma cells is augmented by celastrol through proteasomal degradation of survivin. *Neurosci Lett* **442**:109–113.
- Gong B and Almasan A (2000a) Apo2 ligand/TNF-related apoptosis-inducing ligand and death receptor 5 mediate the apoptotic signaling induced by ionizing radiation in leukemic cells. *Cancer Res* **60**:5754–5760.
- Gong B and Almasan A (2000b) Genomic organization and transcriptional regulation of human Apo2/TRAIL gene. *Biochem Biophys Res Commun* **278**:747–752.
- Jane EP, Premkumar DR, DiDomenico JD, Hu B, Cheng SY, and Pollack IF (2013) YM-155 potentiates the effect of ABT-737 in malignant human glioma cells via survivin and Mcl-1 downregulation in an EGFR-dependent context. *Mol Cancer Ther* **12**:326–338.
- Jane EP, Premkumar DR, and Pollack IF (2011) Bortezomib sensitizes malignant human glioma cells to TRAIL, mediated by inhibition of the NF-kappaB signaling pathway. *Mol Cancer Ther* **10**:198–208.
- Karam JA, Lotan Y, Ashfaq R, Sagalowsky AI, and Shariat SF (2007) Survivin expression in patients with non-muscle-invasive urothelial cell carcinoma of the bladder. *Urology* **70**:482–486.
- Kim S, Kang J, Qiao J, Thomas RP, Evers BM, and Chung DH (2004) Phosphatidylinositol 3-kinase inhibition down-regulates survivin and facilitates TRAIL-mediated apoptosis in neuroblastomas. *J Pediatr Surg* **39**:516–521.
- Kim SH, Ricci MS, and El-Deiry WS (2008) Mcl-1: a gateway to TRAIL sensitization. *Cancer Res* **68**:2062–2064.
- Kim YS, Jin HO, Seo SK, Woo SH, Choe TB, An S, Hong SI, Lee SJ, Lee KH, and Park IC (2011) Sorafenib induces apoptotic cell death in human non-small cell lung cancer cells by down-regulating mammalian target of rapamycin (mTOR)-dependent survivin expression. *Biochem Pharmacol* **82**:216–226.
- Kitchens CA, McDonald PR, Shun TY, Pollack IF, and Lazo JS (2011) Identification of chemosensitivity nodes for vinblastine through small interfering RNA high-throughput screens. *J Pharmacol Exp Ther* **339**:851–858.
- Kobayashi S, Werneburg NW, Bronk SF, Kaufmann SH, and Gores GJ (2005) Interleukin-6 contributes to Mcl-1 up-regulation and TRAIL resistance via an Akt-signaling pathway in cholangiocarcinoma cells. *Gastroenterology* **128**:2054–2065.
- Kummar S, Doroshow JH, Tomaszewski JE, Calvert AH, Lobbezoo M, and Giaccone G; Task Force on Methodology for the Development of Innovative Cancer Therapies (MDICT) (2009) Phase 0 clinical trials: recommendations from the Task Force on Methodology for the Development of Innovative Cancer Therapies. *Eur J Cancer* **45**:741–746.
- Lei Y, Geng Z, Guo-Jun W, He W, and Jian-Lin Y (2010) Prognostic significance of survivin expression in renal cell cancer and its correlation with radioresistance. *Mol Cell Biochem* **344**:23–31.
- Li H, Niederkorn JY, Neelam S, and Alizadeh H (2005) Resistance and susceptibility of human uveal melanoma cells to TRAIL-induced apoptosis. *Arch Ophthalmol* **123**:654–661.
- Meng XW, Lee SH, Dai H, Loegering D, Yu C, Flatten K, Schneider P, Dai NT, Kumar SK, and Smith BD, et al. (2007) Mcl-1 as a buffer for proapoptotic Bcl-2 family members during TRAIL-induced apoptosis: a mechanistic basis for sorafenib (Bay 43-9006)-induced TRAIL sensitization. *J Biol Chem* **282**:29831–29846.
- Nakao K, Hamasaki K, Ichikawa T, Arima K, Eguchi K, and Ishii N (2006) Survivin downregulation by siRNA sensitizes human hepatoma cells to TRAIL-induced apoptosis. *Oncol Rep* **16**:389–392.
- Naumann I, Kappler R, von Schweinitz D, Debatin KM, and Fulda S (2011) Bortezomib primes neuroblastoma cells for TRAIL-induced apoptosis by linking the death receptor to the mitochondrial pathway. *Clin Cancer Res* **17**:3204–3218.
- Oh B, Park S, Pak JH, and Kim I (2012) Downregulation of Mcl-1 by daunorubicin pretreatment reverses resistance of breast cancer cells to TNF-related apoptosis-inducing ligand. *Biochem Biophys Res Commun* **422**:42–47.
- Okada H, Low KL, Kohanbash G, McDonald HA, Hamilton RL, and Pollack IF (2008) Expression of glioma-associated antigens in pediatric brain stem and non-brain stem gliomas. *J Neurooncol* **88**:245–250.
- Okhrimenko H, Lu W, Xiang C, Hamburger N, Kazimirsy G, and Brodie C (2005) Protein kinase C-epsilon regulates the apoptosis and survival of glioma cells. *Cancer Res* **65**:7301–7309.
- Okumura K, Huang S, and Sinicrope FA (2008) Induction of Noxa sensitizes human colorectal cancer cells expressing Mcl-1 to the small-molecule Bcl-2/Bcl-xL inhibitor, ABT-737. *Clin Cancer Res* **14**:8132–8142.
- Opel D, Naumann I, Schneider M, Bertele D, Debatin KM, and Fulda S (2011) Targeting aberrant PI3K/Akt activation by PI103 restores sensitivity to TRAIL-induced apoptosis in neuroblastoma. *Clin Cancer Res* **17**:3233–3247.
- Pollack IF (1994) Brain tumors in children. *N Engl J Med* **331**:1500–1507.
- Pollack IF, Jakacki RI, Blaney SM, Hancock ML, Kieran MW, Phillips P, Kun LE, Friedman H, Packer R, and Banerjee A, et al. (2007) Phase I trial of imatinib in children with newly diagnosed brainstem and recurrent malignant gliomas: a Pediatric Brain Tumor Consortium report. *Neuro-oncol* **9**:145–160.
- Prados MD, Chang SM, Butowski N, DeBoer R, Parvataneni R, Carliner H, Kabuubi P, Ayers-Ringler J, Rabbitt J, and Page M, et al. (2009) Phase II study of erlotinib plus temozolomide during and after radiation therapy in patients with newly diagnosed glioblastoma multiforme or gliosarcoma. *J Clin Oncol* **27**:579–584.
- Premkumar DR, Jane EP, Agostino NR, DiDomenico JD, and Pollack IF (2013) Bortezomib-induced sensitization of malignant human glioma cells to vorinostat-induced apoptosis depends on reactive oxygen species production, mitochondrial dysfunction, Noxa upregulation, Mcl-1 cleavage, and DNA damage. *Mol Carcinog* **52**:118–133.
- Premkumar DR, Jane EP, DiDomenico JD, Vukmer NA, Agostino NR, and Pollack IF (2012) ABT-737 synergizes with bortezomib to induce apoptosis, mediated by Bid cleavage, Bax activation, and mitochondrial dysfunction in an Akt-dependent context in malignant human glioma cell lines. *J Pharmacol Exp Ther* **341**:859–872.
- Raviv Z, Zilberberg A, Cohen S, Reischer-Pelech D, Horrix C, Berger MR, Rosin-Arbesfeld R, and Flescher E (2011) Methyl jasmonate down-regulates survivin

- expression and sensitizes colon carcinoma cells towards TRAIL-induced cytotoxicity. *Br J Pharmacol* **164**:1433–1444.
- Reis FR, Vasconcelos FC, Pereira DL, Moellman-Coelho A, Silva KL, and Maia RC (2011) Survivin and P-glycoprotein are associated and highly expressed in late phase chronic myeloid leukemia. *Oncol Rep* **26**:471–478.
- Ricci MS, Kim SH, Ogi K, Plastaras JP, Ling J, Wang W, Jin Z, Liu YY, Dicker DT, and Chiao PJ, et al. (2007) Reduction of TRAIL-induced Mcl-1 and cIAP2 by c-Myc or sorafenib sensitizes resistant human cancer cells to TRAIL-induced death. *Cancer Cell* **12**:66–80.
- Rieger L, Weller M, Bornemann A, Schabet M, Dichgans J, and Meyermann R (1998) BCL-2 family protein expression in human malignant glioma: a clinical-pathological correlative study. *J Neurol Sci* **155**:68–75.
- Rosato RR, Almenara JA, Coe S, and Grant S (2007) The multikinase inhibitor sorafenib potentiates TRAIL lethality in human leukemia cells in association with Mcl-1 and cFLIPL down-regulation. *Cancer Res* **67**:9490–9500.
- Russo M, Mupo A, Spagnuolo C, and Russo GL (2010) Exploring death receptor pathways as selective targets in cancer therapy. *Biochem Pharmacol* **80**: 674–682.
- Salvesen GS and Duckett CS (2002) IAP proteins: blocking the road to death's door. *Nat Rev Mol Cell Biol* **3**:401–410.
- Satoh T, Okamoto I, Miyazaki M, Morinaga R, Tsuya A, Hasegawa Y, Terashima M, Ueda S, Fukuoka M, and Ariyoshi Y, et al. (2009) Phase I study of YM155, a novel survivin suppressant, in patients with advanced solid tumors. *Clin Cancer Res* **15**: 3872–3880.
- Shankar S, Ganapathy S, and Srivastava RK (2008) Sulforaphane enhances the therapeutic potential of TRAIL in prostate cancer orthotopic model through regulation of apoptosis, metastasis, and angiogenesis. *Clin Cancer Res* **14**: 6855–6866.
- Shin S, Sung BJ, Cho YS, Kim HJ, Ha NC, Hwang JI, Chung CW, Jung YK, and Oh BH (2001) An anti-apoptotic protein human survivin is a direct inhibitor of caspase-3 and -7. *Biochemistry* **40**:1117–1123.
- Srivastava RK (2001) TRAIL/Apo-2L: mechanisms and clinical applications in cancer. *Neoplasia* **3**:535–546.
- Tait SW and Green DR (2010) Mitochondria and cell death: outer membrane permeabilization and beyond. *Nat Rev Mol Cell Biol* **11**:621–632.
- Tanaka K, Iwamoto S, Gon G, Nohara T, Iwamoto M, and Tanigawa N (2000) Expression of survivin and its relationship to loss of apoptosis in breast carcinomas. *Clin Cancer Res* **6**:127–134.
- Tang H, Shao H, Yu C, and Hou J (2011) Mcl-1 downregulation by YM155 contributes to its synergistic anti-tumor activities with ABT-263. *Biochem Pharmacol* **82**: 1066–1072.
- Thaker NG, McDonald PR, Zhang F, Kitchens CA, Shun TY, Pollack IF, and Lazo JS (2010a) Designing, optimizing, and implementing high-throughput siRNA genomic screening with glioma cells for the discovery of survival genes and novel drug targets. *J Neurosci Methods* **185**:204–212.
- Thaker NG, Zhang F, McDonald PR, Shun TY, Lazo JS, and Pollack IF (2010b) Functional genomic analysis of glioblastoma multiforme through short interfering RNA screening: a paradigm for therapeutic development. *Neurosurg Focus* **28**:E4.
- Thaker NG, Zhang F, McDonald PR, Shun TY, Lewen MD, Pollack IF, and Lazo JS (2009) Identification of survival genes in human glioblastoma cells by small interfering RNA screening. *Mol Pharmacol* **76**:1246–1255.
- Troeger A, Siepermann M, Escherich G, Meisel R, Willers R, Gudowius S, Moritz T, Laws HJ, Hanenberg H, and Goebel U, et al. (2007) Survivin and its prognostic significance in pediatric acute B-cell precursor lymphoblastic leukemia. *Haematologica* **92**:1043–1050.
- Wang GQ, Gastman BR, Wieckowski E, Goldstein LA, Rabinovitz A, Yin XM, and Rabinowich H (2001) Apoptosis-resistant mitochondria in T cells selected for resistance to Fas signaling. *J Biol Chem* **276**:3610–3619.
- Wei MC, Zong WX, Cheng EH, Lindsten T, Panoutsakopoulou V, Ross AJ, Roth KA, MacGregor GR, Thompson CB, and Korsmeyer SJ (2001) Proapoptotic BAX and BAK: a requisite gateway to mitochondrial dysfunction and death. *Science* **292**:727–730.
- Wen PY and Kesari S (2008) Malignant gliomas in adults. *N Engl J Med* **359**: 492–507.
- Xia W, Bisi J, Strum J, Liu L, Carrick K, Graham KM, Treece AL, Hardwicke MA, Dush M, and Liao Q, et al. (2006) Regulation of survivin by ErbB2 signaling: therapeutic implications for ErbB2-overexpressing breast cancers. *Cancer Res* **66**: 1640–1647.
- Yang A, Wilson NS, and Ashkenazi A (2010) Proapoptotic DR4 and DR5 signaling in cancer cells: toward clinical translation. *Curr Opin Cell Biol* **22**:837–844.

Address correspondence to: Dr. Ian F. Pollack, Department of Neurosurgery, Children's Hospital of Pittsburgh, 4401 Penn Avenue, Pittsburgh, PA 15224. E-mail: ian.pollack@chp.edu