From the clinic to the lab (and back)—a call for laboratory research to optimize cognitive behavioural treatment of pain

S Van Damme,¹ D J Moore²

¹Department of Experimental-Clinical and Health Psychology, Ghent University, Henri Dunantlaan 2, 9000 Ghent, Belgium

²Centre for Pain Research, School for Health, The University of Bath, Bath, UK

Correspondence to: S Van Damme
stefaan.vandamme@ugent.be

Cite this as: *TBM* 2012;2:102–105
doi: 10.1007/s13142-011-0083-6

ABSTRACT

Cognitive behavioural therapy (CBT) is the dominant psychological treatment for chronic pain. CBT covers a mixture of aims and a broad range of techniques, including attentional control, modification of maladaptive beliefs and coping strategies and flexible goal setting and acceptance. Patients are exposed to varying selections of these strategies, and a major challenge for clinical practice is to determine what works for whom in which context. We propose that incorporating laboratory research into translational behaviour medicine is a critical developmental step that will help optimizing CBT and provide examples of representative experimental research programs.

KEYWORDS

Pain, Psychology, Cognitive behavioural therapy, Experimental research

INTRODUCTION

The biopsychosocial model is widely accepted as the most heuristic approach to the understanding and treatment of chronic pain [6]. Within this approach, cognitive behavioural therapy (CBT) is the dominant psychological treatment. Despite its popularity, a systematic review of randomised controlled trials (RCTs) showed that effects on pain and disability are rather modest [5]. The authors proposed several possible explanations, such as large variations in dose and quality and lack of focus in content and identified treatment outcomes. Particularly striking is the relative lack of specificity of CBT, covering a mixture of aims and a broad range of techniques, including attentional control, modification of maladaptive beliefs and coping strategies and acceptance [23]. Patients are exposed to varying selections of these strategies, and a major challenge for clinical practice is to determine what works for whom in which context.

We propose that incorporating laboratory research into translational behaviour medicine is a critical developmental step that will help fine-tuning and tailoring CBT. Rather than presenting an exhaustive review of contemporary views on chronic

Implications

This paper informs researchers about new developments in research on psychological factors in the experience of pain. Practitioners and policy-makers are informed about how laboratory research is an important step in optimizing cognitive behavioural therapy for chronic pain.

pain, we describe a number of areas within the psychology of pain, argue how laboratory research can help optimizing CBT in each of these areas and provide examples of representative experimental research programs. As an organizing framework, we use the areas previously identified by Morley [22] as targets for CBT: interruption, interference and identity.

INTERRUPTION

Temporarily, interruption of attention by pain can be seen as a beneficial process to warn of potential danger and promote analgesic behaviour [3]. However, this can be very disruptive to daily activities such as work and leisure behaviour. There is a wealth of behavioural, neuropsychological and neurophysiological evidence showing that cognitive functioning is impaired by the presence of chronic pain [7, 21]. Less is known about the underlying mechanisms of pain-related cognitive interruption and the conditions moderating its magnitude. Fundamental research using experimental pain models would allow a more detailed exploration of these issues. Although such research exists, studies have typically focused on simple interruptive effects [3, 15]. It is, however, acknowledged within cognitive psychology that attention has a wide range of separable functions [17]. More precise investigation of which components of the cognitive system are affected by pain may prove valuable for clinical practice. For instance, advances have been made in the development of a standardised test protocol allowing the specific assessment of cognitive dysfunctions [20]. This may be particularly helpful in the evaluation of interventions aimed at reducing the interruptive function of pain.

In addition, there is also a history of research examining the effects of attentional distraction on pain [30], and attentional strategies to manage pain are often part of CBT [24]. The effectiveness of distraction techniques is, however, less clear than originally thought. Experimental research is particularly helpful in identifying when, how and for whom distraction works. For instance, it is well-known that it is more difficult to direct attention away from pain when it is interpreted as threatening, rendering distraction less effective in those who are characterized by a high level of pain catastrophizing [30]. Furthermore, recent advances in experimental distraction research may help further refining and optimizing the use of distraction in clinical practice. One example is a research examining the role of executive functioning abilities in distraction effectiveness. Results from the first studies are promising and suggest that specifically inhibition capability may play an important role in the attentional control of pain [26, 34]. Another example concerns the role of motivational factors. Contemporary theories propose that attention is prioritized to information that is relevant for valuable goals, and that attentional control over pain may be optimized when the distraction has high personal relevance and is related to a valuable personal goal [30]. A recent study experimentally manipulating the motivational value of the distraction task provides preliminary evidence for this idea [33].

INTERFERENCE

Continued interruption leads to interference, meaning failure to complete tasks effectively or task performance at an unsatisfactory level [22]. The extent to which pain interferes with the accomplishment of daily life tasks, often called disability, is the main reason of patients to consult health care providers and is associated with high costs when pain becomes chronic [6]. Patients' beliefs about their pain have been argued to play an essential role in maintaining disability. Specifically, the fear-avoidance (FA) model states that a confrontational style, in which individuals with acute pain gradually resume activities despite pain, leads to recovery, whereas catastrophic beliefs about pain, i.e. interpretation of pain as a sign of serious injury or pathology, lead to excessive fear of pain/injury that gradually extends to avoidance of activities that are presumed to worsen the pain [35]. The FA model has been empirically validated by both clinical and experimental findings [14, 28] and has become one of the most popular biopsychosocial models in pain. Part of this success lies in its conceptual clarity and natural flow from diagnostic information to clinical interventions such as exposure therapy [2].

However, not all chronic pain patients seem to display the typical fear-avoidance pattern [36], and one subgroup even reports task persistence and suppression of pain-related thoughts [10]. It has also

been suggested that fear avoidance may not be a stable behaviour pattern and may be influenced by the motivational context [2]. Specifically, avoiding pain is often only one goal in a dynamic environment with concomitant, competing, goals [11, 29, 37], and the decision to avoid or persist pain-evoking activities may depend on the value of other activated goals [2]. This may have implications for the applicability of exposure as a clinical technique.

An experimental research program testing the boundaries of the fear-avoidance model by systematically investigating determinants of avoidance and persistence behaviour may be an important step in further refining treatment. For example, recent studies have experimentally manipulated goals during pain-provoking tasks and found that the pursuit of a valuable goal reduces avoidance behaviour [12, 31]. More systematic experimental research embedded within a dynamic motivational view on disability, taking into account the goals and values of patients, may help further optimizing the efficacy of therapeutic approaches aimed at increasing functional ability of patients, such as exposure [19].

IDENTITY

Repeated interference with tasks that are essential to achieve valued personal goals affects patients' sense of self [22]. It is therefore likely that patients often perceive pain relief as a necessary step to allow successful pursuit of life goals [2]. In patients with chronic pain, this search for pain relief often becomes the dominant goal in life. Paradoxically, this increase in effort may contribute to the maintenance in suffering. Persistent attempts to control or solve the pain problem when actual control is low and the problem is chronic, often called "misdirected problem solving", might become part of the problem and bring along negative consequences [4]. In such situation, accepting pain may be a more adaptive response, and an extensive body of research has demonstrated that acceptance is associated with better mental and physical well-being in patients with chronic pain [27]. However, note that research about acceptance is dominated by correlational studies in which no conclusions can be drawn about its causal effects.

Of clinical interest are treatments aimed at acceptance and restoration of patients' capacity to live according to their life values despite the presence of pain. Examples of acceptance-based treatments are acceptance and commitment therapy [18, 39], mindfulness-based cognitive therapy [38, 40] and contextual cognitive behavioural therapy [19]. A recent meta-analysis [32] suggests that although acceptance-based interventions are promising, many treatment outcome studies lack (adequate) control groups, and RCTs are small and as yet unconvincing.

In sum, it is still unclear if and by which mechanisms patient functioning is exactly affected

by acceptance or by lack of acceptance, i.e. perseveration in the search for pain relief. Therefore, experimental research to the precise consequences of acceptance-based versus control-based coping strategies is necessary. Recent studies in which participants' goal was to obtain control over a painful stimulus that was actually uncontrollable showed negative consequences such as increase in fear and attentional biases [1, 25]. Other studies have experimentally manipulated acceptance of laboratory-induced pain, and showed that, compared to suppression-based strategies, acceptance resulted in higher pain tolerance and less pain-related distress [8, 16].

CONCLUSION

Reflecting on the evidence reviewed in this essay provides important considerations for the translation of basic scientific research to clinical pain management. Recent theoretical advances suggest a number of pain-cognition relationships to be less universal than originally thought. This has important implications for the utilisation of CBT, and careful examination of pain-cognition interactions under laboratory conditions is essential to allow further optimizing CBT. Individual tailoring of CBT by means of evidence-based selection criteria may be a step in the right direction [9]. This does not necessarily mean that we should abandon group-based interventions. One promising way of adding individual accents while still preserving the benefits of group programs is the implementation of e-health innovations allowing both assessment of patient features requiring specific treatment techniques and delivery of such interventions [13]. To conclude, we recommend that in order to maximize effectiveness of CBT, more use should be made of the advantages of clinically informed experimental research.

- Crombez G, Eccleston C, De Vlieger P, Van Damme S, De Clercq A. Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain. *Pain*. 2008; 137:631-639.
- Crombez G, Eccleston C, Van Damme S, Vlaeyen J, W. S., & Karoly, P. (2011). The fear-avoidance model of chronic pain: the next generation. *Clinical Journal of Pain*, (in press).
- Eccleston C, Crombez G. Pain demands attention: a cognitive affective model of the interruptive function of pain. *Psychol Bull*. 1999; 123:356-366.
- Eccleston C, Crombez G. Worry and chronic pain: a misdirected problem solving model. *Pain*. 2007; 132:233-236.
- Eccleston C, Williams ACD, Morley S. Psychological therapies for the management of chronic pain (excluding headache) in adults. *Cochrane Database Syst Rev*. 2009; 2:CD007407.
- Gatchel RJ, Peng YB, Peters ML, Fuchs PN, Turk DC. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychol Bull*. 2007; 133:581-624.
- Glass JM. Cognitive dysfunction in fibromyalgia syndrome. *Journal of Musculoskeletal Pain*. 2010; 18:367-372.
- Gutierrez O, Luciano C, Rodriguez M, Fink BC. Comparison between an acceptance-based and a cognitive-control-based protocol for coping with pain. *Behav Ther*. 2004; 35:767-783.
- Haldorsen EMH, Grasdahl AL, Skouen JS, Risa AE, Kronholm K, Ursin H. Is there a right treatment for a particular patient group? Comparison of ordinary treatment, light multidisciplinary treatment, and extensive multidisciplinary treatment for long-term sick-listed employees with musculoskeletal pain. *Pain*. 2002; 95:49-63.
- Hasenbring MI, Verbunt JA. Fear-avoidance and endurance-related responses to pain: new models of behavior and their consequences for clinical practice. *Clin J Pain*. 2010; 26:747-753.
- Karoly P, Okun MA, Ruehlman LS, Pugliese JA. The impact of goal cognition and pain severity on disability and depression in adults with chronic pain: an examination of direct effects and mediated effects via pain-induced fear. *Cognitive Therapy and Research*. 2008; 32:418-433.
- Karsdorp PA, Nijst SE, Goossens MEJB, Vlaeyen JWS. The role of current mood and stop rules on physical task performance: an experimental investigation in patients with work-related upper extremity pain. *Eur J Pain*. 2010; 14:434-440.
- Keogh E, Rosser BA, Eccleston C. E-Health and chronic pain management: current status and developments. *Pain*. 2010; 151:18-21.
- Leeuw M, Goossens MEJB, Linton SJ, Crombez G, Boersma K, Vlaeyen JWS. The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med*. 2007; 30:77-94.
- Legrain V, Van Damme S, Eccleston C, Davis KD, Seminowicz DA, Crombez G. A neurocognitive model of attention to pain: behavioral and neuroimaging evidence. *Pain*. 2009; 144:230-232.
- Masedo AI, Esteve MR. Effects of suppression, acceptance and spontaneous coping on pain tolerance, pain intensity and distress. *Behav Res Ther*. 2007; 45:199-2009.
- Mathias JL, Wheaton P. Changes in attention and information-processing speed following severe traumatic brain injury: a meta-analytic review. *Neuropsychology*. 2007; 21:212-223.
- McCracken LM, Guttierrez-Martinez O. Processes of change in psychological flexibility in an interdisciplinary group-based treatment for chronic pain based on acceptance and commitment therapy. *Behav Res Ther*. 2011; 49:267-274.
- McCracken LM, Yang S. The role of values in a contextual cognitive-behavioral approach to chronic pain. *Pain*. 2006; 123:137-145.
- Moore DJ, Keogh E, Eccleston C. (2011). The interruptive effect of pain on attention. *Quarterly Journal of Experimental Psychology*, (in press).
- Moriarty O, McGuire BE, Finn DP. The effect of pain on cognitive function: a review of clinical and preclinical research. *Prog Neurobiol*. 2011; 93:385-404.
- Morley S. Psychology of pain. *Br J Anaesth*. 2008; 101:25-31.
- Morley S. Efficacy and effectiveness of cognitive behavior therapy for chronic pain: progress and some challenges. *Pain*. 2011; 152:S99-S106.
- Morley S, Shapiro DA, Biggs J. Developing a treatment manual for attention management in chronic pain. *Cogn Behav Ther*. 2004; 33:1-11.
- Notebaert L, Crombez G, Vogt J, De Houwer J, Van Damme S, Theeuwes J. Attempts to control pain prioritize attention towards signals of pain: an experimental study. *Pain*. 2011; 152:1068-1073.
- Oosterman JM, Dijkerman HC, Kessels RPC, Scherder EJA. A unique association between cognitive inhibition and pain sensitivity in healthy participants. *Eur J Pain*. 2010; 14:1046-1050.
- Thompson M, McCracken LM. Acceptance and related processes in adjustment to chronic pain. *Curr Pain Headache Rep*. 2011; 15:144-151.
- Turk DC, Wilson HD. Fear of pain as a prognostic factor in chronic pain: conceptual models, assessment, and treatment implications. *Curr Pain Headache Rep*. 2010; 14:88-95.
- Van Damme S, Crombez G, Eccleston C. Coping with pain: a motivational perspective. *Pain*. 2008; 139:1-4.
- Van Damme S, Legrain V, Vogt J, Crombez G. Keeping pain in mind: a motivational account of attention to pain. *Neurosci Biobehav Rev*. 2010; 34:204-213.
- Van Damme S, Van Ryckeghem D, Wyffels F, Van Hulle L, & Crombez G. (2011) No gain without pain: pursuing a competing goal inhibits avoidance behaviour. *Pain*, (in press).
- Veehof MM, Oskam M, Schreurs KMG, Bohlmeijer ET. Acceptance-based interventions for the treatment of chronic pain: a systematic review and meta-analysis. *Pain*. 2011; 152:533-542.
- Verhoeven K, Crombez G, Eccleston C, Van Ryckeghem DML, Morley S, Van Damme S. The role of motivation in distracting attention away from pain: an experimental study. *Pain*. 2010; 149:229-234.
- Verhoeven K, Van Damme S, Eccleston C, Van Ryckeghem D, Legrain V, Crombez G. Distraction from pain and executive functioning: an experimental investigation of inhibition, attentional switching and working memory. *Eur J Pain*. 2011; 15:866-873.

35. Vlaeyen JWS, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain*. 2000; 85:317-332.
36. Vlaeyen JWS, Morley S. Active despite pain: the putative role of stop-rules and current mood. *Pain*. 2004; 110:512-516.
37. Vlaeyen JWS, Morley S. Cognitive and behavioural factors in fibromyalgia: mood, goals, and task performance. *Journal of Musculoskeletal Pain*. 2009; 17:295-301.
38. Vowles KE, Wetherell JL, Sorrell JT. Targeting acceptance, mindfulness, and values-based action in chronic pain: findings of two preliminary trials of an outpatient group-based intervention. *Cognitive and Behavioral Practice*. 2009; 16:49-58.
39. Wicksell RK, Olsson GL, Haeyes SC. Psychological flexibility as a mediator of improvement in acceptance and commitment therapy for patients with chronic pain following whiplash. *Eur J Pain*. 2010; 14:1059.e1-1059.e11.
40. Zautra AJ, Davis MC, Reich JW, et al. Comparison of cognitive behavioural and mindfulness meditation interventions on adaptation to rheumatoid arthritis for patients with and without history of recurrent depression. *J Consult Clin Psychol*. 2008; 76:408-421.