Laryngeal Paraganglioma: A Rare Clinical Entity Managed by Supraselective Embolization and Lateral Pharygotomy

Sudhir M. Naik · Ashok M. Shenoy ·
Purshottam Chavan · Akkamahadevi Patil ·
Sumit Gupta

Received: 25 July 2012 / Accepted: 28 September 2012 / Published online: 6 October 2012
© Association of Otolaryngologists of India 2012

Abstract Laryngeal paragangliomas are benign slow growing tumors with symptoms resembling squamous cell carcinoma. Hoarseness or dysphasia are the commonest presenting symptoms and usually it presents as an submucosal mass on laryngoscopy. Ninety percentage of these tumors occur in the supraglottic larynx and the rest in the glottis and the subglottic region. Functional activity is seen in a few (2.9 %), none are associated with paraneoplastic syndromes. A 35-year-old male presented to us with hoarseness of voice since 4 months duration. Contrast arteriography demonstrated that the left superior thyroid artery supplied >80 % of the blood supply to the laryngeal mass. Supraselective embolization was done from the right femorals under local anaesthesia and sedation which was uneventful. The tumor was excised from lateral pharyngotomy approach with an partial laryngectomy procedure. Microscopy and immune-histochemistry confirmed it to be paraganglioma. Complete surgical resection or partial laryngectomy with meticulous dissection of surrounding tissues and preservation of neurovascular structures gives an excellent prognosis as far as oncological clearance is concerned. Malignant paragangliomas of the larynx are rare and an major meta-analysis is necessary to provide a true biological behaviour of this tumor.

Keywords Paraganglioma · Embolization · Lateral pharygotomy · Digital subtraction angiography

Introduction

Head and neck paragangliomas are rare tumors arising from the neural crest [1, 2]. They are highly vascularized tumors involving the walls of the blood vessels or specific nerves [3–5]. In the larynx they arise from superior or inferior laryngeal paraganglia and are seen in the supraglottis in 90 % of the cases [6]. Since these tumors are from neural crest-derived cells of the parasympathetic nervous system, they usually appear adjacent to nerve structures, most commonly the superior laryngeal nerve or the recurrent laryngeal nerve [7].

The paragangliomas of the larynx show a female preponderance compared to other neuroendocrine tumors [6]. Other neuroendocrine tumors differ in their biologic behavior, management, and prognosis compared to paragangliomas and so immunohistochemical differentiation forms an important criteria [8]. Paragangliomas of larynx are benign submucosal lesion and the management includes a conservative resection [6]. Significant intraoperative blood loss is expected so preoperative transarterial embolization has proved beneficial [9, 10]. Percutaneous direct intratumoral embolization can also be done with same efficacy [11–14]. Supraglottic laryngectomy and lateral pharyngotomy are commonly used approaches for managing these paragangliomas [15–20].

Case Report

A 35-year-old male presented to us with hoarseness of voice since 4 months duration. No history of difficulty in swallowing and any other symptoms. Also on examination a swelling in left side of the neck was seen confirmed on palpation. Indirect laryngoscopic examination revealed a

S. M. Naik (✉) · A. M. Shenoy · P. Chavan · S. Gupta
Department of Head and Neck Oncosurgery, KMIO, Bangalore, India
e-mail: sud223@gmail.com

A. Patil
Department of Pathology, KMIO, Bangalore, India

congested globular flashy mass arising from the left pharyngo-epiglottic fold, extending to the aryepiglottic fold and to the apex of the pyriform fossa with mobile vocal cords.

The patient was a non-smoker and non-drinker also complained of foreign body sensation. Contrast enhanced CT scan was done and an enhancing lesion was seen in the same region with radiological features suggesting of a vascular lesion or a paraganglioma (Figs. 1, 2, 3). Translaryngeal fine needle aspiration cytology revealed cells similar to that of paraganglioma, but a vascular malformation and schwannomas were not ruled out. His blood investigations and urine catecholamine levels were normal. Magnetic resonance angiography revealed a 3–4 cm vascular lesion in the left supraglottic region but the feeder

vessels were not specifically identified (Fig. 4). Chest X-ray was normal and lateral soft tissue neck showed an compromised airway (Fig. 5). So a digital subtraction angiography done revealed the superior thyroid artery to be the feeder and a minor contribution from contralateral superior thyroid was seen.

Digital subtraction angiography demonstrated that the left superior thyroid artery supplied greater than 80 % of the blood supply to the laryngeal mass (Fig. 6). Supraselective embolization was done from the right femorals under ocal anaesthesia and sedation which was uneventfull. Left superior thyroid artery was selectively catheterised using the 4F vertebral glide catheter over the 0.035 Terumo guide wire and embolized using 510–750 sized polyvinyl

Fig. 1 Tumor seen in the coronal and sagittal planes in the supraglottis

Fig. 2 Enhanced lesion seen in the left supraglottis showing a proliferative vascular lesion

Fig. 3 Axial lesions showing the extension of the tumor

Fig. 4 Contrast MRI showing the feeder vessels of the vascular tumor

Fig. 5 A normal chest X-ray and lateral soft tissue shadows showing the airway and the supraglottic shadows

Fig. 6 Digital subtraction angiography showing tumor blush before embolization

alcohol (PVA) particles. A minimal tumor blush was noted after this and the feeders were embolized using a microguide wire with 250–500 PVA particles (Fig. 7). Finally after embolization partial tumor blush was seen with feeders from the proximal part of the left superior thyroid artery. Since these feeders were close to origin of

external carotid artery no attempt was made to block these feeders. There was no other neurological deficit noticed after embolization.

Micro-laryngoscopic evaluation was done in 48 h and the pulsatile vascular lesion was elective posted for partial laryngectomy by lateral pharyngotomy (Fig. 8). Under

Fig. 7 The tumor shadow after embolization

Fig. 8 Micro laryngoscopic view of the tumor with high vascularity

general anaesthesia and endotracheal intubation the larynx was approached by lateral pharyngotomy. The left thyroid cartilage ala was dissected and left superior thyroid artery was ligated and cut along with the superior laryngeal nerve. The subglottis was normal, as was the base of the tongue and pharynx (Fig. 9).

Later the thyroid cartilage was dissected and the tumor excised. The defect closed with pharyngeal constrictors. As this was a partial laryngectomy procedure an elective tracheostomy was done to secure the airway and reduce aspiration and the endotracheal tube extubated. The post operative period uneventful. The temporary tracheostoma

Fig. 9 The tumor being excised by lateral pharyngotomy approach

Fig. 10 Post operative pictures with the temporary tracheostomy

was decannulated and strap closed after a week with swallowing rehabilitation (Fig. 10).

Histopathology revealed features suggestive of paraganglioma and all the lymph nodes had reactive changes. Microscopic examination showed polygonal epithelioid cells arranged in discrete nests (Zellballen pattern) (Fig. 11). Immunohistochemistry results showed the neoplastic cells were cytoplasmic positivity for chromogranin, synaptophysin and sustentacular positivity for S-100 but negative for cytokeratin and epithelial membrane antigen (EMA). So all these features were consistent with paraganglioma. Endothelial

lined vascular channels were showing CD 34 positivity enhancing the cell nest pattern (Figs. 12, 13, 14, 15, 16).

Discussion

Laryngeal paraganglioma was first described in 1955 and since then 85 cases were described in world literature [8]. Ferlito et al. [2, 8, 16] analysed all the reported paragangliomas of the larynx and found 62 case reports genuinely met the criteria for diagnosis.

Fig. 11 Zellballen nests of tumor cells with thin supporting sustentacular cells around each nest in low and high power fields

Fig. 12 The tumor showing cytokeratin negative and the overlying squamous epithelium showing positivity

These are benign slow growing tumors with symptoms resembling squamous cell carcinoma [8]. Hoarseness or dysphasia are the commonest presenting symptoms and usually it presents as an submucosal mass on laryngoscopy [8]. Rarely stridor, dysphagia, foreign body sensation and

Fig. 13 Immunohistochemistry showing cytoplasmic and nuclear S100 reactivity in sustentacular cells

Fig. 14 Tumor showing cytoplasmic positivity for chromogranin

dyspnea are seen and depends on the size and location of the tumor [16]. The majority of these tumors arise in the supraglottic larynx, and 2 % of these are malignant [2, 8, 16].

A histological diagnosis is necessary prior to surgical treatment and repetitive fine needle aspiration biopsies are advised if they are inconclusive [5]. Superficial biopsies are inconclusive as they have intact mucosa, but deeper biopsies can complicate as these lesions are highly vascular [21]. Translaryngeal biopsies are helpful in these conditions as done in our case.

Laryngeal paragangliomas have the same appearance as paragangliomas of other sites on histopathology [22]. They are highly vascular and composed of chief cells and sustentacular cells, arranged in a characteristic alveolar or Zellballen pattern. [22] The chief cells (type I cells,

Fig. 15 Tumor showing cytoplasmic positivity for synaptophysin

Fig. 16 Endothelial lined vascular channels showing CD 34 positive picture

epithelioid cells) are more numerous and contain catecholamine bound neuro-secretory granules and the sustentacular cells (type II cells, supporting cells) are devoid of neurosecretory granules and are characteristically located at the periphery of Zellballen [2]. Zellballen pattern are also seen in typical and atypical carcinoid, malignant melanoma and medullary carcinoma of thyroid and are not specific for paragangliomas [22].

Differentiating laryngeal paragangliomas from carcinoids, atypical carcinoids and other neuroendocrine carcinoma on microscopic examination is not impossible and immunohistochemistry is helpful [22]. The chief cells are immunoreactive to neuroendocrine markers chromogranin, synaptophysin, and neuron-specific enolase and negative for epithelial markers including cytokeratin [1, 20].

Atypical carcinoids of the larynx show immunoreactivity for both neuroendocrine markers and epithelial markers [6].

The sustentacular cells are immunoreactive to S-100, but chief cells are negative for melanoma markers S-100, HMB45, and melan-A, excluding this from melanoma [23]. Non neuroendocrine tumors are excluded by chromogranin positivity and the absence of keratin positivity and presence of S-100 protein-positive sustentacular cells tends to exclude carcinomas [24]. These tumors can be confused on microscopy as carcinoid tumor, malignant melanoma, medullary thyroid carcinoma, and small-cell neuroendocrine carcinoma [24].

S-100 testing is particularly necessary to establish the correct diagnosis as its negative in typical and atypical carcinoid and small-cell neuroendocrine carcinoma [25]. Cytokeratin, epithelial membrane antigen, carcino-embryonic antigen are negative in paragangliomas while positive in carcinoids and small-cell neuroendocrine carcinoma [25]. Chromogranin, synaptophysin, neuron-specific enolase are positive in all the laryngeal neuroendocrine tumors [25]. Galanin, glial fibrillary acidic protein are very specific for laryngeal paragangliomas [25].

The mutations in the SHDB, SDHC, and SHDD genes reflect higher risk for developing malignant paragangliomas whereas VHL and NF1 genes mutation reflect inherited paragangliomas [26, 27]. Rare cases of laryngeal paraganglioma associated with multicentric head and neck paraganglioma syndrome are also reported [28].

Ninety percentage of these tumors occur in the supraglottic larynx and the rest in the glottis and the subglottic region. Functional activity is seen in a few (2.9 %), none are associated with paraneoplastic syndromes [1, 19]. They spread by vocal fold fixation and rarely associated with necrosis and vascular invasion [1, 19]. Some herniate through the thyrohyoid membrane and present as a neck mass [29]. If imaging provide sufficient evidence of probable diagnosis of paraganglioma (or any other vascular tumor), then a biopsy can be omitted and direct surgery following angiography and embolization seems better [5]. Some tumors may be large enough to herniate through the thyrohyoid membrane and present as a neck mass [11].

Contrast-enhanced computerised tomography (CECT neck) and, contrast enhanced magnetic resonance imaging (MRI) rules out less enhancing laryngeal masses [15]. MRI is also useful in detecting lesions that are smaller than 5 mm and to screen for synchronous paragangliomas [30, 31]. Contrast enhanced MRI angiography (MRA) does not increase the diagnostic value of MRI but defines the vascularity of the tumor [32].

Digital subtraction angiography (DSA) is more sensitive in detecting vascularization compared to MRA [10, 33]. Also tumors larger than 2.5 cm, four vessel angiography and

preoperative embolization is useful in minimizing bleeding and injury to surrounding neurovascular structures [10, 33]. Digital subtraction angiography also allows the determination of carotid artery invasion and the performance of balloon occlusion studies to assess collateral cerebral circulation in carotid body paragangliomas [10, 33].

Usually superior thyroid artery is the primary vessel that supplies these tumors, so embolization creates a bloodless surgical field [34]. Preoperative embolization done by superselective catheterization of the supplying branches and transarterial embolization with particulate agents dramatically improves the operative field [35].

Sonography guided or CT guided intratumoral injection of a liquid adhesive like *N*-butyl-2-cyanoacrylate gives better devascularization compared to transarterial embolization and is a safe and effective method of preoperative embolization [11–14]. Percutaneous direct embolization provides an easy access to hypervascular tumor with a higher chance of achieving complete devascularization [12, 36]. Direct embolization is typically carried out with *n*-butyl cyanoacrylate (NBCA) [12, 36]. Percutaneous tumor embolization is easy and safe but complications of glue migration into the intracranial circulation is a fatal complication [12, 36]. Casasco et al. [12, 36] advocated the use of non-detachable balloons to protect the intracranial branches to reduce the risk of glue reflux into the circulation. No other complications are reported in the procedure [12, 36].

Most of the laryngeal paragangliomas are managed surgically and organ preservation partial laryngectomy procedures are becoming the standard protocol [1, 6]. Radical surgeries are not indicated as regional and distant metastasis are rarely documented [1, 6]. Transoral resections are not advised because hemostasis management can prove fatal, so lateral pharyngotomy is preferred depending on the extent and site of the tumor [1, 6]. The endoscopic approach provides limited exposure and access to the lesion [6]. This is not a good approach for vascular tumors, because hemorrhage might obscure visualization of tissue planes [6]. The biological behaviour of the tumor is not extensively defined as the tumor is rare and the malignant potential hardly exist [37]. All these lesions are treated by surgical resection which includes supraglottic or total laryngectomy and endoscopic resection [18, 37]. Inferior laryngeal paragangliomas present in the subglottis having dyspnea and shortness of breath and need careful consideration for airway management pre-operatively [18, 37]. Most of them were managed by total laryngectomy as the primary treatment [18, 37].

Lateral pharyngotomy approach is described for removal of supraglottic submucosal tumors, including paraganglioma [17]. All procedures are aimed at complete tumor removal with minimum functional deficits [17]. This approach can be performed without a tracheostomy for

smaller lesions, but for resection of vascular masses, a tracheotomy should be performed [6]. Also conservation laryngeal approaches are not attempted when cartilage involvement is seen radiographically [18, 37]. Submucosal resection with meticulous dissection results in no postoperative dysphagia and excellent voice preservation [6].

Laryngeal paragangliomas are benign tumors and presence of vascular, capsular or perineural invasion, focal necrosis, mitoses does not indicate metastatic potential [38, 39]. Malignant paragangliomas are reported in 2 % in some series while pathologic findings cannot predict clinical and biological behavior of the tumor [1, 2, 34]. Rare cases of distant metastasis is reported while neck nodes spread are rarely reported and so selective neck node clearance is not warranted [1, 2]. Regular followup is advised as no much literature on the biological behaviour of the tumor is known [38, 39]. A recurrence rates of 17 % is reported in superior laryngeal paragangliomas and 4 % in inferior laryngeal paraganglioma with careful follow-up should be given to the inferior laryngeal cases due to the potential for rapid airway obstruction with recurrent or persistent disease [18, 37]. Local recurrence may be a sign of malignancy [18, 37].

No role of radiotherapy or chemotherapy is used as a laryngeal preservation approach [5]. In cervical paragangliomas external beam radiation provides approximately 90 % tumor control rate [5].

Ferlito et al. [1, 2, 34] suggested the existence of other atypical carcinoid tumors as high as 25 % which were earlier thought as metastasis. They reviewed 65 cases over 35 years and found the mean age to be 44 years, mean duration of symptoms to be 26 months and most of the tumors were excised surgically [1, 2, 34]. They reported no role of radiation or chemotherapy [5].

Conclusion

Early diagnosis of these laryngeal paragangliomas are difficult as they present with nonspecific clinical manifestation. Complete surgical resection or partial laryngectomy with meticulous dissection of surrounding tissues and preservation of neurovascular structures gives an excellent prognosis as far as oncological clearance is concerned. Malignant paragangliomas of the larynx are rare and an major meta-analysis is necessary to provide a true biological behavior of this tumor.

References

1. Ferlito A, Barnes L, Rinaldo A, Gnepp DR, Milroy CM (1998) A review of neuroendocrine neoplasms of the larynx: update on diagnosis and treatment. *J Laryngol Otol* 112:827–834

2. Barnes L (1991) Paraganglioma of the larynx: a critical review of the literature. *ORL J Otorhinolaryngol Relat Spec* 53:220–234
3. Erickson D, Kudva YC, Ebersold MJ et al (2001) Benign paragangliomas: clinical presentation and treatment outcomes in 236 patients. *J Clin Endocrinol Metab* 86:5210–5216
4. Bishop GB Jr, Urist MM, EL Gammal T et al (1992) Paragangliomas of the neck. *Arch Surg* 127:1441–1445
5. Smolarz JR, Hanna EY, Williams MD, Kupferman ME (2010) Paraganglioma of the endolarynx: a rare tumor in an uncommon location: *Head Neck. Oncology* 2:2
6. Del Gaudio JM, Muller S (2004) Diagnosis and treatment of supraglottic laryngeal paraganglioma: report of a case. *Head Neck* 26:94–98
7. Batsakis JG, el-Naggar AK, Luna MA (1992) Neuroendocrine tumors of larynx. *Ann Otol Rhinol Laryngol* 101(8):710–714
8. Ferlito A, Barnes L, Wenig BM (1994) Identification, classification, treatment, and prognosis of laryngeal paraganglioma. Review of the literature and eight new cases. *Ann Otol Rhinol Laryngol* 103:525–536
9. Persky MS, Setton A, Niimi Y et al (2002) Combined endovascular and surgical treatment of head and neck paragangliomas: a team approach. *Head Neck* 24:423–431
10. Tikkakaoski T, Luotonen J, Leinonen S et al (1997) Preoperative embolization in the management of neck paragangliomas. *Laryngoscope* 107:821–826
11. Pierot L, Boulin A, Castings L et al (1994) Embolization by direct puncture of hypervascularised otorhinolaryngology tumours. *Ann Otolaryngol Chir Cervicofac* 111:403–409
12. Casasco A, Herbreteau D, Houdart E et al (1994) Devascularisation of craniofacial tumours by percutaneous tumour puncture. *Am J Neuroradiol* 15:1233–1239
13. Chloupka JC, Mangla S, Huddle DC et al (1999) Evolving experience with direct puncture therapeutic embolization for adjunctive and palliative management of head and neck hypervascular neoplasms. *Laryngoscope* 109:1864–1872
14. Abud DG, Mounayer C, Mounayer C, Benndorf G et al (2004) Intratumoral injection of cyanoacrylate glue in head and neck paragangliomas. *Am J Neuroradiol* 25:1457–1462
15. Sanders KW, Abreo F, Rivera E, Stucker FJ, Nathan CO (2001) A diagnostic and therapeutic approach to paragangliomas of the larynx. *Arch Otolaryngol Head Neck Surg* 127:565–569
16. Myssiorek D, Madnani D, Delacure MD (2001) The external approach for submucosal lesions of the larynx. *Otolaryngol Head Neck Surg* 125:370–373
17. Rubin JS, Silver CE (1992) Surgical approach to submucosal lesions of the supraglottic larynx: the supero-lateral thyrotomy. *J Laryngol Otol* 106:416–419
18. Myssiorek D, Halaas Y, Silver C (2001) Laryngeal and sinonasal paragangliomas. *Otolaryngol Clin North Am* 34:971–982
19. Peterson KL, Fu YS, Calcaterra T (1997) Subglottic paraganglioma. *Head Neck* 19:54–56
20. Wasserman PG, Savargaonkar P (2001) Paragangliomas: classification, pathology, and differential diagnosis. *Otolaryngol Clin North Am* 34:845–862
21. Lustrin ES, Palestro C, Vaheesan K (2001) Radiographic evaluation and assessment of paragangliomas. *Otolaryngol Clin North Am* 34(5):881–906
22. Dhakhwa R, Toran KC, Kafle S, Lakhey M (2012) Paraganglioma of the larynx. *J Pathol Nepal* 2:245–247
23. Martinez-Madrigal F, Bosq J, Micheau C, Nivet P, Luboinski B (1991) Paragangliomas of the head and neck. Immunohistochemical analysis of 16 cases in comparison with neuro-endocrine carcinomas. *Path Res Pract* 187:814–823
24. Khademi B, Taghizadegan L, Gandomi B, Vasei M (2004) Paraganglioma of the larynx: a case report. *IJMS* 29:47–49
25. Ferlito A, Silver CE, Bradford CR, Rinaldo A (2009) Neuroendocrine neoplasms of the larynx: an overview. *Head Neck* 31:1634–1646
26. Jimenez C, Cote G, Arnold A, Gagel RF (2006) Review: should patients with apparently sporadic pheochromocytomas or paragangliomas be screened for hereditary syndromes? *J Clin Endocrinol Metab* 91(8):2851–2858
27. Neumann HP, Pawlu C, Peczkowska M (2004) Distinct clinical features of paraganglioma syndromes associated with SDHB and SDHD gene mutations. *JAMA* 292(8):943–951
28. Rubin AD, Cheng SS, Bradford CR (2005) Laryngeal paraganglioma in a patient with multiple head and neck paragangliomas. *Otolaryngol Head Neck Surg* 132(3):520–522
29. Blanchard CL, Saunders WH (1955) Chemodectoma of the larynx: case report. *AMA Arch Otolaryngol* 61(4):472–474
30. Vogl T, Bruning R, Schedel H, Kang K, Grevers G, Hahn D, Lissner J (1989) Paragangliomas of the jugular bulb and carotid body: MR imaging with short sequences and Gd-DPTA enhancement. *Am J Roentgenol* 153:583–587
31. van Gils APG, van der Mey AGL, Hoogma RP, Sankuijl LA, Maaswinkel-Mooy PD, Falke TH, Pauwels EK (1992) MRI screening of kindred at risk of developing paragangliomas: support for genomic imprinting in hereditary glomus tumours. *Br J Cancer* 65:903–907
32. Vogl TJ, Juergens M, Balzer JO, Mack MG, Bergman C, Grevers G, Lissner J, Felix R (1994) Glomus tumors of the skull base: combined use of MR angiography and spin-echo imaging. *Radiology* 192:103–110
33. van den Berg R, Wasser MN, van Gils AP, van der Mey AG, Hermans J, van Buchem MA (2000) Vascularization of head and neck paragangliomas: comparison of three angiographic techniques with digital subtraction angiography. *Am J Neuroradiol* 21:162–170
34. Myssiorek D, Rinaldo A, Barnes L, Ferlito A (2004) Laryngeal paraganglioma: an updated critical review. *Acta Otolaryngol* 124(9):995–999
35. Rao V, Raghavendra KS, Chavan P, Shenoy AM (2011) Vocal cord paralysis following percutaneous embolization of a vagal paraganglioma. *Int J Head Neck Surg* 2(1):65–66
36. George B, Casasco A, Deffrennes D, Houdart E (1994) Intratumoral embolization of intracranial and extracranial tumors: technical note. *Neurosurgery* 35:771–773
37. Hinojar AG, Prieto JR, Munoz E, Hinojar AA (2002) Relapsing paraganglioma of the inferior laryngeal paraganglion: case report and review of the literature. *Head Neck* 24(1):95–102
38. Ferlito A, Milory CM, Wenig BM et al (1995) Laryngeal paraganglioma versus atypical carcinoid tumor. *Ann Otol Rhinol Laryngol* 104:78–83
39. Kimura N, Chetty R, Capella C et al (2004) Extra-adrenal paraganglioma: carotid body, jugulotympanic, vagal, laryngeal, aortic-pulmonary. In: DeLellis RA, Lloyd RV, Heitz PU, Eng C (eds) *Pathology and genetics. Tumors of endocrine organs*. IARC Press, Lyon, p 16