Published in final edited form as:

Biol Psychiatry. 2013 October 15; 74(8): 591–598. doi:10.1016/j.biopsych.2013.02.025.

White Matter Alterations at 33-Year Follow-Up in Adults with Childhood Attention-Deficit/Hyperactivity Disorder

Samuele Cortese, M.D., Ph.D.^{1,2,3}, Davide Imperati, Ph.D.¹, Juan Zhou, Ph.D.^{1,4}, Erika Proal, Ph.D.^{1,5}, Rachel G. Klein, Ph.D.⁶, Salvatore Mannuzza, Ph.D.⁶, Maria A. Ramos-Olazagasti, Ph.D.¹, Michael P. Milham, M.D., Ph.D.^{7,8}, Clare Kelly, Ph.D.¹, and F. Xavier Castellanos, M.D.^{1,7}

¹Phyllis Green and Randolph Cowen Institute for Pediatric Neuroscience, Child Study Center, NYU Langone School of Medicine, New York, NY, USA

²Child Neuropsychiatry Unit, G. B. Rossi Hospital, Department of Life Science and Reproduction, Verona University, Verona, Italy

³UMR_S INSERM U 930, CNRS ERL 3106, François-Rabelais University, Child Psychiatry Centre, University Hospital, Tours, France

⁴Duke-NUS Graduate Medical School, Neuroscience and Behavioral Disorders Program, & the Agency for Science, Technology and Research, Neuroscience Research Partnership, Singapore

⁵Neuroingenia Clinical and Research Center, México, D.F., México

⁶Anita Saltz Institute for Anxiety and Mood Disorders, Child Study Center, NYU Langone School of Medicine, New York, NY, USA

⁷Nathan Kline Institute for Psychiatric Research, Orangeburg, NY, USA

⁸Center for the Developing Brain, Child Mind Institute, New York, NY, USA

Abstract

Background—Attention-Deficit/Hyperactivity Disorder (ADHD) is increasingly conceived as reflecting altered functional and structural brain connectivity. The latter can be addressed with diffusion tensor imaging (DTI). We examined fractional anisotropy (FA), a DTI index related to white matter structural properties, in adult males diagnosed with ADHD in childhood (probands) and matched comparisons without childhood ADHD. Additionally, we contrasted FA among probands with and without current ADHD in adulthood and comparisons.

© 2013 Society of Biological Psychiatry. Published by Elsevier Inc. All rights reserved.

Corresponding author: Samuele Cortese, M.D., Ph.D., Phyllis Green and Randolph Cowen Institute for Pediatric Neuroscience, NYU Child Study Center, One Park Avenue, 7th Floor, 10016 New York, NY, USA. Phone: +1 646 754 5150; Fax: +1 646 754 5211; samuele.cortese@gmail.com.

Part of this work was presented on May 4, 2012 at the 2012 Annual Meeting of the Society of Biological Psychiatry in Philadelphia, PA.

Financial disclosures:

Dr. Cortese served as scientific consultant for Shire Pharmaceuticals from June 2009 to December 2010. He received support to attend meetings from Eli Lilly and Co. in 2008 and from Shire in 2009–2010. The remaining authors (Drs. Imperati, Zhou, Proal, Klein, Mannuzza, Ramos-Olazagasti, Milham, Kelly, and Castellanos) reported no biomedical financial interests or potential conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Methods—Participants were from an original cohort of 207 boys and 178 male comparisons. At 33-year follow-up, analyzable DTI scans were obtained in 51 probands (41.3 ± 2.8 yrs) and 66 comparisons (41.2 ± 3.1 yrs). Voxel-based FA was computed using tract-based spatial statistics (TBSS), controlling for multiple comparisons.

Results—Probands with childhood ADHD exhibited significantly lower FA than comparisons without childhood ADHD in the right superior and posterior corona radiata, right superior longitudinal fasciculus, and in a left cluster including the posterior thalamic radiation, the retrolenticular part of the internal capsule, and the sagittal stratum ($p < 0.05$, corrected). FA was significantly decreased relative to comparisons in several tracts in both probands with current and remitted ADHD, who did not differ significantly from each other. FA was not significantly increased in probands in any region.

Conclusions—Decreased FA in adults with childhood ADHD regardless of current ADHD may be an enduring trait of ADHD. White matter tracts with decreased FA connect regions involved in high-level as well as sensorimotor functions, suggesting that both types of processes are involved in the pathophysiology of ADHD.

Keywords

ADHD; DTI; Fractional anisotropy; Neuroimaging; Longitudinal follow-up; Pathophysiology

Introduction

Attention-Deficit/Hyperactivity Disorder (ADHD), defined by a persistent and age-inappropriate pattern of inattention, hyperactivity-impulsivity, or both (1), is a common childhood-onset psychiatric condition, with estimated worldwide-pooled prevalence exceeding 5% in school-age children (2). Longitudinal studies have documented that impairing symptoms of ADHD persist into adulthood in a substantial proportion (3). Such symptomatic continuity suggests continuity of pathophysiology, which has been addressed mostly through neuroimaging (4–7). Recently, the pathophysiological conceptualization of ADHD has shifted from models focused primarily on fronto-striatal regions (8) to dysfunction in widely distributed large-scale brain networks (5;9;10). Accordingly, the field has increasingly focused on abnormalities in connectivity, i.e., on white matter properties (9;11;12).

In the present report, we focus on the assessment of white matter using fractional anisotropy (FA), an index obtained with Diffusion Tensor Imaging (DTI), which reflects a complex mixture of tissue properties, including axonal ordering, axonal density, and myelination (13). Although it is inappropriate to interpret FA as a direct measure of white matter integrity, it does reflect physical differences in white matter structure (13), and thus serves as a starting point for exploring structural connectivity.

A meta-analysis of DTI studies in ADHD (14), mostly conducted in children, documented abnormal FA of numerous WM tracts, including the anterior corona radiata, internal capsule, forceps minor, and cerebellar tracts. The smaller DTI literature on adults with ADHD is limited to specific brain regions-of-interest (15–17) or does not correct for multiple comparisons (18). Additionally, DTI studies in adults with ADHD have relied on retrospective recall of childhood symptoms, which can be questionable (19).

We report whole-brain FA analyses corrected for multiple comparisons in the largest sample to date of adults with a childhood diagnosis consistent with DSM-IV combined type ADHD (probands) and prospectively enrolled participants free of ADHD in childhood (comparisons) (20). In our previous study of gray matter structure based on the same cohorts

(21), we conducted analyses based both on original (i.e., childhood) group assignment and on current diagnostic status in adulthood. Similarly, here our first objective was to contrast FA in adulthood between probands with childhood ADHD and comparisons without childhood ADHD. Our prior analyses (21) revealed reduced cortical thickness in probands relative to comparisons in parietal, temporal, frontal, and occipital regions. Analyses of gray matter density using voxel based morphometry (VBM) generally echoed these findings and also revealed decreased gray matter density in caudate, thalamus, and cerebellum. Accordingly, we expected to find alterations of FA in probands with childhood ADHD relative to comparisons without childhood ADHD in tracts connecting the gray matter regions that were abnormal in our previous analyses (21).

As in our prior study (21), the second objective was to contrast probands with persistent ADHD and those in remission to comparisons free of current ADHD symptoms in adulthood and to each other. Analyses based on current diagnosis in adulthood allow comparability with cross-sectional studies of adults with ADHD. They also allowed us to examine FA correlates of persistence versus remission in ADHD for the first time.

Methods and Materials

Participants

The study was approved by the Institutional Review Boards of NYU Langone Medical Center and NYU. Participants provided written informed consent and were compensated for participating.

Probands originally consisted of 207 6- to 12-year-old middle class Caucasian boys referred to a research clinic from 1970 to 1978 (mean $\pm$ SD: 8.3 $\pm$ 1.6 years) (22–24). Inclusion criteria were: school referral because of behavior problems, elevated parent and teacher hyperactivity ratings, behavior problems in multiple settings, IQ $\geq$ 85, and English speaking parents. Children with a pattern of aggressive or antisocial behavior were excluded to rule out comorbid conduct disorder. Psychosis and neurological disorders were also exclusionary. As detailed elsewhere (25), all probands would have met criteria for DSM-IV ADHD combined type. Of 207 ADHD probands, 182 were treated with methylphenidate for an average of 2.2 $\pm$ 1.6 years. Three follow-up waves were conducted, when probands were 18.4 $\pm$ 1.3, 25.0 $\pm$ 1.3, and 41.2 $\pm$ 2.7 years (FU18, FU25, and FU41, respectively). Comparison male subjects (n=178) matched for age, social class, and geographic residence, were recruited at FU18 from the same medical center among children seen for routine physical exams whose record history and interview with parents did not indicate behavior problems during elementary school. At FU41, 135/207 male probands (65%) and 136/178 male comparisons (76%) participated in the follow-up. The Structured Clinical Interview for DSM-IV Axis I Disorders, Non-Patient Edition (SCID-I/NP) (26) was administered by clinicians, inquiring about function during the interval between FU25 and FU41. A special interview, Assessment of Adult Attention Deficit/Hyperactivity Disorder (21;27), was developed for diagnosing DSM-IV ADHD in adults. Current ADHD was defined as meeting DSM-IV criteria during the preceding six months. Participants diagnosed as having ADHD-NOS at FU41 had fewer than the required number of DSM-IV criteria, but reported significant impairment or distress associated with these symptoms.

DTI Acquisition

Anatomic T1-weighted scans were obtained on a 3T Siemens Trio with an 8-channel head coil (probands: n=18, comparisons: n=14, total: n=32) or a 3T Siemens Allegra with a single-channel head coil (probands: n=33, comparisons: n=52, total: n=85) with the following parameters: TR=2100ms; flip angle=12; slice thickness=1.5mm; inversion

time=1100ms; matrix:=192×256; FOV=172.5mm. TE was 3.87ms on the Trio and 3.90ms on the Allegra.

DTI was acquired in one non-weighted and six diffusion-weighted non-collinear directions using an echo-planar sequence with diffusion weighting (b value) of 1000 mm^{-1} . A dual spin echo was used to minimize distortion due to eddy currents. Imaging parameters were: TR=6100ms; flip angle=90; FOV=178×219; matrix=104×128; voxel size=1.71×1.71×4 mm; number of averages=4. TE was 102ms on the Allegra and 92ms on the Trio.

Image Preprocessing

Motion correction was performed by applying 9 degrees of freedom linear registration to a standard MNI template. Eddy-current and EPI distortion-related corrections were also performed. Diffusion gradients were rotated to improve consistency with the motion parameters. Diffusion images were visually inspected for motion effects by two investigators (SC, JZ). Data were discarded if the interslice displacement was more than 3mm. Data for each of the six corrected directions were used to fit the tensor parameters. Diffusion tensors were fitted for each voxel to obtain FA images, which were registered to FMRIB58_FA standard space image with 1mm³ resolution using the nonlinear registration tool FNIRT (28). We then applied tract-based spatial statistics (TBSS) (29) to carry out a voxel-wise analysis of FA data within major WM pathways throughout the brain. TBSS minimizes problems of intersubject registration by first determining a mean FA ‘skeleton,’ representing only the center of major WM fiber tracts, then mapping each participant’s DTI data directly onto the skeleton. Analysis of FA differences is thus restricted to regions that represent, with high confidence, only the center of equivalent WM tracts of each individual. A standard FA template was used, and each participant’s standard space FA data were projected onto the FA template using the standard preprocessing scripts provided with FMRIB Software Library (FSL)’s TBSS and adapted for group features (29). The resultant skeletonized FA images were used for statistical analyses.

Statistical Analyses

Group differences in sample characteristics were tested with independent-samples *t* tests or ² tests. DTI data were analyzed using FSL 4.1.5 (30). We examined voxelwise cross-subject spatial statistics of FA values using permutation-based non-parametric testing (FSL’s RANDOMISE) on the skeletonized FA images. First, we compared probands with childhood ADHD to comparisons without childhood ADHD. Then, to test whether FA differed as a function of current ADHD, we classified probands as to whether they had ADHD at FU41 or not, thus generating two proband subgroups, “probands with persistent ADHD” and “probands with remitted ADHD.” They were contrasted to comparisons who did not meet criteria for ADHD-NOS at FU41 (“non-ADHD comparisons”) (20). Contrasts were: 1) probands with persistent ADHD vs. non-ADHD comparisons; 2) probands with remitted ADHD vs. non-ADHD comparisons; and 3) probands with persistent ADHD vs. probands with remitted ADHD. In each contrast, age and scanner model were covaried. (Supplementary analyses were performed limited to the 85 datasets obtained on the Allegra scanner, to address concerns regarding possible dependence of DTI parameters on scanner type and sequence.) We corrected for multiple comparisons using threshold-free cluster enhancement (TFCE) (31). The Johns Hopkins University DTI-based WM atlas, available in FSL (30), was used to label the WM tracts.

Results

Subjects

A total of 152 participants were scanned at FU41, of whom 144 (61 probands and 83 comparisons) underwent diffusion-weighted scans. DTI data for 10 probands and 17 comparisons failed quality criteria, leaving 51 probands and 66 comparisons with analyzable DTI data. Rates of MRI refusal and failure to schedule or locate subjects did not differ significantly between probands and comparisons (45% vs. 43%). However, a smaller proportion of probands (32%) than comparisons (48%) were scanned. This discrepancy reflects a significantly higher rate of unavoidable factors in probands (i.e., deaths, incarcerations or MRI exclusions) than in comparisons (27% vs. 12%, respectively; $p<0.001$) (20). Within both proband and comparison groups, individuals scanned and those not scanned did not differ significantly on age at referral, childhood IQ, socioeconomic status, Teachers Connors Hyperactivity Factor scores (32), and rates of mental disorders at FU18 (ADHD, Antisocial Personality Disorder, Mood or Anxiety Disorders) (21). However, scanned probands had significantly higher rates of substance use disorders (SUD) at FU18 than not scanned probands (25% vs. 8%; $p=0.02$) (21). Scanned individuals with or without analyzable DTI data did not differ significantly in scanner type ($p=0.99$), age ($p=0.53$), or full scale IQ ($p=0.91$) at FU41.

Fifteen of the 51 probands with analyzable DTI met DSM-IV(TR) criteria for current ADHD: six (11.8%) with inattentive type, six (11.8%) with hyperactive-impulsive type, and three (5.9%) with combined type. Twenty-five probands (49%) were classified as remitters (Table 1). Probands ($n=11$, 21.6%) and comparisons ($n=19$, 28.7%) with ADHD-NOS at FU41 were excluded from subgroup analyses as in our prior study (21). Therefore, we used data from 47 (66 minus 19) non-ADHD comparisons for subgroup analyses (21). Forty-seven probands (92%) had been treated with methylphenidate in childhood. Co-occurring ongoing SUD did not differ between probands with childhood ADHD and comparisons without childhood ADHD (22% vs. 21%, respectively) nor among probands with persistent ADHD, those in remission, or non-ADHD comparisons (Table 1).

DTI Analyses

Table 2 and Figure 1 present FA differences between probands and comparisons. Probands with childhood ADHD ($n=51$) exhibited significantly lower FA than comparisons without childhood ADHD ($n=66$), adjusting for age and scanner model, in two large sets of clusters, one in each hemisphere. The left hemisphere cluster encompassed the sagittal stratum and the retrolenticular part of the internal capsule, including the posterior thalamic radiation. The right hemisphere cluster comprised the superior longitudinal fasciculus II, and the posterior and superior corona radiata.

Probands with persistent ADHD ($n=15$) exhibited significantly lower FA than non-ADHD comparisons ($n=47$) in a right hemisphere tract encompassing the sagittal stratum, external capsule, retrolenticular part of the internal capsule, and the anterior corona radiata (Table 2 and Figure 2). Probands with remitted ADHD ($n=25$) also exhibited significantly lower FA than non-ADHD comparisons in right superior and posterior corona radiata (Table 2 and Figure 3). Probands with persistent ADHD and those who had remitted did not differ significantly in FA when correcting for multiple comparisons or even at $p<0.01$, uncorrected. In no instance did probands exhibit significantly higher FA than comparisons. Analyses limited to the 85 scans obtained using the Allegra scanner revealed similar results, albeit at a somewhat reduced level of statistical significance ($p<0.15$, corrected; see Supplementary Table 1).

Discussion

This is the first DTI study in adults with ADHD established in childhood. As expected, we found significantly decreased FA in probands relative to comparisons, regardless of ADHD diagnosis at mean age 41, in WM tracts connecting gray matter regions that we found to be abnormal in the same cohorts (21). FA in probands with childhood ADHD was decreased relative to comparisons without childhood ADHD in a left hemisphere cluster encompassing the sagittal stratum and the retrolenticular part of the internal capsule and in a right hemisphere cluster comprising the superior longitudinal fasciculus II, the posterior corona radiata, and the superior corona radiata. In subgroups defined by current ADHD diagnoses (i.e., in adulthood), probands with persistent ADHD exhibited significantly reduced FA relative to non-ADHD comparisons in several tracts, two of which overlapped with those found in the all-inclusive analysis, albeit in the opposite (right) hemisphere. Probands with ADHD in remission also had significantly lower FA than non-ADHD comparisons in right hemisphere tracts. Probands with persistent and remitted ADHD did not differ significantly from each other in any tract.

Probands with Childhood ADHD Relative to Comparisons without Childhood ADHD

We found reduced FA in probands in tracts connecting regions involved in higher-level cognitive as well as in sensorimotor functions including, in particular, visual processing. These are:

Left sagittal stratum—This tract contains fibers of the inferior fronto-occipital fasciculus (IFO), which connects the lateral aspects of the frontal and occipital lobes as well as prefrontal cortex to auditory (BA 22) and visual association cortex (BA 20–21). The IFO is implicated in attention set-shifting abilities (33), which are deficient in ADHD (34–36). FA in the IFO was reduced in a study of adults with ADHD (18). Additionally, the sagittal stratum contains fibers of the inferior longitudinal fasciculus (ILF), which was reported abnormal in children with ADHD (37). The ILF relays visual information from occipital to temporal cortex, subserving visual perception and object recognition (38). Alterations in this tract may underlie visual memory deficits in ADHD (39).

Left retrolenticular part of internal capsule (including posterior thalamic radiation)—Consistent with findings in children with ADHD (40), probands had reduced FA in the retrolenticular part of internal capsule (RIC), mostly within the posterior thalamic radiation, which contains fibers of the optic radiation carrying visual information from the lateral geniculate nucleus to the occipital lobe (41). This finding further underscores the possible role of dysfunctions within the visual system in ADHD. Although largely overlooked in the ADHD literature, partly because of the focus on fronto-striatal dysfunctions (8), anomalies of the visual system have been reported in structural (42) and functional (43) MRI studies. Additionally, event-related electrophysiological and MRI studies concur in suggesting a possible deficit in early visual processing in ADHD (44;45).

Right superior longitudinal fasciculus (SLF) II—The SLF II is a pathway connecting dorsolateral prefrontal regions and caudal-inferior parietal lobe (46) that has been found to be abnormal in adults (17) and children (37;47) with ADHD. Visual perceptual information carried by the SLF II from the parietal lobe to the prefrontal cortex allows prefrontal cortex to regulate focusing attention in space (46). The SLF II also subserves spatial working memory (48), a consistent executive function deficit in children (49;50) and adults (51) with ADHD.

Also in the right hemisphere, we found two clusters with reduced FA in probands in the *superior* and *posterior corona radiata*, consistent with results in children with ADHD (52). The superior and posterior corona radiata include descending sensorimotor fibers contributing to the corticospinal tract (53;54). Alterations in these tracts may underpin sensorimotor deficits in ADHD (55).

FA in the anterior corona radiata, found reduced in children with ADHD (52;56–58), did not differ significantly in probands vs. comparison at $p < 0.05$, corrected. However, at a lower threshold of $p = 0.12$, corrected, a significant FA reduction in the anterior corona radiata in probands vs. comparisons emerged (data not shown). Discrepancies between our findings and previous results may be ascribable to different analytical procedure (i.e., TBSS was not implemented in early studies). We note that a prior study using TBSS in children with ADHD found increased, rather than reduced FA in several tracts, which was speculatively linked to decreased neuronal branching in ADHD (37).

Relationship of White Matter Findings to Cortical Thickness and VBM Findings in this Study Cohort: The white matter tracts that differentiated probands from comparisons connect brain regions that we found to be abnormal in cortical thickness and VBM (21). Specifically, probands had relatively: reduced cortical thickness in right dorsolateral (right middle frontal gyrus, BA9) and right inferior parietal regions, which are connected by the right SLF II; reduced cortical thickness in the right (as well as left) precentral gyrus (BA6), from which descending fibers of the superior and posterior corona radiata depart; reduced cortical thickness in the left frontal pole and reduced gray matter density in the left middle temporal gyrus (BA21), connected by the IFO; reduced cortical thickness and gray matter density in left temporal and occipital regions, connected by the ILF; and reduced gray matter density in the left temporo-occipital cortex (BA37; involved in visual recognition), as well as right occipital areas 18 and 19, which are targets of the posterior thalamic radiation. In contrast with our finding of reduced VBM in cerebellum (21), and reports of abnormal FA in cerebellar peduncles in children with ADHD (57;59;60), we did not find significant alterations in tracts connected with cerebellar regions. The only other voxel-wise DTI study in ADHD adults (18) also failed to find significant alterations in cerebellar-related tracts.

Study Findings in Relation to Neural Models of ADHD: Summarizing, we found evidence of WM alterations in tracts connecting regions involved in higher-level cognitive functions as well as sensory and motor functions. In response to an early model of ADHD highlighting executive functions (61), imaging studies have tended to focus on prefrontal and related regions and ignore those involved in more basic sensory and motor processes. In particular, abnormalities in visual areas have been overlooked, even when abnormal results were reported, e.g., (62;63). A recent meta-analysis (43) of functional MRI studies of ADHD documented ADHD-related abnormalities in systems subserving higher-level cognitive functions, such as the frontoparietal, dorsal attention, and default network, as well as regions underpinning sensory (including visual) and motor functions.

Our findings of significant FA differences in ADHD probands relative to comparisons, regardless of current ADHD diagnosis, support the interpretation that WM alterations may represent an enduring neurobiological trait independent of syndromic remission. Since this is a cross-sectional DTI assessment embedded in a prospective clinical follow-up, we cannot establish whether significant reductions in FA reflect recent, or early and enduring alterations in ADHD. The latter seems most likely in light of prior results of FA deficits in children with ADHD (52), and the lack of differences between probands with remitted and persistent ADHD.

Probands with Persistent ADHD Relative to Non-ADHD Comparisons

The analysis in probands with childhood ADHD vs. comparisons without childhood ADHD revealed FA differences in both hemispheres. By contrast, individuals with persistent ADHD, relative to non-ADHD comparisons, showed reduced FA only in the right hemisphere. However, analyses with a relaxed statistical threshold ($p < 0.09$, corrected, data not shown), revealed that these tracts also tended to be abnormal in the left hemisphere. Thus, hemispheric differences likely reflect threshold effects. At $p = 0.05$, corrected, reduced FA in probands with persistent ADHD vs. non-ADHD comparisons was found in two clusters, i.e., sagittal stratum and the retrolenticular part of the internal capsule, which had been detected in the left hemisphere in the all-inclusive analysis.

Three additional clusters of reduced FA in probands with persistent ADHD were found in the right anterior corona radiata and external capsule. Several groups have reported FA reductions in the anterior corona radiata in ADHD (52;56–58). The anterior corona radiata contains WM fibers connecting the anterior cingulate cortex (ACC) to the striatum (64). Abnormalities of the dorsal ACC in ADHD have been found in both structural (65–67) and functional data (68;69). The dorsal ACC subserves multiple functions which are altered in ADHD such as attention, target detection, response selection/ inhibition, error detection, and motivation (70). Finally, external capsule FA was reduced in a study of adolescents with very low birth weight and ADHD (71). Consistent with findings from the contrast “probands with childhood ADHD vs. comparisons without childhood ADHD,” in no instance did probands with persistent ADHD at FU41 have significantly higher FA than non-ADHD comparisons.

Because of methodological differences, comparisons with prior studies of FA in adults with ADHD are not straightforward. Makris et al. (17) found significantly lower FA values in the cingulum bundle and in the SLF II in adults with ADHD vs. comparisons; Dransdahl et al. (15) showed ADHD-related reduction of FA in the isthmus/splenium of the corpus callosum; Konrad et al. (16) reported lower FA in the left inferior longitudinal fasciculus (ILF) in adults with ADHD vs. comparisons. However, all three studies used a region-of-interest approach, whereas we did not limit our investigation to specific regions. In an earlier study, Konrad et al. (18) reported reduced FA bilaterally in medial orbitofrontal white matter and in the right anterior cingulate bundle, and elevated FA bilaterally in temporal WM structures in the ADHD group. However, differently from ours, these results were uncorrected for multiple comparisons.

Probands with Remitted ADHD Relative to Non-ADHD Comparisons and Probands with Persistent ADHD

This study provided the first opportunity to investigate FA in relation to ADHD remission. Probands with remitted ADHD exhibited significantly lower FA than non-ADHD comparisons in right superior and posterior corona radiata. These two tracts were also found when contrasting probands with childhood ADHD and comparisons without childhood ADHD. However, probands with remitted ADHD did not differ significantly from probands with current ADHD even though the former had less than a total of two ADHD symptoms on average (data not shown). This suggests that decreased FA in certain white matter tracts predominantly reflects childhood ADHD status, independently of current diagnostic status, presumably reflecting the influence of genetic factors.

Limitations

We were able to analyze imaging data for only 25% of the original cohort of ADHD probands and 37% of comparison subjects. However, probands and comparisons studied were representative of the original sample, and the probands studied did not differ

significantly from lost subjects on nearly all clinical and demographic variables, except for significantly higher rates of SUD at FU18 in scanned probands. However, probands and comparisons who were scanned had nearly identical rates of current SUD at FU41. Second, our subjects were exclusively Caucasian males, since the number of originally diagnosed females with ADHD was too small ($n=19$) for meaningful study. Thus, our results may not generalize to women, or to other racial or ethnic groups. However, this constraint avoided potential confounds from possible sex, ethnic, or socioeconomic differences. Exclusion of conduct disorder comorbidity in childhood also averted confusion as to the origin of the FA deficits we found. Third, our DTI acquisition protocol included only six directions, which was standard when the study was planned. However, imprecision in FA values resulting from the use of only six directions of diffusion sensitization would affect FA measures in comparisons and probands equally and should not have biased group differences. For logistic reasons, we used two different scanners. Fortunately, group representation did not differ significantly across the scanners and we performed all analyses using scanner model as a covariate. We also confirmed that the same pattern of results held up when analyses were limited to the single scanner on which most studies were performed. Fourth, we cannot comment on the effects of stimulant treatment in childhood on FA, since all but four of the probands received stimulant treatment in childhood for an average of ~2 years. However, prior studies have failed to detect stimulant effects on FA (47;59;72). Finally, despite the substantial overall size of the sample, sub-analyses based on current diagnosis were based on small groups, which limited statistical power. Thus the lack of differences between probands in remission from those with persistent ADHD should be viewed as suggestive.

Conclusions

We found evidence of decreased FA, reflecting altered white matter properties, in adults with childhood ADHD, regardless of presence or absence of current ADHD, suggesting that FA alteration may be an enduring trait related to ADHD. The implicated WM tracts connect regions involved in high-level as well as sensorimotor functions, suggesting that both types of processes may be involved in the pathophysiology of ADHD.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work has been supported by R01MH018579 (to R. Klein) and R01DA016979 and T32MH-067763 (to F. X. Castellanos). Dr. Cortese was supported by a grant from the European Union Commission (Marie Curie Actions, International Outgoing Fellowships for Career Development, Grant# 253103).

References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4. Washington DC: 2000. Text revision
2. Polanczyk G, de Lima MS, Horta BL, Biederman J, Rohde LA. The worldwide prevalence of ADHD: a systematic review and meta-regression analysis. *Am J Psychiatry*. 2007; 164(6):942–948. [PubMed: 17541055]
3. Mannuzza S, Klein RG, Moulton JL III. Persistence of Attention-Deficit/Hyperactivity Disorder into adulthood: what have we learned from the prospective follow-up studies? *J Atten Disord*. 2003; 7(2):93–100. [PubMed: 15018358]
4. Castellanos FX. Toward a pathophysiology of attention-deficit/hyperactivity disorder. *Clin Pediatr (Phila)*. 1997; 36(7):381–393. [PubMed: 9241475]

5. Castellanos FX, Proal E. Large-scale brain systems in ADHD: beyond the prefrontal-striatal model. *Trends Cogn Sci.* 2012; 16(1):17–26. [PubMed: 22169776]
6. Durston S. A review of the biological bases of ADHD: what have we learned from imaging studies? *Ment Retard Dev Disabil Res Rev.* 2003; 9(3):184–195. [PubMed: 12953298]
7. Shaw P, Rabin C. New insights into attention-deficit/hyperactivity disorder using structural neuroimaging. *Curr Psychiatry Rep.* 2009; 11(5):393–398. [PubMed: 19785981]
8. Barkley RA. Behavioral inhibition, sustained attention, and executive functions: constructing a unifying theory of ADHD. *Psychol Bull.* 1997; 121(1):65–94. [PubMed: 9000892]
9. Konrad K, Eickhoff SB. Is the ADHD brain wired differently? A review on structural and functional connectivity in attention deficit hyperactivity disorder. *Hum Brain Mapp.* 2010; 31(6):904–916. [PubMed: 20496381]
10. Makris N, Biederman J, Monuteaux MC, Seidman LJ. Towards conceptualizing a neural systems-based anatomy of attention-deficit/hyperactivity disorder. *Dev Neurosci.* 2009; 31(1–2):36–49. [PubMed: 19372685]
11. Liston C, Malter CM, Teslovich T, Levenson D, Casey BJ. Atypical prefrontal connectivity in attention-deficit/hyperactivity disorder: pathway to disease or pathological end point? *Biol Psychiatry.* 2011; 69(12):1168–1177. [PubMed: 21546000]
12. Rubia K, Halari R, Cubillo A, Mohammad AM, Brammer M, Taylor E. Methylphenidate normalises activation and functional connectivity deficits in attention and motivation networks in medication-naïve children with ADHD during a rewarded continuous performance task. *Neuropharmacology.* 2009; 57(7–8):640–652. [PubMed: 19715709]
13. Jones DK, Knosche TR, Turner R. White matter integrity, fiber count, and other fallacies: The do's and don'ts of diffusion MRI. *Neuroimage.* in press.
14. van Ewijk H, Heslenfeld DJ, Zwiers MP, Buitelaar JK, Oosterlaan J. Diffusion tensor imaging in attention deficit/hyperactivity disorder: A systematic review and meta-analysis. *Neurosci Biobehav Rev.* 2012; 36(4):1093–106. [PubMed: 22305957]
15. Dransdahl M, Westerhausen R, Haavik J, Hugdahl K, Plessen KJ. Adults with attention-deficit/hyperactivity disorder - a diffusion-tensor imaging study of the corpus callosum. *Psychiatry Res.* 2012; 201(2):168–173. [PubMed: 22386969]
16. Konrad A, Dielentheis TF, Masri DE, Dellani PR, Stoeter P, Vucurevic G, et al. White matter abnormalities and their impact on attentional performance in adult attention-deficit/hyperactivity disorder. *Eur Arch Psychiatry Clin Neurosci.* 2012; 262(4):351–60. [PubMed: 21879383]
17. Makris N, Buka SL, Biederman J, Papadimitriou GM, Hodge SM, Valera EM, et al. Attention and executive systems abnormalities in adults with childhood ADHD: A DT-MRI study of connections. *Cereb Cortex.* 2008; 18(5):1210–1220. [PubMed: 17906338]
18. Konrad A, Dielentheis TF, El MD, Bayerl M, Fehr C, Gesierich T, et al. Disturbed structural connectivity is related to inattention and impulsivity in adult attention deficit hyperactivity disorder. *Eur J Neurosci.* 2010; 31(5):912–919. [PubMed: 20374289]
19. Mannuzza S, Klein RG, Klein DF, Bessler A, ShROUT P. Accuracy of adult recall of childhood attention deficit hyperactivity disorder. *Am J Psychiatry.* 2002; 159(11):1882–1888. [PubMed: 12411223]
20. Klein RG, Mannuzza S, Olazagasti MA, Roizen E, Hutchison JA, Lashua EC, et al. Clinical and Functional Outcome of Childhood Attention-Deficit/Hyperactivity Disorder 33 Years Later. *Arch Gen Psychiatry.* in press.
21. Proal E, Reiss PT, Klein RG, Mannuzza S, Gotimer K, Ramos-Olazagasti MA, et al. Brain gray matter deficits at 33-year follow-up in adults with attention-deficit/hyperactivity disorder established in childhood. *Arch Gen Psychiatry.* 2011; 68(11):1122–1134. [PubMed: 22065528]
22. Gittelman R, Mannuzza S, Shenker R, Bonagura N. Hyperactive boys almost grown up. I. Psychiatric status. *Arch Gen Psychiatry.* 1985; 42(10):937–947. [PubMed: 4037987]
23. Mannuzza S, Klein RG, Konig PH, Giampino TL. Hyperactive boys almost grown up. IV. Criminality and its relationship to psychiatric status. *Arch Gen Psychiatry.* 1989; 46(12):1073–1079. [PubMed: 2589922]

24. Mannuzza S, Klein RG, Bessler A, Malloy P, LaPadula M. Adult outcome of hyperactive boys. Educational achievement, occupational rank, and psychiatric status. *Arch Gen Psychiatry*. 1993; 50(7):565–576. [PubMed: 8317950]
25. Mannuzza S, Klein RG, Moulton JL III. Lifetime criminality among boys with attention deficit hyperactivity disorder: a prospective follow-up study into adulthood using official arrest records. *Psychiatry Res*. 2008; 160(3):237–246. [PubMed: 18707766]
26. First, M.; Spitzer, R.; Gibbon, M.; Williams, J. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Non-patient Edition (SCID-I/NP). New York: Biometrics Research, New York State Psychiatric Institute; 2002.
27. Mannuzza S, Castellanos FX, Roizen ER, Hutchison JA, Lashua EC, Klein RG. Impact of the impairment criterion in the diagnosis of adult ADHD: 33-year follow-up study of boys with ADHD. *J Atten Disord*. 2011; 15(2):122–129. [PubMed: 20378923]
28. Rueckert D, Sonoda LI, Hayes C, Hill DL, Leach MO, Hawkes DJ. Nonrigid registration using free-form deformations: application to breast MR images. *IEEE Trans Med Imaging*. 1999; 18(8): 712–721. [PubMed: 10534053]
29. Smith SM, Johansen-Berg H, Jenkinson M, Rueckert D, Nichols TE, Miller KL, et al. Acquisition and voxelwise analysis of multi-subject diffusion data with tract-based spatial statistics. *Nat Protoc*. 2007; 2(3):499–503. [PubMed: 17406613]
30. FMRIB Software Library; Oxford, UK: <http://www.fmrib.ox.ac.uk/fsl/> [Last access: 08.08.2012.]
31. Smith SM, Nichols TE. Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage*. 2009; 44(1):83–98. [PubMed: 18501637]
32. Conners CK. Clinical use of rating scales in diagnosis and treatment of attention-deficit/hyperactivity disorder. *Pediatr Clin North Am*. 1999; 46(5):857–70. vi. [PubMed: 10570692]
33. Kvickstrom P, Eriksson B, van WD, Latt J, Elfgrén C, Nilsson C. Selective frontal neurodegeneration of the inferior fronto-occipital fasciculus in progressive supranuclear palsy (PSP) demonstrated by diffusion tensor tractography. *BMC Neurol*. 2011; 11:13. [PubMed: 21269463]
34. Helenius P, Laasonen M, Hokkanen L, Paetau R, Niemivirta M. Impaired engagement of the ventral attentional pathway in ADHD. *Neuropsychologia*. 2011
35. Nigg JT, Stavro G, Ettenhofer M, Hambrick DZ, Miller T, Henderson JM. Executive functions and ADHD in adults: evidence for selective effects on ADHD symptom domains. *J Abnorm Psychol*. 2005; 114(4):706–717. [PubMed: 16351391]
36. Nigg JT, Casey BJ. An integrative theory of attention-deficit/hyperactivity disorder based on the cognitive and affective neurosciences. *Dev Psychopathol*. 2005; 17(3):785–806. [PubMed: 16262992]
37. Silk TJ, Vance A, Rinehart N, Bradshaw JL, Cunnington R. White-matter abnormalities in attention deficit hyperactivity disorder: a diffusion tensor imaging study. *Hum Brain Mapp*. 2009; 30(9):2757–2765. [PubMed: 19107752]
38. Tusa RJ, Ungerleider LG. The inferior longitudinal fasciculus: a reexamination in humans and monkeys. *Ann Neurol*. 1985; 18(5):583–591. [PubMed: 4073852]
39. Shang CY, Gau SS. Visual memory as a potential cognitive endophenotype of attention deficit hyperactivity disorder. *Psychol Med*. 2011; 1–12. [PubMed: 21835095]
40. Peterson DJ, Ryan M, Rimrodt SL, Cutting LE, Denckla MB, Kaufmann WE, et al. Increased regional fractional anisotropy in highly screened attention-deficit hyperactivity disorder (ADHD). *J Child Neurol*. 2011; 26(10):1296–1302. [PubMed: 21628699]
41. Sherbondy AJ, Dougherty RF, Napel S, Wandell BA. Identifying the human optic radiation using diffusion imaging and fiber tractography. *J Vis*. 2008; 8(10):12–11. [PubMed: 19146354]
42. Ahrendts J, Rusch N, Wilke M, Philipsen A, Eickhoff SB, Glauche V, et al. Visual cortex abnormalities in adults with ADHD: A structural MRI study. *World J Biol Psychiatry*. 2011; 12(4):260–70. [PubMed: 20879808]
43. Cortese S, Kelly C, Chabernaud C, Proal E, Di MA, Milham MP, et al. Toward Systems Neuroscience of ADHD: A Meta-Analysis of 55 fMRI Studies. *Am J Psychiatry*. 2012; 169(10): 1038–1055. [PubMed: 22983386]

44. Nazari MA, Berquin P, Missonnier P, Aarabi A, Debatisse D, de BA, et al. Visual sensory processing deficit in the occipital region in children with attention-deficit/hyperactivity disorder as revealed by event-related potentials during cued continuous performance test. *Neurophysiol Clin.* 2010; 40(3):137–149. [PubMed: 20513613]
45. Yong-Liang G, Robaey P, Karayanidis F, Bourassa M, Pelletier G, Geoffroy G. Stimulus-response incompatibility effects on event-related potentials in children with attention-deficit hyperactivity disorder. *Brain Cogn.* 2000; 43(1–3):211–215. [PubMed: 10857696]
46. Makris N, Kennedy DN, McInerney S, Sorensen AG, Wang R, Caviness VS Jr, et al. Segmentation of subcomponents within the superior longitudinal fascicle in humans: a quantitative, in vivo, DT-MRI study. *Cereb Cortex.* 2005; 15(6):854–869. [PubMed: 15590909]
47. Hamilton LS, Levitt JG, O'Neill J, Alger JR, Luders E, Phillips OR, et al. Reduced white matter integrity in attention-deficit hyperactivity disorder. *Neuroreport.* 2008; 19(17):1705–1708. [PubMed: 18841089]
48. Petrides M. Lateral prefrontal cortex: architectonic and functional organization. *Philos Trans R Soc Lond B Biol Sci.* 2005; 360(1456):781–795. [PubMed: 15937012]
49. Martinussen R, Hayden J, Hogg-Johnson S, Tannock R. A meta-analysis of working memory impairments in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry.* 2005; 44(4):377–384. [PubMed: 15782085]
50. Willcutt EG, Doyle AE, Nigg JT, Faraone SV, Pennington BF. Validity of the executive function theory of attention-deficit/hyperactivity disorder: a meta-analytic review. *Biol Psychiatry.* 2005; 57(11):1336–1346. [PubMed: 15950006]
51. McLean A, Dowson J, Toone B, Young S, Bazanis E, Robbins TW, et al. Characteristic neurocognitive profile associated with adult attention-deficit/hyperactivity disorder. *Psychol Med.* 2004; 34(4):681–692. [PubMed: 15099422]
52. Nagel BJ, Bathula D, Herting M, Schmitt C, Kroenke CD, Fair D, et al. Altered white matter microstructure in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry.* 2011; 50(3):283–292. [PubMed: 21334568]
53. Dougherty RF, Ben-Shachar M, Deutsch GK, Hernandez A, Fox GR, Wandell BA. Temporal-callosal pathway diffusivity predicts phonological skills in children. *Proc Natl Acad Sci U S A.* 2007; 104(20):8556–8561. [PubMed: 17483487]
54. Giorgio A, Watkins KE, Chadwick M, James S, Winmill L, Douaud G, et al. Longitudinal changes in grey and white matter during adolescence. *Neuroimage.* 2010; 49(1):94–103. [PubMed: 19679191]
55. Mostofsky SH, Rimrod SL, Schafer JG, Boyce A, Goldberg MC, Pekar JJ, et al. Atypical motor and sensory cortex activation in attention-deficit/hyperactivity disorder: a functional magnetic resonance imaging study of simple sequential finger tapping. *Biol Psychiatry.* 2006; 59(1):48–56. [PubMed: 16139806]
56. Davenport ND, Karatekin C, White T, Lim KO. Differential fractional anisotropy abnormalities in adolescents with ADHD or schizophrenia. *Psychiatry Res.* 2010; 181(3):193–198. [PubMed: 20153608]
57. Kobel M, Bechtel N, Specht K, Klarhofer M, Weber P, Scheffler K, et al. Structural and functional imaging approaches in attention deficit/hyperactivity disorder: does the temporal lobe play a key role? *Psychiatry Res.* 2010; 183(3):230–236. [PubMed: 20702071]
58. Pavuluri MN, Yang S, Kamineni K, Passarotti AM, Srinivasan G, Harral EM, et al. Diffusion tensor imaging study of white matter fiber tracts in pediatric bipolar disorder and attention-deficit/hyperactivity disorder. *Biol Psychiatry.* 2009; 65(7):586–593. [PubMed: 19027102]
59. Ashtari M, Kumra S, Bhaskar SL, Clarke T, Thaden E, Cervellione KL, et al. Attention-deficit/hyperactivity disorder: a preliminary diffusion tensor imaging study. *Biol Psychiatry.* 2005; 57(5):448–455. [PubMed: 15737658]
60. Bechtel N, Kobel M, Penner IK, Klarhofer M, Scheffler K, Opwis K, et al. Decreased fractional anisotropy in the middle cerebellar peduncle in children with epilepsy and/or attention deficit/hyperactivity disorder: a preliminary study. *Epilepsy Behav.* 2009; 15(3):294–298. [PubMed: 19362604]

61. Barkley RA. Attention-deficit/hyperactivity disorder, self-regulation, and time: toward a more comprehensive theory. *J Dev Behav Pediatr.* 1997; 18(4):271–279. [PubMed: 9276836]
62. Dillo W, Goke A, Prox-Vagedes V, Szyck GR, Roy M, Donnerstag F, et al. Neuronal correlates of ADHD in adults with evidence for compensation strategies--a functional MRI study with a Go/No-Go paradigm. *Ger Med Sci.* 2010; 8:Doc09. [PubMed: 20421953]
63. Schneider MF, Krick CM, Retz W, Hengesch G, Retz-Junginger P, Reith W, et al. Impairment of fronto-striatal and parietal cerebral networks correlates with attention deficit hyperactivity disorder (ADHD) psychopathology in adults - a functional magnetic resonance imaging (fMRI) study. *Psychiatry Res.* 2010; 183(1):75–84. [PubMed: 20558047]
64. Wakana S, Jiang H, Nagae-Poetscher LM, van Zijl PC, Mori S. Fiber tract-based atlas of human white matter anatomy. *Radiology.* 2004; 230(1):77–87. [PubMed: 14645885]
65. Amico F, Stauber J, Koutsouleris N, Frodl T. Anterior cingulate cortex gray matter abnormalities in adults with attention deficit hyperactivity disorder: a voxel-based morphometry study. *Psychiatry Res.* 2011; 191(1):31–35. [PubMed: 21129938]
66. Makris N, Seidman LJ, Valera EM, Biederman J, Monuteaux MC, Kennedy DN, et al. Anterior cingulate volumetric alterations in treatment-naïve adults with ADHD: a pilot study. *J Atten Disord.* 2010; 13(4):407–413. [PubMed: 20008822]
67. Seidman LJ, Valera EM, Makris N, Monuteaux MC, Boriol DL, Kelkar K, et al. Dorsolateral prefrontal and anterior cingulate cortex volumetric abnormalities in adults with attention-deficit/hyperactivity disorder identified by magnetic resonance imaging. *Biol Psychiatry.* 2006; 60(10):1071–1080. [PubMed: 16876137]
68. Bush G, Frazier JA, Rauch SL, Seidman LJ, Whalen PJ, Jenike MA, et al. Anterior cingulate cortex dysfunction in attention-deficit/hyperactivity disorder revealed by fMRI and the Counting Stroop. *Biol Psychiatry.* 1999; 45(12):1542–1552. [PubMed: 10376114]
69. Tian L, Jiang T, Wang Y, Zang Y, He Y, Liang M, et al. Altered resting-state functional connectivity patterns of anterior cingulate cortex in adolescents with attention deficit hyperactivity disorder. *Neurosci Lett.* 2006; 400(1–2):39–43. [PubMed: 16510242]
70. Sheth SA, Mian MK, Patel SR, Asaad WF, Williams ZM, Dougherty DD, et al. Human dorsal anterior cingulate cortex neurons mediate ongoing behavioural adaptation. *Nature.* 2012; 488(7410):218–221. [PubMed: 22722841]
71. Skranes J, Vangberg TR, Kulseng S, Indredavik MS, Evensen KA, Martinussen M, et al. Clinical findings and white matter abnormalities seen on diffusion tensor imaging in adolescents with very low birth weight. *Brain.* 2007; 130(Pt 3):654–666. [PubMed: 17347255]
72. de Zeeuw P, Mandl RC, Hulshoff Pol HE, van EH, Durston S. Decreased frontostriatal microstructural organization in attention deficit/hyperactivity disorder. *Hum Brain Mapp.* 2012; 33(8):1941–1951. [PubMed: 21826757]

Figure 1.

Regions where probands with childhood ADHD ($n=51$) had significantly lower FA than comparisons without childhood ADHD ($n=66$) (TFCE $p<0.05$, corrected). The mean FA skeleton is represented in green. Significantly different clusters have been thickened with the FMRIB Software Library (FSL) option “tbss_fill” (30) (in a yellow-red scale) for visualization. The clusters are labeled according to the *John Hopkins University DTI*-based white-matter atlas (ICBM-DTI-81), available in FSL (30). Although slices show all significant clusters, arrows indicate the specific cluster of interest for each column. PCR R: Posterior Corona Radiata Right; PTR L: Posterior Thalamic Radiation Left; RIC L: Retrolenticular Part of Internal Capsule Left; SCR R: Superior Corona Radiata Right; SLF R: Superior Longitudinal Fasciculus Right.

Figure 2.

Regions where probands with persistent ADHD ($n=15$) had significantly lower FA than non-ADHD comparisons ($n=47$) (TFCE $p<0.05$, corrected). The mean FA skeleton is represented in green. Significantly different clusters have been thickened with the FMRIB Software Library (FSL) option “tbss_fil” (30) (in a yellow-red scale) for visualization. The clusters are labeled according to the *John Hopkins University DTI*-based white-matter atlas (ICBM-DTI-81), available in FSL (30)

ACR R: Anterior Corona Radiata Right; ECR R: External Capsule Right; RIC R: Retrolenticular part of Internal Capsule Right; SS R: Sagittal Stratum Right.

Figure 3.

Regions where probands with remitted ADHD (n= 15) had significantly lower FA than non-ADHD comparisons (n=47) (TFCE $p < 0.05$, corrected). The mean FA skeleton is represented in green. Significantly different clusters have been thickened with the FMRIB Software Library (FSL) option “tbss_fill” (30) (in a yellow-red scale) for visualization. The clusters are labeled according to the *John Hopkins University DTI*-based white-matter atlas (ICBM-DTI-81), available in FSL (30)

PCR R: Posterior Corona Radiata Right; SCR R: Superior Corona Radiata Right.

Table 1
Demographic and clinical characteristics of participants with analyzable DTI data at follow-up at mean age 41 (FU41).

	Prob. <i>N</i> =51 <i>Mean (SD) or N</i> (%)	Comp. <i>N</i> =66 <i>Mean (SD) or N</i> (%)	ADHD+ <i>N</i> =15 <i>Mean (SD) or N</i> (%)	ADHD- <i>N</i> =25 <i>Mean (SD) or N</i> (%)	Non-ADHD Comp.¹ <i>N</i> =47 <i>Mean (SD) or N</i> (%)	<i>p</i> value (2-sided)	<i>p</i> value (2-sided)	<i>Post-hoc LSD</i>
Age at FU41	41.3 (2.8)	41.2 (3.1)	41.8 (3.0)	41.3 (2.6)	41.1 (3.0)	0.81	0.74	
SES at FU41	3.4 (1.1)	2.4 (1.0)	3.7 (1.2)	3.2 (1.1)	2.4 (1.1)	<0.001	<0.001	Comp.> ADHD-; <i>p</i> =0.01 Comp.> ADHD+; <i>p</i> <0.001
Comorbid SUDs [*]	11 (21.6)	14 (21.2)	4 (26.7)	5 (20.0)	8 (17.0)	0.59	0.71	
Full scale IQ at FU41 ^{**}	101.3 (13.7)	108.9 (16.0)	99.3 (13.0)	103.8 (13.1)	111.1 (14.3)	0.01	0.01	Comp.> ADHD-; <i>p</i> =0.08 Comp.> ADHD+; <i>p</i> =0.04
Type of scan						0.09	0.25	
Siemens Allegra	33 (64.7)	52 (78.8)	9 (60.0)	18 (72.0)	38 (80.9)			
Siemens Trio	18 (35.3)	14 (21.2)	6 (40.0)	7 (28.0)	9 (19.1)			

¹ Excluding those (n=19) with ADHD-NOS at FU 41.

^{*} Definite diagnoses, current;

^{**} Data available for 46 probands and 61 comparisons. "ADHD+" and "ADHD-" refer to ADHD probands with persistent and remitted ADHD, respectively. Prob.= Probands; Comp.= Comparisons. LSD=Least Significant Difference. SUDs=Substance Use Disorders.

Table 2

Significant Fractional Anisotropy (FA) clusters between Probands and Comparisons ($p < 0.05$, corrected) *.

Probands with childhood ADHD < Comparisons without childhood ADHD						
Cluster	White matter tract	N voxels	MNI coordinates (peak voxel)			P value
			X	Y	Z	
1	Posterior thalamic radiation L (including optic radiation)	205	-40	-42	-3	3.7 0.02
2a	Retrolenticular part of internal capsule L	94	-40	-35	-2	4.1 0.02
2b	Sagittal stratum L (including inferior longitudinal fasciculus and inferior fronto-occipital fasciculus)	129	-41	-38	-6	3.4 0.02
3	Superior longitudinal fasciculus II R	95	30	-40	34	3.3 0.02
4	Posterior corona radiata R	147	23	-42	36	3.0 0.02
5	Superior corona radiata R	94	21	-15	41	2.8 0.03
Probands with persistent ADHD < Non-ADHD Comparisons						
Cluster	White matter tract	N voxels	MNI coordinates (peak voxel)			P value
			X	Y	Z	
1	Sagittal stratum R (including inferior longitudinal fasciculus and inferior fronto-occipital fasciculus)	266	36	-21	-6	2.8 0.03
2	External capsule R	263	34	3	-10	2.9 0.03
3	Retrolenticular part of internal capsule R	38	40	-32	-3	2.7 0.04
4	Anterior corona radiata R	13	23	28	-4	3.2 0.05
Probands with remitted ADHD < Non-ADHD Comparisons						
Cluster	White matter tract	N voxels	MNI coordinates (peak voxel)			P value
			X	Y	Z	
1	Posterior corona radiata R	123	20	-44	38	4.3 0.02
2	Superior corona radiata R	72	22	-28	39	3.3 0.03

* controlling for age and scanner type