Published in final edited form as:

Curr Top Med Chem. 2011 ; 11(9): 1103–1127.

Design, Synthesis and Interaction at the Vesicular Monoamine Transporter-2 of Lobeline Analogs: Potential Pharmacotherapies for the Treatment of Psychostimulant Abuse

Peter A. Crooks*, Guangrong Zheng, Ashish P. Vartak, John P. Culver, Fang Zheng, David B. Horton, and Linda P. Dwoskin

Department of Pharmaceutical Sciences, College of Pharmacy, University of Kentucky, Lexington, KY 40536-0082, USA

Abstract

The vesicular monoamine transporter-2 (VMAT2) is considered as a new target for the development of novel therapeutics to treat psychostimulant abuse. Current information on the structure, function and role of VMAT2 in psychostimulant abuse are presented. Lobeline, the major alkaloidal constituent of *Lobelia inflata*, interacts with nicotinic receptors and with VMAT2. Numerous studies have shown that lobeline inhibits both the neurochemical and behavioral effects of amphetamine in rodents, and behavioral studies demonstrate that lobeline has potential as a pharmacotherapy for psychostimulant abuse. Systematic structural modification of the lobeline molecule is described with the aim of improving selectivity and affinity for VMAT2 over neuronal nicotinic acetylcholine receptors and other neurotransmitter transporters. This has led to the discovery of more potent and selective ligands for VMAT2. In addition, a computational neural network analysis of the affinity of these lobeline analogs for VMAT2 has been carried out, which provides computational models that have predictive value in the rational design of VMAT2 ligands and is also useful in identifying drug candidates from virtual libraries for subsequent synthesis and evaluation.

Keywords

Vesicular monoamine transporter-2 (VMAT2); psychostimulant abuse; lobeline; lobeline analogs; computational modeling

1. VMAT2

1.1. Introduction

The accumulation and storage of the monoamines serotonin, dopamine (DA), norepinephrine, and histamine in the vesicles of neurons is essential to the regulatory processes governing available cytosolic concentrations of these endogenous neurotransmitters [1]. The vesicular monoamine transporter (VMAT) exists as two distinct isoforms VMAT1 and VMAT2 [2–5]. VMAT2 is expressed in sympathetic postganglionic neurons as well as adult human and monoaminergic neurons of the central nervous system, whereas VMAT1 is primarily expressed in neuroendocrine cells including chromaffin cells of the adrenal medulla and enterochromaffin cells of the intestine [1, 6–10]. However, recent

studies have shown that VMAT1 is also expressed in brain [11]. Both isoforms of VMAT are expressed during the embryonic growth and development phase in mammals [12]. Studies utilizing transfected Chinese hamster ovary (CHO) cells expressing either VMAT1 or VMAT2 revealed differences in both substrate recognition and activity of inhibitors. For example, VMAT2 has greater affinity for the monoamines, and greater sensitivity to inhibition by tetrabenazine (TBZ) compared to VMAT1 [1, 4].

The natural alkaloid reserpine **1**, Fig. (1) and the synthetic compound TBZ **2**, Fig. (1) are both considered classical inhibitors of VMAT2 [13]. Reserpine inhibits monoamine transport into synaptic storage vesicles as well as chromaffin granules by interaction with VMAT [14, 15]. Reserpine binds to the amine recognition site on VMAT2 and TBZ binds to an allosteric site on this protein that is distinct from the substrate binding site [15–17].

1.2. The Role of VMAT2 in Psychostimulant Abuse

The psychostimulant amphetamine and other structurally-related compounds, including methamphetamine, interact with VMAT2 and the DA transporter (DAT) to increase synaptic DA concentrations leading to reward and drug-seeking behavior [18–23]. These psychostimulants inhibit DA uptake into and promote DA release from the synaptic vesicles via its interaction with VMAT2 [24, 25]. Amphetamine enters the vesicle by diffusing across the vesicular membrane or by uptake via VMAT2 with a resulting decrease in pH gradient across the vesicular membrane and a reduction in the free energy available for DA transport into the vesicle [24–27]. Amphetamine accumulating inside the vesicle also competes with monoamine neurotransmitters for available protons, as monoamines are normally found as cationic at the low pH found in the vesicle, but the increase in pH due to the influx of amphetamine results in monoamines reverting to their non-ionized form, resulting in diffusion of neutral mono-amines out of the vesicle [27]. Studies utilizing VMAT2 knockout mice demonstrate reduced drug-seeking behavior compared to wild-type in locomotor response studies that included a array of drugs, including amphetamine and cocaine.[28, 29]. Collectively, these results support the validity of VMAT2 serving as a target for the development of therapeutic agents that have the ability to augment cessation of psychostimulant abuse [30].

1.3. VMAT2 Structure

VMAT2 is a member of the major facilitator superfamily (MFS) of transporters [36]. The computationally predicted molecular structure of VMAT2 is thought to incorporate 12 putative transmembrane domains (TMDs) Fig. (2) based on hydrophobicity plots of the amino acid sequence of the protein [31]. Both the N- and C- termini are cytosolic, and a hydrophobic, N-glycosylated loop is believed to be located between TMDs 1 and 2 Fig. (2) left, projecting into the vesicle lumen. [32].

Structural biology studies have identified specific amino acid residues in the VMAT2 protein that are likely important sites for ligand binding and monoamine transport. Mutagenesis experiments indicate that the negatively charged aspartate 33 residue in TMD1 and serine residues 180 and 183 in TMD3 of VMAT2 are essential for substrate recognition. Substrate-recognition is believed to result interaction of the protonated amino group of the substrate with aspartate 33 and interaction of the catechol or indole hydroxyl groups with serine residues 180 and 183 [32]. Further, lysine 139 in TMD2 and aspartate 427 in TMD11 of VMAT2 form an ion-pair and provide a structural foundation for high affinity substrate recognition [33]. Experiments utilizing chimeras of VMAT1 and VMAT2 show that two regions of the protein, incorporating TMD5 through TMD8 Region B, Fig. (2), and TMD9 through TMD12 Region C, Fig. (2), liaise to confer TBZ and histamine with high affinity at VMAT2. Similar experiments reveal that the region that includes TMD3 and TMD4 Region

A, Fig. (2) confers serotonin affinity, but does not impart affinity for histamine or sensitivity to TBZ [30]. Studies employing photoaffinity labelling of purified VMAT2 from rat brain suggest that TMD1 and TMD10/TMD11 may be adjacent to one another, and likely interact and thereby affecting function [34]. Cysteine mutagenesis and derivatization experiments with human VMAT2 (hVMAT2) have demonstrated that cysteine residues 439, 476, and/or 497, and possibly cysteine 126 and/or 333, are critical for [³H]TBZ binding. Moreover, modification of a combination of three or more of the cysteine residues 126, 333, 439, 476, and 497 alters [³H]TBZ binding characteristics [35]. Furthermore, preincubation with [³H]TBZ, even at low concentrations, effectively blocked subsequent modification at cysteine 439 in TMD 11, suggesting that cysteine 439 is positioned at the TBZ binding site on VMAT2. Several studies provide support for location of the TBZ binding site on VMAT2 at TMD 11, as well as suggest contributions from TMDs 1, 2 and 12 [36]. More recently, comparative modeling studies were conducted utilizing high resolution structures of proteins from the MFS. These studies predict tertiary structures of the rVMAT2 protein allowing for high structural conservation of the MFS proteins, despite low rates of sequence homology. These models also lend further credence to the idea that TMD 2 and 11 are likely involved in the TBZ binding site, and this hypothesis is further supported by recent mutagenesis studies using functionally expressed rVMAT2 in *S. cerevisiae* cells [37].

2. LOBELINE

(–)-Lobeline lobeline, 2*R*,6*S*,10*S*-lobeline, **3**, Fig. (3) is the major alkaloidal constituent of *Lobelia inflata*, which is also known as Indian tobacco. The name “Indian tobacco” most likely originated from the fact that Native Americans smoked the dried leaves of the plants (Millspaugh 1974). In 1838, Proctor prepared a liquid alkaloid extract of the plant and named it lobeline (Millspaugh 1974). Historically, the alkaloidal extract of *L. inflata* has been used as an expectorant, emetic, anti-asthmatic, anti-spasmodic, respiratory stimulant, general muscular relaxant, diaphoretic, diuretic, and stimulant and has been used to treat narcotic overdose. In the plant, lobeline is both the most abundant and the most pharmacologically active constituent of more than 20 structurally-related piperidine alkaloids, which include lobelanine (**4**), nor-lobelanine (**5**), lobelanidine (**6**), and nor-lobelanidine (**7**) Fig. (3) (Felpin and Lebreton 2004). Lobeline was initially isolated (Wieland 1921) and characterized (Wieland, Schopf *et al.* 1925; Wieland and Dragendorff 1929) by Wieland, who subsequently reported a total synthesis to confirm its structure (Wieland and Drishaus 1929; Wieland, Koschara *et al.* 1929). Numerous total syntheses of lobeline have been reported either by the racemic approach (Scheuing and Winterhalder 1929; Schopf and Lehmann 1935) or by the asymmetric approach (Schoepf and Mueller 1965; Compere, Marazano *et al.* 1999; Felpin and Lebreton 2002; Klingler and Sobotta 2006; Birman, Jiang *et al.* 2007).

Both the free base and the hydrochloride and sulfate salt forms of lobeline are stable when in crystalline form. However, in solution, due to the presence of the β-aminoketone moiety in the molecule, lobeline readily undergoes a pH-dependent epimerization at C2 of the piperidine ring to yield a mixture of *cis*-lobeline (**3**) and *trans*-lobeline **8**, Fig. (4) (Compere, Marazano *et al.* 1999; Felpin and Lebreton 2002; Zheng, Dwoskin *et al.* 2004). The epimerization is believed to occur via the transient retro-aza Michael addition product **9** Fig. (4). β-Aminoketone compounds often exhibit stability problems, such as epimerization and elimination, due to their susceptibility to retro-aza Michael addition. Retro-aza Michael addition often occurs under acidic conditions (Vazquez, Galindo *et al.* 2001; Davis, Zhang *et al.* 2005); however, epimerization does not occur in aqueous solutions of lobeline at pH 3.0. Thus, the epimerization process can be terminated by decreasing the pH of the aqueous media to below 3.0, and this ‘quench’ technique has been used to determine the time course and kinetics of lobeline epimerization under a variety of conditions utilizing HPLC as an

analytical methodology for determining epimer ratio (unpublished data). At room temperature, epimerization of lobeline is curtailed when the *cis:trans* ratio is ~65:35 in aqueous solutions. In a chloroform solution, epimerization stops when a *cis:trans* ratio of 46:54 is reached (unpublished data). Interestingly, the *cis:trans* lobeline ratio in human plasma after sublingual administration of lobeline sulfate is about 1:19. Surprisingly, this ratio is reversed to 16:1 *cis:trans* lobeline, in rat plasma after sublingual administration of lobeline sulfate (Crooks *et al.*, unpublished work).

Due to the presence of the β -aminoketone moiety, an additional stability problem associated with lobeline is decomposition via a retro-Mannich reaction which affords ace-tophenone (**10**) and reactive intermediates **11a**, **11b**, and **11c** Fig. (5) in an aqueous solution (Crooks *et al.* 2010). As illustrated in Fig. (5), the nitrogen lone pair must be available for the retro-Mannich reaction to occur, which is consistent with experimental observations, demonstrating that this degradation pathway does not occur in acidic media.

2.1. Lobeline Pharmacology

The pharmacology of lobeline has been reviewed previously (McCurdy, Miller *et al.* 2000; Dwoskin and Crooks 2002; Felpin and Lebreton 2004; Zheng, Dwoskin *et al.* 2006). Lobeline exhibits high affinity for nicotinic acetylcholine receptors (nAChRs) (Court, Perry *et al.* 1994; Damaj, Patrick *et al.* 1997; Flammia, Dukat *et al.* 1999; Miller, Crooks *et al.* 2004), and has many nicotine-like effects, such as tachycardia and hypertension (Korczyn, Bruderma.I *et al.* 1969), nausea (Bevan and Hughes 1966), anxiolytic activity (Brioni, Oneill *et al.* 1993), improvement in learning and memory (Decker, Majchrzak *et al.* 1993), hyperalgesia (Hamann and Martin 1994), and analgesia after intrathecal administration (Damaj, Patrick *et al.* 1997). Consequently, lobeline has been categorized as a nicotinic receptor agonist. However, lobeline has no obvious structural resemblance to nicotine, and structure-activity relationships (SARs) do not suggest a common pharmacophore (Barlow and Johnson 1989). Further, lobeline and nicotine have different effects in both behavioral and neurochemical studies, suggesting that they do not act *via* a common mechanism (Dwoskin and Crooks 2002). Specifically, in contrast to nicotine, lobeline only marginally supports self-administration in mice (Rasmussen and Swedberg 1998) and does not support self-administration in rats (Harrod, Dwoskin *et al.* 2003). In contrast to nicotine, lobeline does not produce conditioned place preference. Also, chronic lobeline treatment does not increase locomotor activity in rats (Fudala and Iwamoto 1986; Stolerma, Garcha *et al.* 1995). Conversely, lobeline attenuates the hyperactivity induced by repeated nicotine administration in rats (Miller, Harrod *et al.* 2003). Furthermore, lobeline inhibits nicotine-evoked [³H]dopamine (DA) overflow from rat striatal slices (IC₅₀ ~1 μ M), suggesting that the alkaloid acts as an antagonist at nAChRs mediating nicotine-evoked DA release (Miller, Crooks *et al.* 2000). Lobeline binds to $\alpha 4\beta 2^*$ nAChRs with high affinity (K_i = 4 nM), and also inhibits nicotine-evoked ⁸⁶Rb⁺ efflux from rat thalamic synaptosomes (IC₅₀ = 0.7 μ M), indicating that lobeline is an antagonist at $\alpha 4\beta 2^*$ nAChRs (Miller, Crooks *et al.* 2000). Additionally, lobeline has also been reported to be an antagonist (IC₅₀ = 8.5 μ M) at human $\alpha 7^*$ nAChRs expressed in *Xenopus* oocytes (Briggs and McKenna 1998). These results suggest that lobeline acts as a potent, but nonselective, nAChR antagonist.

In addition to the interaction of lobeline with nAChRs, lobeline interacts with VMAT2 and DAT. Lobeline inhibits [³H]dihydrotetrabenazine (DTBZ) binding to VMAT2 (IC₅₀ = 0.90 μ M) and inhibits [³H]DA uptake into rat striatal vesicle preparations (IC₅₀ = 0.88 μ M) (Teng, Crooks *et al.* 1997; Teng, Crooks *et al.* 1998). In addition, lobeline inhibits [³H]DA uptake into rat striatal synaptosomes (IC₅₀ = 80 μ M) via DAT, but with 100-fold lower affinity compared to its inhibition of VMAT2 (Teng, Crooks *et al.* 1997). In a VMAT2 and DAT co-expressed cell system, lobeline has been shown to evoke [³H]DA release through

an interaction with VMAT2, but not with DAT (Wilhelm, Johnson *et al.* 2004), which was accompanied by a decrease in methamphetamine-evoked [^3H]DA release (Wilhelm, Johnson *et al.* 2008).

Numerous studies have shown that lobeline inhibits both the neurochemical and behavioral effects of amphetamine in rodents. Lobeline inhibits amphetamine-evoked DA release from superfused rat striatal slices (Miller, Crooks *et al.* 2001) and decreases amphetamine- and methamphetamine-induced hyperactivity in rats and mice (Miller, Crooks *et al.* 2001), and decreases methamphetamine self-administration in rats without acting as a substitute reinforcer (Harrod, Dwoskin *et al.* 2001; Harrod, Dwoskin *et al.* 2003). Pre-treatment with lobeline attenuates the intensity of methamphetamine-induced stereotypy in adolescent mice (Tatsuta, Kitanaka *et al.* 2006). Lobeline also attenuates cocaine self-administration in rats. The effects of lobeline on cocaine-induced hyperactivity in rats is complex. Acutely, lobeline did not decrease the effect of cocaine on hyperactivity. However, repeated administration of lobeline attenuated cocaine-induced hyperactivity and prevented the development of sensitization to cocaine. Interestingly, repeated administration of a lower dose of lobeline (0.3 mg/kg) increased cocaine (10 mg/kg)-induced hyperactivity (Polston, Cunningham *et al.* 2006). These behavioral studies demonstrate that lobeline has potential as a pharmacotherapy for psychostimulant abuse. In addition, since lobeline attenuates methamphetamine-induced decreases in rat striatal VMAT2 protein, this suggests a possible neuroprotective role against methamphetamine toxicity (Eyerman and Yamamoto 2005). Lobeline is currently undergoing Phase 2 clinical evaluation to determine its effectiveness in humans as a therapeutic agent for methamphetamine abuse.

Lobeline has recently been shown to interact with opiate receptors, and has been suggested as a potential therapeutic for treatment of opiate abuse [77].

3. LOBELINE INTERACTION WITH VMAT2

Lobeline is believed to inhibit the rewarding effects of psychostimulants by altering synaptic DA concentration via interactions with VMAT2 and DAT [30]. Since the affinity of lobeline at VMAT2 is 100-fold higher than at DAT and lobeline inhibits amphetamine evoked DA release in the concentration range which inhibits VMAT2, but not DAT, it has been proposed that the interaction of lobeline with VMAT2 is the primary mechanism responsible for lobeline-induced inhibition of the behavioral effects of psychostimulants. Thus, VMAT2 is currently being investigated as a novel target for the development of pharmacotherapies to treat psychostimulant addiction (Dwoskin and Crooks 2002). In this respect, a mechanism of action for lobeline has been proposed, i.e. lobeline interacts with the TBZ binding site on VMAT2 to inhibit dopamine uptake and promote dopamine release from storage vesicles within dopaminergic presynaptic terminals (Dwoskin and Crooks 2002).

4. STRUCTURAL MODIFICATION OF LOBELINE

Lobeline represents a novel structural entity as a VMAT2 ligand even though it is a molecule that interacts with multiple receptors and transporters. Systematic structural modification of lobeline has been carried out with the aim of improving selectivity and affinity for VMAT2 over other neurotransmitter transporters and nAChRs.

4.1. Modification on the 10-Hydroxyl Group of Lobeline

Jones oxidation (Wieland, Koschara *et al.* 1929) and acid-catalyzed dehydration (Ebnoter 1958; Flammia, Dukat *et al.* 1999) of lobeline afforded lobelanine (**4**) and ketoalkene (**12**), respectively (Scheme 1). Compared to lobeline, both compounds **4** and **12** exhibited a robust 110-9280-fold reduction in affinity for the [^3H]nicotine ([^3H]NIC) binding site, probing

$\alpha 4\beta 2^*$ nAChRs, a slight reduction in affinity for the [^3H]methyllycaconitine ([^3H]MLA) binding site, probing $\alpha 7^*$ nAChRs, and a slight increase in affinity for the [^3H]methoxytetrabenazine ([^3H]MTBZ) binding site, probing VMAT2 (Table 1) (Miller, Crooks *et al.* 2004). These results demonstrate that the 10-hydroxyl group is important for $\alpha 4\beta 2^*$ nAChRs interaction, but not for $\alpha 7^*$ nAChR nor VMAT2 interaction.

Acylation or sulfonation of the 10-hydroxyl group of lobeline provided a series of lobeline carboxylic and sulfonic esters which exhibited interesting biological activity (Scheme 2). All the lobeline sulfonate derivatives (**13a–k**), including both aliphatic and aromatic sulfonates, exhibited similar binding profiles to lobeline at $\alpha 4\beta 2^*$, $\alpha 7^*$ nAChRs and VMAT2 (Table 2). However, within the lobeline ester series, aromatic esters (**14a–c**) exhibited 400–4830-fold lower affinity for $\alpha 4\beta 2^*$ nAChRs compared to lobeline, while the aliphatic esters **14d–g** exhibited only 15–125-fold lower affinity for $\alpha 4\beta 2^*$ nAChRs (Table 2). In addition, all the carboxylic esters (**14a–g**) were equipotent to lobeline at VMAT2 (Table 2) [85]. Compounds **13a–13k** and **14a–14g** generally exhibited poor affinity for $\alpha 7$ nAChRs. Thus, esterification of the lobeline molecule with sulphonic acids affords analogs with high affinity for $\alpha 4\beta 2$ nAChRs and comparable affinity to lobeline at VMAT2.

4.2. Modification on the 8-Ketone Moiety in the Lobeline Molecule

The carboxyl group of lobeline was stereoselectively reduced to lobelanidine **6**, Fig. (3) by using LAH (Ebnother 1958) or NaBH_4 , or eliminated to form a series of *des*-keto analogs (**15a–d** and **16a–d**) (Scheme 3) (Zheng, Horton *et al.* 2006). Compound **15a** was prepared by Clemmensen reduction of lobeline (Flammia, Dukat *et al.* 1999). Oxidation of **15a** to compound **17**, an enantiomer of ketoalkene **12**, by Jones oxidation, followed by NaBH_4 reduction yielded a mixture of **15a** and **15b**; this which could be easily separated by silica gel column chromatography. Similarly, NaBH_4 reduction of ketoalkene **12** afforded compounds **15c** and **15d**, which were obtained in an isomerically pure form using silica gel chromatography. Catalytic hydrogenation of the unsaturated compounds **15a**, **15b**, **15c**, and **15d** afforded the corresponding hydrolic compounds **16a**, **16b**, **16c**, and **16d**, respectively.

Compared to lobeline, the dihydroxyl compound, lobelanidine (**6**) exhibited 200-fold lower affinity for $\alpha 4\beta 2^*$ nAChRs, slightly higher affinity at $\alpha 7^*$ nAChRs, and 5-fold lower affinity at VMAT2 (Table 1) (Miller, Crooks *et al.* 2004). *Des*-keto lobeline analogs displayed diminished affinity at $\alpha 4\beta 2^*$ and $\alpha 7^*$ nAChRs, except for compounds **15c** and **16c**, which had slightly higher affinity than lobeline at $\alpha 7^*$ nAChRs. The *des*-keto analogs had similar affinities for VMAT2 compared to lobeline (i.e., within one-order of magnitude); with compound **15b** exhibiting the highest affinity of the series ($K_i = 0.59 \mu\text{M}$) (Table 3) (Zheng, Horton *et al.* 2006).

4.3. DES-Oxygen Lobeline Analogs

From the above structure-activity relationships, both the hydroxyl group and the carboxyl group of lobeline clearly are important for the high affinity for $\alpha 4\beta 2^*$ nAChRs, but not important for VMAT2 interaction. A series of fully defunctionalized lobeline analogs, in which both of these two functional groups were removed from the molecule were synthesized and evaluated for affinity at VMAT2 assessed by inhibition of [^3H]DTBZ binding to rat synaptic vesicle membranes (Zheng, Dwoskin *et al.* 2005). The synthesis of the defunctionalized analogs **18a–c**, **19a–c**, and **20a–d** is illustrated in Scheme 4. Briefly, *cis*-lobeline (**3**) was treated with K_2CO_3 /methanol to yield a mixture of the two C2 epimers, **3** and **8**, which was then converted into a mixture of lobelanidine isomers via reduction with NaBH_4 . Treatment of the lobelanidine isomeric mixture with 85% phosphoric acid afforded a separable mixture of *meso*-transdiene (MTD, **18a**), and (–)-*trans*-transdiene ((–)-TTD, **18b**). The (+)-TTD (**18c**) isomer was separated from a mixture of **18a** and **18c**, obtained via

successive oxidation and C6-epimerization of **15a**. Catalytic hydrogenation of **18a**, **18b**, and **18c** afforded lobeline (**19a**), (–)-2S, 6S-*trans*-lobeline ((–)-*trans*-lobeline, **19b**), and (+)-2R, 6R-*trans*-lobeline ((+)-*trans*-lobeline, **19c**), respectively. Simultaneous reduction of the double bond and the carbonyl group in the ketoalkene mixture (**12**) or the ketoalkene mixture (**17**) by catalytic hydrogenation each afforded compounds **21** (as a mixture of 4 isomers) and **22** (as a mixture of 4 isomers). Dehydration of **21** and **22** with 85% H₃PO₄, respectively, followed by silica gel column chromatography separation, yielded compounds **20a**, **20b**, **20c**, and **20d**, respectively.

Results from the pharmacological analysis of this series of defunctionalized analogs are shown in Table 4. MTD (**18a**), the fully deoxygenated, unsaturated *meso*-compound, has no affinity at either $\alpha 4\beta 2^*$ or $\alpha 7^*$ nAChRs, but had comparable affinity at VMAT2 ($K_i = 9.88 \mu\text{M}$) to lobeline. Reduction of the double-bonds in MTD afforded lobeline (**19a**, $K_i = 0.97 \mu\text{M}$ at VMAT2), which was more selective for VMAT2 but only 3-fold more potent than lobeline at VMAT2. Reduction of either the C7 or the C9 double bond in MTD afforded enantiomers **20c** and **20a**, respectively, both of which had no affinity at either $\alpha 4\beta 2^*$ or $\alpha 7^*$ nAChRs, but both isomers had 4-fold higher affinity at VMAT2 compared to MTD. The *trans* analogs of MTD (**18b** and **18c**) had no affinity at either $\alpha 4\beta 2^*$ or $\alpha 7^*$ nAChRs and exhibited comparable affinity for VMAT2 with respect to MTD. Surprisingly, no difference in affinity between these two *trans* enantiomers was observed at VMAT2. These results indicate a surprising lack of stereochemical sensitivity at the ligand recognition site at VMAT2 with these lobeline analogs because changes in the stereochemistry of the substituents at C2 and C6 of the piperidine ring from *cis* to *trans* within the MTD series (i.e., **18a** to **18b** and **18c**) had no effect on affinity for VMAT2. The *trans* analogs of lobeline, **19b** and **19c**, exhibited a 5–6-fold decrease in affinity at VMAT2 compared to lobeline (**19a**). Again, the *trans* enantiomers **19b** and **19c** exhibited comparable affinities at VMAT2. Taken together, these data indicate that the VMAT2 binding site can accept major stereochemical changes to the MTD and lobeline molecules at the C2 and C6 piperidino ring carbons.

Similarly, the enantiomers **20b** and **20d** showed low affinity at $\alpha 4\beta 2^*$ or $\alpha 7^*$ nAChRs, but interestingly exhibited very different affinities for VMAT2. That is compound **20b** had a similar affinity for VMAT2 compared to MTD, whereas **20d** exhibited low affinity for VMAT2. These results suggest a somewhat complex structure-activity relationship within these structurally-related analogs, in that the VMAT2 recognition site appears to accommodate three mono double bond isomers, with different stereochemistry at C2 and C6, but does not recognize one specific stereochemical form **20d**, which has the 2*R*, 6*R* stereochemical configuration.

4.4. Fragmented Lobeline Analogs

(–)-Sedamine (**29**) and compound **32**, which represent the hydroxyl containing fragment and the keto containing fragment of lobeline, were synthesized to determine if the structure of the whole lobeline molecule is required for potent VMAT2 interaction. (–)-Sedamine was synthesized via a key ring opening reaction of the TBS protected lobeline (**24**) to afford compound **25**. The double bond in compound **25** was then cleaved and the resulting aldehyde was reduced by NaBH₄ to afford compound **26**. Bromination of compound **26**, followed by removal of the Troc group, cyclization, and removal of the TBS group, yielded the final product (Scheme 5) (Zheng, Dwoskin *et al.* 2004). Compound **32** was synthesized via initial Sonogashira cross-coupling of 2-bromopyridine and phenylacetylene to form compound **30**. Compound **30** was then hydrolyzed to the keto compound **31** with sulfuric acid. Hydrogenation of compound **31** over Adams' catalyst followed by *N*-methylation afforded the final product (Scheme 6) (Zheng, Dwoskin *et al.* 2005). Compounds **29** and **32**

exhibited low affinity at $\alpha 4\beta 2^*$ nAChRs, $\alpha 7^*$ nAChRs and VMAT2 (Table 4) (Zheng, Dwoskin *et al.* 2005), suggesting that both the C2 and C6 side chains on the piperidine ring of lobeline are essential for VMAT2 affinity.

5. STRUCTURAL MODIFICATION OF LOBELANE

Systematic structural modification of the lobeline molecule provided lobelane (**19a**, Scheme 4), which exhibited high affinity (K_i of 0.97 μ M) for the DTBZ binding site, on VMAT2, with low affinity for the $\alpha 4\beta 2^*$ and $\alpha 7^*$ nAChRs (Zheng, Dwoskin *et al.* 2005). Compared to lobeline, lobelane appears to be slightly more potent and selective at VMAT2, and lobelane is also effective in decreasing methamphetamine self-administration, suggesting that it may have potential as a novel treatment for methamphetamine abuse (Neugebauer, Harrod *et al.* 2007). Thus, systematic structural modifications based on lobelane molecule have been carried out in order to identify more selective and potent analogs at VMAT2, as well as analogs with improved physicochemical properties compared to lobelane.

5.1. Fragmented Lobelane Analogs

Similar to the lobeline series, structural fragments of MTD and lobelane (compounds **33** and **34**, respectively; Scheme 6), were also prepared in order to determine if the whole molecule was required for potent VMAT2 interaction. Compounds **33** and **34** were synthesized from compound **32** as shown in Scheme 6 (Zheng, Dwoskin *et al.* 2005). Interestingly, compound **34**, in which one of the piperidine ring side chains of lobelane has been removed, showed 3-fold higher affinity at $\alpha 4\beta 2^*$ nAChRs compared to lobelane, and exhibited no affinity at $\alpha 7^*$. Compound **33**, in which a *trans*-double bond has been introduced into the molecule, showed no affinity at either $\alpha 4\beta 2^*$ or $\alpha 7^*$ nAChRs. Both of the fragmented compounds (**33** and **34**) exhibited low affinity at VMAT2 (Table 4) (Zheng, Dwoskin *et al.* 2005). These results suggest that both the C2 and C6 side chains of the piperidine ring of MTD and/or lobelane are essential for VMAT2 affinity and selectivity. Compounds **35** and **36** Fig. (6), two other fragmented compounds in which the piperidine ring has been truncated, were also synthesized (Zheng, Dwoskin *et al.* 2005). Compounds **35** and **36** exhibited no affinity at either $\alpha 4\beta 2^*$ or $\alpha 7^*$ nAChRs and had affinity at VMAT2 similar to that of lobelane (Table 4) (Zheng, Dwoskin *et al.* 2005). These results indicate that the nitrogen and two phenyl rings of lobelane are likely the essential pharmacophoric elements for recognition by the VMAT2 binding site. Interestingly, two open ring compounds, **37** and **38** Fig. (6), which are by-products from the hydrogenation reaction of **18b** and **18c** to form **19b** and **19c**, respectively, showed comparable affinity for VMAT2 compared to lobelane (Table 4) (Zheng, Dwoskin *et al.* 2005).

5.2. Modification of the Phenyl Rings of Lobelane

A series of lobelane analogs, in which the two phenyl rings were replaced by substituted phenyl rings or other aromatic ring systems, was synthesized (Scheme 7 and 8). The syntheses of analogs **40–60b** (as shown in Table 5) were initiated from condensation reactions between 2,6-lutidine and appropriate aldehydes to form the conjugated products of general structure **39**. Compounds of general structure **39** were reduced to the corresponding *nor*-compounds by Adams' catalyzed hydrogenation. *N*-Methylation was then carried out using NaCNBH₃/(CH₂O)_n to afford the corresponding *N*-methyl compounds (Scheme 7) (Zheng, Dwoskin *et al.* 2005; Zheng, Dwoskin *et al.* 2005). All the compounds synthesized in this series (Table 5) exhibited low affinity for both $\alpha 4\beta 2^*$ and $\alpha 7^*$ nAChRs. With the exception of compounds **51a** and **54a**, which exhibited low affinity at VMAT2, all other compounds exhibited comparable affinity for VMAT2 relative to lobelane, i.e., within about one-order of magnitude. The 3-fluoro, 2-methoxy, and 3-methoxy, substituted analogs (**42b**, **44b**, and **45b**, respectively) had the highest affinity (K_i = 0.43–0.57 μ M) at VMAT2 among

the *mono*-substituted analogs series (Zheng, Dwoskin *et al.* 2005). Among the di-substituted series, 3,4-methylenedioxy, 3,5-difluoro, 2,3-dimethoxy, and 2,5-dimethoxy substituted analogs (**47b**, **56b**, **59b**, and **60b**, respectively) showed the highest affinity ($K_i = 0.29$ – 0.62 μ M) at VMAT2 (Culver *et al.*, 2010). Interestingly, there were no marked differences in VMAT2 affinity between analogs bearing either electron donating or withdrawing substituents. Also, the *N*-methyl analogs generally had higher affinity at VMAT2 compared to their corresponding *nor*-analogs.

The syntheses of analogs **63**–**67**, in which the two phenyl rings of lobelane have been replaced by two hetero-aromatic rings, were initiated from condensation reaction between *N*-methylated 2,6-lutidine (**61**) and appropriate aldehydes to form compound **62**. Adams' catalyzed hydrogenation of compound **62** afforded the final products (Scheme 8) (Vartak *et al.*, 2010). These analogs displayed affinities that were 5 to 10-fold lower at VMAT2 compared to lobelane, with the exception of 2-quinolinyll compound **65**, which showed similar affinity to lobelane at VMAT2 (Table 5; Vartak *et al.*, 2010). It would appear that a neutral aryl substituent is highly preferred by the putative binding site on VMAT2, since only the quinolyl compounds **65** and **66**, which can potentially remain uncharged at physiological pH, retained the affinity of lobelane toward VMAT2.

Two asymmetric lobelane analogs **70** and **71**, which incorporate non-identical aromatic moieties at the C2, C6 of the piperidine ring, were also prepared. A Wittig reaction of 2-bromo-6-pyridincarboxaldehyde with benzyl ylid, followed by Sonogashira coupling with 1-ethynynaphthalene or 3-ethynylanisole furnished the key intermediate, compound **69**. Adams' catalyzed hydrogenation of compound **69** followed by *N*-methylation provided the final products (Scheme 9; Crooks *et al.*, 2010). Both compounds **70** and **71** exhibited similar affinity at VMAT2 when compared to that of lobelane, and slightly lower affinity at VMAT2 compared to their corresponding symmetric *bis*-1-naphthyl analog (**53b**) and *bis*-3-anisolyll analog (**45b**) (Table 5).

5.3. Isomerized Lobelane Analogs

To further understand the importance of the juxtaposition and stereochemistry of the piperidino C2 and C6 substituents in the lobelane molecule, and the importance of the intramolecular distance between the nitrogen atom and the phenyl rings of lobelane, compounds **74a**–**76a**, **74b**–**76b**, **74c**–**76c**, **79a**, **79b**, **80a**, and **80b** were synthesized (Zheng, Dwoskin *et al.* 2005). In these compounds, the C2, C6 phenethyl groups of lobelane were moved to the N1, C2; N1, C3; N1, C4; C3, C5; or C2, C4 positions around the piperidine ring. Analogues with one side chain attached to the nitrogen atom of the piperidine ring, i.e., **74a**–**76a**, **74b**–**76b**, and **74c**–**76c**, were synthesized from initial condensation reaction of 2, 3, or 4-picoline with benzaldehyde. Adams' reduction of the resulting conjugated products **72a**–**72c**, followed by *N*-alkylation afforded compounds **74a**–**74c** and **75a**–**75c**. Reduction of the double bond in compounds **74a**, **74b**, and **74c** by catalytic hydrogenation afforded compounds **76a**, **76b**, and **76c**, respectively (Scheme 10). The synthesis of compounds **80a**, and **80b** was similar to the synthesis of *nor*-lobelane and lobelane, except that 2,4-lutidine was used as a starting material instead of 2,6-lutidine (Scheme 11). Similar procedures were also employed for the synthesis of compounds **79a** and **79b**. In the condensation reaction, *N*-benzylidene-4-chloroaniline was used as an activated form of benzaldehyde, due to the low activity of the 3-methyl group of the pyridinyl moiety (Scheme 11). Most of these compounds showed low affinity at either $\alpha_4\beta_2^*$ or α_7^* nAChRs. All of these analogs retained affinity at VMAT2, but were 2–10-fold less potent than lobelane (Table 6). Analogues bearing a side chain at the piperidino C3 position, i.e., **74b**, **75b**, **76b**, **79a** and **79b**, exhibited 3–10-fold lower potency at VMAT2 compared with lobelane. Thus, surprisingly, the position of the piperidine nitrogen atom relative to the C2 and C6 side chains does not

appear to be critical for interaction with the VMAT2 binding site. The most potent compounds at VMAT2 in this series were **74c** and **80a** ($K_i = 1.87$ and $1.36 \mu\text{M}$, respectively), in which the two side chains are located on the piperidino N1, C4 or C2, C4 positions. These results also indicate that the VMAT2 binding site can tolerate changes in distance between the piperidine nitrogen and the two phenyl rings.

Based on the interesting data obtained on compound **74c**, a series of N1, C4 lobelane analogs, in which one of the phenyl rings was replaced by an anisoyl or dimethoxyphenyl ring, was synthesized (Scheme 12). The 2-methoxy and 3-methoxy substituted analogs, **84a** and **84b**, exhibited similar affinity ($K_i = 0.30$ and $0.63 \mu\text{M}$, respectively) at VMAT2 compared to lobelane and **74c** (Table 6) (Culver *et al.*, 2010).

A series of isomerized MTD analogs, including **86a–88a**, **86b–88b** and **89–93**, in which the C2, C6 side chains in MTD were moved to C3, C5, while the carbon numbers between nitrogen and the aromatic rings were retained, was also synthesized. These analogs generally displayed similar affinity at VMAT2 compared with MTD. Compounds **86c** and **86d**, two isomerized lobelane analogs containing a linker unit that is one carbon shorter between the piperidine and the phenyl rings than compound **79b**, displayed a 3-fold increase in affinity at VMAT2 compared with compound **79b** (Table 6) (Horton *et al.*, submitted).

5.4. Modification on the two Ethylene Groups of Lobelane

To determine the optimal bridge lengths between the piperidine ring and each of the two phenyl rings in lobelane, a series of lobelane homologs, in which the two ethylene groups connecting the two phenyl rings to the C-2 and C-6 positions of the piperidine ring in lobelane have been replaced by carbon linkers of a range of different length (0–3 carbons), i.e., combinations of 0,0-, 0,1-, 0,2-, 0,3-, 1,1-, 1,2-, 1,3-, 2,3-, and 3,3-carbons on each side of piperidine ring, were synthesized as ligands for VMAT2 (Table 7)[90]. The removal of the two-carbon bridge on both sides of the piperidine ring of lobelane to generate compound **94** eliminated binding at VMAT2. Compounds **95b**, **96b**, and **97b**, which contain no carbon bridge on one side, and a one-, two, or three-carbon bridge, respectively, on the other side of the piperidine ring, had 14–40-fold lower affinity at VMAT2 compared to lobelane. The reduction of one methylene unit on each side of the two-carbon bridge of lobelane to afford compound **98b** resulted in a 20-fold decrease in VMAT2 affinity compared to lobelane. Increasing the length of one side of the bridge in compound **98b** by one or two methylene units to afford compounds **99b** and **100b** afforded an improvement in affinity ($K_i = 2.77$ and $1.62 \mu\text{M}$, respectively), at VMAT2 compared to **98b**, similar to lobelane. The addition of a methylene unit to one side or both sides of the piperidine ring in lobelane to produce longer homologs, compounds **101b** and **102b**, respectively, had no effect on VMAT2 affinity compared to lobelane. Compound **101b** ($K_i = 0.88 \mu\text{M}$) had similar potency to lobelane. Compounds **95a–102a**, exhibited affinities at VMAT2 within an order-of-magnitude compared to their corresponding methylated analogs **95b–102b**, which is consistent with the SAR observed in the previous lobelane analog series [90]. These results indicate that retention of binding affinity at VMAT2 found for lobelane, structurally-related homologs require linker lengths between the central piperidine and two phenyl rings to be no shorter than one methylene group on one side of the piperidine ring and no shorter than two methylene groups on the other side of the piperidine ring. The optimal methylene unit lengths are a combination of two and three methylene units.

5.5. Modification on the *n*-Methyl Group of Lobelane

A series of modifications on the nitrogen atom of the piperidine ring of lobelane was carried out (Fig. (7)). *Nor*-lobelane (**40**, $K_i = 2.31 \mu\text{M}$), *N*-ethyl lobelane (**103a**, $K_i = 3.41 \mu\text{M}$) or *N*-*n*-propyl lobelane (**103b**, $K_i = 1.87 \mu\text{M}$) exhibited similar affinity for VMAT2 compared to

lobelane (Zheng, Dwoskin *et al.* 2005). Replacing the *N*-methyl group in lobelane with a dimethylhydrazine group afforded compound **104**, which exhibited no affinity at VMAT2. Replacing the *N*-methyl group in lobelane with an acetonitrile, an ethylamine, or an ethyldimethylamine group afforded compound **105**, **106**, or **107**, respectively, which exhibited 5–20-fold lower affinity at VMAT2 compared to lobelane. Interestingly, compound **108**, which contains a 1,2(*R*)-dihydroxypropyl group on the piperidine nitrogen, displayed similar affinity ($K_i = 0.56 \mu\text{M}$) at VMAT2 (Table 8) compared to lobelane (Zheng *et al.*, 2010). Importantly, **108** is a more water soluble analog of lobelane, with obvious advantages for clinical development.

5.6. Modification on the Piperidine Ring of Lobelane

5.6.1. Introduction of an Oxygen Atom at the C4 of Piperidine Ring—By introduction of an oxygen atom (ketone or hydroxyl group) at the C4 position of the piperidine ring of lobelane, analogs **109–113** incorporate some structural features of the TBZ and DTBZ molecules, which were thought to improve affinity at VMAT2 Fig. (8). Interestingly, all the analogs in this series displayed more than 10-fold lower affinity at VMAT2 when compared to lobelane, indicating that lobelane and TBZ may bind differently at VMAT2 (Table 8) (Zheng *et al.*, submitted).

5.6.2. Replacing the Piperidine Ring in Lobelane with a Pyrrolidine Ring—Subtle conformational changes caused by contraction of the central piperidine ring to the 5-membered pyrrolidine can be expected to affect the 3-dimensional projection of the C2, C6 substituents, as well as the direction of the *N*-substituent. Such compounds were prepared via the addition of Grignard reagents to Katritzky's bicyclic hemiaminal (**114**; Scheme 14), followed by elaboration of the separable diastereomers (**115** and **116**) into the requisite pyrrolidine compounds (**117–120**) (Katritzky, Cui *et al.* 1999). Enantiomers of compounds **115**, **116**, **119** and **120** (compounds **121**, **122**, **123** and **124**, respectively) were similarly obtained from the epimer of compound **114**.

The *nor*-analog (**117**) exhibited an order-of-magnitude higher affinity at VMAT2 than its *N*-methylated analog (**118**; Table 9), which is in sharp contrast to the higher affinity of the *N*-methyl analog usually observed in the corresponding piperidine (lobelane) series. *N*-Substitutions as large as a phenethyl substituent (compounds **115**, **116**, **121**, **122**) did not decrease binding affinity within this series of pyrrolidine compounds [92]. The relative stereochemistry of substitutions at the C2 and C5 positions also seems to be of little consequence with regard to affinity for VMAT. The overall affinity of the pyrrolidine compounds, with the exception of **118**, does not vary substantially within the series. Clearly, replacement of the central piperidine ring with a pyrrolidine ring does not afford compounds with enhanced affinity when compared to lobelane.

5.6.3. Replacing the Piperidine Ring in Lobelane with a Piperizine Ring—Further modification on the piperidine ring of lobelane has been carried out by replacing the piperidine ring with a piperizine ring (compounds **125** and **126**), which can be achieved by employment of a similar approach as described in Scheme 7 (Scheme 15). Compounds **125** and **126** were 36- and 6-fold less potent at VMAT2, respectively, compared to lobelane (Table 9).

5.6.4. Replacing the Piperidine Ring in Lobelane with a Tropane Ring—Compounds **128a**, **128b**, **128c**, **129a**, **129b**, **129c**, and **131**, in which the more conformationally-restricted tropane ring was substituted for the piperidine ring in lobelane, were synthesized (Zheng, Dwoskin *et al.* 2005). The general methodology for the preparation of these analogs is summarized in Scheme 16. Compounds **127a–127c** and **130**

were obtained by aldol condensation of tropinone with benzaldehyde, 4-fluorobenzaldehyde, 4-methoxybenzaldehyde or *trans*-cinnamaldehyde. Clemmensen reduction of compounds **127a**, **127b** or **127c** afforded compounds **128a**, **128b** or **128c**, respectively. Compounds **129a**, **129b**, and **129c** were each obtained by catalytic hydrogenation of the corresponding precursor molecules, compounds **128a**, **128b** and **128c**. Clemmensen reduction of compound **130**, followed by catalytic hydrogenation afforded compound **131**.

The trop-2-ene compounds **128a** and **128b** exhibited similar affinity ($K_i = 1.30 \mu\text{M}$ and $1.38 \mu\text{M}$, respectively) at VMAT2 to that of lobelane (Table 9). Interestingly, compounds **129a** and **129b**, which are saturated analogs of **128a** and **128b**, respectively, had no affinity (both with $K_i > 100 \mu\text{M}$) for VMAT2. These results suggest that the double bond in compounds **128a** and **128b** is important for VMAT2 binding. In this respect, the double bond affects both the configuration of the tropane ring and the orientations of two side chains at C-2 and C-4. In particular, the molecules of **128a** and **128b** are more extended than the corresponding reduced analogs, i.e., compounds **129a** and **129b**. Compound **128c**, in which an electron donating methoxy group was introduced into the *para* position of each of the two phenyl rings, exhibited lower potency ($K_i = 4.80 \mu\text{M}$) at VMAT2 compared to either lobelane, compound **128a** or compound **128b**. Surprisingly, in contrast to compounds **129a** and **129b**, when the double bond in compound **128c** was reduced to afford the tropane analog, compound **129c**, the affinity of compound **129c** ($K_i = 3.88 \mu\text{M}$) at VMAT2 was retained. Thus, the electron density in the phenyl rings may influence binding at VMAT2. However, compound **129c** is more extended than either **129a** or **129b**, because of the presence of the *para*-methoxy groups. Thus, full extension of the molecule in these tropane analogs also may be important for VMAT2 binding. Compound **131**, which incorporates an increased distance between the phenyl rings and the C2, C4 atoms of the tropane ring compared to compound **129a**, had a K_i value of $3.95 \mu\text{M}$ at VMAT2. Similar to compound **129c**, compound **131** is a more extended molecule than compound **129a**, which may be the reason for the observed increase in VMAT2 affinity (Zheng, Dwoskin *et al.* 2005).

6. COMPUTATIONAL NEURAL NETWORK ANALYSIS OF THE AFFINITY OF LOBELINE ANALOGS FOR VMAT2

Towards the discovery of more potent, novel and selective lobeline analogs as ligands for VMAT2, we considered computational modeling as a valuable aid in drug design and optimization. In this respect, the nature of the interaction of these novel ligands with binding site(s) on VMAT2 is not known due to the lack of a crystal structure of this protein. Thus, a structure-based drug design approach is not available. On the other hand, artificial neural network (ANN) approaches, such as back-propagation network to data analysis, has received much attention over the last decade. This artificial system emulates the function of the brain, in which a very high number of information-processing neurons are interconnected and are known for their ability to model a wide set of functions, including linear and non-linear functions, without knowing in advance the analytic forms[94]. By use of an ANN that simulates the processes of brain neurons, one can train the network to solve many types of problems. The rapid advancement of computing systems in the past 20 years led to the success of this machine-learning approach in various engineering, business, and medical applications. One of the best known and widely used applications is NETalk [95], an application of an ANN for machine reading of text. Most recently, the ANN approach also has been applied in a variety of biomedical areas, which includes analysis of appendicitis [96], cancer imaging extraction and classification [97], and AIDS research and therapy [98]. Relevant to this chapter, this approach has also been used in drug design and discovery, in such areas as prediction of molecular bioactivity [99–102]. Pharmaceutical applications utilizing this approach include pharmaceutical production development [103],

pharmacodynamic modeling [104], and mapping dose-effect relationships to pharmacological response. [105].

In our recent study, the ANN approach was used to build a quantitative SAR (QSAR) model on a set of 104 TBZ and lobeline analogs with known affinity for VMAT2 [103]. This is the first QSAR modeling study that has addressed the interaction of a small library of ligands at the VMAT2 binding site. Most of the entities in the library were lobeline analogs. TBZ and its analogs were introduced into the dataset as they represent ligands having high affinity for VMAT2. The affinity of the lobeline analogs were measured using [³H]DTBZ as a probe, thus the same bioactivity endpoint was used for the entire set of 104 compounds in this QSAR study. With high activity analogs in a training set, the trained model would have a better chance of identifying the features that lead to high affinity novel analogs. The goal of the QSAR work were: (i) to extract the relevant descriptors to establish the QSAR of the library of ligands, (ii) to establish the high predictive power of ANN modeling of this library of compounds, and (iii) to develop insights regarding the relationship between the descriptors of the compounds and their affinity for VMAT2.

The QSAR study revealed that while the linear partial least squares model built from the same dataset is predictive, the fully interconnected three-layer neural network model trained with the back-propagation procedure was superior learning the correct association between a set of relevant descriptors of compounds and the $\log(1/K_i)$ for VMAT2 [106]. The optimal ANN architecture was determined as 11:3:1, i.e. eleven input, three hidden and one output neuron(s). The eleven related descriptors (shown in Table 10) constructed the input vector for the input neurons of the ANN. These descriptors were composed of topological, geometrical, GETAWAY, aromaticity, and WHIM descriptors. Evaluation of the contributions of the descriptors to the QSAR reflected the importance of atomic distribution in the molecules, molecular size, and steric effects of the ligand molecules when interacting with their target binding site on VMAT2. Statistical analysis for the trained ANN indicated that the computed activities were in excellent agreement with the experimentally observed values ($r^2 = 0.91$, $\text{rmsd} = 0.225$; predictive $q^2 = 0.82$, $\text{loormsd} = 0.316$). The generated models were further tested by use of an external prediction set of 15 molecules. The nonlinear ANN model had $r^2 = 0.93$ and root-mean-square errors of 0.282 compared with the experimentally measured activities of the test set Fig. (9). Furthermore, the stability test of the model with regard to data division was performed. The dataset of 104 molecules was randomly divided into two sets; one set of 89 molecules for training and leave-one-out validation, and the other set of 15 molecules for external testing. Among each division, the dissimilarity of one test set from the test set of another division was greater than 90%. This process was randomly repeated five times. The same eleven descriptors, learning epochs (200,000) and ANN architecture were used to train and test the ANN models. The data in Table 11 shows that the trained 11-3-1 ANN models are stable with regard to the data division, indicating that the generated model is reliable and predictive.

The developed models are expected to be valuable in the rational design of chemical modifications of the next-generation of VMAT2 ligands in order to identify the most likely candidates for synthesis and discovery of new lead compounds. One of the most important attributes of ANN is the ability to make predictions on new molecules with accuracy similar to that within the learning set. With a system that was theoretically validated to estimate binding affinity (K_i values) for known compounds with reasonable accuracy, we predicted the K_i values of various new (virtual) compounds designed by intuitive modification of the structures of the compounds in the ANN training set. Insights obtained from the theoretical predictions included the importance of the aromatic methoxy groups, and the optimal number of these groups in the molecule, which were determined to be a factor leading to high affinity for VMAT2. For example, for molecules with the general structure shown in

Table 12, the $IC_{50}(\text{exp.})$ values of four molecules (ID 1–4) that had been synthesized and evaluated experimentally (out of 104 new structures utilized for model generation), were inserted into the training set for the ANN to ‘learn’. Compounds 5–16 were novel molecules in which binding affinities were predicted by the generated models. According to the results predicted by the trained model, it was shown that when $R_0 = H$, if R_1, R_2, R_3 and $R_4 = H$ (compound 1), the computationally fitted binding affinity for the molecule was $4.90 \mu\text{M}$. However, if one of the R_1, R_2, R_3 and/or R_4 groups was CH_3O , the affinity of the compound was predicted to be improved, and if two of the R_1, R_2, R_3 and R_4 groups were CH_3O , the affinity of the compound would be improved by one-order-of magnitude compared to the predicted binding affinity for compound 1. When R_0 was OH , and R_1, R_2, R_3 and R_4 all were H (compound 9), the binding affinity of the molecule was predicted to be $4.62 \mu\text{M}$. Compounds 10–12, which contain one aromatic methoxy substituent, are predicted to have their affinity improved similar to compounds 2–4. Compounds 13–16, which contain two aromatic methoxy substituents, are predicted to exhibit a two-fold increase in binding affinity at VMAT2 compared to those of compounds 5–8. Compounds 6–9 were subsequently synthesized and their binding affinities were measured experimentally. The experimental results for compounds 7–9 shown in Table 12 are consistent with the computational predictions, but those for compound 6 are not. Clearly, the position of the methoxy group in the aromatic ring has a significant influence on K_i value; however, the model is not sensitive to this variable, since the information relating the importance of methoxy to binding activity was determined by topological descriptors. In this situation, the 3D effect of the methoxy group was not reflected. However, in other cases the generated model does predict the difference in binding affinity of a molecule with different configuration. Table 13 shows that compounds **132–135**, which constitute 4 different stereoisomers of the same structure, have different binding affinities for VMAT2. The experiment measurements for compounds **132** and **135** were consistent with theoretical predictions [106].

The accuracy of a QSAR model should be determined by statistical analysis with a much greater number (>500) of analogs than the limited number in our practical application of the currently generated models. For the VMAT2 binding assay, three compounds (**6, 7** and **8**, Table 12) were among those predicted to be lead compounds (i.e. binding affinity was predicted to be less than $1 \mu\text{M}$) based on computational analysis, and were subsequently synthesized. Two of these compounds (**7** and **8**) had experimental binding affinities ($0.62 \mu\text{M}$ and $0.56 \mu\text{M}$, respectively) that were close to the original best leads ($0.43 \mu\text{M}$ and $0.58 \mu\text{M}$). Three compounds (**132** and **135**, Table 13, and **9**, Table 12) were predicted to be inactive from computational models, and indeed were inactive when evaluated experimentally. These results indicate that the generated computational models have predictive value in the rational design of chemical modifications of VMAT2 ligands, and can be utilized to identify the most likely candidates for synthesis and discovery of new lead compounds. A larger data base needs to be examined to form conclusions with statistical significance.

References

1. Erickson JD, Schafer MK, Bonner TI, Eiden LE, Weihe E. Distinct pharmacological properties and distribution in neurons and endocrine cells of isoforms of the human vesicular monoamine transporter. *Proc Natl Acad Sci.* 1996; 93:5166–5171. [PubMed: 8643547]
2. Erickson JD, Eiden L. Functional identification and molecular cloning of a human brain vesicle monoamine transporter. *J Neurochem.* 1993; 61:2314–2317. [PubMed: 8245983]
3. Erickson JD, Eiden LE, Hoffman BJ. Expression cloning of a reserpine-sensitive vesicular monoamine transporter. *Proc Natl Acad Sci.* 1992; 89:10993–10997. [PubMed: 1438304]

4. Zheng G, Dwoskin LP, Crooks PA. Vesicular monoamine transporter-2: Role as a novel target for drug development. *AAPS J.* 2006; 8(4):E682–E692. [PubMed: 17233532]
5. Peter D, Jimenez J, Liu Y, Kim J, Edwards RH. The chromaffin granule and synaptic vesicle amine transporters differ in substrate recognition and sensitivity to inhibitors. *J Biol Chem.* 1994; 269:7231–7237. [PubMed: 8125935]
6. Peter D, Liu Y, Sternini C, de Giorgio R, Brecha N, Edwards RH. Differential expression of two vesicular monoamine transporters. *J Neurosci.* 1995; 15:6179–6188. [PubMed: 7666200]
7. Weihe E, Schafer MK, Erickson JD, Eiden LE, et al. Localization of vesicular monoamine transporter isoforms (VMAT1 and VMAT2) to endocrine cells neurons in rat. *J Mol Neurosci.* 1994; 5:149–164. [PubMed: 7654518]
8. Krejci E, Gasnier B, Botton D, Isambert MF, Sagné C, Gagnon J, Massoulié J, Henry JP. Expression and regulation of the bovine vesicular monoamine transporter gene. *FEBS Lett.* 1993; 335:27–32. [PubMed: 7902299]
9. Stern-Bach Y, Keen JN, Bejerano M, Steiner-Mordoch S, Wallach M, Findlay JB, Schuldiner S. Homology of a vesicular amine transporter to a gene conferring resistance to 1-methyl-4-phenylpyridinium. *Proc Natl Acad Sci USA.* 1992; 89:9730–9733. [PubMed: 1357668]
10. Howell M, Shirvan A, Stern-Bach Y, Steiner-Mordoch S, Strasser JE, Dean GE, Schuldiner S. Cloning and functional expression of a tetrabenazine sensitive vesicular monoamine transporter from bovine chromaffin granules. *FEBS Lett.* 1994; 338:16–22. [PubMed: 8307150]
11. Lohoff FW, Dahl JP, Ferraro TN, Arnold SE, Gallinat J, Sander T, Berrettini WH. Variations in the vesicular mono-amine transporter 1 gene (VMAT1/SLC18A1) are associated with bipolar I disorder. *Neuropsychopharmacology.* 2006; 31:2739–2747. [PubMed: 16936705]
12. Hansson SR, Mezey E, Hoffman BJ. Ontogeny of vesicular monoamine transporter mRNAs VMAT1 and VMAT2, II. Expression in neural crest derivatives and their target sites in the rat. *Dev Brain Res.* 1998; 110:159–174. [PubMed: 9733958]
13. Pletscher A. Effect of neuroleptics and other drugs on monoamine uptake by membranes of adrenal chromaffin granules. *Br J Pharmacol.* 1977; 59:419–424. [PubMed: 584682]
14. Darchen F, Scherman D, Henry JP. Reserpine binding to bovine chromaffin granule membranes. *Mol Pharmacol.* 1984; 25:113–122. [PubMed: 6708929]
15. Scherman D, Henry JP. Radioligands of the vesicular monoamine transporter and their use as markers of monoamine storage vesicles. *Biochem Pharmacol.* 1989; 38:2395–2404. [PubMed: 2667522]
16. Henry JP, Scherman D. Radioligands of the vesicular monoamine transporter and their use as markers of monoamine storage vesicles. *Biochem Pharmacol.* 1989; 38:2395–2404. [PubMed: 2667522]
17. Scherman D, Jaudon P, Henry JP. Characterization of the monoamine carrier of chromaffin granule membrane by binding of [2-³H]dihydrotetrabenazine. *Proc Natl Acad Sci USA.* 1983; 80:584–588. [PubMed: 6572908]
18. Amara SG, Sonders MS. Neurotransmitter transporters as molecular targets for addictive drugs. *Drug Alcohol Depend.* 1998; 51:87–96. [PubMed: 9716932]
19. Wise RA, Bozarth MA. A psychostimulant theory of addiction. *Psychol Rev.* 1987; 94:469–492. [PubMed: 3317472]
20. Koob GF. Neural mechanisms of drug reinforcement. *Ann N Y Acad Sci.* 1992; 654:171–191. [PubMed: 1632582]
21. Fleckenstein AE, Hanson GR. Impact of psychostimulants on vesicular monoamine transporter function. *Eur J Pharmacol.* 2003; 479:283–289. [PubMed: 14612158]
22. Riddle EL, Fleckenstein AE, Hanson GR. Role of monoamine transporters in mediating psychostimulant effects. *AAPS J.* 2005; 7:E847–E851. [PubMed: 16594636]
23. Brown JM, Hanson GR, Fleckenstein AE. Regulation of the vesicular monoamine transporter-2: A novel mechanism for cocaine and other psychostimulants. *J Pharmacol Exp Ther.* 2001; 296:762–767. [PubMed: 11181904]
24. Sulzer D, Maidment NT, Rayport S. Amphetamine and other weak bases act to promote reverse transport of dopamine in ventral midbrain neurons. *J Neurochem.* 1993; 60:527–535. [PubMed: 8419534]

25. Sulzer D, Chem TK, Lau YY, Kristensen H, Rayport S, Ewing A. Amphetamine redistributes dopamine from synaptic vesicles to the cytosol and promotes reverse transport. *J Neurosci.* 1995; 15:4102–4108. [PubMed: 7751968]
26. Johnson RG. Accumulation of biological amines into chromaffin granules: A model for hormone and neurotransmitter transport. *Physiol Rev.* 1988; 68:232–307. [PubMed: 2892215]
27. Sulzer D, Rayport S. Amphetamine and other psychostimulants reduce pH gradients in midbrain dopaminergic neurons and chromaffin granules: A mechanism of action. *Neuron.* 1990; 5:797–808. [PubMed: 2268433]
28. Takahashi N, Miner LL, Sora I, Ujike H, Revay RS, Kostic V, Jackson-Lewis V, Przedborski S, Uhl GR. VMAT2 knockout mice: heterozygotes display reduced amphetamine-conditioned reward, enhanced amphetamine locomotion and enhanced MPTP toxicity. *Proc Natl Acad Sci USA.* 1997; 94:9938–9943. [PubMed: 9275230]
29. Wang Y, et al. Knockout of the vesicular monoamine transporter 2 gene results in neonatal dath and supersensitivity to cocaine and amphetamine. *Neuron.* 1997; 19:1285–1296. [PubMed: 9427251]
30. Dwoskin LP, Crooks PA. A Novel mechanism of action and potential use for lobeline as a treatment for psychostimulant abuse. *Biochem Pharmacol.* 2002; 63:89–98. [PubMed: 11841781]
31. Peter D, Vu T, Edwards RH. Chimeric vesicular monoamine transporters identify structural domains that influence substrate affinity and sensitivity to tetrabenazine. *J Biol Chem.* 1996; 271:2979–2986. [PubMed: 8621690]
32. Yelin, R.; Schuldiner, S. Vesicular neurotransmitter transporters: Pharmacology, biochemistry, and molecular analysis. In: Reith, M., editor. *Neurotransmitter Transporters; Structure, Function, and Regulation.* 2. Humana Press; Totowa: 2002. p. 313-354.
33. Merickel A, et al. Identification of residues involved in substrate recognition by a vesicular monoamine transporter. *J Biol Chem.* 1995; 270:25798–25804. [PubMed: 7592763]
34. Merickel A, Kaback HR, Edwards RH. Charged residues in transmembrane domains II and XI of a vesicular monoamine transporter form a chard pair that promotes high affinity substrate recognition. *J Biol Chem.* 1997; 272:5403–5408. [PubMed: 9038139]
35. Sievert MK, Ruoho AE. Peptide mapping of the [¹²⁵I]iodoazidoketanserin and [¹²⁵I]2-N-[(3'-iodo-4'-azidophenyl)propionyl]tetrabenazine binding sites for the synaptic vesicle monoamine transporter. *J Biol Chem.* 1997; 272:26049–26055. [PubMed: 9325342]
36. Thiriot DS, Ruoho AE. Mutagenesis and deriviatization of human vesicle monoamine transporter 2 (VMAT2) cysteines identifies transporter domains involved in tetrabenazine binding and substrate transport. *J Biol Chem.* 2001; 276:27304–27315. [PubMed: 11375404]
37. Vardy E, Arkin IT, Gottschalk KE, Kaback HR, Schuldiner S. Structural conservation in the major facilitator superfamily as revealed by comparative modeling. *Protein Sci.* 2004; 13:1982–1840.
38. Gros Y, Schuldiner S. Directed evolution reveals hidden properties of VMAT, a neurotransmitter transporter. *J Biol Chem.* 2009 Paper in Press: Manuscript M109.081216.
39. Millspaugh, CF. *American Medicinal Plants: An Illustrated and Descriptive Guide to Plants Indigenous to and Naturalized in the United States Which Are Used in Medicine.* Dover; New York: 1974. *Lobelia Inflata*; p. 385-388.
40. Felpin FX, Lebreton J. History, chemistry and biology of alkaloids from *Lobelia inflata*. *Tetrahedron.* 2004; 60:10127–10153.
41. Wieland H. Alkaloids of the *Lobelia* plant. I. (Preliminary Communication.). *Ber Dtsch Chem Ges.* 1921; 54B:1784–1788.
42. Wieland H, Schopf C, Hermesen W. *Lobelia* Alkaloids. Ii. *Ann Chem.* 1925; 444:40–68.
43. Wieland H, Dragendorff O. *Lobelia* alkaloids. Iii. Constitution of the *Lobelia* alkaloids. *Liebigs Ann Chem.* 1929; 473:83–101.
44. Wieland H, Drishaus I. *Lobelia* Alkaloids. Iv. Synthesis of *Lobelia* alkaloids. *Liebigs Ann Chem.* 1929; 473:102–118.
45. Wieland H, Koschara W, Dane E. *Lobelia* alkaloids. V. Several bases accompanying lobeline and the reciprocal relations of the *Lobelia* alkaloids. *Liebigs Ann Chem.* 1929; 473:118–126.
46. Scheuing G, Winterhalder L. Synthesis of *Lobelia* alkaloids. *Justus Liebigs Ann Chem.* 1929; 473:126–136.

47. Schopf C, Lehmann G. Syntheses and transformations of natural products under physiological conditions (model experiments on the question of the biogenesis of natural products). Iv. Syntheses of tropinone, pseudopelletierine, lobelanine and related alkaloids under physiological conditions. *Liebigs Ann Chem.* 1935; 518:1–37.
48. Schoepf C, Mueller E. Absolute configuration of (–)-lobeline and of its reduction products. *Liebigs Ann Chem.* 1965; 687:241–250.
49. Compere D, Marazano C, Das BC. Enantioselective access to Lobelia alkaloids. *J Org Chem.* 1999; 64:4528–4532.
50. Felpin FX, Lebreton J. A highly stereoselective asymmetric synthesis of (–)-lobeline and (–)-sedamine. *J Org Chem.* 2002; 67:9192–9199. [PubMed: 12492320]
51. Klingler, F-D.; Sobotta, R. US. Boehringer Ingelheim Pharma GmbH & Co; KG, Germany: 2006. p. 4
52. Birman VB, Jiang H, Li X. Enantioselective synthesis of lobeline via nonenzymatic desymmetrization. *Org Lett.* 2007; 9:3237–3240. [PubMed: 17658751]
53. Zheng G, Dwoskin LP, Crooks PA. Indirect trapping of the retroconjugate addition reaction intermediate involved in the epimerization of lobeline: Application to the synthesis of (–)-sedamine. *J Org Chem.* 2004; 69:8514–8517. [PubMed: 15549833]
54. Vazquez E, Galindo A, Gnecco D, Bernes S, Teran JL, Enriquez RG. Unexpected retro-Michael reaction of (–)-(1's,4as,8ar)- and (+)-(1's,4ar,8as)-4a-ethyl-1-(1-phenylethyl)octahydroquinolin-7-ones. *Tetrahedron: Asymmetry.* 2001; 12:3209–3211.
55. Davis FA, Zhang J, Li Y, Xu H, DeBrosse C. Asymmetric synthesis of 2,4,5-trisubstituted piperidines from sulfinimine-derived D-amino B-ketoesters. formal synthesis of pseudodistomin B triacetate. *J Org Chem.* 2005; 70:5413–5419. [PubMed: 15989321]
56. McCurdy CR, Miller RL, Beach JW. Lobeline: A natural product with high affinity for neuronal nicotinic receptors and a vast potential for use in neurological disorders. *Biolog Active Nat Prod.* 2000:151–162.
57. Court JA, Perry EK, Spurdin D, Lloyd S, Gillespie JI, Whiting P, Barlow R. Comparison of the binding of nicotinic agonists to receptors from human and rat cerebral cortex and from chick brain (alpha(4)beta(2)) transfected into mouse fibroblasts with ion-channel activity. *Brain Res.* 1994; 667:118–122. [PubMed: 7534607]
58. Damaj MI, Patrick GS, Creasy KR, Martin BR. Pharmacology of lobeline, a nicotinic receptor ligand. *J Pharmacol Exp Ther.* 1997; 282:410–419. [PubMed: 9223582]
59. Flammia D, Dukat M, Damaj MI, Martin B, Glennon RA. Lobeline: Structure-affinity investigation of nicotinic acetylcholine receptor binding. *J Med Chem.* 1999; 42:3726–3731. [PubMed: 10479304]
60. Miller DK, Crooks PA, Zheng GR, Grinevich VP, Norrholm SD, Dwoskin LP. Lobeline analogs with enhanced affinity and selectivity for plasmalemma and vesicular monoamine transporters. *J Pharmacol Exp Ther.* 2004; 310:1035–1045. [PubMed: 15121762]
61. Korczyn AD, Bruderma I, Braun K. Cardiovascular effects of lobeline. *Archives Internationales De Pharmacodyn Et De Therap.* 1969; 182:370–375.
62. Bevan JA, Hughes T. Inhibition of gastric contraction following stimulation of intrathoracic sensory endings. *Arch Int Pharmacodyn Ther.* 1966; 161:334–342. [PubMed: 5926875]
63. Brioni JD, Oneill AB, Kim DJB, Decker MW. Nicotinic receptor agonists exhibit anxiolytic-like effects on the elevated plus-maze test. *Eur J Pharmacol.* 1993; 238:1–8. [PubMed: 8405072]
64. Decker MW, Majchrzak MJ, Arneric SP. Effects of lobeline, a nicotinic receptor agonist, on learning and memory. *Pharmacol Biochem Behav.* 1993; 45:571–576. [PubMed: 8332618]
65. Hamann SR, Martin WR. Hyperalgesic and analgesic actions of morphine, U50-488, naltrexone, and (–)-lobeline in the rat brainstem. *Pharmacol Biochem Behav.* 1994; 47:197–201. [PubMed: 8115422]
66. Barlow RB, Johnson O. Relations between structure and nicotine-like activity - X-ray crystal-structure analysis of (–)-cytisine and (–)-lobeline hydrochloride and a comparison with (–)-nicotine and other nicotine-like compounds. *Br J Pharmacol.* 1989; 98:799–808.

67. Rasmussen T, Swedberg MDB. Reinforcing effects of nicotinic compounds: Intravenous self-administration in drug-naïve mice. *Pharmacol Biochem Behav.* 1998; 60:567–573. [PubMed: 9632242]
68. Harrod SB, Dwoskin LP, Green TA, Gehrke BJ, Bardo MT. Lobeline does not serve as a reinforcer in rats. *Psychopharmacol.* 2003; 165:397–404.
69. Fudala PJ, Iwamoto ET. Further studies on nicotine-induced conditioned place preference in the rat. *Pharmacol Biochem Behav.* 1986; 25:1041–1049. [PubMed: 3786357]
70. Stolerman IP, Garcha HS, Mirza NR. Dissociations between the locomotor stimulant and depressant effects of nicotinic agonists in rats. *Psychopharmacology.* 1995; 117:430–437. [PubMed: 7604144]
71. Miller DK, Harrod SB, Green TA, Wong MY, Bardo MT, Dwoskin LP. Lobeline attenuates locomotor stimulation induced by repeated nicotine administration in rats. *Pharmacol Biochem Behav.* 2003; 74:279–286. [PubMed: 12479946]
72. Miller DK, Crooks PA, Dwoskin LP. Lobeline inhibits nicotine-evoked [³H]dopamine overflow from rat striatal slices and nicotine-evoked ⁸⁶Rb⁽⁺⁾ efflux from thalamic synaptosomes. *Neuropharmacology.* 2000; 39:2654–2662. [PubMed: 11044735]
73. Briggs CA, McKenna DG. Activation and inhibition of the human alpha 7 nicotinic acetylcholine receptor by agonists. *Neuropharmacology.* 1998; 37:1095–1102. [PubMed: 9833639]
74. Teng LH, Crooks PA, Sonsalla PK, Dwoskin LP. Lobeline and nicotine evoke [³H]overflow from rat striatal slices preloaded with [³H]dopamine: Differential inhibition of synaptosomal and vesicular [³H]dopamine uptake. *J Pharmacol Exp Ther.* 1997; 280:1432–1444. [PubMed: 9067333]
75. Teng L, Crooks PA, Dwoskin LP. Lobeline displaces [³H]dihydrotetrabenazine binding and releases [³H]dopamine from rat striatal synaptic vesicles: Comparison with D-amphetamine. *J Neurochem.* 1998; 71:258–265. [PubMed: 9648873]
76. Wilhelm CJ, Johnson RA, Lysko PG, Eshleman AJ, Janowsky A. Effects of methamphetamine and lobeline on vesicular monoamine and dopamine transporter-mediated dopamine release in a cotransfected model system. *J Pharmacol Exp Ther.* 2004; 310:1142–1151. [PubMed: 15102929]
77. Wilhelm CJ, Johnson RA, Eshleman AJ, Janowsky A. Lobeline effects on tonic and methamphetamine-induced dopamine release. *Biochem Pharmacol.* 2008; 75:1411–1415. [PubMed: 18191815]
78. Miller DK, Lever JR, Rodvelt KR, Baskett JA, Will MJ, Kracke GR. Lobeline, a potential pharmacotherapy for drug addiction, binds to mu opioid receptors and diminishes the effects of opioid receptor agonists. *Drug Alcohol Depend.* 2007; 89:282–291. [PubMed: 17368966]
79. Miller DK, Crooks PA, Teng LH, Witkin JM, Munzar P, Goldberg SR, Acri JB, Dwoskin LP. Lobeline inhibits the neurochemical and behavioral effects of amphetamine. *J Pharmacol Exper Ther.* 2001; 296:1023–1034. [PubMed: 11181937]
80. Harrod SB, Dwoskin LP, Crooks PA, Klebaur JE, Bardo MT. Lobeline attenuates D-methamphetamine self-administration in rats. *J Pharmacol Exp Ther.* 2001; 298:172–179. [PubMed: 11408539]
81. Tatsuta T, Kitanaka N, Kitanaka J, Morita Y, Takemura M. Lobeline attenuates methamphetamine-induced stereotypy in adolescent mice. *Neurochem Res.* 2006; 31:1359–1369. [PubMed: 17053968]
82. Neugebauer NM, Dwoskin LP, Bardo MT. Lobeline attenuates cocaine self-administration in rats. *Behav Pharmacol.* 2007; 18:S26–S26.
83. Polston JE, Cunningham CS, Rodvelt KR, Miller DK. Lobeline augments and inhibits cocaine-induced hyperactivity in rats. *Life Sci.* 2006; 79:981–990. [PubMed: 16765386]
84. Eyerman DJ, Yamamoto BK. Lobeline attenuates methamphetamine-induced changes in vesicular monoamine transporter 2 immunoreactivity and monoamine depletions in the striatum. *J Pharmacol Exp Ther.* 2005; 312:160–169. [PubMed: 15331654]
85. Ebnöther A. Alkaloids. IX. Mutarotation of lobeline. cis-trans isomers in the Lobelia alkaloid series. *Helv Chim Acta.* 1958; 41:386–396.

86. Hojahmat M, Horton DB, Norrholm SD, Miller DK, Grinevich VP, Deaciuc AG, Dwoskin LP, Crooks PA. Lobeline esters as novel ligands for neuronal nicotinic acetylcholine receptors and neurotransmitter transporters. *Bioorg Med Chem Lett*. 2010; 18:640–649.
87. Zheng G, Dwoskin LP, Deaciuc AG, Norrholm SD, Crooks PA. Defunctionalized lobeline analogues: Structure-activity of novel ligands for the vesicular monoamine transporter. *J Med Chem*. 2005; 48:5551–5560. [PubMed: 16107155]
88. Zheng G, Horton DB, Deaciuc AG, Dwoskin LP, Crooks PA. Des-keto lobeline analogs with increased potency and selectivity at dopamine and serotonin transporters. *Bioorg Med Chem Lett*. 2006; 16:5018–5021. [PubMed: 16905316]
89. Zheng G, Dwoskin LP, Deaciuc AG, Zhu J, Jones MD, Crooks PA. Lobelane analogues as novel ligands for the vesicular monoamine transporter-2. *Bioorg Med Chem*. 2005; 13:3899–3909. [PubMed: 15911306]
90. Neugebauer NM, Harrod SB, Stairs DJ, Crooks PA, Dwoskin LP, Bardo MT. Lobelane decreases methamphetamine self-administration in rats. *Eur J Pharmacol*. 2007; 571:33–38. [PubMed: 17612524]
91. Zheng G, Dwoskin LP, Deaciuc AG, Crooks PA. Synthesis and evaluation of a series of homologues of lobelane at the vesicular monoamine transporter-2. *Bioorg Med Chem*. 2008; 18:6509–6512.
92. Katritzky AR, Cui XL, Yang B, Steel PJ. Asymmetric syntheses of 2-substituted and 2,5-disubstituted pyrrolidines from (3*s*,5*r*,7*ar*)-5-(benzotriazol-1-yl)-3-phenyl[2,1-*b*]oxazopyrrolidine. *J Org Chem*. 1999; 64:1979–1985. [PubMed: 11674292]
93. Varkat AP, Nickell JR, Chagkutip J, Dwoskin LP, Crooks PA. Pyrrolidine Analogs of lobeline: relationship of Affinity for the dihydrotetrabenazine binding site with function of the vesicular monoamine transporter (VMAT2). *J Med Chem*. 2009; 52:7878–7882. [PubMed: 19691331]
94. Zheng G, Dwoskin LP, Deaciuc AG, Crooks PA. Synthesis and evaluation of a series of tropane analogues as novel vesicular monoamine transporter-2 ligands. *Bioorg Med Chem Lett*. 2005; 15:4463–4466. [PubMed: 16112864]
95. Rich, E.; Knight, K. *Artificial Intelligence*. 2. McGraw-Hill; New York: 1991.
96. Sejnowski TJ, Rosenberg CR. *Complex systems*. 1987; 1:145.
97. Shah DR, Dev VR, Brown CA. The utilization of a neural network system in the diagnosis of appendicitis. *M D Comput: Comput Med Pract*. 1999; 16:65–67.
98. Seker H, Odetayo MO, Petrovic D, Naguib RNG, Bartoli C, Alasio L, Lakshmi MS, Sherbet GV. Assessment of nodal involvement and survival analysis in breast cancer patients using image cytometric data: statistical, neural network and fuzzy approaches. *Anticancer Res*. 2002; 22:433–438. [PubMed: 12017328]
99. Sardari S, Sardari D. Applications of artificial neural networks in AIDS research and therapy. *Curr Pharm Des*. 2002; 8:659–670. [PubMed: 11945163]
100. Ochoa C, Chana A, Stud M. Applications of neural networks in the medicinal chemistry field. *Curr Med Chem– Central Nervous System Agents*. 2001; 1:247–256.
101. Kaiser KLE. *Quant Struct-Act Relat*. 2003; 22:1.
102. Zheng F, Bayram E, Sumithran SP, Ayers JT, Zhan CG, Schmitt JD, Dwoskin LP, Crooks PA. QSAR modeling of *mono*- and *bis*-quaternary ammonium salts that act as antagonists at neuronal nicotinic acetylcholine receptors mediating dopamine release. *Bioorg Med Chem*. 2006; 14:3017–3037. [PubMed: 16431111]
103. Katritzky AR, Pacureanu LM, Dobchev DA, Fara DC, Duchowicz PR, Karelson M. QSAR modeling of the inhibition of glycogen synthase kinase-3. *Bioorg Med Chem*. 2006; 14:4987–5002. [PubMed: 16650999]
104. Huuskonen J, Salo M, Taskinen J. Neural network modeling for estimation of the aqueous solubility of structurally related drugs. *J Pharm Sci*. 1997; 86:450–454. [PubMed: 9109047]
105. Li H, Yap CW, Xue Y, Li ZR, Ung CY, Han LY, Chen YZ. Statistical learning approach for predicting specific pharmacodynamic, pharmacokinetic, or toxicological properties of pharmaceutical agents. *Drug Dev Res*. 2005; 66:245–259.

106. Mager DE, Shirey JD, Cox D, Fitzgerald DJ, Abernethy DR. Mapping the dose-effect relationship of orbofiban from sparse data with an artificial neural network. *J Pharm Sci.* 2005; 94:2475–2486. [PubMed: 16200613]
107. Zheng F, Zheng GR, Deaciuc AG, Zhan CG, Dwoskin LP, Crooks PA. Computational neural network analysis of the affinity of lobeline and tetrabenazine analogs for the vesicular monoamine transporter-2. *Bioorg Med Chem.* 2007; 15:2975–2992. [PubMed: 17331733]

Fig. 1.
Chemical structures of reserpine (**1**) and tetrabenazine (**2**).

Fig. 2.
VMAT2 Predicted Secondary Structure (Copyright permission requested, need to login to JBC, create account, costs \$86.25. **NEED COPYRIGHT AUTHORIZATION**) Peter, D. *et al.* J. Biol. Chem. 1996; 271: 2979–2986

Fig. 3.
Chemical structures of lobeline and other structurally-related alkaloids in *L. inflata*.

Fig. 4.
Lobeline epimerization.

Fig. 5.
Retro-Mannich reaction of lobeline.

Fig. 6.
Structures of compounds **35**, **36**, **37**, and **38**.

Fig. 7.
Structures of compounds **103a–b**, and **104–108**.

- 109** 2,6-trans, R = H, X,Y = O
110 2,6-trans, R = CH₃, X,Y = O
111 2,6-trans, R = H, X = OH, Y = H
112 2,6-cis, R = H, X,Y = O
113 2,6-cis, R = H, X = OH, Y = H

Fig. 8.
Structures of compounds **109–113**.

Fig. 9.
The calculated versus experimental activity data for the training (shown in squares), LOO cross-validation (shown in triangles) and test set runs (shown in diamonds) for the best NN11-3-1 QSAR model with VMAT2 ligands. The solid line represents a perfect correlation.

Scheme 1.
Synthesis of lobelanine and ketoalkene.

Scheme 2.
Synthesis of lobeline esters and sulfonates.

Scheme 3.
Synthesis of *des*-keto analogs of lobeline.

Scheme 4.
Synthesis of fully defunctionalized lobeline analogs.

Scheme 5.
Synthesis of (-)-sedamine.

Scheme 6.
Synthesis of compounds **32**, **33**, and **34**.

Scheme 7.
Synthesis of compounds **40–60b** in Table 5.

Scheme 8. Synthesis of compounds **63–68** in Table 5.

Scheme 9. Synthesis of compounds **70** and **71**.

Scheme 10.
Synthesis of compounds **74a–74c**, **75a–75c**, and **76a–76c**.

Scheme 11.
Synthesis of compounds **79a**, **79b**, **80a**, and **80b**.

Scheme 12.
Synthesis of compounds **83a-d**, and **84a-d**.

Scheme 13.

Synthesis of compounds **86a–d**, **87a**, **87b**, **88a**, **88b** and **89–93**.

Scheme 14.
 Synthesis of compounds **115**–**124**.

Scheme 15.
Synthesis of compounds **125** and **126**.

Scheme 16.
Synthesis of compounds **128a–c**, **129a–c**, and **131**.

Table 1

$\alpha 4\beta 2^*$, $\alpha 7^*$ nAChR, and VMAT2 Binding Affinity of Compounds 3, 4, 6, and 12.

Compd	K_i , $\mu\text{M} \pm \text{SEM}$		
	[^3H]NIC Binding $\alpha 4\beta 2^*$ nAChR ^a	[^3H]MLA Binding $\alpha 7^*$ nAChR ^a	[^3H]MTBZ Binding VMAT2 ^a
3	0.004	11.6	5.46
4	37.1	22.7	2.41
6	3.10	3.29	25.9
12	0.44	34.2	1.35

^a n = 3–6/compound/assay

Table 2 $\alpha 4\beta 2^*$, $\alpha 7^*$ nAChR, and VMAT2 Binding Affinity of Lobeline Carboxylic and Sulfonic Esters.

Compd	K_i , $\mu\text{M} \pm \text{SEM}$		
	[^3H]NIC Binding $\alpha 4\beta 2^*$ nAChRs ^a	[^3H]MLA Binding $\alpha 7^*$ nAChRs ^a	[^3H]DTBZ Binding VMAT2 ^a
3	0.004 $\pm$ 0.0001	6.26 $\pm$ 1.30	2.76 $\pm$ 0.64
13a	0.005 $\pm$ 0.001	18.0 $\pm$ 4.56	10.8 $\pm$ 4.64
13b	0.008 $\pm$ 0.001	34.8 $\pm$ 11.3	3.50 $\pm$ 0.80
13c	0.016 $\pm$ 0.001	> 100	3.24 $\pm$ 0.28
13d	0.012 $\pm$ 0.001	24.2 $\pm$ 1.59	3.00 $\pm$ 0.48
13e	0.013 $\pm$ 0.002	53.8 $\pm$ 10.4	4.23 $\pm$ 1.08
13f	0.017 $\pm$ 0.002	> 100	4.12 $\pm$ 1.15
13g	0.010 $\pm$ 0.001	18.5 $\pm$ 3.06	2.48 $\pm$ 0.33
13h	0.013 $\pm$ 0.001	46.6 $\pm$ 5.22	3.98 $\pm$ 0.74
13i	0.015 $\pm$ 0.002	47.7 $\pm$ 7.03	2.09 $\pm$ 0.30
13j	0.011 $\pm$ 0.001	41.0 $\pm$ 15.4	1.98 $\pm$ 0.65
13k	0.016 $\pm$ 0.006	25.6 $\pm$ 3.12	2.69 $\pm$ 0.63
14a	1.62 $\pm$ 0.30	> 100	1.76 $\pm$ 0.54
14b	2.87 $\pm$ 0.52	> 100	1.53 $\pm$ 0.38
14c	19.3 $\pm$ 8.80	18.4 $\pm$ 0.42	2.98 $\pm$ 0.21
14d	0.07 $\pm$ 0.01	> 100	8.05 $\pm$ 1.06
14e	0.15 $\pm$ 0.04	8.86 $\pm$ 2.36	4.75 $\pm$ 0.92
14f	0.50 $\pm$ 0.08	> 100	3.38 $\pm$ 0.98
14g	0.06 $\pm$ 0.005	> 100	2.61 $\pm$ 0.52

^a n = 3–6/compound/assay

Table 3 $\alpha 4\beta 2^*$, $\alpha 7^*$ nAChR, and VMAT2 Binding Affinity of *des*-keto Lobeline Analogs

Compd	K_i , $\mu\text{M} \pm \text{SEM}$		
	$[^3\text{H}]\text{NIC}$ Binding $\alpha 4\beta 2^*$ nAChR ^a	$[^3\text{H}]\text{MLA}$ Binding $\alpha 7^*$ nAChR ^a	$[^3\text{H}]\text{DTBZ}$ Binding VMAT2 ^a
15a	9.75 $\pm$ 0.91	> 100	6.44 $\pm$ 0.54
15b	> 100	> 100	0.59 $\pm$ 0.15
15c	4.19 $\pm$ 0.80	1.70 $\pm$ 0.32	5.16 $\pm$ 0.30
15d	> 100	> 100	6.06 $\pm$ 0.45
16a	1.77 $\pm$ 0.61	39.3 $\pm$ 12.9	3.09 $\pm$ 0.41
16b	> 100	> 100	6.60 $\pm$ 2.96
16c	2.36 $\pm$ 0.18	1.21 $\pm$ 0.09	1.98 $\pm$ 0.31
16d	33.6 $\pm$ 8.54	> 100	3.01 $\pm$ 0.44

^a n = 3–6/compound/assay

Table 4

$\alpha 4\beta 2^*$, $\alpha 7^*$ nAChR, and VMAT2 Binding Affinity of Compounds 18a–c, 19a–c, 20a–d, 29, and 32–38.

Compd	$K_i, \mu\text{M} \pm \text{SEM}$		
	$[^3\text{H}]\text{NIC}$ binding $\alpha 4\beta 2^*$ nAChR ^a	$[^3\text{H}]\text{MLA}$ binding $\alpha 7^*$ nAChR ^a	$[^3\text{H}]\text{DTBZ}$ binding VMAT2 ^a
3	0.004 $\pm$ 0.000	6.26 $\pm$ 1.30	2.76 $\pm$ 0.64
18a	11.6 $\pm$ 2.01	>100	9.88 $\pm$ 2.22
18b	>100	>100	19.4 $\pm$ 1.25
18c	>100	>100	7.09 $\pm$ 2.42
19a	14.9 $\pm$ 1.67	26.0 $\pm$ 6.57	0.97 $\pm$ 0.19
19b	>100	25.3 $\pm$ 4.27	5.32 $\pm$ 0.45
19c	>100	40.0 $\pm$ 14.1	6.46 $\pm$ 1.70
20a	>100	>100	2.50 $\pm$ 0.23
20b	>100	55.1 $\pm$ 13.3	5.27 $\pm$ 1.32
20c	>100	>100	2.67 $\pm$ 0.56
20d	>100	25.2 $\pm$ 3.00	> 100
29	> 100	> 100	> 100
32	25.9 $\pm$ 1.62	> 100	> 100
33	> 100	> 100	> 100
34	4.52 $\pm$ 0.58	> 100	> 100
35	> 100	> 100	2.37 $\pm$ 0.55
36	> 100	> 100	3.07 $\pm$ 1.72
37	>100	>100	5.21 $\pm$ 1.23
38	>100	>100	3.96 $\pm$ 0.80

^a n = 3–6/compound/assay

Table 5

$\alpha 4\beta 2^*$, $\alpha 7^*$ nAChR, and VMAT2 Binding Affinity of Compounds 41–60, 63–67, 70 and 71.

Compd	Ar (Het)	R	K_i , $\mu\text{M} \pm \text{SEM}$		
			$[^3\text{H}]\text{NIC}$ binding $\alpha 4\beta 2^*$ nAChR ^a	$[^3\text{H}]\text{MLA}$ binding $\alpha 7^*$ nAChR ^a	$[^3\text{H}]\text{DTBZ}$ binding VMAT2 ^a
lobelane	phenyl	CH ₃	14.9 $\pm$ 1.67	26.0 $\pm$ 6.57	0.97 $\pm$ 0.19
nor-lobelane	phenyl	H	> 100	> 100	2.31 $\pm$ 0.21
41a	2-fluorophenyl	H	> 100	> 100	1.60 $\pm$ 0.10
41b	2-fluorophenyl	CH ₃	> 100	> 100	1.07 $\pm$ 0.07
42a	3-fluorophenyl	H	> 100	49.2 $\pm$ 21.3	1.60 $\pm$ 0.08
42b	3-fluorophenyl	CH ₃	> 100	16.6 $\pm$ 3.47	0.57 $\pm$ 0.07
43a	4-fluorophenyl	H	> 100	> 100	1.25 $\pm$ 0.08
43b	4-fluorophenyl	CH ₃	> 100	> 100	0.98 $\pm$ 0.31
44a	2-anisoyl	H	> 100	> 100	1.87 $\pm$ 0.19
44b	2-anisoyl	CH ₃	> 100	29.7 $\pm$ 7.73	0.58 $\pm$ 0.04
45a	3-anisoyl	H	> 100	> 100	1.67 $\pm$ 0.07
45b	3-anisoyl	CH ₃	> 100	25.1 $\pm$ 2.23	0.43 $\pm$ 0.03
46a	4-anisoyl	H	> 100	> 100	3.32 $\pm$ 0.82
46b	4-anisoyl	CH ₃	> 100	> 100	1.73 $\pm$ 0.20
47a	3,4-methylenedioxy-phenyl	H	> 100	> 100	2.34 $\pm$ 0.16
47b	3,4-methylenedioxy-phenyl	CH ₃	> 100	18.2 $\pm$ 6.87	0.52 $\pm$ 0.25
48a	4-methylphenyl	H	> 100	> 100	3.23 $\pm$ 0.10
48b	4-methylphenyl	CH ₃	> 100	> 100	4.36 $\pm$ 0.18
49a	3-trifluoromethylphenyl	H	> 100	> 100	9.90 $\pm$ 2.01
49b	3-trifluoromethylphenyl	CH ₃	> 100	> 100	1.51 $\pm$ 0.07
50a	2,4-dichlorophenyl	H	> 100	> 100	1.32 $\pm$ 0.18
50b	2,4-dichlorophenyl	CH ₃	> 100	> 100	1.04 $\pm$ 0.06
51a	4-phenylphenyl	H	> 100	> 100	> 100
51b	4-phenylphenyl	CH ₃	> 100	> 100	10.7 $\pm$ 6.60
52a	4-acetoxyphenyl	H	> 100	> 100	5.96 $\pm$ 1.76

Compd	Ar (Het)	R	K_i , μ M $\pm$ SEM		
			[³ H]NIC binding $\alpha 4\beta 2^*$ nAChR ^a	[³ H]MLA binding $\alpha 7^*$ nAChR ^a	[³ H]DTBZ binding VMAT2 ^a
52b	4-hydroxyphenyl	H	> 100	> 100	5.26 $\pm$ 0.47
53a	1-naphthyl	H	> 100	> 100	4.68 $\pm$ 0.70
53b	1-naphthyl	CH ₃	> 100	> 100	0.63 $\pm$ 0.16
54a	2-naphthyl	H	> 100	> 100	> 100
54b	2-naphthyl	CH ₃	> 100	> 100	2.03 $\pm$ 0.45
55	3-chlorophenyl	H	-	-	3.88 $\pm$ 0.43
56a	3,5-difluorophenyl	H	-	-	5.24 $\pm$ 1.81
56b	3,5-difluorophenyl	CH ₃	-	-	0.29 $\pm$ 0.04
57a	3,5-dimethoxyphenyl	H	-	-	2.57 $\pm$ 0.48
57b	3,5-dimethoxyphenyl	CH ₃	-	-	2.87 $\pm$ 0.83
58a	3,4-dimethoxyphenyl	H	-	-	29.0 $\pm$ 5.00
58b	3,4-dimethoxyphenyl	CH ₃	-	-	5.93 $\pm$ 1.40
59a	2,3-dimethoxyphenyl	H	-	-	2.08 $\pm$ 0.10
59b	2,3-dimethoxyphenyl	CH ₃	-	-	0.62 $\pm$ 0.02
60a	2,5-dimethoxyphenyl	H	-	-	2.70 $\pm$ 0.70
60b	2,5-dimethoxyphenyl	CH ₃	-	-	0.56 $\pm$ 0.08
63	2-pyridinyl	-	-	-	5.87 $\pm$ 1.72
64	3-pyridinyl	-	-	-	33.3 $\pm$ 16.3
65	2-quinolinyl	-	-	-	0.96 $\pm$ 0.37
66	4-quinolinyl	-	-	-	2.64 $\pm$ 1.41
67	2-(N-methyl)pyrrolyl	-	-	-	9.21 $\pm$ 1.65
70	1-naphthyl	-	-	-	1.02 $\pm$ 0.30
71	1-anisoly	-	-	-	0.83 $\pm$ 0.23

^a n =3–6/compound/assay

Table 6

$\alpha 4\beta 2^*$, $\alpha 7^*$ nAChR, and VMAT2 Binding Affinity of 74a–76a, 74b–76b, 79a, 79b, 80a, 80b, 83a–d, 84a–d, 86a–d, 87a, 87b, 88a, 88b and 89–93.

Compd	K_i , $\mu\text{M} \pm \text{SEM}$		
	$[^3\text{H}]\text{NIC}$ Binding $\alpha 4\beta 2^*$ nAChR ^a	$[^3\text{H}]\text{MLA}$ Binding $\alpha 7^*$ nAChR ^a	$[^3\text{H}]\text{DTBZ}$ Binding VMAT2 ^a
lobelane	14.9 ± 1.67	26.0 ± 6.57	0.97 ± 0.19
74a	> 100	66.9 ± 27.4	3.32 ± 0.74
75a	> 100	> 100	3.11 ± 0.51
76a	> 100	9.02 ± 1.54	3.35 ± 0.80
74b	> 100	> 100	3.95 ± 0.61
75b	> 100	> 100	10.5 ± 1.58
76b	> 100	> 100	4.04 ± 1.76
74c	7.62 ± 1.47	> 100	1.87 ± 0.25
75c	> 100	> 100	3.28 ± 1.65
76c	> 100	> 100	2.47 ± 0.70
79a	> 100	> 100	6.11 ± 0.92
79b	> 100	> 100	11.7 ± 0.65
80a	> 100	> 100	1.36 ± 0.11
80b	> 100	> 100	2.62 ± 0.51
83a	-	-	3.35 ± 0.78
83b	-	-	3.17 ± 0.58
83c	-	-	3.36 ± 0.72
83d	-	-	5.29 ± 0.96
84a	-	-	0.30 ± 0.11
84b	-	-	0.63 ± 0.27
84c	-	-	2.68 ± 0.44
84d	-	-	3.62 ± 0.59
86a	> 100	> 100	12.3 ± 4.70
86b	> 100	> 100	31.8 ± 5.84
86c	> 100	> 100	2.57 ± 0.63
86d	> 100	> 100	3.92 ± 0.19
87a	> 100	> 100	3.42 ± 0.26
87b	> 100	> 100	16.0 ± 2.83
88a	> 100	> 100	41.0 ± 14.0
88b	> 100	> 100	4.60 ± 1.70
89	> 100	> 100	7.27 ± 2.28
90	> 100	> 100	10.4 ± 2.62
91	> 100	> 100	91.3 ± 26.2
92	> 100	> 100	15.5 ± 1.61
93	> 100	> 100	32.6 ± 6.79

^a
n =3–6/compound/assay

Table 7

VMAT2 binding affinity of compounds 94–102.

Compd	K_i , $\mu\text{M} \pm \text{SEM}$
	$[^3\text{H}]\text{DTBZ}$ binding, VMAT2 ^a
	R	m	n	
lobelane	CH ₃	2	2	0.97 $\pm$ 0.19
94	CH ₃	0	0	>100
95a	H	0	1	24.1 $\pm$ 6.10
95b	CH ₃	0	1	13.6 $\pm$ 8.14
96a	H	0	2	28.4 $\pm$ 7.30
96b	CH ₃	0	2	16.1 $\pm$ 4.07
97a	H	0	3	28.6 $\pm$ 2.44
97b	CH ₃	0	3	62.5 $\pm$ 16.6
98a	H	1	1	7.68 $\pm$ 1.44
98b	CH ₃	1	1	10.5 $\pm$ 3.36
99a	H	1	2	12.2 $\pm$ 0.19
99b	CH ₃	1	2	2.77 $\pm$ 1.77
100a	H	1	3	1.13 $\pm$ 0.28
100b	CH ₃	1	3	1.62 $\pm$ 0.39
101a	H	2	3	4.60 $\pm$ 0.56
101b	CH ₃	2	3	0.88 $\pm$ 0.30
102a	H	3	3	4.50 $\pm$ 1.02
102b	CH ₃	3	3	1.00 $\pm$ 0.23

^a n = 3–6/compound

Table 8

VMAT2 Binding Affinity of

Compd	[³ H]DTBZ binding, VMAT2 Ki, $\mu\text{M} \pm \text{SEM}$
lobelane	0.97 ± 0.19^a
<i>nor</i> -lobelane	2.31 ± 0.21
103a	3.41 ± 0.67
103b	1.87 ± 0.25
104	> 100
105	22.6 ± 0.23
106	5.59 ± 0.94
107	9.59 ± 1.47
108	0.56 ± 0.08
109	12.7 ± 1.09
110	15.5 ± 3.90
111	17.7 ± 4.66
112	25.0 ± 3.08
113	24.9 ± 3.07

Table 9

VMAT2 Binding Affinity of Pyrrolidino Analogs 115–126, 128a–128c, 129a–129c, and 131

Compd	$K_i, \mu\text{M} \pm \text{SEM}$
	[³ H]DTBZ binding, VMAT2 ^a
lobelane	0.97 ± 0.19
<i>nor</i> -lobelane	2.31 ± 0.21
115	2.75 ± 0.63
116	4.79 ± 0.71
117	1.50 ± 0.22
118	14.5 ± 2.13
119	3.02 ± 0.17
120	8.80 ± 2.30
121	5.05 ± 1.05
122	4.34 ± 0.71
123	3.40 ± 0.30
124	3.97 ± 0.64
125	36.7 ± 20.3
126	6.09 ± 0.19
128a	1.30 ± 0.21
128b	1.38 ± 0.20
128c	4.80 ± 1.70
129a	> 100
129b	> 100
129c	3.88 ± 0.90
131	3.95 ± 0.54

^a
n = 3–6/compound

Table 10

Brief Description of the Descriptors Used in the PLS and Nonlinear Neural Network Analyses.

Descriptor	Definition	Type ^a
Ram	Ramification index	1
PW5	Path/walk 5-Randic shape index.	1
LP1	Lovasz-Pelikan index [leading eigenvalue].	1
SEige	Eigenvalue sum from electronegativity weighted distance matrix.	1
VEp2	Average eigenvector coefficient sum from polarizability weighted distance matrix.	1
DISPm	d COMMA2 value/weighted by atomic masses.	2
G(N..N)	Sum of geometrical distances between N..N.	2
H7m	H autocorrelation of lag 7/weighted by atomic masses.	3
RCON	Randic-type R matrix connectivity.	3
R1p+	R maximal autocorrelation of lag 1/ weighted by atomic polarizabilities.	3
HOMT	HOMA (harmonic oscillator model of aromaticity index) total.	4

a1. Topological; 2. Geometrical; 3. GETAWAY; 4. Aromaticity indices; 5. WHIM

^a_n = 3–6/compound

Table 11

Stability Analysis of the Predictive Models Built from the Set of 104 Tetrabenazine and Lobeline Analogs

No.	NN11-3-1 Model		
	Train r^2 (rmsd)	LOO q^2 (rmsd)	Test r^2 (rmsd)
1	0.91 (0.225)	0.82 (0.316)	0.93 (0.282)
2	0.93 (0.207)	0.83 (0.330)	0.89 (0.341)
3	0.93 (0.220)	0.81 (0.358)	0.92 (0.303)
4	0.92 (0.235)	0.84 (0.339)	0.88 (0.353)
5	0.93 (0.223)	0.80 (0.370)	0.94 (0.246)
Avg.	0.92 (0.222)	0.82 (0.343)	0.91 (0.305)