Complementation Test of *Rpe65* Knockout and *Tvrm148*

Charles B. Wright,¹ Micah A. Chrenek,¹ Stephanie L. Foster,¹ Todd Duncan,²
T. Michael Redmond,² Machel T. Pardue,^{1,3} Jeffrey H. Boatright,¹ and John M. Nickerson¹

¹Department of Ophthalmology, School of Medicine, Emory University, Atlanta, Georgia

²Laboratory of Retinal Cell and Molecular Biology, National Eye Institute, National Institutes of Health, Bethesda, Maryland

³Rehabilitation Research and Development Center of Excellence, Atlanta VA Medical Center, Decatur, Georgia

Correspondence: John M. Nickerson, Department of Ophthalmology, Emory University, B5602, 1365B Clifton Road NE, Atlanta, GA 30322; litjn@emory.edu.

Submitted: May 1, 2013
Accepted: June 10, 2013

Citation: Wright CB, Chrenek MA, Foster SL, et al. Complementation test of *Rpe65* knockout and *tvrm148*. *Invest Ophthalmol Vis Sci*. 2013;54:5111–5122. DOI:10.1167/iov.13-12336

PURPOSE. A mouse mutation, *tvrm148*, was previously reported as resulting in retinal degeneration. *Tvrm148* and *Rpe65* map between markers D3Mit147 and D3Mit19 on a genetic map, but the physical map places RPE65 outside the markers. We asked if *Rpe65* or perhaps another nearby gene is mutated and if the mutant reduced 11-*cis*-retinal levels. We studied the impact of the *tvrm148* mutation on visual function, morphology, and retinoid levels.

METHODS. Normal phase HPLC was used to measure retinoid levels. *Rpe65*^{+/+}, *tvrm148*/*+* (*T*^{+/+}), *tvrm148/tvrm148* (*T*^{-/-}), *RPE65*^{KO/KO} (*Rpe65*^{-/-}), and *Rpe65*^{T/-} mice visual function was measured by optokinetic tracking (OKT) and electroretinography (ERG). Morphology was assessed by light microscopy and transmission electron microscopy (TEM). qRT-PCR was used to measure *Rpe65* mRNA levels. Immunoblotting measured the size and amount of RPE65 protein.

RESULTS. The knockout and *tvrm148* alleles did not complement. No 11-*cis*-retinal was detected in *T*^{-/-} or *Rpe65*^{-/-} mice. Visual acuity in *Rpe65*^{+/+} and *T*^{+/+} mouse was ~0.382 c/d, but 0.037 c/d in *T*^{-/-} mice at postnatal day 210 (P210). ERG response in *T*^{-/-} mice was undetectable except at bright flash intensities. Outer nuclear layer (ONL) thickness in *T*^{-/-} mice was ~70% of *Rpe65*^{+/+} by P210. *Rpe65* mRNA levels in *T*^{-/-} mice were unchanged, yet 14.5% of *Rpe65*^{+/+} protein levels was detected. Protein size was unchanged.

CONCLUSIONS. A complementation test revealed the RPE65 knockout and *tvrm148* alleles do not complement, proving that the *tvrm148* mutation is in *Rpe65*. Behavioral, physiological, molecular, biochemical, and histological approaches indicate that *tvrm148* is a null allele of *Rpe65*.

Keywords: RPE65/*Rpe65*, visual cycle, mutation

Recently a catalog of over 160 mouse mutations that cause retinal disease was published.¹ This collection included a brief description of *tvrm148*, a putative novel autosomal recessive mutation in the *Rpe65* gene. The *tvrm148* lesion was mapped between markers D3Mit147 (at 74.02 cM, corresponding to Chr3:148745409-148745542) and D3Mit19 (at 87.60 cM, corresponding to Chr3:157983038-157983067), approximately 9,237,658 bases apart, according to the most recent mouse genome sequence assembly, GRCm38 (Fig. 1A). This region includes 21 validated genes, but surprisingly, does not include *Rpe65*. Further, *Rpe65* is physically located at Chr3:159599181-159625307 by sequence analysis (Fig. 1B). This is approximately 1.6 Mb further telomeric from the distal end of the mapped disease locus (that is, from D3Mit19; Fig. 1B). The region from D3Mit19 to the physical location of the *Rpe65* gene contains another four candidate genes. *Rpe65* is just distal to these genes. Thus, *Rpe65* is not one of the 25 known genes in the mapped locus (Fig. 1B). From these two observations arises the potential complication that the gene harboring the *tvrm148* lesion is unknown among 25 candidates.

Another potential complication of the *tvrm148* assignment is that it is a missense mutation in *Rpe65* (a single base mutation resulting in the change of a wild type Phe codon to a

Ser codon). It is often problematic to assign disease causation to missense mutations in the absence of experimental data; after all, chimpanzee contains a Tyr residue at the same position¹ and presumably retains function. For instance, a counter-hypothesis is that *tvrm148* is a mutation in a gene close to *Rpe65*, and due to their proximity, a neutral missense mutation in *Rpe65* might be mistaken for the causative lesion. Because the putative causative mutation being a missense mutation that has not been previously detected in mouse, dog, or man, we thought an independent approach to the mutation assignment and additional experiments of *tvrm148* as a candidate mutation in *Rpe65* were warranted.

A straightforward way to formally resolve the question of whether *tvrm148* is a mutation in the *Rpe65* gene is to perform a complementation test using a deletion mutation solely within the *Rpe65* gene. Here we used the *Rpe65* knockout mouse.² This knockout is well characterized, has been backcrossed against wild type C57BL/6J for 10 generations, and the knockout of the *Rpe65* gene is well understood both genetically² and biochemically.² The *Rpe65* gene was partially deleted in the construction of the knockout,² and the resulting gene fragment no longer exhibits any *Rpe65* mRNA² or protein² expression or retinoid isomerase enzymatic activity.²

FIGURE 1. *Rpe65* resides outside the markers D3Mit147 and D3Mit19 on the physical map of chromosome 3. (A) Simplified genetic map of chromosome 3 showing the positions of D3Mit147, D3Mit19, and *Rpe65*. (B) Ensembl genome browser map of the critical region neighboring the D3Mit147 and D3Mit19 markers based on the GRCh38 build of the mouse genome released in December 2011.

The fundamental principles of the complementation test are basic in genetics. Here we perform this test asking if the mutation in *tvrm148* is in the same gene as the *Rpe65* knockout. We compared the phenotypes of the homozygous *tvrm148* mutation, *tvrm148/Rpe65* knockout heterozygotes, and homozygous *Rpe65*^{KO} to wild-type C57BL/6J by several histological, behavioral, and biochemical tests. We conclude that *tvrm148* does not complement the knockout allele, thus proving unequivocally that this mutation is within the *Rpe65* gene. Given this, we hypothesized that *tvrm148* should manifest as a retinoid isomerase mutation. This was confirmed on molecular, biochemical, morphological, and functional levels.

METHODS

Experimental Animals

C57BL/6J (*Rpe65*^{+/+}) mice served as wild type in all experiments. Mutant *tvrm148* (*T*^{-/-}) mutant mice were induced by ENU mutagenesis at The Jackson Laboratory¹ (Bar Harbor, ME) and were purchased for this study; *T*^{+/-} mice were bred by crossing *T*^{-/-} mice with *Rpe65*^{+/+} mice. *Rpe65*^{KO} (*Rpe65*^{-/-}) mice were transferred from the National Eye Institute to Emory University, where they were bred and maintained. *T*^{-/-} and *Rpe65*^{-/-} mice were bred together to create *T*^{+/-} mice for the complementation test. Mice were provided food and water ad libitum and maintained in a 12:12 light-dark cycle. Mice were sacrificed with CO₂ gas, and all procedures and care were approved by Emory's Institutional Animal Care Use Committee and all animals were treated in accordance with the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research.

Computer Predictions

Web-based computer prediction programs were used to predict the effect a putative F229S amino acid substitution could have on RPE65 protein (Sorting Intolerant from Tolerant,

SIFT³⁻⁷; Polymorphism Phenotyping, polyPhen 2.0⁸⁻¹⁰; Protein Analysis Through Evolutionary Relationships, PANTHER¹¹; MutationTaster¹²⁻¹⁹; PMUT²⁰; Screening for Non-Acceptable Polymorphisms, SNAP²¹; SNPs3D²²⁻²⁴; Predictor of Human Deleterious Single Nucleotide Polymorphisms, PhD-SNP²⁵⁻²⁸; and Protein Variation Effect Analyzer, PROVEAN²⁹).

Electroretinography (ERG)

Mice were dark-adapted overnight prior to ERG measurements. Scotopic recordings were performed after mice were dark-adapted, and photopic recordings were performed after mice had light adapted for 10 minutes under a constant illumination of 1.46 log cd·m⁻². Measurements were recorded with a commercial ERG system (UTAS-E3000; LKC, Gaithersburg, MD)³⁰⁻³³ on a range of light intensities from 3 × 10⁻⁴ cd·s·m⁻² to 137 cd·s·m⁻². The b-wave amplitudes were measured from the trough of the a-wave to the peak of the positive wave; if the a-wave was not present, b-wave amplitudes were measured from the baseline to the top of the positive wave.³⁴⁻³⁶

Retinoid Analysis

All steps were performed in a darkroom under dim red light. RPE/choroid samples were removed from eyecups and homogenized in 200 μL PBS (pH 7.4, 150 mM NaCl, 1.06 mM KH₂PO₄, 5.60 mM Na₂HPO₄) in a disposable homogenizer. Homogenate (10 μL) was taken for the protein assay (Microplate Pierce Coomassie [Bradford] Plus Protein Assay; Thermo Fisher Scientific Inc., Rockford, IL). The remaining homogenate was transferred to a 15 mL screw-top polypropylene tube (BD Falcon; BD Biosciences, San Jose, CA), and 400 μL methanol was added. Retinoids were extracted two times by adding 1.5 mL hexane and vortex mixing at top speed for 1 minute. Samples were centrifuged at 1424g for 3 minutes. Hexane was drawn off the upper phase and transferred to a fresh 15 mL screw-top polypropylene tube (BD Falcon; BD Biosciences). Pooled hexane extracts were dried in an evaporation system (TurboVap LV; Zymark Corporation, Hopkinton, MA). Samples were redissolved in 100 μL hexane by vortex mixing for 1

minute followed by centrifugation at 1424g for 3 minutes. A 50 μ L aliquot of each sample was run on a 5- μ m particle normal phase column (LiChrospher Si-60; EMD Millipore, Billerica, MA) on an isocratic HPLC (Agilent 1100/1200; Agilent Technologies, Santa Clara, CA) at 1.2 mL/min using HEDO (hexane with 11.2% [vol/vol] ethyl acetate; 2% [vol/vol] dioxane; and 1.4% [vol/vol] octanol) mobile phase.³⁷ Data were analyzed using commercial software (ChemStation32; Agilent Technologies). The remaining portion of mutant samples was subjected to saponification with alkaline hydrolysis in ethanolic potassium hydroxide and assayed for isomeric retinols contained in the retinyl esters.³⁷ For analysis of retinas, a pair of mouse retinas were homogenized in 0.5 mL of 50 mM MOPS, pH 6.5, 10 mM NH_2OH , and then transferred to a 15 mL glass screw-cap tube. A further 0.5 mL of 50 mM MOPS, pH 6.5, 10 mM NH_2OH was added along with 1 mL ethanol. The tubes were capped, mixed, and incubated at room temperature in the dark for 30 minutes. Retinoids were extracted two times by addition of 4 mL hexane, vortex mixing and centrifugation, and the hexane upper phases were combined and dried. The dried samples were redissolved in 100 μ L hexane. Retinal oxime standards were prepared following Garwin and Saari.³⁸ Retinal oxime standards and samples (20 μ L) were separated using a 5- μ m column (EMD Millipore) running HEDO mobile phase (see above) at 0.7 mL/min.

Optokinetic Tracking (OKT)

Visual acuities of *Rpe65*^{+/+}, *T*^{+/+}, *T*^{-/-}, and *Rpe65*^{-/-} mice were measured at 100% contrast from postnatal day 30 (P30) to P210 using a virtual optomotor system (OptoMotry 1.7.4; Cerebral Mechanics, Vancouver, British Columbia, Canada).^{39,40}

Histology and Morphometrics

Rpe65^{+/+}, *T*^{+/+}, and *T*^{-/-} mice were killed at P60 and P210. Eyes were enucleated and fixed in glutaraldehyde (2.5% [wt/vol] in 0.1 M sodium cacodylate buffer, pH 7.4; Electron Microscopy Sciences, Hatfield, PA) for 4 days at 4°C. After fixation, the eyes were rinsed three times in 0.1 M sodium cacodylate buffer (Electron Microscopy Sciences). The superior portion of each eye was marked by leaving a small corneal flap after dissecting away the remaining cornea and lens. Eyes were embedded in LX-112 (Ladd Research Industries, Williston, VT) and sectioned at 1- μ m thickness on a vertical meridian through the optic nerve head before staining with toluidine blue. Retina cross-sections were imaged at $\times 20$ magnification (Nikon Diaphot 300; Nikon Corporation, Tokyo, Japan) and photomerged with a graphics editing program (Adobe Photoshop CS2; Adobe Systems, Mountain View, CA). Images were imported into commercial image analysis software (Longboard V7.5; ImagingPlanet, Goleta, CA) for quantitative measures of retina layer thicknesses. Measurements were taken at 500- μ m increments superior and inferior to the optic nerve. Nuclei were counted as the number of nuclei on a line perpendicular to the RPE moving through the outer nuclear layer (ONL). ONL, inner segment (IS), and outer segment (OS) thickness were measured on a line perpendicular to the RPE moving through those respective layers. Cone nuclei were counted in 200- μ m areas 0.25 mm, 1 mm, and 2 mm superior and inferior of the optic nerve from retina sections as previously described.^{41,42} For transmission electron microscopy (TEM), gray-silver thin sections (700 angstroms) were postfixed in 1% (wt/vol) osmium tetroxide and stained with 2% (wt/vol) uranyl acetate and Reynold's lead citrate, and images of *Rpe65*^{+/+}, *T*^{+/+}, and *T*^{-/-} RPE were collected at P60 and P210 at $\times 2900$ and $\times 10,000$ magnification.

Quantitative Real Time Polymerase Chain Reaction (qRT-PCR)

Rpe65 mRNA levels were measured by qRT-PCR with P60 *Rpe65*^{+/+} and *T*^{-/-} samples. RNA was isolated from RPE/choroid tissue placed in Trizol (Invitrogen, Grand Island, NY) and extracted using an RNA purification kit (RNeasy Mini Kit; Qiagen, Valencia, CA). *Rpe65* primers (Qiagen) were used to amplify *Rpe65* mRNA with a one-step reaction (Quantitect SYBR Green RT-PCR kit; Qiagen) under these conditions: 30 minutes at 50°C, 15 minutes at 95°C, 40 cycles at 94°C for 15 seconds, and 55°C for 15 seconds and 72°C for 40 seconds, followed by melt curve analysis. Four technical repeats for each reaction were performed in each run of the instrument (iCycler; Bio-Rad, Hercules, CA), and *Rpe65* amplification was normalized to 18S RNA; 18S primers were: forward 5'-GTTGGTTTTCGGAAGTGGAGGC-3' and reverse 5'-GTCGGCATCGTTTATGGTCG-3'. There were five mice from different litters per sample group.

Immunoblotting

RPE/choroid tissue was isolated from P60 *Rpe65*^{+/+}, *T*^{+/+}, and *T*^{-/-} mice in RIPA buffer (Teknova, Hollister, CA). Samples were homogenized by vortex mixing vigorously with stainless steel set screws for 5 minutes. Following homogenization, samples were centrifuged at 16,000g for 1 minute. Supernatants were collected and protein concentration determined via BCA protein assay (Novagen, Darmstadt, Germany). Protein (1 μ g) was resolved on a 10% gel (Bis-Tris Criterion Gel; Bio-Rad) and transferred to a nitrocellulose membrane (Bio-Rad) with a wet transfer system (Bio-Rad). Western blocking reagent (Roche Diagnostics, Indianapolis, IN) was used. A custom-made N-terminal specific antibody to mouse *Rpe65* was made by raising antiserum against amino acids 1-44 of RPE65 mouse protein. Primary antibody incubation was performed at a 1:5000 dilution for 1 hour at room temperature. A goat anti-rabbit-HRP conjugate secondary antibody (Invitrogen) was incubated at a 1:5000 dilution at room temperature for 1 hour. Chemiluminescent detection (ECL Plus; GE Healthcare BioSciences Corp., Piscataway, NJ) was used for band visualization. Densitometry of immunoblots with the same protein samples run on different gels ($n = 3$) was performed in a graphics editing program (Adobe Photoshop CS2; Adobe Systems).

Statistical Analysis

Graphs were constructed in spreadsheet software (Excel 2008; Microsoft Inc., Redmond, WA). All graphs represent the data as means plus or minus standard deviations. Visual acuity, ERG, and morphometric data were analyzed by two-way repeated measures ANOVA (RM-ANOVA) with post hoc Student-Newman-Keuls testing. qRT-PCR and densitometry data were analyzed by one-way ANOVA with post hoc Student-Newman-Keuls testing (SigmaPlot 12; Systat Software Inc., San Jose, CA). Sample sizes are provided in the results. All groups contain at least five mice from different litters.

RESULTS

Predictions of *Tvrm148* Mutation Pathogenicity

Web-based algorithms were used to predict whether the *tvrm148* mutation that was previously reported¹ could have a potentially harmful effect on RPE65 protein function and could thus have the previously reported effect¹ on visual function. The prediction programs SIFT,³⁻⁷ polyPhen 2.0,⁸⁻¹⁰

PANTHER,¹¹ MutationTaster,^{12–19} PMUT,²⁰ SNAP,²¹ SNPs3D,^{22–24} PhD-SNP,^{25–28} and PROVEAN²⁹ were used to predict if the putative *Rpe65* mutation could negatively impact visual function in mice. With the exception of PROVEAN, all programs predicted the *tvrm148* mutation is responsible for RPE65 dysfunction (Table 1). DNA sequencing from *T*^{−/−} mice confirmed the presence of the *tvrm148* mutation that was previously reported.¹ There were no other sequence differences in any of the other exons or regions sequenced comparing *tvrm148* to the C57BL/6J reference sequence. See Supplementary Material S1 for a description of the primers and PCR products that were sequenced, and methods used.

Complementation Test

A complementation test was performed to determine whether the causative mutation driving visual function loss in *T*^{−/−} mice resides in the *Rpe65* locus. By breeding *T*^{−/−} mice and *Rpe65*^{−/−} mice together, *Rpe65*^{+/−} offspring were produced. *Rpe65*^{+/+}, *Rpe65*^{−/−}, *T*^{−/−}, and *Rpe65*^{+/−} mice were then screened by ERG at 137 cd·s·m^{−2} at P30 to assess visual function (Fig. 2). All mutant mice had significantly lower responses ($P < 0.001$) than *Rpe65*^{+/+} mice ($264.7 \pm 56.4 \mu\text{V}$ [$n = 6$]; Fig. 2), but the a-wave responses of *Rpe65*^{−/−}, *Rpe65*^{+/−}, and *Rpe65*^{+/−} mice were not significantly different when compared with each other alongside wild type mice (*Rpe65*^{+/+} mice, $24.6 \pm 16.2 \mu\text{V}$ [$n = 7$]); *T*^{−/−} mice, $6.1 \pm 6.7 \mu\text{V}$ ($n = 5$); *Rpe65*^{+/−} mice, $35.1 \pm 16.4 \mu\text{V}$ ($n = 9$).

Retinoid Levels in *Tvrm148*, Knockout, and Normal Mouse Eyes

The fact that mutant *T*^{−/−} mice did not have appreciable a-wave responses suggested they lack the ability to produce sufficient amounts of 11-*cis*-retinol (the enzymatic product of RPE65) to support proper photoreceptor function.¹ *Rpe65*^{−/−} mice have also been shown to harbor excessive amounts of retinyl ester,² and we sought to determine if *T*^{−/−} mice shared this canonical chronic loss of 11-*cis*-retinal and increase in retinyl ester content in the RPE/choroid. *Rpe65*^{−/−} and *T*^{−/−} mice had elevated retinyl esters, while 11-*cis* retinal was not detected in *Rpe65*^{−/−} or *T*^{−/−} retinas (Table 2). The *T*^{−/−} samples had 22× more retinyl esters than *Rpe65*^{+/+} samples. Retinyl ester content in *Rpe65*^{−/−} and *T*^{−/−} mice did not differ significantly from one another. Retinol isomers in all samples were below the level of detection prior to saponification of mutant samples. For the saponified portion, only all-*trans*-retinol was detected in the mutant samples and no quantification was attempted.

Inheritance of *Tvrm148*

We sought to determine how the *tvrm148* allele is inherited in mice (alternatively recessive or dominant, autosomal or sex-linked—data not shown). ERG measurements on dark- and light-adapted *Rpe65*^{+/+}, *T*^{+/−}, and *T*^{−/−} mice were performed at P60 and P180 (Fig. 3). While *T*^{+/−} mice did not show any reduction in dark-adapted b-wave amplitudes, *T*^{−/−} mice showed depressed dark-adapted b-wave amplitudes at both ages (Figs. 3A, 3C). At P60, in scotopic conditions ($0.01551 \text{ cd} \cdot \text{s} \cdot \text{m}^{-2}$), *T*^{−/−} mice had no measurable ($0 \pm 0 \mu\text{V}$, $P < 0.001$) b-wave response, whereas *Rpe65*^{+/+} and *T*^{+/−} mice had an average b-wave amplitude of $197.9 \pm 62.4 \mu\text{V}$ and $221.8 \pm 59.6 \mu\text{V}$, respectively. At the brightest flash intensity tested, $137 \text{ cd} \cdot \text{s} \cdot \text{m}^{-2}$, *T*^{−/−} mice had an average b-wave response of $132.4 \pm 52.4 \mu\text{V}$ in comparison to $648.2 \pm 196.9 \mu\text{V}$ and $654.3 \pm 177.9 \mu\text{V}$ in *Rpe65*^{+/+} and *T*^{+/−} mice, respectively (Fig. 3A). By P180, *T*^{−/−} mice had an average scotopic ($0.01551 \text{ cd} \cdot \text{s} \cdot \text{m}^{-2}$) b-wave amplitude of $0 \pm 0 \mu\text{V}$ compared with $216.0 \pm 40.6 \mu\text{V}$ in *Rpe65*^{+/+} mice and $191.0 \pm 39.1 \mu\text{V}$ in *T*^{+/−} mice; at 137

TABLE 1. Prediction of Putative *Tvrm148* Mutation on RPE65 Function

Program	Predicted Damaging Effect?
SIFT	Yes
PolyPHEN 2.0	Yes
PANTHER	Yes
MutationTaster	Yes
PMUT	Yes
SNAP	Yes
SNPs3D	Yes
PhD-SNP	Yes
PROVEAN	No

Multiple computer programs were used to predict whether or not the *tvrm148* mutation, which is predicted to produce an F229S amino acid substitution in RPE65 protein, could in fact be pathogenic with respect to visual function in mice.

$\text{cd} \cdot \text{s} \cdot \text{m}^{-2}$, *T*^{−/−} mice had an average b-wave amplitude of $69.1 \pm 30.2 \mu\text{V}$ compared with $526.1 \pm 75.1 \mu\text{V}$ in *Rpe65*^{+/+} mice and $541.3 \pm 68.6 \mu\text{V}$ in *T*^{+/−} mice (Fig. 3C).

Light-adapted b-wave response was also significantly reduced in *T*^{−/−} mice (Figs. 3B, 3D). In response to the brightest flash intensity ($75 \text{ cd} \cdot \text{s} \cdot \text{m}^{-2}$) at P60, *T*^{−/−} mice had an average b-wave response of $140.4 \pm 58.0 \mu\text{V}$ compared with $220.4 \pm 61.5 \mu\text{V}$ in *Rpe65*^{+/+} mice and $241.9 \pm 76.0 \mu\text{V}$ in *T*^{+/−} mice (Fig. 3B), which decreased at P180 in *T*^{−/−} mice to $91.6 \pm 19.5 \mu\text{V}$ compared with $196.5 \pm 42.8 \mu\text{V}$ in *Rpe65*^{+/+} mice and $217.5 \pm 29.8 \mu\text{V}$ in *T*^{+/−} mice at P180 (Fig. 3D).

Visual Acuity Affected by Null Mutations

The optokinetic tracking testing, a behavioral measure of visual acuity, was used to determine the effects of the *tvrm148* mutation on the visual guided activity of the mouse. A virtual optomotor system^{39,40} was used to measure visual acuity in *Rpe65*^{+/+}, *T*^{+/−}, and *T*^{−/−} mice. *Rpe65*^{+/+} and *T*^{+/−} mice maintained a constant high visual acuity ($0.381 \pm 0.002 \text{ c/d}$) with little variability from P30 through the end of the study at P210 (Fig. 4). *T*^{+/−} mice had no detectable reduction in visual acuity. *T*^{−/−} mice, on the other hand, had a significant reduction in visual acuity at P30 ($0.318 \pm 0.003 \text{ c/d}$), and continued to lose visual acuity until the end of the study (Fig. 4). At P210, only a few *T*^{−/−} mice retained measurable visual acuity, which produced the average value of $0.037 \pm 0.045 \text{ c/d}$ (Fig. 4).

Comparison of Slow Loss of Visual Acuity in *T*^{−/−} and *Rpe65*^{−/−} Mice

The visual acuities of *Rpe65*^{+/+}, *T*^{+/−}, and *Rpe65*^{−/−} mice were compared at P30, P120, and P210 to determine if *Rpe65*^{−/−} and *T*^{−/−} mice behaved similarly in early, middle, and late stages of the retinal degeneration (Fig. 5). The slow loss of visual acuity in *T*^{−/−} and *Rpe65*^{−/−} mice was consistent at each age examined. At P30, *T*^{−/−} visual acuity was $0.318 \pm 0.003 \text{ c/d}$, and *Rpe65*^{−/−} visual acuity was $0.318 \pm 0.003 \text{ c/d}$. At P120, *T*^{−/−} visual acuity was $0.225 \pm 0.010 \text{ c/d}$, and *Rpe65*^{−/−} visual acuity was $0.245 \pm 0.007 \text{ c/d}$. At P210, *T*^{−/−} visual acuity was $0.037 \pm 0.045 \text{ c/d}$, and *Rpe65*^{−/−} visual acuity was $0.000 \pm 0.000 \text{ c/d}$.

Histology and Morphometrics Changes—Slight Losses in Mutants

We sought to determine if loss of visual function correlated with a significant loss of retina layer thickness. Eyes were

FIGURE 2. The *tvrm148* and *Rpe65*^{KO} alleles do not complement. Dark-adapted a-wave amplitudes from a flash stimulus of 137 cd-s/m² from *Rpe65*^{+/+} (*n* = 6), *Rpe65*^{-/-} (*n* = 7), *T*^{-/-} (*n* = 5), and *Rpe65*^{T/-} (*n* = 9) mice at P30 are shown. *Rpe65*^{-/-}, *T*^{-/-}, and *Rpe65*^{T/-} mice have significantly lower a-wave amplitudes than *Rpe65*^{+/+} mice, but a-wave amplitudes from *Rpe65*^{-/-}, *T*^{-/-}, and *Rpe65*^{T/-} mice do not differ significantly from one another, showing the two mutant alleles do not complement one another. **P* < 0.001 comparing *Rpe65*^{+/+} with each of *Rpe65*^{-/-}, *T*^{-/-}, and *Rpe65*^{T/-} mice. Data are presented as mean ± SD.

harvested from P60, P120, P180, and P210 from *Rpe65*^{+/+}, *T*^{+/+}, and *T*^{-/-} mice, and representative sections from P60 and P210 are shown (Fig. 6). The *T*^{-/-} retina was thinner than *Rpe65*^{+/+} or *T*^{+/+} retinas (Fig. 6). Retina architecture was largely intact even at P210, despite some significant ONL thinning (35.9 ± 2.9 μm in *T*^{-/-} mice compared with 50.9 ± 4.5 μm in *Rpe65*^{+/+} mice) 500 μm on the superior side of the optic nerve (*P* < 0.001). Some lipid droplets appear in the RPE of *T*^{-/-} mice (Fig. 7). Images of *Rpe65*^{+/+}, *T*^{+/+}, and *T*^{-/-} retinas taken at P120 and P180 look similar to P60 and P210 (data not shown).

Quantitative measurements of retina cross-sections from P60 and P210 eyes revealed reduced nuclei counts (Figs. 7A, 7B), thinning of the outer nuclear layer (ONL; Figs. 7C, 7D); IS (Figs. 7E, 7F); and OS (Figs. 7G, 7H) in *T*^{-/-} mice compared with *T*^{+/+} and *Rpe65*^{+/+} mice (Fig. 7). *T*^{-/-} retina thinning was apparent as early as P60. The ONLs of *Rpe65*^{+/+} and *T*^{+/+} mice were 11 to 12 nuclei thick at both P60 and P210, but *T*^{-/-} mice had an ONL only 8 to 9 nuclei thick at P60 and 7 to 8 nuclei thick at P210 (Figs. 7A, 7B). By P60, *T*^{-/-} retina had lost a significant number of cone nuclei (Fig. 8A); *Rpe65*^{+/+} retina had 11.8 ± 3.4 cones per 200-μm segment, but *T*^{-/-} retina had only 6.3 ± 2.3 cone nuclei per 200-μm segment on the superior side of the optic nerve and 3.2 ± 1.4 cone nuclei per 200-μm segment on the inferior side (Fig. 8A). By P210, *Rpe65*^{+/+} retina had 10.0 ± 1.8 cone nuclei per 200-μm

segment, and *T*^{-/-} retina had only 2.7 ± 1.4 and 2.0 ± 0.9 cone nuclei per 200-μm segment on the superior and inferior sides of the retina, respectively (Fig. 8B).

Retinyl ester accumulations in the RPE are known to correlate with the formation of lipid droplets in *RPE65*-deficient mice,^{2,43,44} and we decided to examine the RPE of *T*^{-/-} mice to see if lipid droplet formation in these mice occurred consistent with other *RPE65* mutant models. TEM images of *Rpe65*^{+/+}, *T*^{+/+}, and *T*^{-/-} RPE show mutant mice have relatively normal morphology with the exception of lipid droplets that are present (Fig. 9). Lipid droplets were detected qualitatively in the RPE in *T*^{-/-} mice at P60 (Fig. 9C), and lipid droplets appeared larger by P210 (Fig. 9F). No lipid droplets were detected in *Rpe65*^{+/+} or *T*^{+/+} RPE at P60 or P210 (Figs. 9A, 9B, 9D, 9E, 9G).

RNA Quantitation of *Tvrm148* Mutation

RNA was isolated from the RPE of *Rpe65*^{+/+} and *T*^{-/-} mice to determine if the mutant mice expressed similar levels of *Rpe65* mRNA. qRT-PCR results (Fig. 10) indicated insignificant differences in steady-state levels of *Rpe65* mRNA in *T*^{-/-} mice compared to *Rpe65*^{+/+} mice (*P* = 0.617); *T*^{-/-} mice express *Rpe65* mRNA at 91.3 ± 34.4% of the level in *Rpe65*^{+/+} mice.

RPE65 Protein Expression in *T*^{-/-} Mice

Immunoblotting for RPE65 protein showed the size and abundance of RPE65 protein in *Rpe65*^{+/+}, *T*^{+/+}, and *T*^{-/-} mice (Fig. 11A). Immunoblots for *Rpe65* showed a reduction in RPE65 protein amount in *T*^{+/+} and *T*^{-/-} mice (Fig. 10A). Densitometry analysis was performed on three immunoblots (Fig. 11B). *T*^{+/+} mice expressed RPE65 protein at 59.7 ± 4.7% of *Rpe65*^{+/+} levels, and *T*^{-/-} mice expressed RPE65 protein at 14.5 ± 1.5% of *Rpe65*^{+/+} levels (Fig. 11B). There did not appear to be any difference in the size of RPE65 protein in *T*^{-/-} mice despite a reduction in the amount of protein (Fig. 11A).

DISCUSSION

The *Rpe65*^{KO} and *tvrm148* mutations do not complement one another, proving that the *tvrm148* mutation occurs in the *Rpe65* locus and is the causative lesion driving visual function loss (Fig. 2; Table 1). The complementation test with the knockout and *tvrm148* alleles definitively proves that there is no other mutation contributing to the loss of visual function (Fig. 2). This genetic test also resolves concerns associated with discrepancies we found between the genetic and physical maps of chromosome 3 in the region where the *Rpe65* locus and D3Mit147 and D3Mit19 markers reside (Fig. 1).

Retinoid analyses of *Rpe65*^{+/+}, *Rpe65*^{-/-}, and *T*^{-/-} mice, showing undetectable amounts of 11-*cis*-retinal (downstream enzymatic product) in retina but large amounts of retinyl esters (enzymatic substrate) in RPE/choroid samples from mutant

TABLE 2. Retinoid Isomerase Activity Is Abolished in *Rpe65*^{-/-} and *T*^{-/-} Mice

Genotype*	11- <i>cis</i> -Retinal, pmol/eye†	All- <i>trans</i> -Retinal, pmol/eye†	Retinyl Ester Content, pmol/mg protein‡
<i>Rpe65</i> ^{+/+}	765.9 ± 70.1	105 ± 13.5	63.40 ± 38.30
<i>Rpe65</i> ^{-/-}	None detected	None detected	1826.63 ± 654.34
<i>T</i> ^{-/-}	None detected	None detected	1385.77 ± 629.77

11-*cis*-retinal, all-*trans*-retinal and retinyl esters from *Rpe65*^{+/+}, *Rpe65*^{-/-}, and *T*^{-/-} retina and RPE/choroid samples, respectively. *n* ≥ 3 for each group. Data presented as mean ± SD.

* Number of animals analyzed per genotyped ≥ 3.

† The lower limit of detection in this assay was 1 pmol.

‡ The lower limit of detection in this assay was 8.8 pmol.

FIGURE 3. *T*^{-/-} mice, but not *T*^{+/-} mice, have a loss of ERG response. Dark- and light-adapted b-wave amplitudes, respectively, from *Rpe65*^{+/+}, *T*^{+/-}, and *T*^{-/-} mice at P60 (A, B) and P180 (C, D) are shown. **P* < 0.05 *T*^{-/-} compared with *Rpe65*^{+/+}. ***P* < 0.001 *T*^{-/-} compared with *Rpe65*^{+/+}, significance determined through two-way repeated measures ANOVA with post hoc Student-Newman-Keuls testing. There was no difference between *Rpe65*^{+/+} and *T*^{+/-} ERG responses. Data are presented as absolute values (mean ± SD).

FIGURE 4. *T*^{-/-} mice, but not *T*^{+/-} mice, lose visual acuity with age. Visual acuities of *Rpe65*^{+/+} (solid line); *T*^{+/-} (dotted line; largely hidden behind the solid back line); and *T*^{-/-} mice (dashed line) from P30 to P210 are represented on the same graph. ***P* < 0.001 *T*^{-/-} compared with *Rpe65*^{+/+}, significance determined through two-way repeated measures ANOVA with post hoc Student-Newman-Keuls testing. Comparison of *T*^{+/-} with *Rpe65*^{+/+} does not reach statistical significance between P30 and P210, and visual acuity measures overlap through the duration of the study. Data are presented as mean ± SD.

FIGURE 5. *T*^{-/-} mice have residual visual function similar to *Rpe65*^{-/-} mice. Comparison of visual phenotype measured by visual acuity. *T*^{-/-} mice have a residual visual function similar to *Rpe65*^{-/-} mice. Visual acuities of *Rpe65*^{+/+} (black bar); *T*^{-/-} (gray bar); and *Rpe65*^{-/-} (white bar) mice at P30, P120, and P210 are shown on the same graph. Data are presented as mean ± SD. ***P* < 0.001 compared to wild type. #*P* < 0.05 *T*^{-/-} compared with *Rpe65*^{-/-}. ##*P* < 0.001 *T*^{-/-} compared with *Rpe65*^{-/-}.

FIGURE 6. $T^{-/-}$ mice have a thinner retina than $T^{+/-}$ or $Rpe65^{+/+}$ counterparts. Bright-field images of retina cross-sections stained with toluidine blue are shown. $T^{-/-}$ mice experience slight retina thinning but retain normal retina architecture as they age. Representative images of $Rpe65^{+/+}$, $T^{+/-}$, and $T^{-/-}$ retina 1 mm superior of the optic nerve at P60 (A–C) and P210 (D–F) are shown. INL, inner nuclear layer; GCL, ganglion cell layer.

mice, confirmed that the *tvrm148* mutation results in a biochemically null RPE65 protein product that lacks retinoid isomerase activity (Table 2). Previous reports of mouse *Rpe65* mutations^{1,2,43,44} show $Rpe65^{KO}$, $Rpe65^{rd12}$, and $Rpe65^{R91W}$ mice have undetectable scotopic responses at dim flash intensities and bright flash intensities in addition to reduced photopic responses^{1,2,43,44} and these previous findings support the importance of RPE65 in 11-*cis*-retinal production and photoreceptor function. Consistent with other *Rpe65* mutants,^{1,2,43,44} $T^{-/-}$ mice at P30 and P180 had poor dark-mediated vision but relatively appreciable light-mediated vision at early ages (Fig. 3). LCA2 patients suffer from a similar pathology early in their degenerations,⁴⁵ so we sought to determine if $T^{-/-}$ and $Rpe65^{-/-}$ mice conform with clinical observations of human patients. $T^{-/-}$ mice, but not $T^{+/-}$ mice, lose visual acuity through P210 (indicating the *tvrm148* mutation inherits in a recessive manner) but have appreciable visual function at early ages (Fig. 4). $Rpe65^{-/-}$ mice have visual acuity losses that parallel with $T^{-/-}$ mice and do not significantly differ from one another (Fig. 5). This is the first reported progressive visual acuity loss in *Rpe65* mutant mice and goes well beyond previous work reporting RPE65 visual acuity.⁴⁶

Human LCA2 patients^{45,47,48} have mostly intact retina architecture even at late ages,^{45,47,48} so we sought to qualitatively and quantitatively examine the retina for similar phenotypes. The architecture of the retina in $T^{-/-}$ mice relative to $Rpe65^{+/+}$ and $T^{+/-}$ mice was strikingly normal considering the visual function deficits (Fig. 6), but these findings were consistent with published literature on RPE65-deficient mouse models.^{1,2,43,44} Droplets (Fig. 9) in the RPE in $T^{-/-}$ mice are similar to reports in $Rpe65^{KO}$, $Rpe65^{rd12}$, and $Rpe65^{R91W}$ mice^{2,43,44} that are the result of retinyl ester accumulation.² $T^{+/-}$ and $T^{-/-}$ mice experienced significant reductions in

RPE65 protein level compared to $Rpe65^{+/+}$ (Fig. 11) even though there was no reduction in *Rpe65* mRNA levels in $T^{-/-}$ mice compared with $Rpe65^{+/+}$ (Fig. 10). $T^{-/-}$ mice had no detectable 11-*cis*-retinal in neural retina and a 22-fold increases in retinyl esters in RPE samples (Table 2), further indicating the mutant protein is a null mutation that possesses no measurable enzymatic activity.

The visual acuity of $T^{-/-}$ mice is surprisingly high (almost normal) at early ages (Fig. 4) considering the lack of 11-*cis*-retinal generation in RPE65-deficient mice^{2,43}; neither $Rpe65^{KO}$ nor $Rpe65^{rd12}$ mice have detectable amounts of 11-*cis*-retinal in the retina.^{2,43} $Rpe65^{R91W}$ mice have ~5% the wild type amount of 11-*cis*-retinal in the retina, and this residual enzymatic product is sufficient to preserve vision significantly longer than what is seen in other *Rpe65* mutant mice.⁴⁴ We show *tvrm148* mice lack 11-*cis*-retinal altogether (Table 2). The residual visual function in $T^{-/-}$ mice may correspond to residual rod function, not cone function^{42,49} because in the absence of 11-*cis*-retinal, cone opsins in mice mislocalize,^{50,51} and it has been proposed this cone opsin mislocalization may have a toxic effect on cone cells.⁵¹ Work with *rd12* mice shows most cones are lost from the retina by P30.⁴⁶ This is in contrast to most other retinal degenerations reported in mice in which visual function loss is driven primarily by rod photoreceptor loss.⁴⁶ Visual acuity measurements are performed in photopic conditions that are dominated by cone photoreceptors in C57BL/6J mice,⁵² so it may at first seem counterintuitive that the visual acuity measurements in $T^{-/-}$ mice probably rely on residual rod photoreceptor function. Previous work shows the greatly altered kinetics of rod photoreceptors in *Rpe65* knockout mice cause them to function exclusively in photopic conditions because of reduced photon quantum-catch capability.⁵³ This, coupled with the fact that $Rpe65^{-/-}::Rho^{-/-}$ (*Rhodopsin*; pure cone function) mice do not retain visual acuity,⁴⁶ suggests the residual visual acuity in $T^{-/-}$ mice is driven primarily by rod function. Consistent with previous work,^{46,51,54} $T^{-/-}$ mice lost 59.9% of cones by P60 (Fig. 8); furthermore, many remaining cones are not functional at that age in $Rpe65^{-/-}$ mice, although they can be restored to activity by reintroduction of 11-*cis*-retinal.⁵⁵

The source and identity of chromophore for rod photoreceptors in $T^{-/-}$ mice, though, is unclear at this time. However, $Rpe65^{-/-}$ mice may generate and utilize 9-*cis*-retinal to form isorhodopsin.⁵⁶ $R91W$ *Rpe65* knock-in mice generate a small but significant amount of 11-*cis*-retinal.⁴⁴ It is possible *Rpe65* mutant mice could obtain enough 11-*cis*-retinal for residual visual function from photoisomerization.⁵⁷ More recently, evidence is emerging that Müller cells may actually provide small amounts of 11-*cis*-retinal to the retina,^{58–60} and although this mechanism would not be sufficient to preserve cones, we speculate residual visual function in *Rpe65* mutant mice may be aided by this putative cone visual cycle.

Low RPE65 protein levels in $T^{-/-}$ mice could be the result of reduced translation of the *Rpe65* transcript in $T^{-/-}$ mice or the result of RPE65 protein instability. Recent work concerning the three dimensional structure of *B. taurus* RPE65 protein^{61–64} provides insight into reasons why the RPE65^{F229S} mutant protein may be nonfunctional and possibly unstable. The $Rpe65^{tvrm148}$ mutation resides in exon 7 of RPE65¹ and results in a substitution of Ser for Phe at amino acid position 229. Because Phe 229 is conserved in RPE65 from a wide variety of species,¹ it is possible that a missense mutation at this position could be pathogenic even though the residue is not thought to participate in the active site of RPE65.⁶¹ The pathogenicity (Table 2) of the F229S substitution might be attributed to two other factors. First, RPE65 is a seven-bladed β -propeller that folds to form a hydrophobic tunnel leading to the active site. Mutations interfering with β -propeller folding may perturb the

FIGURE 7. *T*^{-/-} mouse retina layers progressively thin with age. Quantitative measurements every 500 μ m superior and inferior of the optic nerve in *Rpe65*^{+/+} (solid line); *T*^{+/-} (dotted line); and *T*^{-/-} (dashed line) retina were taken for the number of stacked nuclei in the ONL (A, B); ONL thickness (C, D); IS thickness (E, F); and OS thickness (G, H) at P60 (*Rpe65*^{+/+} *n* = 11, *T*^{+/-} *n* = 8, *T*^{-/-} *n* = 12) and P210 (*Rpe65*^{+/+} *n* = 7, *T*^{+/-} *n* = 6, *T*^{-/-} *n* = 14). **P* < 0.05 *T*^{-/-} compared with *Rpe65*^{+/+}. ***P* < 0.001 *T*^{-/-} compared with *Rpe65*^{+/+}. +*P* < 0.05 *Rpe65*^{+/+} compared with *T*^{+/-}, significance determined through two-way repeated measures ANOVA with post hoc Student-Newman-Keuls testing. Data are presented as mean $\pm$ SD.

FIGURE 8. *T*^{-/-} mice have significant cone nuclei loss at P60 and P210. Cone nuclei were counted at distances of 0.25, 1, and 2 mm superior and inferior of the optic nerve in a 200- μm segment of the retina at (A) P60 and (B) P210. ***P* < 0.001 *Rpe65*^{+/+} compared with *T*^{-/-}, significance determined through two-way repeated measures ANOVA with post hoc Student-Newman-Keuls testing. Data are presented as mean $\pm$ SD.

FIGURE 9. TEM shows lipid droplets indicative of retinyl ester accumulation in the RPE of *T*^{+/-} and *T*^{-/-} mice. TEM was performed on sections of RPE 1000 μm superior to the optic nerve in *Rpe65*^{+/+}, *T*^{+/-}, and *T*^{-/-} mice at P60 and P210. (A-C) TEM images of RPE from *Rpe65*^{+/+}, *T*^{+/-}, and *T*^{-/-} mice at P60, respectively, at $\times 2900$ magnification are shown. (D-F) TEM images of *Rpe65*^{+/+}, *T*^{+/-}, and *T*^{-/-} mice at P210, respectively, at $\times 2900$ magnification are shown. (G-I) TEM images of *Rpe65*^{+/+}, *T*^{+/-}, and *T*^{-/-} mice at P210, respectively, at $\times 10,000$ magnification are shown. N, nucleus; L, lipid droplet.

FIGURE 10. $T^{-/-}$ mice express *Rpe65* mRNA at the same level as *Rpe65* $^{+/+}$ mice under steady-state conditions. Whole cell RNA from P60 *Rpe65* $^{+/+}$ and $T^{-/-}$ RPE/choroid extracts was harvested for qRT-PCR. Steady-state *Rpe65* mRNA levels in *Rpe65* $^{+/+}$ mice ($n = 4$) and $T^{-/-}$ mice ($n = 5$) show no significant difference ($P = 0.617$).

formation of the hydrophobic tunnel or the appropriate shape of the active site itself.⁶¹ A comparison of the location of many of the known pathogenic mutations in human RPE65 with the crystal structure of bovine RPE65 confirm many of these missense mutations reside between the β -propellers of the protein.⁶¹ Phe 229 resides between the second and third β -propellers of RPE65, and because of the different chemical properties of the phenylalanine (large, aromatic, hydrophobic) and serine (small, polar, hydrophilic) side chains, it would seem possible this amino acid substitution could affect protein folding or shape. The fact that the *tvrm148* mutation abolishes 11-*cis*-retinal production, it would seem possible this residue is near the active site of the enzyme and could affect substrate binding and catalysis.

The *Rpe65* tvrm148 mouse serves as an interesting contrast to other models of RPE65 deficiency, specifically the *Rpe65* R91W

mouse. The *Rpe65* tvrm148 mutation occurs in a region of *Rpe65* that is rich in pathogenic mutations in the human homolog.^{53,65-71} The *Rpe65* R91W mouse, a knock-in model for the most common missense mutation seen in humans, retains residual RPE65 enzymatic activity.⁴⁴ The *Rpe65* tvrm148 mouse, though, is the first model of RPE65 deficiency to present a null phenotype while retaining the ability to produce a mutant RPE65 protein (albeit low; Fig. 11). Because this model has a missense mutation in a region of the protein known for recessively inheriting, disease-causing missense mutations in human subjects while still retaining a null phenotype, the *Rpe65* tvrm148 model may have unique utility for studying future treatments for LCA2.

Future work with this mutant mouse model could involve *Rpe65* gene augmentation to see if visual function rescue in these mice resemble rescue seen in other RPE65-deficient mice.⁵⁶ Other future work could examine cone photoreceptor loss in $T^{-/-}$ mice to see if the loss resembles that seen in other RPE65-deficient mice.^{49,72} Like the knockout mouse, the *tvrm148* mutation acts as an autosomal recessive, null mutation. Compared with the knockout, almost all other properties were very similar based on visual optokinetic testing (Fig. 4) and ERGs (Fig. 2). Molecular properties of the *tvrm148* mRNA are remarkable in its nearly normal abundance (Fig. 10), but with little accumulated protein under steady-state conditions (Fig. 11). We might speculate that the mRNA is processed normally but that the protein is unstable and rapidly degraded.

In summary, a complementation test proves that *tvrm148* is a mutation in the *Rpe65* gene. This mutation results in a null *Rpe65* mouse that makes no 11-*cis*-retinal. The *tvrm148* mice have a visual phenotype resembling that found in *Rpe65* KO mice, losing visual acuity and ERGs in a similar manner. Protein is present in low amounts and lacks activity, despite wild type levels of mRNA. This work suggests that the *tvrm148* mouse may serve as a model for LCA2 and can thus help in the understanding of the basic biology of the visual cycle and aid in the future development of treatment strategies for the disease. Although our work does not resolve the apparent difference in the genetic map and current gene sequence assembly, it does demonstrate the *tvrm148* mutation as a causative lesion in *Rpe65*.

FIGURE 11. Immunoblots and densitometry for RPE65 in *Rpe65* $^{+/+}$, $T^{+/-}$, and $T^{-/-}$ mice show mutant mice have reduced protein levels. Protein extracts harvested from P60 *Rpe65* $^{+/+}$ and $T^{-/-}$ RPE. (A) Immunoblotting for RPE65 in *Rpe65* $^{+/+}$, $T^{+/-}$, and $T^{-/-}$ protein extracts showed a decrease in RPE65 levels in $T^{+/-}$ and $T^{-/-}$ mice but no change in protein size. Gapdh is shown as a loading control. (B) Densitometry of immunoblots showed $60 \pm 4.7\%$ *Rpe65* $^{+/+}$ levels of RPE65 protein in $T^{+/-}$ mice and $14.5 \pm 1.5\%$ protein levels in $T^{-/-}$ mice. Data are presented as percentage change from *Rpe65* $^{+/+}$ (mean $\pm$ SD).

Acknowledgments

Supported by National Institutes of Health (NIH) Grants P30EY006360, R01EY016470, T32EY007092, R01EY014026, R01EY016435, R01EY021592, and a grant from the Abraham J. & Phyllis Katz Foundation (JHB), an unrestricted Departmental Award from Research to Prevent Blindness, the Intramural Research Program of the National Eye Institute, NIH (Z01EY000444). The authors alone are responsible for the content and writing of the paper.

Disclosure: **C.B. Wright**, None; **M.A. Chrenek**, None; **S.L. Foster**, None; **T. Duncan**, None; **T.M. Redmond**, None; **M.T. Pardue**, None; **J.H. Boatright**, None; **J.M. Nickerson**, None

References

- Won J, Shi LY, Hicks W, et al. Mouse model resources for vision research. *J Ophthalmol*. 2011;2011:391384.
- Redmond TM, Yu S, Lee E, et al. Rpe65 is necessary for production of 11-cis-vitamin A in the retinal visual cycle. *Nat Genet*. 1998;20:344–351.
- Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc*. 2009;4:1073–1081.
- Ng PC, Henikoff S. Predicting the effects of amino acid substitutions on protein function. *Ann Rev Genomics Hum Genet*. 2006;7:61–80.
- Ng PC, Henikoff S. SIFT: predicting amino acid changes that affect protein function. *Nucleic Acids Res*. 2003;31:3812–3814.
- Ng PC, Henikoff S. Accounting for human polymorphisms predicted to affect protein function. *Nucleic Acids Res*. 2003;31:3812–3814.
- Ng PC, Henikoff S. Predicting deleterious amino acid substitutions. *Genome Res*. 2001;11:863–874.
- Adzhubei IA, Schmidt S, Peshkin L, et al. A method and server for predicting damaging missense mutations. *Nat Methods*. 2010;7:248–249.
- Ramensky V, Bork P, Sunyaev S. Human non-synonymous SNPs: server and survey. *Nucleic Acids Res*. 2002;30:3894–3900.
- Sunyaev SR, Eisenhaber F, Rodchenkov IV, Eisenhaber B, Tumanyan VG, Kuznetsov EN. PSIC: profile extraction from sequence alignments with position-specific counts of independent observations. *Protein Eng*. 1999;12:387–394.
- Mi H, Muruganujan A, Thomas PD. PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucleic Acids Res*. 2013;41:D377–D386.
- Flicek P, Aken BL, Beal K, et al. Ensembl 2008. *Nucleic Acids Res*. 2008;36:D707–D714.
- Gasteiger E, Gattiker A, Hoogland C, Ivanyi I, Appel RD, Bairoch A. ExPASy: The proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res*. 2003;31:3784–3788.
- Grantham R. Amino acid difference formula to help explain protein evolution. *Science*. 1974;185:862–864.
- Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution rates on mammalian phylogenies. *Genome Res*. 2010;20:110–121.
- Abecasis GR, Auton A, Brooks LD, et al. An integrated map of genetic variation from 1,092 human genomes. *Nature*. 2012;491:56–65.
- Reese MG, Eeckman FH, Kulp D, Haussler D. Improved splice site detection in Genie. *J Comput Biol*. 1997;4:311–323.
- Altschul SE, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*. 1990;215:403–410.
- Tabaska JE, Zhang MQ. Detection of polyadenylation signals in human DNA sequences. *Gene*. 1999;231:77–86.
- Ferrer-Costa C, Gelpi JL, Zamakola L, Parraga I, de la Cruz X, Orozco M. PMUT: a web-based tool for the annotation of pathological mutations on proteins. *Bioinformatics*. 2005;21:3176–3178.
- Bromberg Y, Rost B. SNAP: predict effect of non-synonymous polymorphisms on function. *Nucleic Acids Res*. 2007;35:3823–3835.
- Yue P, Li Z, Moulton J. Loss of protein structure stability as a major causative factor in monogenic disease. *J Mol Biol*. 2005;353:459–473.
- Yue P, Moulton J. Identification and analysis of deleterious human SNPs. *J Mol Biol*. 2006;356:1263–1274.
- Yue P, Melamud E, Moulton J. SNPs3D: candidate gene and SNP selection for association studies. *BMC Bioinformatics*. 2006;7:166.
- Capriotti E, Calabrese R, Casadio R. Predicting the insurgence of human genetic diseases associated to single point protein mutations with support vector machines and evolutionary information. *Bioinformatics*. 2006;22:2729–2734.
- Capriotti E, Fariselli P, Calabrese R, Casadio R. Predicting protein stability changes from sequences using support vector machines. *Bioinformatics*. 2005;21(suppl 2):ii54–ii58.
- Capriotti E, Fariselli P, Casadio R. I-Mutant2.0: predicting stability changes upon mutation from the protein sequence or structure. *Nucleic Acids Res*. 2005;33:W306–W310.
- Altschul SE, Madden TL, Schaffer AA, et al. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*. 1997;25:3389–3402.
- Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. Predicting the functional effect of amino acid substitutions and indels. *PLoS One*. 2012;7:e46688.
- Chang B, Hawes NL, Pardue MT, et al. Two mouse retinal degenerations caused by missense mutations in the beta-subunit of rod cGMP phosphodiesterase gene. *Vision Res*. 2007;47:624–633.
- Jablonski MM, Dalke C, Wang X, et al. An ENU-induced mutation in Rslh causes disruption of retinal structure and function. *Mol Vis*. 2005;11:569–581.
- Pardue MT, McCall MA, LaVail MM, Gregg RG, Peachey NS. A naturally occurring mouse model of X-linked congenital stationary night blindness. *Invest Ophthalmol Vis Sci*. 1998;39:2443–2449.
- Pardue MT, Phillips MJ, Yin H, et al. Neuroprotective effect of subretinal implants in the RCS rat. *Invest Ophthalmol Vis Sci*. 2005;46:674–682.
- Akula JD, Hansen RM, Martinez-Perez ME, Fulton AB. Rod photoreceptor function predicts blood vessel abnormality in retinopathy of prematurity. *Invest Ophthalmol Vis Sci*. 2007;48:4351–4359.
- Akula JD, Mocko JA, Moskowitz A, Hansen RM, Fulton AB. The oscillatory potentials of the dark-adapted electroretinogram in retinopathy of prematurity. *Invest Ophthalmol Vis Sci*. 2007;48:5788–5797.
- Weymouth AE, Vingrys AJ. Rodent electroretinography: methods for extraction and interpretation of rod and cone responses. *Prog Retin Eye Res*. 2008;27:1–44.
- Landers GM, Olson JA. Rapid, simultaneous determination of isomers of retinal, retinal oxime and retinol by high-performance liquid chromatography. *J Chromatogr*. 1988;438:383–392.
- Garwin GG, Saari JC. High-performance liquid chromatography analysis of visual cycle retinoids. *Methods Enzymol*. 2000;316:313–324.
- Douglas RM, Alam NM, Silver BD, McGill TJ, Tschetter WW, Prusky GT. Independent visual threshold measurements in the

- two eyes of freely moving rats and mice using a virtual-reality optokinetic system. *Vis Neurosci.* 2005;22:677-684.
40. Prusky GT, Alam NM, Beekman S, Douglas RM. Rapid quantification of adult and developing mouse spatial vision using a virtual optomotor system. *Invest Ophthalmol Vis Sci.* 2004;45:4611-4616.
 41. Carter-Dawson LD, LaVail MM. Rods and cones in the mouse retina. I. Structural analysis using light and electron microscopy. *J Comp Neurol.* 1979;188:245-262.
 42. Seeliger MW, Grimm C, Stahlberg F, et al. New views on RPE65 deficiency: the rod system is the source of vision in a mouse model of Leber congenital amaurosis. *Nat Genet.* 2001;29:70-74.
 43. Pang JJ, Chang B, Hawes NL, et al. Retinal degeneration 12 (rd12): a new, spontaneously arising mouse model for human Leber congenital amaurosis (LCA). *Mol Vis.* 2005;11:152-162.
 44. Samardzija M, von Lintig J, Tanimoto N, et al. R91W mutation in *Rpe65* leads to milder early-onset retinal dystrophy due to the generation of low levels of 11-cis-retinal. *Hum Mol Genet.* 2008;17:281-292.
 45. Jacobson SG, Aleman TS, Cideciyan AV, et al. Defining the residual vision in Leber congenital amaurosis caused by RPE65 mutations. *Invest Ophthalmol Vis Sci.* 2009;50:2368-2375.
 46. Cachafeiro M, Bemelmans AP, Canola K, et al. Remaining rod activity mediates visual behavior in adult *Rpe65*^{-/-} mice. *Invest Ophthalmol Vis Sci.* 2010;51:6835-6842.
 47. Caruso RC, Aleman TS, Cideciyan AV, et al. Retinal disease in *Rpe65*-deficient mice: comparison to human leber congenital amaurosis due to RPE65 mutations. *Invest Ophthalmol Vis Sci.* 2010;51:5304-5313.
 48. Pasadhika S, Fishman GA, Stone EM, et al. Differential macular morphology in patients with RPE65-, CEP290-, GUCY2D-, and AIPL1-related Leber congenital amaurosis. *Invest Ophthalmol Vis Sci.* 2010;51:2608-2614.
 49. Chen Y, Moiseyev G, Takahashi Y, Ma JX. Impacts of two point mutations of RPE65 from Leber's congenital amaurosis on the stability, subcellular localization and isomerohydrolase activity of RPE65. *FEBS Lett.* 2006;580:4200-4204.
 50. Rohrer B, Lohr HR, Humphries P, Redmond TM, Seeliger MW, Crouch RK. Cone opsin mislocalization in *Rpe65*^{-/-} mice: a defect that can be corrected by 11-cis retinal. *Invest Ophthalmol Vis Sci.* 2005;46:3876-3882.
 51. Zhang T, Zhang N, Baehr W, Fu Y. Cone opsin determines the time course of cone photoreceptor degeneration in Leber congenital amaurosis. *Proc Natl Acad Sci U S A.* 2011;108:8879-8884.
 52. Umino Y, Solessio E, Barlow RB. Speed, spatial, and temporal tuning of rod and cone vision in mouse. *J Neurosci.* 2008;28:189-198.
 53. Ablonczy Z, Crouch RK, Goletz PW, et al. 11-cis-retinal reduces constitutive opsin phosphorylation and improves quantum catch in retinoid-deficient mouse rod photoreceptors. *J Biol Chem.* 2002;277:40491-40498.
 54. Cachafeiro M, Bemelmans A-P, Canola K, et al. Remaining rod activity mediates visual behavior in adult *Rpe65*^{-/-} mice. *Invest Ophthalmol Vis Sci.* 2010;51:6835-6842.
 55. Tang PH, Wheless L, Crouch RK. Regeneration of photopigment is enhanced in mouse cone photoreceptors expressing RPE65 protein. *J Neurosci.* 2011;31:10403-10411.
 56. Fan J, Rohrer B, Moiseyev G, Ma JX, Crouch RK. Isorhodopsin rather than rhodopsin mediates rod function in RPE65 knockout mice. *Proc Natl Acad Sci U S A.* 2003;100:13662-13667.
 57. Van Hooser JP, Liang Y, Maeda T, et al. Recovery of visual functions in a mouse model of Leber congenital amaurosis. *J Biol Chem.* 2002;277:19173-19182.
 58. Wang JS, Kefalov VJ. The cone-specific visual cycle. *Prog Retin Eye Res.* 2011;30:115-128.
 59. Takahashi Y, Moiseyev G, Chen Y, Nikolaeva O, Ma JX. An alternative isomerohydrolase in the retinal Muller cells of a cone-dominant species. *FEBS J.* 2011;278:2913-2926.
 60. Kaylor JJ, Yuan Q, Cook J, et al. Identification of DES1 as a vitamin A isomerase in Muller glial cells of the retina. *Nat Chem Biol.* 2013;9:30-36.
 61. Kiser PD, Golczak M, Lodowski DT, Chance MR, Palczewski K. Crystal structure of native RPE65, the retinoid isomerase of the visual cycle. *Proc Natl Acad Sci U S A.* 2009;106:17325-17330.
 62. Golczak M, Kiser PD, Lodowski DT, Maeda A, Palczewski K. Importance of membrane structural integrity for RPE65 retinoid isomerization activity. *J Biol Chem.* 2010;285:9667-9682.
 63. Kiser PD, Palczewski K. Membrane-binding and enzymatic properties of RPE65. *Prog Retin Eye Res.* 2010;29:428-442.
 64. Kiser PD, Farquhar ER, Shi W, Sui X, Chance MR, Palczewski K. Structure of RPE65 isomerase in a lipidic matrix reveals roles for phospholipids and iron in catalysis. *Proc Natl Acad Sci U S A.* 2012;109:E2747-E2756.
 65. Gu SM, Thompson DA, Srikumari CR, et al. Mutations in RPE65 cause autosomal recessive childhood-onset severe retinal dystrophy. *Nat Genet.* 1997;17:194-197.
 66. Nikolaeva O, Moiseyev G, Rodgers KK, Ma JX. Binding to lipid membrane induces conformational changes in RPE65: implications for its isomerohydrolase activity. *Biochem J.* 2011;436:591-597.
 67. Marlhens F, Bareil C, Griffioen JM, et al. Mutations in RPE65 cause Leber's congenital amaurosis. *Nat Genet.* 1997;17:139-141.
 68. Morimura H, Fishman GA, Grover SA, Fulton AB, Berson EL, Dryja TP. Mutations in the RPE65 gene in patients with autosomal recessive retinitis pigmentosa or leber congenital amaurosis. *Proc Natl Acad Sci U S A.* 1998;95:3088-3093.
 69. Philp AR, Jin M, Li S, et al. Predicting the pathogenicity of RPE65 mutations. *Hum Mutat.* 2009;30:1183-1188.
 70. Stone EM. Leber congenital amaurosis - a model for efficient genetic testing of heterogeneous disorders: LXIV Edward Jackson Memorial Lecture. *Am J Ophthalmol.* 2007;144:791-811.
 71. Sundaresan P, Vijayalakshmi P, Thompson S, Ko AC, Fingert JH, Stone EM. Mutations that are a common cause of Leber congenital amaurosis in northern America are rare in southern India. *Mol Vis.* 2009;15:1781-1787.
 72. Pang J, Boye SE, Lei B, et al. Self-complementary AAV-mediated gene therapy restores cone function and prevents cone degeneration in two models of *Rpe65* deficiency. *Gene Ther.* 2010;17:815-826.