Published in final edited form as:

J Comp Neurol. 2013 October 1; 521(14): . doi:10.1002/cne.23340.

Amygdala projections to the lateral bed nucleus of the stria terminalis in the Macaque: comparison with ventral striatal afferents

Danielle M. deCampo¹ and Julie L. Fudge^{1,2,*}

¹Institutional Affiliations: Department of Neurobiology and Anatomy

²Department of Psychiatry

Abstract

The lateral bed nucleus of the stria terminalis (BSTL) is involved in mediating anxiety-related behaviors to sustained aversive stimuli. The BSTL forms part of the central extended amygdala, a continuum composed of the BSTL, the amygdala central nucleus, and cell columns running between the two. The central subdivision (BSTLcn), and the juxtacapsular subdivision (BSTLj) are two BSTL regions that lie above the anterior commissure, near the ventral striatum. The amygdala, a heterogeneous structure that encodes emotional salience, projects to both the BSTL and ventral striatum. We placed small injections of retrograde tracers into the BSTL, focusing on the BSTLcn and BSTLj, and analyzed the distribution of labeled cells in amygdala subregions. We compared this to the pattern of labeled cells following injections into the ventral striatum. All retrograde results were confirmed by anterograde studies. We found that the BSTLcn receives stronger amygdala inputs relative to the BSTLj. Furthermore, the BSTLcn is defined by inputs from the corticoamygdaloid transition area and central nucleus, while the BSTLj receives inputs mainly from the magnocellular accessory basal and basal nucleus. In the ventral striatum, the dorsomedial shell receives inputs that are similar, but not identical, to inputs to the BSTLcn. In contrast, amygdala projections to the ventral shell/core are similar to projections to the BSTLj. These findings indicate that the BSTLcn and BSTLj receive distinct amygdala afferent inputs and that the dorsomedial shell is a transition zone with the BSTLcn, while the ventral shell/core are transition zones with the BSTLj.

Keywords

extended amygdala; corticoamygdaloid transition region; juxtacapsular; oval; amygdalopiriform transition area; dorsomedial shell; anxiety

© 2013 Wiley Periodicals, Inc.

*Correspondence University of Rochester Medical Center 601 Elmwood Avenue Rochester, NY 14642 P: 585 273 2028 F: 585 756 5334 Julie_Fudge@urmc.rochester.edu.

ROLE OF AUTHORS

All authors had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. J.F. and D.D. designed the experiments and wrote the manuscript. J.F. and D.D. performed the surgeries and D.D. did the analysis with assistance from J.F.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

INTRODUCTION

The bed nucleus of the stria terminalis (BST) forms part of the extended amygdala, a forebrain continuum that mediates autonomic, motor and neuroendocrine responses to emotionally relevant stimuli. The ‘central extended amygdala’ is comprised of the lateral bed nucleus of the stria terminalis (BSTL), the central nucleus of the amygdala (CeN) and columns of cells running between the two structures, both within the stria terminalis and in the subpallidal region (central sublenticular extended amygdala, SLEAc) (Alheid and Heimer, 1988). The BSTL and CeN are each composed of multiple subdivisions whose internal organization is similar, mirroring one another in terms of cell type and histochemistry (Heimer et al., 1999; Martin et al., 1991). The main components of the BSTL are the large central subdivision (BSTLcn), the thin capsule that surrounds it (BSTLc), the juxtacapsular subdivision (BSTLj), and the posterior subdivision (BSTLP). The BSTLD is a cellular matrix surrounding the BSTLcn and BSTLc, and is often seen just beneath the ventromedial caudate nucleus.

Mounting evidence indicates that the CeN and the BSTL mediate related, but distinct responses to aversive stimuli. While the CeN mediates rapid behavioral and autonomic fear responses to immediate threats and conditioned cues, the BSTL mediates sustained fear responses (likened to anxiety) to threat cues maintained over long periods of time, or cues signaling a potential threat (Fendt et al., 2003; Sullivan et al., 2004; Walker and Davis, 1997; Zimmerman and Maren, 2011). Recent studies in primates and humans support this distinction, showing that the BSTL is involved in ‘threat monitoring’ and may be responsible for individual differences in anxiety (Duvarci et al., 2009; Fox et al., 2008; Somerville et al., 2010).

The BSTL resides behind the ventral striatum, where it merges with the ventral striatal shell. The ventral striatum is classically divided into shell and core subterritories based on histochemical, connectional, and pharmacologic features. The shell, which is directly adjacent to the BST, is further subdivided into dorsomedial, ventral and lateral components based on behavioral and pharmacologic studies (Ikemoto et al., 2005; Kelley, 1999; Meredith et al., 1996; Reynolds and Berridge, 2002). In rodents, some authors have noted similarities between the dorsomedial shell and the central extended amygdala with respect to cell morphology, histochemical features and connections, and have described this region as a transition zone ((Alheid and Heimer, 1988; Heimer et al., 1997) but see (Zahm, 1998)). Across species, both the BSTL and ventral striatum receive strong inputs from the amygdala (Fudge et al., 2004; Fudge et al., 2002; Novotny, 1977).

The amygdala is a complex structure formed by multiple subnuclei. Specific amygdala subnuclei send differential projections to the ventral striatal subterritories, along a gradient from the dorsomedial shell to the ventrolateral shell and core (Friedman et al., 2002; Fudge et al., 2002; Russchen et al., 1985b). Previously, we noted that amygdala inputs to the dorsomedial shell in the monkey are distinct from inputs to more ventral and lateral shell regions, with inputs to the dorsomedial shell arising more from the corticomедial amygdala and corticoamygdaloid transition area (CTA), in contrast to the well-described basal and accessory basal nucleus inputs to the ventral and lateral shell and core (Fudge et al., 2002; Russchen et al., 1985b).

Given the importance of the BSTL and ventral striatum for emotion-related behaviors, we had two goals for this study. The first was to determine the organization of amygdala afferents to BSTL subterritories in nonhuman primates using retrograde and anterograde tracing studies. We hypothesized that discrete subdivisions of the BSTL receive differential amygdala inputs just as specific subdivisions of the ventral striatum receive differential

amygdala afferent projections. In addition, because of the BSTL's close proximity to the ventral striatum, the second goal of the study was to compare amygdaloid projections to the main BSTL subdivisions with those of adjacent striatal territories. We focused specifically on the BSTLcn and the BSTLJ, which lie in close proximity to the ventral striatum and are thought to play a role in responses to stressful stimuli.

MATERIALS AND METHODS

Injection sites

For the retrograde studies, we placed small injections (40 nL) of the tracers, Lucifer yellow conjugated to dextran amine (LY; 10%, Molecular Probes, Eugene, OR), tetramethylrhodamine, conjugated to dextran amine ('fluoruby', FR; 4%, Molecular Probes), fluorescein conjugated to dextran amine (FS; 10%, Molecular Probes), or wheat-germ agglutinin-horse radish peroxidase (WGA; 10%, Sigma, St. Louis, MO) diluted in 0.1M PO₄ into the BST and ventral striatum. In a separate group of animals, anterograde injections into the amygdala were placed using 40 nL of the tracers, LY, FR, FS, or 200 nL of tritiated amino acids (AA) (1:1 ratio of tritiated proline and leucine: PerkinElmer, Boston, MA). Previous studies have indicated that there is no cross-reactivity of antibodies to FR, FS, WGA and LY (Haber et al., 2000). In our experience FR, FS, LY and AA have similar anterograde properties, and are sensitive indicators of terminal fields when survival is between 10–14 days. In the retrograde studies FR, FS, and LY have similar retrograde properties. However, WGA injected into similar regions results in relatively more labeled cells, albeit in a similar distribution.

Surgery

Nineteen Old World male primates (*Macaca fascicularis*, n = 18; *Macaca nemestrina*, n = 1) (2–3 years old) weighing between 2.5 kg – 4.9 kg were used in this study (World Wide Primates, Tallahassee, FL, USA; Three Springs Laboratories, Pekaski, PA; Labs of Virginia, Yemassee, SC; Alpha Genesis, Yemassee, SC). All experiments were carried out in accordance with National Institute of Health guidelines. Experimental design and techniques were aimed at minimizing animal use and suffering and were reviewed by the University of Rochester Committee on Animal Research. To minimize animal use, several of these injections were performed as parts of other studies (Fudge et al., 2004; Fudge et al., 2012; Fudge and Tucker, 2009). Stereotaxic coordinates were determined prior to injection using MRI imaging (3Telsa, coronal sections, 0.8 mm thick, 0.1 mm apart).

Initial anesthesia was administered by intramuscular injections of ketamine hydrochloride (10mg/kg). Animals were intubated, and a deep plane of anesthesia was induced by isoflurane gas, maintained as needed during surgery. A craniotomy was performed and the bone flap removed to visualize cortical surface landmarks. Small deposits of tracer were pressure-injected over 10–15 min into the BSTL or ventral striatum using a 0.5 µL Hamilton syringe with an attached glass pipette (tip diameter at 40 µm) (Hamilton Company, Reno, NV). The syringe was left in place for an additional 20 minutes to prevent leakage of tracer up the needle track. Only one injection for each type of tracer was used per animal. Following injection placement, the bone flap was replaced, and overlying muscles and skin sutured. Anterograde injections of tracer into various amygdala subnuclei were performed as part of previous studies, and were used for comparison (Fudge et al., 2012; Fudge and Tucker, 2009).

Two weeks after surgery, animals were deeply anesthetized and killed by perfusion through the heart with 0.9% saline containing 0.5 ml of heparin sulfate (200 ml/min for 10 minutes), followed by cold 4% paraformaldehyde in a 0.1 M phosphate buffer/30% sucrose solution

(100 ml/min for 1 h). The brain was extracted from the skull, placed in a fixative overnight, and then put through increasing gradients of sucrose (10%, 20%, 30%). Brains were cut on a freezing microtome (40 μ m) and all sections were stored in cryoprotectant solution (30% ethylene glycol and 30% sucrose in 0.1 M phosphate buffer) at -20°C (Rosene et al., 1986).

Immunocytochemistry (Table 1)

Characterization of antibodies—The antibodies used for detection of the specific tracer molecules have been documented in numerous tract-tracing studies (Lanciego and Wouterlood, 2011). The pattern of labeled cells and fibers in the amygdala and BST, respectively, for all tracer/antibody combinations was similar to previously reported results (e.g. (Fudge et al., 2002; Grove, 1988; Ottersen and Ben-Ari, 1979; Russchen et al., 1985a) see also *Technical considerations*, Discussion). The pattern of calbindin-D28K (CaBP) expression was identical to published reports for our regions of interest as well as in other brain areas such as the substantia nigra (Cote et al., 1991; Meredith et al., 1996). Finally, the pattern of enkephalin, somatostatin, and neurotensin immunoreactivity in the basal forebrain matched previously published results in human and monkey (Fudge and Haber, 2001; Heimer et al., 1999; Mai et al., 1987; Martin et al., 1991).

Tracers—For retrograde analyses, 1:8 sections through the brain were immunostained for tracer. Sections were rinsed in 0.1 M phosphate buffer (pH, 7.2) with 0.3 Triton-X (PB-TX) and treated with peroxidase inhibitor for 5 min, followed by more rinses in 0.1 M PB-TX. Sections were then preincubated in a blocking solution of 10% normal goat serum (NGS) in 0.1 M PB-TX for 30 min. Tissue was placed in primary antisera to LY (1:2000, Molecular Probes, rabbit), FS (1:2000, Molecular Probes, rabbit), FR (1:1000, Molecular Probes, rabbit), and WGA (1:2000, Sigma, rabbit) for 96 h at 4°C . After thorough rinsing with 0.1 M PB-TX, and preincubation with 10% NGS-PB-TX, sections were incubated in biotinylated anti-rabbit secondary antibody (1:200; Vector, Burlingame, CA). Tracers were visualized using avidin-biotin reaction (Vector ABC Standard kit, Burlingame, CA). Additional compartments for each case were processed for tracer and counterstained for cresyl violet. Detection of these tracer molecules in tract-tracing has been extensively documented (see *Technical Considerations*, Discussion).

Enkephalin, somatostatin, neurotensin and CaBP-immunoreactivity (IR)—To identify the relative location of the BSTL injection sites, the distribution of retrogradely labeled cells in specific amygdala nuclei, and the distribution of anterogradely labeled fibers in BSTL subdivisions, we used a variety of histochemical stains in adjacent sections, including cresyl violet, acetylcholinesterase (AChE), enkephalin (ENK), neurotensin (NT) and somatostatin (ST). ENK, ST, NT, and AChE were used to distinguish BSTL and amygdala subnuclei (Amaral and Bassett, 1989; Heimer et al., 1999; Martin et al., 1991; Walter et al., 1991). For example, ST-IR was used to distinguish the BSTLcn, where it is high relative to the adjacent BSTLj. Both NT and ST demarcated the SLEAc from the neighboring globus pallidus (see Results). CaBP-IR was chosen to identify the boundaries of the globus pallidus versus the SLEAc, and the boundary between shell and core subterritories of the ventral striatum (Chang and Kuo, 1991; Cote et al., 1991; Meredith et al., 1996).

For visualization of neuropeptide and CaBP proteins, sections were rinsed in 0.1 M PB-TX, preincubated in 10% NGS-PB-TX as described above, and then incubated for 96 h at 4°C in CaBP (Sigma, 1: 10,000, mouse), ENK (Immunostar, 1:1000, rabbit), ST (Immunostar, 1:1000, rabbit), or NT (Immunostar, 1:1000, rabbit). Sections were then rinsed, blocked with 10% NGS-TX, and incubated with the appropriate secondary biotinylated mouse or rabbit antibody. Following rinsing, the molecules were visualized using avidin-biotin reaction as

described above. AChE staining was performed using the Geneser technique (Geneser-Jensen and Blackstad, 1971).

Autoradiography

Tissue sections were mounted on chrome alum gelatin-coated slides, dehydrated in increasing gradients of alcohol, and cleared in xylene for 48 h. Slides were then dipped in Kodak NTB2 photographic emulsion (Kodak, Rochester, NY) and exposed for 4–6 months in a light-tight box at -20°C . Slides were then developed in cold Kodak D19 developer (15°C) for 2.5 min, fixed, washed and counterstained with cresyl violet.

Analysis

Retrograde injection sites—The relative position of the retrograde injection sites was confirmed by charting the injection site and an adjacent neuropeptide-immunoreactive or AChE-stained section. We also analyzed the localization of the injection site by inspecting the pattern of projections to or from other brain regions based on the literature. For example, injection sites that included the BSTLcn resulted in anterogradely labeled fibers in the parabrachial nucleus and paraventricular nucleus of the hypothalamus, while injection sites in the BSTLJ had few to no anterogradely labeled fibers in these areas, and resulted in projections to the dorsal tier of the substantia nigra, pars compacta (Dong et al., 2001b; Fudge and Haber, 2001; Moga et al., 1989). Injection sites in the BSTL that spread into the overlying striatum resulted in labeled fibers in the globus pallidus, and were classified as such.

The distribution of retrogradely labeled cells in specific amygdala nuclei was charted for each case individually using Neurolucida (Microbrightfield, Rutland, VT) under 10x objective with brightfield illumination. The boundaries of various amygdaloid subregions, along with identifying landmarks such as blood vessels and fiber tracts were then hand-drawn from adjacent sections stained with cresyl violet or AChE using a Jena macroprojector. Boundary charts were converted into digital form using a drawing tablet interfaced with the program Adobe Illustrator CS2 (Adobe Systems, San Jose, CA), and carefully aligned with cell distribution charts, matching blood vessels and fiber tracts for reference. Additional sections counterstained with cresyl violet were used to confirm these data.

To compare the relative distribution of labeled cells in specific amygdala nuclei resulting from different injection sites, we created composite charts using transparent layers in Adobe Illustrator CS2. Although WGA results in relatively more labeled cells compared to similar injections of LY, FS, and FR, the distribution pattern is the same. Digital charts of retrogradely labeled cells and nuclear boundaries were matched for rostrocaudal level across cases, and then carefully overlaid in Adobe Illustrator CS2 (Adobe Systems, San Jose CA). Landmarks such as fiber tracts, as well as subnuclear boundaries previously charted from AChE-stained adjacent sections, were used for alignment.

Anterograde injection sites—The location of anterograde injection sites in specific amygdala nuclei was determined using adjacent sections stained for AChE and cresyl violet (deCampo and Fudge, 2012; Fudge et al., 2012; Fudge and Tucker, 2009). The distribution of labeled fibers in the BSTL subdivisions was determined using a 20x objective in conjunction with brightfield and darkfield illumination. Hand-drawn charts were created, scanned into the computer using Adobe Photoshop CS2, and imported into an Adobe Illustrator CS2 format. Adjacent sections immunoreacted for BSTL subdivision-specific markers were used to localize labeled fibers within specific subdivisions.

Digital photography for brightfield and darkfield images was done with an Insight 4 Megasample Camera mounted on the microscope using SPOT imaging software 5.0 (Sterling Heights, Michigan, USA). Captured images were optimized in Adobe Photoshop using the brightness/contrast tool.

RESULTS

Nomenclature of the BST subdivisions (Figure 1)

We used classification schema by Heimer and others, which recognizes two major divisions, the lateral (BSTL) and medial (BSTM) in monkeys and humans (De Olmos, 2004; Heimer et al., 1999; Lesur et al., 1989; Martin et al., 1991). The distribution and intensity of markers used in these primate studies and the present results are similar. The focus of this study, the BSTL, is comprised of several subdivisions that are distinguished by histochemical properties. Ventrally, the BSTL extends below the globus pallidus and is contiguous with the ‘central division of the sublenticular extended amygdala’ (SLEAc).

BSTLcn, BSTLc, BSTLD—The BSTLD is the rostral-most subdivision of the BSTL, in which the BSTLcn, the BSTLc, and a few smaller ‘accessory’ islands are embedded. The BSTLc, which surrounds the BSTLcn, is composed of cells embedded in fibers of the stria terminalis (Figure 1A, D, G, J). The BSTLcn is an oval cellular mass that extends along the entire rostrocaudal extent of the BSTL, although it is most clearly seen at central to caudal levels (Figure 1C, F, I, L). The BSTLcn exhibits low to moderate staining for AChE, and is surrounded by the BSTLc which, because of its fibrous components, is AChE-poor (Figure 1A–C). In contrast, in ENK preparations, the BSTLcn stains prominently as an oval shaped mass surrounded by thick, ENK-positive fibers of the BSTLc (Figure 1D–F). Similarly, NT and ST immunostaining reveal the BSTLcn as a strongly labeled oval mass containing numerous labeled cells and fibers, encapsulated by a darkly stained BSTLc (Figure 1G–I, J–L).

BSTLJ—The BSTLJ is located between the internal capsule laterally, and the BSTLcn medially. It merges with the overlying BSTLD and striatum adjacent to the internal capsule. It is easily distinguished from the BSTLcn by high AChE activity and distinct immunostaining characteristics. Visualized with AChE staining, the BSTLJ stains darkly as a narrow strip, similar to the overlying striatal mass (Figure 1A–C). In contrast to the BSTLcn, ENK-IR, ST-IR and NT-IR are low in the BSTLJ (Figure 1D–L).

BSTLP—The BSTLP extends ventrally from the BSTLcn to the anterior commissure, and continues caudally to merge with the SLEAc. Supracommissurally, the BSTLP is situated between the BSTM and the BSTLcn (Figure 1B, E, H, K), and exhibits moderate staining for AChE, ST, and NT (Figure 1A–C, G–I, J–L). In ENK preparations, the BSTLP also has moderate staining (Figure 1E), which decreases caudally (Figure 1F).

SLEAc—The SLEAc is a column of cells ventral to the globus pallidus (known as the ‘lenticular nucleus’ in older literature) that links the BSTL with the CeN. The SLEAc appears to extend from the BSTLP and like it, is also clearly visualized with NT staining, appearing as a column of cells sweeping laterally through the basal forebrain (Figure 1J–L). Unlike the BSTLP however, the SLEAc also stains darkly for ENK. The SLEAc is easily differentiated from the adjacent globus pallidus using CaBP staining, which is low in the SLEAc but moderate in the globus pallidus (not shown).

Amygdaloid subnuclei (Figure 2)

Amygdala subnuclei are described according to Price and colleagues (Price et al., 1987), with two exceptions. We refer to the transition region between the medial basal nucleus, the periamygdaloid cortex, and the entorhinal cortex as the ‘corticoamygdaloid transition area’ (CTA), a name given by Crosby and Humphrey (Crosby and Humphrey, 1941). For the CeN, we use the naming schema of de Olmos (De Olmos, 1990).

The CeN forms the caudal pole of the central extended amygdala. It is located in the dorsal amygdala and merges rostrally with the anterior amygdaloid area (AAA), and the SLEAc (Figure 2B–D). The CeN is subdivided into medial (CeM) and lateral divisions, the latter comprised of the lateral central (CeLcn) and lateral capsular (CeLc) subdivisions. The CeLcn is an oval region with low AChE activity, and is surrounded by fibers and cells of the CeLc (Figure 2F–H). The CeM is a funnel shaped area that sweeps medially around the CeLcn and contains moderate AChE activity, distinguishing it from the CeLcn (Figure 2F–H). In AChE staining, as well as with histochemical staining with neuropeptide markers (not shown), the CeM and CeLcn mirror the pattern in the BSTLP and BSTLcn, respectively (Figure 1A–C).

The remaining amygdala subnuclei are divided into ‘deep’ and ‘superficial’ nuclear groups. The ‘deep’ nuclear group includes the lateral, basal, and accessory basal nuclei. The basal nucleus extends along the rostrocaudal extent of the amygdala, and is subdivided into magnocellular (Bmc), intermediate (Bi) and parvicellular (Bpc) subdivisions. AChE activity is highest in the Bmc and decreases gradually in the Bi and Bpc (Figure 2E–G), paralleling the gradual decrease in pyramidal cell size from the Bmc to Bpc (Figure 2B, C). At all rostrocaudal levels, the basal nucleus is intimately associated with the paralaminar nucleus (PL), a sheet of heterogeneous cells that wraps around the Bpc rostrally and caudally (Figure 2A–D, I). The PL breaks up into cell clusters and merges with the Bpc and periamygdaloid cortex (PAC) medially, at the level of the amygdaloid fissure (semiannular sulcus) (Figure 2I, arrowhead). This confluence creates a highly undifferentiated cortical bridge between the Bpc laterally, the entorhinal cortex ventrally and the PAC dorsally (Crosby and Humphrey, 1941; deCampo and Fudge, 2012; Rosene and Van Hoesen, 1987) (Figure 2B–C, F–G, I asterisk). We refer to this region as the ‘corticoamygdaloid transition area (CTA)’, to emphasize the lack of clear boundaries that exist between the PAC, entorhinal cortex, and the Bpc and PL.

The last of the deep nuclei, the accessory basal nucleus, is composed of the magnocellular (ABmc), parvicellular (ABpc), and superficial (ABs) subdivisions. The ABs has also been referred to as the rostral most extension of the amygdalohippocampal area (Amaral et al., 1992; De Olmos, 2004) (Figure 2G). The ABmc and ABs are both marked by high AChE staining, while the ABpc has low levels of AChE (Figure 2F–H).

The superficial amygdala includes the anterior (CoA) and posterior (CoP) cortical nuclei, the medial nucleus (M), nucleus of the lateral olfactory tract (NLOT), the periamygdaloid cortex (PAC), and amygdalohippocampal area (AHA). All of these regions except the NLOT and the AHA have low AChE activity (Figure 2E–H).

Intercalated islands are collections of small, tightly packed cell clusters interspersed in fiber bundles separating and surrounding the major amygdala subdivisions. They are best seen with Nissl staining under high power magnification (not shown).

Retrograde cases

BSTL injections

Relative placement of retrograde injections in BSTL subdivisions (Figure 3A, Table 2):

There were a total of seven injections in the BSTL. Three injections were centered in the BSTLcn (J24FS, J28WGA, J29WGA), with some encroachment on adjacent subdivisions but no leakage into the striatum (Figure 3A, light gray). One small injection landed in the BSTLD, centered between the BSTLcn and BSTLJ, with slight encroachment on the BSTLP (J28LY, light gray). Three cases were centered in the BSTLJ (J28FS, J28FR, J29FR), none of which spread into the BSTLcn or BSTLP (Figure 3A, dark gray). However, two of these injections spread into the overlying ventromedial caudate nucleus (J28FR, J29FR).

Distribution of retrogradely labeled cells in the amygdala

BSTLcn, BSTLc, and BSTLD: Cases J29WGA (Figure 4A), J24FS, J28WGA, J28LY (not shown): The BSTLcn was involved in the injection site in all cases except J28LY. In these instances, the distribution of retrogradely labeled cells was similar, and representative case J29WGA is described in detail (Figure 4A). The majority of labeled cells were concentrated in the CTA and medial Bpc, the ABs, and both the CeM and CeLcn (Figure 5A–F). There were also a few labeled cells interspersed in the fibers of the CeLc. The relatively high number of labeled cells in the CTA was particularly striking, appearing as a dark band of cells, sweeping from the medial Bpc towards the semiannular sulcus (Figure 5G–I, arrowheads).

Moderate numbers of labeled cells occupied the Bi and Bmc, and were aligned along the medial border of the basal nucleus. Moderate numbers of retrogradely labeled cells were also found in the ABpc and ABmc. Labeled cells in the ABpc were found along its ventral border, adjacent to cells in the Bi and Bpc. Low concentrations of retrogradely labeled cells were found in the PAC, with a few scattered retrogradely labeled cells found in the CoP and M.

In contrast, a tiny injection placed mainly in the BSTLD, and involving little of the BSTLcn (case J28LY), had relatively fewer labeled cells in the amygdala. These predominated in the ABmc, Bi, and Bpc, similar to injections into the BSTLJ (described below).

BSTLJ: Cases J28FS (Figure 4B), J29FR, J28FR (not shown): The distribution of labeled cells was similar among all cases involving the BSTLJ, including cases in which the injection site spread dorsally into the striatum. Case J28FS, with an injection confined to the BSTLJ, is described in detail (Figure 4B). Overall, injections in the BSTLJ resulted in a more restricted distribution of labeled cells than cases involving injections in the BSTLcn. In the BSTLJ, the majority of labeled cells were in the ABmc (see also Figure 5G–I), followed by moderate numbers of labeled cells in the Bmc, Bi and the medial-central Bpc (Figure 4B). Throughout the basal nucleus, labeled cells were dispersed medially and centrally, with a few labeled cells in the ventrolateral Bpc. There were scattered labeled cells in the medial Bpc/CTA, CeM, PAC, AAA and AHA, and few to no cells in the remaining nuclei.

Ventral Striatal Injections

Ventral striatum injection site placement (Figure 3B, Table 3): There were seven injections placed at different positions in the ventral striatum. Four injections were centered in the shell of the ventral striatum (J24WGA, J26FS, J27FS, J27FR). Case J24WGA was centered in the dorsomedial shell with slight spread into the ventromedial caudate (Figure 3B, light gray). Cases J26FS, J27FS and J27FR were confined to the ventral shell (Figure

3B, gray). There was also one injection site confined to the core of the ventral striatum (J13WGA) (Figure 3B black), one injection in the ventromedial caudate nucleus (J24FR) and one injection into the central caudate nucleus (J25FS) (Figure 3B, white).

Distribution of labeled cells in the amygdala

Dorsomedial shell: Case J24WGA (Figure 6A): High concentrations of retrogradely labeled cells were found in the CTA, medial Bpc, ABmc, ABs, and AHA. Moderate numbers of cells were present in the CoP, CeM, M, the PAC and the ABpc. There were few to no cells in the CeLcn, CeLc or Bmc.

Ventral shell: Cases J27FS (Figure 6B), J26FS, J27FR (not shown): Retrogradely labeled cells following all ventral shell injections were mainly confined to ABmc and central Bpc and Bi. There were a few scattered cells found in the medial Bpc and ABpc. A few labeled cells in the CeM were seen after the more ventral injection sites (J26FS and J27FR), but not in case J27FS. Overall, the pattern of labeled cells in the amygdala was similar to the distribution of labeled cells seen after BSTLJ injections.

Ventral core: Case J13WGA (not shown): Compared to the ventral shell injections, an injection placed in the core resulted in relatively fewer labeled cells, but in the same distribution, mainly in the ABmc and Bi. A few scattered cells were also found in the Bmc, Bpc and CTA.

Ventromedial and central caudate nucleus: Cases J24FR, J25 FS (not shown): Compared to other striatal injections, the concentration of labeled cells following caudate injections was less than that seen after other ventral striatal injections. Labeled cells were mainly found in the ABmc at the midlevel of the amygdala, with a few scattered FR-positive cells found in the Bmc, Bi and L.

Anterograde cases

Injection site placement in the amygdala subregions (Figure 7): We used 20 injections of bidirectional tracer into the amygdala to assess the pattern of labeled fibers within specific BSTL subdivisions and the ventral striatum. There were eight injection sites in the basal nucleus: three were in the Bmc (J12FR, J12LY, J16LY), and five in the Bpc (J20LY, J15LY, J14FS, J15FS, J14FR). In the Bmc, injections in cases J12FR and J16LY were located medially within the Bmc, with the injection in case J16LY located dorsal to that in J12FR. Case J12LY had an injection that was mostly in the lateral Bmc, but also included part of the CeM. In the Bpc, three injections (J20LY, J15FS, J14FS, asterisks) were located in the vicinity of the medial Bpc/CTA. The injections in cases J15FS and J14FS were caudal to the injection in case J20LY. Case J15LY had an injection in the central Bpc, and the injection site in case J14FR was located in the ventrolateral Bpc. Five injections were placed in the accessory basal nucleus: two were confined to the ABmc (J8LY and J12FS), one was confined to the ABpc (J18FR), and two larger injections encompassed both the ABmc and ABpc (J21FR, J20FS). In the ABmc injections, the injection in case J8LY was centered in the dorsal aspect of the ABmc, while in case J12FS, a smaller injection site was in the caudal ABmc. There were four injections in the CeN: one site was confined to the CeM (J9LY), two were confined to the CeLcn (J9FS, J1LY) and one injection included both the CeM and CeLcn (J1FR). All four injections included the CeLc but did not encroach on the amygdalostratial area. Two injections were located in the M (J19LY, J16FR). One injection site also encompassed the AHA and ABs (J15AA). A large injection site covered the majority of the L, with spread to surrounding white matter (J17 LY).

Distribution of anterogradely labeled fibers in the BSTL and ventral striatum

Bmc: Cases J12FR (Figure 8), J12LY, J16LY (not shown): The concentration of labeled fibers in the BSTL was highest following the two more medially placed injections, consistent with the distribution of labeled cells seen in retrograde cases. Case J12FR is described in detail. Labeled fibers were most densely distributed to the BSTLJ, and there was a light concentration of labeled fibers in the BSTLP (Figure 8A, B). There was a conspicuous absence of labeled fibers in the BSTLcn (or BSTLc), that stood out as a fiber-poor oval structure (Figure 8C). The absence of labeled fibers in the BSTLcn indicates that any retrogradely labeled cells seen in the Bmc following retrograde injections likely resulted from tracer uptake in the BSTLJ or BSTLP.

In the ventral striatum, there were high concentrations of labeled fibers in the ventral and lateral shell and small patches of labeled fibers in the core of the ventral striatum (Figure 8D) and ventromedial caudate nucleus (not shown). The dorsomedial shell was relatively devoid of labeled fibers.

Medial Bpc/CTA: Cases J20LY (Figure 9), J15FS, J14FS (not shown): The concentration of labeled fibers in the BSTL subdivisions was high for all injections involving the medial Bpc/CTA. However, the pattern of distribution among subregions depended on the rostrocaudal location of the injection. Case J20LY, the most rostral injection, is described first. Labeled fibers were found in all subdivisions of the BSTL, with highest concentrations in the BSTLcn (Figure 9A, B). There were many labeled fibers in the BSTLD, at the junction of the ventral striatum. At the level of the anterior commissure (Figure 9A), moderate concentrations of labeled fibers overlapped the BSTLcn, including the BSTLc, and BSTLP. There were a concentrated number of LY-positive fibers in the BSTLJ at this level. As the decussation of the anterior commissure reached its caudal extent, labeled fibers were most densely concentrated in the BSTLcn, with a moderate concentration of labeled fibers in the BSTLJ and a light concentration of labeled fibers in the BSTLP (Figure 9B, C). At this level, varicose fibers in the SLEAc were moderately concentrated. Passing fibers, seen following all anterograde injections, and depicted in the photomicrograph (Figure 9C, asterisks), were not charted. Interestingly, injections in the caudal medial Bpc/CTA (cases J15FS and J14FS) resulted in a more restricted pattern of labeled fibers in the BSTL. In these cases, there were many labeled fibers in the BSTLcn and BSTLc, moderate concentrations of labeled fibers in the BSTLP, but virtually no labeled fibers in the BSTLJ. This suggests that the caudal medial Bpc/CTA projection is relatively confined to the BSTLcn and BSTLP, while more rostral CTA has a broader input that includes the BSTLJ.

The pattern of labeled fibers in the ventral striatum also depended on the rostrocaudal location of the medial Bpc/CTA injections. Labeled fibers were confined to the dorsomedial shell following caudal medial Bpc/CTA injections (cases J15FS and J14FS), whereas anterogradely labeled fibers resulting from the more rostral medial Bpc/CTA (case J20LY) occupied the dorsomedial shell with additional patches of labeled fibers in ventral shell and core rostrally (Figure 9D), and ventromedial caudate nucleus and ventromedial putamen caudally (not shown).

Central Bpc: Case J15 LY (not shown): This injection site was centered in the central Bpc. In contrast to cases involving the medial Bpc/CTA, high concentrations of labeled fibers occupied only the ventrolateral edge of the BSTLcn and BSTLc. There were high concentrations of thin, varicose labeled fibers packed in the BSTLJ. The BSTLP and contiguous SLEAc also contained moderate concentrations of labeled fibers, similar to the more medial injections involving the CTA.

In the ventral striatum, there were many labeled fibers in the ventral shell, but no labeled fibers in the dorsomedial shell. The core of the ventral striatum had a light distribution of labeled fibers and there were no labeled fibers in the caudate nucleus.

Lateral Bpc: Case J14FR (not shown): Labeling in the BSTL was negligible, however, a few scattered fibers could be seen in the SLEAc. Labeled fibers in the ventral striatum were confined to the ventrolateral shell.

ABmc: Cases J8LY (Figure 10), J12FS, J20FS, J21FR (not shown): In cases involving the ABmc alone (J8LY and J12FS), there was a dense distribution of finely beaded fibers in the BSTLJ (Figure 10A, B). Posterior to the anterior commissure, moderate numbers of labeled fibers were also found in the BSTLP (Figure 10B, C). These labeled fibers were continuous with ones that extended through the basal forebrain in the SLEAc. The BSTLcn was relatively devoid of labeled fibers following these injection sites (Figure 10 B, C), and there was a very light distribution of labeled fibers in the BSTLc.

Cases J21FR and J20FS were relatively larger tear-shaped injections that involved the ABmc and extended into the lateral ABpc. In both cases, anterogradely labeled fibers were predominantly found in the BSTLJ, however, there was also moderate numbers of labeled fibers in the BSTLcn and BSTLc. The BSTLP contained many labeled fibers only in case J20FS. All injections that included the ABmc resulted in labeled fibers in the ventral shell and core of the striatum (Figure 10D), and the ventromedial caudate (not shown),

ABpc: Case J18FR (not shown): Case J18FR was centered in the medial aspect of the ABpc, and had no involvement of the ABmc. There were many FR-positive fibers overlying the BSTLcn and the BDTLc. Unlike cases that included the ABmc in the site, there were few to no labeled fibers in the BSTLJ. The BSTLP contained moderate concentrations of labeled fibers. The ventral striatum contained scattered labeled fibers in the shell.

CeM: Cases J9LY (Figure 11A-D), and J1FR (not shown): Cases J9LY and J1FR had a similar distribution of labeled fibers. The fact that case J1FR, which involved both CeM and CeLcn, was similar to J9LY, a small injection confined to the CeM, suggested a relatively restricted input from the CeLcn. Case J9LY, in which the injection site was confined to the CeM, is described. In the ventral striatum (Figure 11A), light patches of labeled fibers were seen in the dorsomedial and ventral shell. There were no labeled fibers in the core or caudate nucleus. Behind the caudal shell, in the rostral BSTL, there were many labeled fibers in the BSTLcn. At caudal levels, there were many densely packed, very fine varicose-labeled fibers in the BSTLcn, BSTLJ, and BSTLP. The BSTLc also contained fine, labeled fibers. Labeled fibers in the BSTLP were continuous with high concentrations of thin, beaded fibers sweeping under the globus pallidus, in the SLEAc (Figure 11D).

CeLcn: Cases J9 FS, J1LY (Figure 11E-H): Compared to cases with injections encompassing the CeM (above), labeled fibers in these cases were relatively confined to the BSTLcn, with only a few labeled fibers in the BSTLc, and in both cases, the labeled fibers were of relatively lower density. There were relatively few labeled fibers in the BSTLJ or BSTLP, although there were moderate concentrations of labeled fibers in the SLEAc. There were no labeled fibers in the ventral striatum.

AHA: Cases J15 AA (not shown): Tritiated amino acids were used for this large injection, which encompassed the AHA and ABs. Dense concentrations of silver grains were found in both BSTM and BSTL. While the BSTM contained the highest concentration of labeled fibers, the BSTLP also contained many labeled fibers, with a lighter concentration of AA-labeled fibers in the BSTLJ. While the main BSTLcn was conspicuously devoid of labeled

fibers, a small accessory island beneath the BSTLcn was densely overlaid with labeled fibers. In the basal forebrain, labeled fibers extended in two, dorsal and ventral streams, corresponding to the location of the SLEAc and medial sublenticular extended amygdala (SLEAm), respectively.

In contrast to the broad distribution of labeled fibers in the BSTL and BSTM, the distribution of labeled fibers in the striatum was confined to the tip of dorsomedial shell.

M: Cases J16FR, J19LY (not shown): While the majority of labeled fibers following these injections were in the BSTM, the lateral septum and the SLEAm, there were a few labeled fibers in the BSTLcn and BSTLP and SLEAc. There were no fibers in the BSTLJ. In the ventral striatum, there was a light concentration of labeled fibers in the dorsomedial shell and occasional labeled fibers in the ventral shell.

L: Case J17 LY (not shown): There were no labeled fibers in the BSTL and only scattered labeled fibers the striatum.

DISCUSSION

This is the first study in the monkey detailing amygdala inputs to BSTL subdivisions using both retrograde and anterograde tracing methods. We also compared how amygdala inputs to the BSTLcn and BSTLJ, related to projections to the ventral striatum, a region considered to be a transition zone with the BSTL (Alheid et al., 1995; Alheid and Heimer, 1988; Heimer et al., 1997). We first found that the main afferent inputs to the BSTL as a whole originate from the medial Bpc/CTA, CeN, basal nucleus and accessory basal nucleus, consistent with previous findings in rats and primates (Aggleton et al., 1987; De Olmos, 1990; Dong et al., 2001a; Krettek and Price, 1978; Price and Amaral, 1981; Price et al., 1987; Russchen et al., 1985b; Weller and Smith, 1982). These findings extend previous studies in primates by showing that specific amygdala inputs vary according to relative placement of retrograde injections in the BSTLcn or BSTLJ. Anterograde injections in specific amygdala nuclei confirmed that there are distinct subcircuits that project to the BSTLcn and BSTLJ.

The BSTLcn receives its strongest projections from the medial Bpc/CTA and CeN, while the medial Bpc/CTA and CeN provide only light inputs to the BSTLJ. The BSTLcn receives input from the CeM, CeLcn and CeLc. The projection from the CeLcn is relatively specific, projecting only to the BSTLcn, and avoiding other BSTL subdivisions. Finally, the ABpc also has a relatively specific input to the BSTLcn. In contrast, projections to the BSTLJ preferentially originate from the ABmc and basal nucleus outside the CTA region (Figure 12). Anterograde injections confirmed that the ABmc mainly innervates the BSTLJ, avoiding the BSTLcn. This indicates that retrogradely labeled cells in the ABmc following injections centered in the BSTLcn resulted from some spread into the BSTLJ or BSTLP.

Comparison of medial Bpc injections involving the CTA (J20LY, J14FS, J15FS) indicates that this region consistently, and strongly projects to the BSTLcn. In contrast to the caudal medial Bpc/CTA (cases J14FS and J15FS), the rostral medial Bpc/CTA (case J20LY) projects additionally to the BSTLJ. Furthermore, as the injection site moves laterally into the Bpc proper (case J15LY), the BSTLJ remains strongly innervated, while input to the BSTLcn diminishes. The rostral medial Bpc/CTA thus appears to be a transition region between the caudal medial Bpc/CTA and Bpc proper.

The BSTLJ is also distinguished from the BSTLcn in receiving no input from the CeLcn. The CeLcn thus has a relatively focused input to the BSTLcn, a finding noted in the rat

(Petrovich and Swanson, 1997). This projection links the BSTLDCn with its ‘mirror’ subdivision at the caudal pole of the extended amygdala, the CeLCn.

The BSTLc is a relatively thin structure in the primate, and therefore could not be isolated in retrograde studies. For the most part, anterograde injections indicate that the BSTLc contained labeled fibers whenever there were inputs to the BSTLCn, with the medial Bpc/CTA and CeM providing the strongest projections. An injection into the ABpc also resulted in labeled fibers in the BSTLc, as well as BSTLCn.

Based on anterograde results, the BSTLP receives the broadest inputs of all BSTL subdivisions, with afferent projections from the CeM, AHA/ABs subnuclei, ABmc, ABpc, central Bpc and medial/CTA regions. In addition, anterogradely labeled fibers in the BSTLP were consistently seen in conjunction with patches of labeled fibers in the contiguous SLEAc, which share a similar histochemical profile.

Our current results, along with our previous study examining amygdala inputs to the CeN and its transition zone, provide further support for the organization of the central extended amygdala into ‘symmetrical’ BSTL and CeN subdivisions (present study, Fudge and Tucker, 2009). In addition to matching histochemical profiles (Heimer et al., Martin 1991, present findings), the CeLCn and BSTLCn each receive their main inputs from the medial Bpc and CTA regions. Furthermore, the BSTLCn, but not the BSTLJ, receives CeLCn input, suggesting a unique relationship between these two ‘symmetrical’ structures. One possible exception to the overall ‘symmetry’ of amygdala inputs to BSTL and CeN subdivisions may be the relatively stronger input from the ABpc to the BSTLCn compared to the CeLCn since our anterograde injection in the ABpc resulted in relatively more labeled fibers in the BSTLCn than the CeLCn (Fudge and Tucker, 2009). However, more injections are needed to confirm this relationship. The broad amygdala inputs to the BSTLP roughly parallel those to the CeM, its counterpart in the caudal pole of the central extended amygdala (Heimer et al., 1999). The BSTLJ, which we have suggested in the present study to serve as a transition zone with the ventral shell, resemble amygdala projections to transition zones between the CeN and the shell, including the lateral amygdalostratial area and the lateral interstitial nucleus of the posterior limb of the anterior commissure (Fudge and Tucker, 2009).

Comparison with previous studies of amygdala-BSTL connectivity in rodents

Our results are generally consistent with rodent findings (Dong et al., 2001a; Krettek and Price, 1978; Price et al., 1987; Shammah-Lagnado et al., 1996; Sun et al., 1991; Weller and Smith, 1982). In Krettek and Price’s classic rodent study focusing on amygdala inputs to the basal forebrain, anterograde injections in the CTA (known as the ‘amygdalopiriform transition area’ or ‘postpiriform transition area’ in rats), the Bpc (referred to as the ‘posterior basolateral amygdala’), the accessory basal nucleus (referred to as the ‘basomedial nucleus’), and the CeN resulted in strong anterograde labeling in the BSTL region (Krettek and Price, 1978). These general results are consistent with our data in monkeys.

Many retrograde and anterograde studies later extended Krettek and Price’s findings in rats, showing subdivision-specific projections (Dong et al., 2001a; Jolkkonen et al., 2001; McDonald, 1991; McDonald, 1998; McDonald et al., 1999; Petrovich et al., 1996; Petrovich and Swanson, 1997; Shammah-Lagnado et al., 1996; Shammah-Lagnado and Santiago, 1999; Sun et al., 1991; Weller and Smith, 1982). In a comprehensive review based on a series of anterograde studies of amygdala outputs, Swanson and colleagues concluded that there were differential inputs to the rodent BSTLJ and the BSTLCn (referred to as the ‘oval nucleus’)(Dong et al., 2001a). Specifically, they found that the Bpc (referred to as the ‘posterior basolateral amygdala’, BLAp) provides the main source of inputs to the BSTLJ, while the CeLCn, CeM and CTA (‘postpiriform transition region’) provide the main source

of inputs to the ‘oval nucleus’. They also noted that the region of the AHA (‘posterior nucleus’) and PAC (‘cortical nucleus, posterior division’) largely avoid the BSTLcn (‘oval nucleus’) but target aspects of the Krettek and Price’s BSTM (‘principal nucleus’) and dorsomedial shell (Canteras et al., 1992); findings that are broadly consistent with our data in the monkey.

With respect to the CTA-like region in rodents, works by Dong and Shammah-Lagnado report that the CTA (the ‘rostromedial postpiriform transition area’ and ‘medial amygdalopiriform area’, respectively) sends strong inputs to both the BSTLJ and BSTLcn (Dong et al., 2001a; Shammah-Lagnado and Santiago, 1999). We also found strong projections to the BSTLcn and BSTLJ following injections involving the rostromedial CTA (case J20LY). However, more caudal injection sites that included the CTA (cases J14FS and J15FS) exclusively innervated the BSTLcn.

We found that the ABmc provides specific inputs to the monkey BSTLJ, while in rats, the ‘posterior basomedial nucleus’ (sometimes assumed to be a homologue of the AB) does not project there (Dong et al., 2001a; Petrovich et al., 1996). In the rat, both anterior and posterior basomedial amygdala project strongly to the BSTL (Petrovich et al., 1996), although it is unclear the extent to which either has a clear homologue in primates (Canteras and Swanson, 1992; Crosby and Humphrey, 1941; Fudge and Tucker, 2009; Petrovich et al., 1996). Therefore, differences between our results and those from rodent studies may reflect species differences as previously noted for the amygdalostratial (Fudge et al., 2004; Fudge et al., 2002) and amygdala-CeN paths (Fudge and Tucker, 2009).

The primate CeN also has several discontinuities with the rat in its cytoarchitectural and connectional features (Fudge and Tucker, 2009; Price and Amaral, 1981; Price et al., 1987). Thus, while our data is generally consistent with the rodent data of Sun et al. (Sun et al., 1991) and Petrovich (Petrovich and Swanson, 1997)-- namely, that the CeLcn (analogous to the rodent ‘CeL’) and CeM both provide input to the BSTLcn (‘anterolateral bed nucleus subdivision’ and ‘oval’ nucleus, respectively)-- the relatively large ‘intermediate’ subdivision of the rat CeN, which also projects to the BSTL, has not been demonstrated in the primate.

Differences in details of the amygdala-BSTL path between rats and monkeys may also result from a lack of strict homology of BSTL subdivisions. For example, the ‘oval’ subnucleus of the BSTL (Ju et al., 1989), also termed the ‘dorsal, lateral subdivision’ (Alheid et al., 1995), has been assumed to be homologous to the primate BSTLcn. While both structures contain high concentrations of NT- and CCK-positive cells and fibers, the ‘oval’ subdivision of the rat also has high SP-IR, a feature that is absent in the primate (Heimer et al., 1999; Ju et al., 1989; Martin et al., 1991)(present results). In fact, the primate BSTLcn is actually identified by low SP immunoreactivity, underscoring important caveats in making cross-species comparisons.

Amygdala inputs across the BSTLcn/BSTLJ and ventral striatum subdivisions

In primates, amygdala inputs to the ventral striatum are relatively better-characterized than those to the BSTL (Friedman et al., 2002; Fudge et al., 2002; Russchen et al., 1985b). Due to the close anatomic and connectional relationships shared by the ventral striatum and the BSTL in rats (e.g. (Zahm, 1998)), we examined the organization of amygdala inputs to each of these structures. The ABmc and basal nucleus are the main source of inputs to the ventral striatum as a whole, including the ventromedial caudate nucleus, consistent with previous studies (Friedman et al., 2002; Fudge et al., 2002; Russchen et al., 1985b) (Table 4). In this study, we replicated and extended our previous results, showing that the ‘transitional’ nature of the shell between the BSTL and striatum can be appreciated in inputs from specific

amygdaloid nuclei. For example, the shell is distinguished from the core and ventromedial caudate nucleus by light to moderate inputs from the CeM, which also target the BSTL. The dorsomedial shell is further distinguished by additional inputs from the medial Bpc/CTA, which are also an important afferent source for the BSTLcn. The dorsomedial shell is also unique in receiving inputs that target the medial extended amygdala, including those from the AHA, a finding also noted in the rodent (Brog et al., 1993; Canteras et al., 1992). This indicates that the dorsomedial shell is a site where afferents common to the BSTL, BSTM and striatum converge.

There is a shift in the topography of amygdala inputs as the projection to the striatum moves ventrally from the ventromedial caudate nucleus to the core of the ventral striatum, and then to the ventral and dorsomedial shell. In addition to 'striatal'-like inputs from the basal nucleus and ABmc that are typical of the ventromedial caudate nucleus and ventral striatal core, the shell receives additional projections that innervate the BSTL, including additional inputs from the CeN, the PAC, and the medial Bpc/CTA. Within the shell, there is a medial to lateral topography paralleling that in the BSTLcn and BSTLJ, respectively. While the BSTLcn and dorsomedial shell receive strong inputs from the medial Bpc/CTA and relatively fewer inputs from the ABmc, Bpc, and Bi, the BSTLJ and ventral shell/core primarily receive inputs from these latter subnuclei (Figure 12, Table 4). Interestingly, when both the BSTLcn and BSTLJ were strongly labeled, for example, following an anterograde injection in the rostrocentral CTA (case 20LY, Figure 9), both the dorsomedial and ventral shell/core contained patches of anterogradely labeled fibers. As noted, the dorsomedial shell also receives some inputs that innervate the BSTM, such as the AHA and M. In sum, as amygdala inputs move ventrally in the striatum and into the BSTL, the projection shifts to involve increasingly transitional regions of the amygdala such as the CTA, ABs, and AHA as well as the CeN.

The connectional similarities observed across BSTL and ventral striatal subdivisions are also seen in the intrinsic histochemical organization of both structures. Specifically, the BSTLJ/ventral shell have relatively high levels of the striatal-like marker, AChE, compared to the BSTLcn/dorsomedial shell. Conversely, high levels of the classic extended amygdala peptides ST and NT are expressed in the dorsomedial shell and BSTLcn, but not the BSTLJ/ventral shell. The connectional and histochemical gradients noted in this study are supported by the original work of Heimer and colleagues that proposed that the dorsomedial shell, particularly its 'cone' region, is more closely associated with the extended amygdala (Alheid and Heimer, 1988). 'Striatal' properties of the BSTLJ observed in this study are also consistent with earlier reports of the BSTLJ in rodents. For example, McDonald commented on the 'striatal-like' nature of the spiny neurons confined to the BSTLJ in Golgi impregnation specimens (McDonald, 1983), a finding echoed by more recent studies (Heimer et al., 1999; Larriva-Sahd, 2004; Martin et al., 1991).

Technical considerations

An advantage of retrograde studies is the ability to provide a sensitive overview of afferent structures that can influence specific brain regions. Within subjects, a single injection also enables analysis of the relative number of labeled cells projecting to a region. Key concerns, however, are the encroachment of tracer on nearby structures, and the possibility of tracer uptake and transport by fibers of passage, leading to false-positive results (Halperin and LaVail, 1975; Nance and Burns, 1990). Both issues are particularly relevant for injections in structures such as the BSTL, which is small and lies in, or near, passing fiber tracts. Because injections which cause necrosis or tissue damage at the site are most prone to fibers of passage problems, we used slow, pressure injections of tracer (Nance and Burns, 1990; Schmued et al., 1990; Vercelli et al., 2000). We also eliminated cases where there was obvious tissue damage and nonspecific labeling, and then remaining cases were examined

for evidence of uptake by fibers of passage. For example, fibers leaving the amygdala and ‘passing through’ the BSTL go on to innervate the cortex and mediodorsal thalamus (Aggleton and Mishkin, 1984; Miyashita et al., 2007; Porrino et al., 1981; Russchen et al., 1987). Retrograde labeling in the amygdala resulting from uptake by these fibers of passage would therefore be expected to also label these brain regions. While we noted retrogradely labeled cells in the cortex, their distribution was confined to areas previously noted to innervate the BSTL (Freedman et al., 2000; Ghashghaei and Barbas, 2001). There were no retrogradely labeled cells in broader cortical regions connected with the amygdala proper, or the mediodorsal thalamus. Finally, to check for fiber of passage issues, we examined cases where the injection target was “missed” and an injection placed in nearby fiber tracts, such as the anterior commissure or internal capsule. These injection sites resulted in very few retrogradely labeled cells in any brain region, suggesting that the fiber of passage problem was negligible.

Retrograde studies of complex structures require anterograde studies to confirm the specificity of nuclei-specific afferent paths. Indeed, our anterograde studies were critical for clarifying projections to the BSTLcn versus BSTLJ. Anterograde studies are also important since the density of retrogradely labeled cells does not necessarily correspond to the density of the terminal field. Because anterograde tracers are also theoretically subject to a “fibers of passage” problem, as well as false labeling of axon collaterals, careful analysis in conjunction with retrograde studies is required. Since injections within the lateral nucleus, basal nucleus subdivisions, and accessory basal nucleus resulted in different patterns of labeling not only in the BSTL subdivisions, but also in various predicted terminal fields, false positive labeling by passing fibers or axon collateral was thought to be minimal.

Functional considerations

In this study, we show that the amygdala sends differential projections to the BSTLJ and BSTLcn, suggesting subtle differences in the microcircuitry carrying information to each region. The density and specificity of inputs from the medial Bpc/CTA to the BSTLcn and dorsomedial shell is particularly striking. The CTA is a poorly understood part of the amygdala, and due to its relatively undifferentiated nature, has received multiple names including, the ‘amygdalohippocampal transition area’, ‘postpiriform transition area’, ‘amygdalopiriform transition area’, and amygdaloentorhinal field (Insausti et al., 1997; Jolkkonen et al., 2001; Santiago and Shammah-Lagnado, 2005; Shammah-Lagnado and Santiago, 1999). We have previously emphasized the primate CTA’s continuity with the medial PL, which flows into the CTA dispersing into streams and islands of cells (deCampo and Fudge, 2012). We previously found that the CTA and contiguous PL is the only amygdala subregion that projects to the CeLcn (Fudge and Tucker, 2009). Thus, the CTA is a common and specific input to both lateral core subdivisions at each end of the central extended amygdala continuum (BSTLcn and CeLcn), adding a common connective feature to their histochemical and cellular similarities (Heimer et al., 1999).

In monkeys, the CTA is unique in serving as a crossroads between the so-called ‘olfactory’ (corticomedial) amygdala and ‘sensory’ (basolateral) amygdala. While the corticomedial amygdala is best known for receiving olfactory information from the piriform cortex and olfactory bulb (Krettek and Price, 1977), in primates, this projection tapers off in the region of the CTA (Price’s ‘PACs’) (Carmichael et al., 1994). The CTA region in monkeys is also characterized by abundant ‘sensory association’ inputs from the lateral nucleus of the amygdala (Pitkanen and Amaral, 1998), which processes higher level visual, auditory, and tactile information on which primates are relatively dependent compared to macrosmatic species (Turner et al., 1980; Webster et al., 1993). The insula, which mediates sensations derived from inside the body such as taste, olfaction, and bodily sensations (Carmichael et al., 1994; Craig, 2002; Critchley et al., 2004), also has a strong input to the CTA region

(Carmichael and Price, 1996; Hoistad and Barbas, 2008; Mufson et al., 1981)(unpublished results). Insula projections suggest that internal sensory states, rather than simple olfactory inputs, are important drivers of the CTA responses in monkeys. Finally, the medial Bpc/CTA receives some of the strongest hippocampal inputs in the amygdala (Fudge et al., 2012; Rosene and Van Hoesen, 1977; Saunders et al., 1988). These inputs can be thought of as conveying temporal and spatial coordinates of past and present experiences (Naya and Suzuki, 2011; Paz et al., 2010; Squire, 1993), enabling interpretation of the environment. Thus, with converging inputs from exteroceptive and interoceptive sensory centers and the hippocampus, the CTA is positioned to convey information on potentially dangerous environments based on past experience to modulate the BSTL. Recognizing complex environments, and their salience, may be advantageous, enabling the organism to enter a state of defensive preparedness manifested by behavior changes (hypervigilance) and autonomic arousal levels. These functions are associated with BST activation in response to danger (Alvarez et al., 2011; Somerville et al., 2010; Walker and Davis, 1997).

In humans, Alvarez and colleagues demonstrated heightened BST activity in normal subjects upon exposure to a dangerous virtual environment (a scene associated with unpredicted shock). BST activity was associated with heightened subjective fear ratings and autonomic arousal, as well as activation in the ventral amygdala and hippocampus (Alvarez et al., 2011). This finding suggests that contextual knowledge from the hippocampus and amygdala are important for protective behaviors associated with BST activity, including threat monitoring and elevated arousal levels (Mobbs et al., 2010; Somerville et al., 2010). Although functional imaging techniques are still unable to resolve specific BSTL subdivisions, our connectional data suggest that the BSTLcn, with its prominent afferents from the medial Bpc/CTA and CeN, is the main subdivision mediating these responses.

In contrast to the BSTLcn, the BSTLJ, like the ventral striatum, receives its main amygdala inputs from the ABmc and basal nucleus. In rodents, the BSTLJ has efferents typical of the of the ventral striatum, avoiding the paraventricular nucleus, parabrachial nucleus and the nucleus of the solitary tract; regions typically associated with the extended amygdala outputs (Dong et al., 2000; Moga et al., 1989; Zahm et al., 1999). Pharmacologic studies implicate both the BSTLJ and ventral striatum in the negative affective responses of drug and alcohol withdrawal (review, (Koob and Volkow, 2010))(Francesconi et al., 2009; Gracy et al., 2001; Szucs et al., 2012; Veinante et al., 2003; Walker et al., 2000), supporting functional similarities between the two structures.

Conclusions

The BSTLcn and BSTLJ are prominent subdivisions of the BSTL that are histochemically distinct, and sit directly behind the shell of the ventral striatum. The BSTL has been relatively neglected in primate tract-tracing work despite its importance in threat monitoring and anxiety-like responses across species. We found that at a microscopic level, the BSTLcn and BSTLJ receive distinct afferent inputs from specific amygdala subnuclei, with the medial Bpc/CTA and CeN targeting the BSTLcn, and the ABmc and basal nucleus forming most of the afferent path to the BSTLJ. The rostrocentral medial Bpc/CTA and CeM send projections to both subdivisions. The ventral striatum receives many of these same inputs, which are topographically organized such that the dorsomedial shell shares inputs with the BSTLcn and the ventral shell/core shares inputs with the BSTLJ. This arrangement not only supports the idea of the shell as a transition with extended amygdala structures, but also describes distinctions between the dorsomedial and ventral shell/core subregions and their relationship to the BSTL.

Acknowledgments

We are grateful for the technical assistance of Nanette Alcock and Daniel Tylee in performing histochemical work. We also thank Nancy Bui for her help in digitizing hand-drawn charts.

NIMH: RO1 MH63291-09 (JLF), F30MH096502 (DMD)

ABBREVIATIONS

AAA	anterior amygdaloid area
ABmc	accessory basal nucleus, magnocellular subdivision
ABpc	accessory basal nucleus, parvicellular subdivision
ABs	accessory basal nucleus, superficial subdivision
ac	anterior commissure
AChE	acetylcholinesterase
AHA	amygdalohippocampal area
Astr	amygdalostratial area
Bi	basal nucleus, intermediate subdivision
Bmc	basal nucleus, magnocellular subdivision
Bpc	basal nucleus, parvicellular subdivision
BST	bed nucleus stria terminalis
BSTL	lateral bed nucleus stria terminalis
BSTLD	lateral bed nucleus stria terminalis, dorsal subdivision
BSTLc	lateral bed nucleus stria terminalis, capsular subdivision
BSTLcn	lateral bed nucleus stria terminalis, central subdivision
BSTLJ	lateral bed nucleus stria terminalis, juxtacapsular subdivision
BSTM	medial bed nucleus stria terminalis
BSTLP	lateral bed nucleus stria terminalis, posterior subdivision
C	caudate nucleus
CaBP	calbindin
CeLc	central nucleus, lateral capsular division
CeLcn	central nucleus, lateral core subdivision
CeM	central nucleus, medial subdivision
CeN	central nucleus
CoA	anterior cortical nucleus
CoP	posterior cortical nucleus
CTA	corticoamygdaloid transition area
DM shell	dorsomedial shell
ec	external capsule
ENK	enkephalin

FR	fluororuby
FS	fluorescein
GP	globus pallidus
GPe	external globus pallidus
GPI	internal globus pallidus
H	hippocampus
ic	internal capsule
L	lateral nucleus
LV	lateral ventricle
LY	Lucifer yellow
M	medial nucleus
NLOT	nucleus of the lateral olfactory tract
NT	neurotensin
P	putamen
PAC	periamygdaloid cortex, subdivisions I-III
PACs	periamygdaloid cortex, sulcal division
PC	piriform cortex
PL	paralamina nucleus
S	shell
SLEAc	central subdivision of the sublenticular extended amygdala
SLEAm	medial subdivision of the sublenticular extended amygdala
ST	somatostatin
VP	ventral pallidum
V. shell	ventral shell
WGA	wheat-germ agglutinin-horse radish peroxidase

Literature cited

- Aggleton JP, Friedman DP, Mishkin M. A comparison between the connections of the amygdala and hippocampus with the basal forebrain in the macaque. *Exp Brain Res.* 1987; 67(3):556–568. [PubMed: 2820778]
- Aggleton JP, Mishkin M. Projections of the amygdala to the thalamus in the cynomolgus monkey. *J Comp Neurol.* 1984; 222:56–68. [PubMed: 6321564]
- Alheid, GF.; DeOlmos, CA.; Beltramino, CA. Amygdala and extended amygdala. In: Paxinos, G., editor. *The Rat Nervous System*. San Diego: Academic Press; 1995. p. 495-578.
- Alheid GF, Heimer L. New perspectives in basal forebrain organization of special relevance for neuropsychiatric disorders: the striatopallidal, amygdaloid, and corticopetal components of substantia innominata. *Neuroscience.* 1988; 27:1–39. [PubMed: 3059226]
- Alvarez RP, Chen G, Bodurka J, Kaplan R, Grillon C. Phasic and sustained fear in humans elicits distinct patterns of brain activity. *Neuroimage.* 2011; 55(1):389–400. [PubMed: 21111828]

- Amaral DG, Bassett JL. Cholinergic innervation of the monkey amygdala: An immunohistochemical analysis with antisera to choline acetyltransferase. *J Comp Neurol.* 1989; 281:337–361. [PubMed: 2703552]
- Amaral, DG.; Price, JL.; Pitkanen, A.; Carmichael, ST. Memory, and Mental Dysfunction. Wiley-Liss, Inc; 1992. Anatomical organization of the primate amygdaloid complex. The Amygdala: Neurobiological Aspects of Emotion; p. 1-66.
- Brog JS, Salyapongse A, Deutch AY, Zahm DS. The patterns of afferent innervation of the core and shell in the “accumbens” part of the rat ventral striatum: Immunohistochemical detection of retrogradely transported fluoro-gold. *J Comp Neurol.* 1993; 338:255–278. [PubMed: 8308171]
- Canteras NS, Simerly RB, Swanson LW. Connections of the posterior nucleus of the amygdala [published erratum appears in *J Comp Neurol* 1993 Feb 22;328(4):604]. *J Comp Neurol.* 1992; 324(2):143–179. [PubMed: 1430327]
- Canteras NS, Swanson LW. Projections of the ventral subiculum to the amygdala, septum, and hypothalamus: a PHAL anterograde tract-tracing study in the rat. *J Comp Neurol.* 1992; 324(2): 180–194. [PubMed: 1430328]
- Carmichael ST, Clugnet MC, Price JL. Central olfactory connections in the macaque monkey. *J Comp Neurol.* 1994; 346:403–434. [PubMed: 7527806]
- Carmichael ST, Price JL. Limbic connections of the orbital and medial prefrontal cortex in macaque monkeys. *J Comp Neurol.* 1996; 363:615–641. [PubMed: 8847421]
- Chang HT, Kuo H. Relationship of calbindin D-28k and cholinergic neurons in the nucleus basalis of Meynert of the monkey and the rat. *Brain Res.* 1991; 549:141–145. [PubMed: 1893247]
- Cote P-Y, Sadikot AF, Parent A. Complementary distribution of calbindin D-28k and parvalbumin in the basal forebrain and midbrain of the squirrel monkey. *European J Neurosci.* 1991; 3:1316–1329. [PubMed: 12106229]
- Craig AD. How do you feel? Interoception: the sense of the physiological condition of the body. *Nature Reviews Neuroscience.* 2002; 3(8):655–666.
- Critchley HD, Wiens S, Rotshtein P, Ohman A, Dolan RJ. Neural systems supporting interoceptive awareness. *Nat Neurosci.* 2004; 7(2):189–195. [PubMed: 14730305]
- Crosby EC, Humphrey T. Studies of the vertebrate telencephalon: the nuclear pattern of the anterior olfactory nucleus, tuberculum olfactorium and the amygdaloid complex in adult man. *J Comp Neurol.* 1941; 74:309–347.
- De Olmos, J. The Amygdala. In: Paxinos, G.; Mai, JK., editors. *The Human Nervous System.* 2. San Diego: Elsevier Academic Press; 2004.
- De Olmos, JS. The Amygdala. In: Paxinos, G., editor. *The Human Nervous System.* San Diego: Academic Press; 1990. p. 583-710.
- deCampo DM, Fudge JL. Where and what is the paralaminar nucleus? A review on a unique and frequently overlooked area of the primate amygdala. *Neurosci Biobehav Rev.* 2012; 36:520–535. [PubMed: 21906624]
- Dong H, Petrovich GD, Swanson LW. Organization of projections from the juxtacapsular nucleus of the BST: a PHAL study in the rat. *Brain Res.* 2000; 859(1):1–14. [PubMed: 10720609]
- Dong HW, Petrovich GD, Swanson LW. Topography of projections from amygdala to bed nuclei of the stria terminalis. *Brain Res - Brain Res Reviews.* 2001a; 38(1–2):192–246.
- Dong HW, Petrovich GD, Watts AG, Swanson LW. Basic organization of projections from the oval and fusiform nuclei of the bed nuclei of the stria terminalis in adult rat brain. *J Comp Neurol.* 2001b; 436(4):430–455. [PubMed: 11447588]
- Duvarci S, Bauer EP, Pare D. The bed nucleus of the stria terminalis mediates inter-individual variations in anxiety and fear. *J Neurosci.* 2009; 29(33):10357–10361. [PubMed: 19692610]
- Fendt M, Endres T, Apfelbach R. Temporary inactivation of the bed nucleus of the stria terminalis but not of the amygdala blocks freezing induced by trimethylthiazoline, a component of fox feces. *J Neurosci.* 2003; 23(1):23–28. [PubMed: 12514197]
- Fox AS, Shelton SE, Oakes TR, Davidson RJ, Kalin NH. Trait-like brain activity during adolescence predicts anxious temperament in primates. *PLoS One.* 2008; 3(7):e2570. [PubMed: 18596957]
- Francesconi W, Berton F, Repunte-Canonigo V, Hagihara K, Thurbon D, Lekic D, Specio SE, Greenwell TN, Chen SA, Rice KC, Richardson HN, O'Dell LE, Zorrilla EP, Morales M, Koob

- GF, Sanna PP. Protracted withdrawal from alcohol and drugs of abuse impairs long-term potentiation of intrinsic excitability in the juxtacapsular bed nucleus of the stria terminalis. *J Neurosci.* 2009; 29(17):5389–5401. [PubMed: 19403807]
- Freedman LJ, Insel TR, Smith Y. Subcortical projections of area 25 (subgenual cortex) of the macaque monkey. *J Comp Neurol.* 2000; 421(2):172–188. [PubMed: 10813780]
- Friedman DP, Aggleton JP, Saunders RC. Comparison of hippocampal, amygdala, and perirhinal projections to the nucleus accumbens: combined anterograde and retrograde tracing study in the Macaque brain. *J Comp Neurol.* 2002; 450(4):345–365. [PubMed: 12209848]
- Fudge JL, Breitbart MA, McClain C. Amygdaloid inputs define a caudal component of the ventral striatum in primates. *J Comp Neurol.* 2004; 476(4):330–347. [PubMed: 15282709]
- Fudge JL, deCampo DM, Becoats KT. Revisiting the hippocampal-amygdala pathway in primates: association with immature-appearing neurons. *Neuroscience.* 2012; 212:104–119. [PubMed: 22521814]
- Fudge JL, Haber SN. Bed nucleus of the stria terminalis and extended amygdala inputs to dopamine subpopulations in primates. *Neuroscience.* 2001; 104(3):807–827. [PubMed: 11440812]
- Fudge JL, Kunishio K, Walsh P, Richard C, Haber SN. Amygdaloid projections to ventromedial striatal subterritories in the primate. *Neuroscience.* 2002; 110(2):257–275. [PubMed: 11958868]
- Fudge JL, Tucker T. Amygdala projections to central amygdaloid nucleus subdivisions and transition zones in the primate. *Neuroscience.* 2009; 159(2):819–841. [PubMed: 19272304]
- Geneser-Jensen FA, Blackstad TW. Distribution of acetyl cholinesterase in the hippocampal region of the guinea pig. *Z Zellforsch.* 1971; 114:460–481. [PubMed: 5550728]
- Ghashghaei HT, Barbas H. Neural interaction between the basal forebrain and functionally distinct prefrontal cortices in the rhesus monkey. *Neuroscience.* 2001; 103(3):593–614. [PubMed: 11274781]
- Gracy KN, Dankiewicz LA, Koob GF. Opiate withdrawal-induced fos immunoreactivity in the rat extended amygdala parallels the development of conditioned place aversion. *Neuropsychopharmacology.* 2001; 24(2):152–160. [PubMed: 11120397]
- Grove EA. Neural associations of the substantia innominata in the rat: afferent connections. *J Comp Neurol.* 1988; 277:315–346. [PubMed: 2461972]
- Haber SN, Fudge JL, McFarland N. Striatonigrostriatal pathways in primates form an ascending spiral from the shell to the dorsolateral striatum. *J Neurosci.* 2000; 20(6):2369–2382. [PubMed: 10704511]
- Halperin JJ, LaVail JH. A study of the dynamics of retrograde transport and accumulation of horseradish peroxidase in injured neurons. *Brain Res.* 1975; 100(2):253–269. [PubMed: 53088]
- Heimer L, Alheid GF, de Olmos JS, Groenewegen HJ, Haber SN, Harlan RE, Zahm DS. The Accumbens: Beyond the core-shell dichotomy. *J Neuropsychiatry Clin Neurosci.* 1997; 9 (3):354–381. [PubMed: 9276840]
- Heimer, L.; De Olmos, JS.; Alheid, GF.; Person, J.; Sakamoto, N.; Shinoda, K.; Marksteiner, J.; Switzer, RC. The human basal forebrain. Part II. In: Bloom, FE.; Bjorkland, A.; Hokfelt, T., editors. *Handbook of Chemical Neuroanatomy.* Amsterdam: Elsevier; 1999. p. 57–226.
- Hoistad M, Barbas H. Sequence of information processing for emotions through pathways linking temporal and insular cortices with the amygdala. *Neuroimage.* 2008; 40(3):1016–1033. [PubMed: 18261932]
- Ikemoto S, Qin M, Liu ZH. The functional divide for primary reinforcement of D-amphetamine lies between the medial and lateral ventral striatum: is the division of the accumbens core, shell, and olfactory tubercle valid? *J Neurosci.* 2005; 25(20):5061–5065. [PubMed: 15901788]
- Insausti R, Herrero MT, Witter MP. Entorhinal cortex of the rat: cytoarchitectonic subdivisions and the origin and distribution of cortical efferents. *Hippocampus.* 1997; 7(2):146–183. [PubMed: 9136047]
- Jolkkonen E, Miettinen R, Pitkanen A. Projections from the amygdalo-piriform transition area to the amygdaloid complex: a PHA-I study in rat. *J Comp Neurol.* 2001; 432(4):440–465. [PubMed: 11268008]

- Ju G, Swanson LW, Simerly RB. Studies on the cellular architecture of the bed nuclei of the stria terminalis in the rat: II. Chemoarchitecture. *J Comp Neurol.* 1989; 280(4):603–621. [PubMed: 2468695]
- Kelley AE. Functional specificity of ventral striatal compartments in appetitive behaviors. *Ann N Y Acad Sci.* 1999; 877:71–90. [PubMed: 10415644]
- Koob GF, Volkow ND. Neurocircuitry of addiction. *Neuropsychopharm.* 2010; 35(1):217–238.
- Krettek JE, Price JL. Projections from the amygdaloid complex and adjacent olfactory structures to the entorhinal cortex and to the subiculum in the rat and cat. *J Comp Neurol.* 1977; 172(4):723–752. [PubMed: 838896]
- Krettek JE, Price JL. Amygdaloid projections to subcortical structures within the basal forebrain and brainstem in the rat and cat. *J Comp Neurol.* 1978; 178:225–254. [PubMed: 627625]
- Lanciego JL, Wouterlood FG. A half century of experimental neuroanatomical tracing. *J Chem Neuroanat.* 2011; 42(3):157–183. [PubMed: 21782932]
- Larriva-Sahd J. Juxtacapsular nucleus of the stria terminalis of the adult rat: extrinsic inputs, cell types, and neuronal modules: a combined Golgi and electron microscopic study. *J Comp Neurol.* 2004; 475(2):220–237. [PubMed: 15211463]
- Lesur A, Gaspar P, Alvarez C, Berger B. Chemoanatomic compartments in the human bed nucleus of the stria terminalis. *Neuroscience.* 1989; 32(1):181–194. [PubMed: 2573858]
- Mai JK, Triepel J, Metz J. Neurotensin in the human brain. *Neuroscience.* 1987; 22(2):499–524. [PubMed: 3670596]
- Martin LJ, Powers RE, Dellovade TL, Price DL. The bed nucleus-amygdala continuum in human and monkey. *J Comp Neurol.* 1991; 309:445–485. [PubMed: 1918444]
- McDonald AJ. Neurons of the bed nucleus of the stria terminalis: a golgi study in the rat. *Brain Res Bull.* 1983; 10(1):111–120. [PubMed: 6824959]
- McDonald AJ. Topographical organization of amygdaloid projections to the caudatoputamen, nucleus accumbens, and related striatal-like areas of the rat brain. *Neuroscience.* 1991; 44(1):15–33. [PubMed: 1722890]
- McDonald AJ. Cortical pathways to the mammalian amygdala. *Prog Neurobiol.* 1998; 55(3):257–332. [PubMed: 9643556]
- McDonald AJ, Shammah-Lagnado SJ, Shi C, Davis M. Cortical afferents to the extended amygdala. *Annals of the New York Academy of Sciences.* 1999; 877:309–338. [PubMed: 10415657]
- Meredith GE, Pattiselanno A, Groenewegen HJ, Haber SN. Shell and core in monkey and human nucleus accumbens identified with antibodies to calbindin-D28k. *J Comp Neurol.* 1996; 365:628–639. [PubMed: 8742307]
- Miyashita T, Ichinohe N, Rockland KS. Differential modes of termination of amygdalothalamic and amygdalocortical projections in the monkey. *J Comp Neurol.* 2007; 502(2):309–324. [PubMed: 17348015]
- Mobbs D, Yu R, Rowe JB, Eich H, FeldmanHall O, Dagleish T. Neural activity associated with monitoring the oscillating threat value of a tarantula. *Proc Natl Acad Sci U S A.* 2010; 107(47):20582–20586. [PubMed: 21059963]
- Moga MM, Saper CB, Gray TS. Bed nucleus of the stria terminalis: cytoarchitecture, immunohistochemistry, and projection to the parabrachial nucleus in the rat. *J Comp Neurol.* 1989; 283(3):315–332. [PubMed: 2568370]
- Mufson EJ, Mesulam MM, Pandya DN. Insular interconnections with the amygdala in the rhesus monkey. *Neuroscience.* 1981; 6:1231–1248. [PubMed: 6167896]
- Nance DM, Burns J. Fluorescent dextrans as sensitive anterograde neuroanatomical tracers: applications and pitfalls. *Brain Res Bull.* 1990; 25(1):139–145. [PubMed: 1698517]
- Naya Y, Suzuki WA. Integrating what and when across the primate medial temporal lobe. *Science.* 2011; 333(6043):773–776. [PubMed: 21817056]
- Novotny GEK. A direct ventral connection between the bed nucleus of the stria terminalis and the amygdaloid complex in the monkey (*Macaca fascicularis*). *J Hirnforsch.* 1977; 18:271–284. [PubMed: 411830]

- Ottersen OP, Ben-Ari Y. Afferent connections to the amygdaloid complex of the rat and cat. I. Projections from the thalamus. *J Comp Neurol.* 1979; 187:401–424. [PubMed: 489786]
- Paz R, Gelbard-Sagiv H, Mukamel R, Harel M, Malach R, Fried I. A neural substrate in the human hippocampus for linking successive events. *Proc Natl Acad Sci U S A.* 2010; 107(13):6046–6051. [PubMed: 20231430]
- Petrovich GD, Risold PY, Swanson LW. Organization of projections from the basomedial nucleus of the amygdala: A PHAL study in the rat. *J Comp Neurol.* 1996; 374:387–420. [PubMed: 8906507]
- Petrovich GD, Swanson LW. Projections from the lateral part of the central amygdalar nucleus to the postulated fear conditioning circuit. *Brain Res.* 1997; 763(2):247–254. [PubMed: 9296566]
- Pitkanen A, Amaral DG. Organization of the intrinsic connections of the monkey amygdaloid complex: projections originating in the lateral nucleus. *J Comp Neurol.* 1998; 398(3):431–458. [PubMed: 9714153]
- Porrino LJ, Crane AM, Goldman-Rakic PS. Direct and indirect pathways from the amygdala to the frontal lobe in rhesus monkeys. *J Comp Neurol.* 1981; 198:121–136. [PubMed: 6164704]
- Price JL, Amaral DG. An autoradiographic study of the projections of the central nucleus of the monkey amygdala. *J Neurosci.* 1981; 1:1242–1259. [PubMed: 6171630]
- Price, JL.; Russchen, FT.; Amaral, DG. The limbic region. II. The amygdaloid complex. In: Hokfelt, BT.; Swanson, LW., editors. *Handbook of Chemical Neuroanatomy.* Amsterdam: Elsevier; 1987. p. 279-381.
- Reynolds SM, Berridge KC. Positive and negative motivation in nucleus accumbens shell: bivalent rostrocaudal gradients for GABA-elicited eating, taste “liking”/“disliking” reactions, place preference/avoidance, and fear. *J Neurosci.* 2002; 22(16):7308–7320. [PubMed: 12177226]
- Rosene, D.; Van Hoesen, G. The hippocampal formation of the primate brain. In: Jones, E.; Peters, A., editors. *Cerebral Cortex: Further Aspects of Cortical Function, Including Hippocampus.* New York: Plenum Press; 1987. p. 345-456.
- Rosene DL, Roy NJ, Davis BJ. A cryoprotection method that facilitates cutting frozen sections of whole monkey brains for histological and histochemical processing without freezing artifact. *J Histochem Cytochem.* 1986; 34(10):1301–1315. [PubMed: 3745909]
- Rosene DL, Van Hoesen GW. Hippocampal efferents reach widespread areas of cerebral cortex and amygdala in the rhesus monkey. *Science.* 1977; 198:315–317. [PubMed: 410102]
- Russchen FT, Amaral DG, Price JL. The afferent connections of the substantia innominata in the monkey, *Macaca fascicularis*. *J Comp Neurol.* 1985a; 242:1–27. [PubMed: 3841131]
- Russchen FT, Amaral DG, Price JL. The afferent input to the magnocellular division of the mediodorsal thalamic nucleus in the monkey, *Macaca fascicularis*. *J Comp Neurol.* 1987; 256:175–210. [PubMed: 3549796]
- Russchen FT, Bakst I, Amaral DG, Price JL. The amygdalostriatal projections in the monkey. An anterograde tracing study. *Brain Res.* 1985b; 329:241–257. [PubMed: 3978445]
- Santiago AC, Shammah-Lagnado SJ. Afferent connections of the amygdalopiriform transition area in the rat. *J Comp Neurol.* 2005; 489(3):349–371. [PubMed: 16025448]
- Saunders RC, Rosene DL, Van Hoesen GW. Comparison of the efferents of the amygdala and the hippocampal formation in the rhesus monkey: II. Reciprocal and non-reciprocal connections. *J Comp Neurol.* 1988; 271:185–207. [PubMed: 2454247]
- Schmued L, Kyriakidis K, Heimer L. In vivo anterograde and retrograde axonal transport of the fluorescent rhodamine-dextran-amine, Fluoro-Ruby, within the CNS. *Brain Res.* 1990; 526(1): 127–134. [PubMed: 1706635]
- Shammah-Lagnado SJ, Alheid GF, Heimer L. Efferent connections of the caudal part of the globus pallidus in the rat. *J Comp Neurol.* 1996; 376:489–507. [PubMed: 8956113]
- Shammah-Lagnado SJ, Santiago AC. Projections of the amygdalopiriform transition area (APir). A PHA-L study in the rat. *Ann N Y Acad Sci.* 1999; 877:655–660. [PubMed: 10415677]
- Somerville LH, Whalen PJ, Kelley WM. Human bed nucleus of the stria terminalis indexes hypervigilant threat monitoring. *Biol Psychiatry.* 2010; 68(5):416–424. [PubMed: 20497902]
- Squire LR. The hippocampus and spatial memory. *Trends Neurosci.* 1993; 16(2):56–57. [PubMed: 7680500]

- Sullivan GM, Apergis J, Bush DEA, Johnson LR, Hou M, LeDoux JE. Lesions in the bed nucleus of the stria terminalis disrupt corticosterone and freezing responses elicited by a contextual but not by a specific cue-conditioned fear stimulus. *Neuroscience*. 2004; 128(1):7–14. [PubMed: 15450349]
- Sun N, Roberts L, Cassell MD. Rat central amygdaloid nucleus projections to the bed nucleus of the stria terminalis. *Brain Res Bull*. 1991; 27(5):651–662. [PubMed: 1721859]
- Szucs A, Berton F, Sanna PP, Francesconi W. Excitability of jcBNST neurons is reduced in alcohol-dependent animals during protracted alcohol withdrawal. *PLoS One*. 2012; 7(8):e42313. [PubMed: 22927925]
- Turner BM, Mishikin M, Knapp M. Organization of the amygdalopetal projections from modality-specific cortical association areas in the monkey. *J Comp Neurol*. 1980; 191:515–543. [PubMed: 7419732]
- Veinante P, Stoeckel ME, Lasbennes F, Freund-Mercier MJ. c-Fos and peptide immunoreactivities in the central extended amygdala of morphine-dependent rats after naloxone-precipitated withdrawal. *Eur J Neurosci*. 2003; 18(5):1295–1305. [PubMed: 12956728]
- Vercelli A, Repici M, Garbossa D, Grimaldi A. Recent techniques for tracing pathways in the central nervous system of developing and adult mammals. *Brain Res Bull*. 2000; 51(1):11–28. [PubMed: 10654576]
- Walker DL, Davis M. Double dissociation between the involvement of the bed nucleus of the stria terminalis and the central nucleus of the amygdala in startle increases produced by conditioned versus unconditioned fear. *J Neurosci*. 1997; 17(23):9375–9383. [PubMed: 9364083]
- Walker JR, Ahmed SH, Gracy KN, Koob GF. Microinjections of an opiate receptor antagonist into the bed nucleus of the stria terminalis suppress heroin self-administration in dependent rats. *Brain Res*. 2000; 854(1–2):85–92. [PubMed: 10784110]
- Walter A, Mai JK, Lanta L, Gorcs T. Differential distribution of immunohistochemical markers in the bed nucleus of the stria terminalis in the human brain. *J Chem Neuroanat*. 1991; 4(4):281–298. [PubMed: 1718318]
- Webster MJ, Bachevalier J, Ungerleider LG. Subcortical connections of inferior temporal areas TE and TEO in macaque monkeys. *J Comp Neurol*. 1993; 335(1):73–91. [PubMed: 8408774]
- Weller KL, Smith DA. Afferent connections to the bed nucleus of the stria terminalis. *Brain Res*. 1982; 232(2):255–270. [PubMed: 7188024]
- Zahm DS. Is the caudomedial shell of the nucleus accumbens part of the extended amygdala? A consideration of connections. *Crit Rev Neurobiol*. 1998; 12(3):245–265. [PubMed: 9847057]
- Zahm DS, Jensen SL, Williams ES, Martin JR 3rd. Direct comparison of projections from the central amygdaloid region and nucleus accumbens shell. *Eur J Neurosci*. 1999; 11(4):1119–1126. [PubMed: 10103108]
- Zimmerman JM, Maren S. The bed nucleus of the stria terminalis is required for the expression of contextual but not auditory freezing in rats with basolateral amygdala lesions. *Neurobiol Learn Mem*. 2011; 95(2):199–205. [PubMed: 21073972]

Figure 1. Adjacent sections through the BST processed for AChE (panels A–C), ENK (panels D–F), ST (panels G–I), and NT (panels J–L). Sections are arranged in rostral to caudal order from left to right. Scale bar = 1mm.

Figure 2. Coronal sections through the rostral to caudal primate amygdala stained using cresyl violet (A–D), with adjacent sections processed for AChE activity (E–H). Scale bar = 2.5 mm. (I) Higher power photomicrograph of the primate CTA, at the level of panel C (boxed area). The asterisk is centered on the CTA, which is situated in the semiannual sulcus (arrowhead) between the tri-laminar PAC and the six-layered entorhinal cortex. Scale bar = 1 mm.

Figure 3.

(A) Schematics and representative photomicrographs of retrograde injections into the BSTL. Injections were classified as centered mainly in the BSTLcn (light gray; J24FS, J28LY, J29WGA, J28WGA) or the BSTLJ (dark gray; J28FR, J28FS, J29FR). Representative injection sites are shown with adjacent sections processed for AChE to delineate boundaries of the BSTLcn (case J28WGA) and BSTLJ (case J28LY), respectively. (B) Schematics of injections into subregions of the ventral striatum including the dorsomedial shell (light gray; J24WGA), ventral shell (dark gray; J27FS, J26FS, J27FR), core (black; J13WGA), and the centromedial and ventromedial caudate nucleus (white; J25FS, J24FR). Photomicrographs of several representative injections are shown. Scale bars = 2 mm.

Figure 4.

Charts of retrogradely labeled cells through the rostrocaudal amygdala (left to right) following tracer injections in the BSTLcn (case J29WGA) (A) and the BSTLJ (J28FS) (B), arranged in rostrocaudal order. 1 dot represents 1 cell.

Figure 5.

Photomicrographs of retrogradely labeled cells in the amygdala following injections in the BSTLcn (A–B, D–E) or BSTLJ (G–H). (A–B) Low and high power photomicrographs of labeled cells in the CeLcn (level shown in C) following injection centered on the BSTLcn. (D–E) Photomicrographs of labeled cells in the medial Bpc/CTA (level shown in F) after the same injection. (G–H) Image of labeled cells in the ABmc resulting from an injection centered on the BSTLJ (level in I). Scale bars = 50 μ m.

Figure 6.

Charts of retrogradely labeled cells through the rostrocaudal amygdala (left to right) following tracer injections in the dorsomedial shell (J24WGA) (A) and the ventral shell (J27FS) (B). 1 dot represents 1 cell.

A. Rostrocentral Amygdala

B. Caudal Amygdala

Figure 7. Schematic of anterograde injection sites in the amygdala at mid- (A) and caudal (B) levels. Pink= accessory basal nucleus; green= basal nucleus (asterisk denotes inclusion of CTA); blue=lateral nucleus; yellow= central nucleus; orange= medial nucleus; purple = amygdalohippocampal area/ABs. See text for details of placement.

Figure 8.

Distribution of anterogradely labeled fibers in the BSTL (A–C) and shell (D) following an injection in the Bmc (J12FR). (C) Darkfield photomicrograph of the boxed area in (B) showing labeled fibers in the BSTLJ, BSTLP and SLEAc, with an absence of fibers in the BSTLcn. Asterisk indicates passing fibers, and dotted oval indicates boundary of the BSTLcn. Scale bar = 2mm.

Figure 9.

Distribution of anterogradely labeled fibers in the BSTL (A–C), core and shell (D), following an injection in rostral medial Bpc/CTA (J20LY). (C) Darkfield photomicrograph of the boxed area in (B) showing labeled fibers in the BSTL subdivisions and SLEAc. Asterisks indicate passing fibers and dotted oval indicates boundaries of the BSTLDcn. Scale bar = 2mm.

Figure 10.

Distribution of anterogradely labeled fibers in the BSTL (A–C) and shell following an injection in the ABmc (J8LY). (C) Darkfield photomicrograph of the boxed area in (B) showing labeled fibers in the BSTLJ, BSTLP and SLEAc, with an absence of fibers in the BSTLcn. Asterisk indicates passing fibers and dotted oval indicates boundaries of the BSTLDcn. Scale bar = 2mm.

Figure 11.

Distribution of anterogradely labeled fibers following injections in the CeM and the CeLcn. (A–D) Anterogradely labeled fibers in the shell (A) and BSTL subdivisions (B–C) following an injection centered in the CeM. (D) Darkfield photomicrograph of the boxed area in (C) showing labeled fibers in the BSTLcn, BSTLP, BSTLJ and SLEAc. (E–H) Distribution of anterogradely labeled fibers in the BSTL subdivisions (E–G) following an injection in the CeLcn (J9FS). (H) Darkfield photomicrograph of the boxed area in (G), showing fibers in the BSTLcn and SLEAc.

Figure 12.

Composite charts of retrogradely labeled cells resulting from injections centered on the BSTL and ventral striatal subregions. (A–B) Injections centered in the BSTLcn (green) and BSTLJ (magenta) result in differential distributions of retrogradely labeled cells in amygdala subregions. (C–D) Injections into specific subregions of the ventral striatum also result in differential distributions of labeled cells in the amygdala subregions. Dorsomedial shell (green); ventral shell (magenta); core (light pink); ventromedial caudate nucleus (black). Comparisons across BSTL and ventral striatal cases shows that the distribution following BSTLcn injections is most similar to that following an injection into the dorsomedial shell (green), while injections in the BSTLJ (magenta) result in a pattern of labeled cells that is similar to that of the ventral shell (magenta) and core (light pink).

Table 1

Primary and secondary antibodies.

Primary Antibody	Immunogen	Source (cat.no.)	Working dilution	Secondary antibody
Rabbit polyclonal anti-Lucifer yellow	Lucifer yellow conjugated to dextran amine	Invitrogen/Molecular Probes (A-5750)	1:2000	Goat anti-rabbit IgG, 1:200, Vector Laboratories
Rabbit polyclonal anti-fluorescein	Fluorescein conjugated to dextran amine	Invitrogen/Molecular Probes (A-889)	1:2000	Goat anti-rabbit IgG, 1:200, Vector Laboratories
Rabbit polyclonal anti-fluororuby	Fluororuby conjugated to dextran amine (tetramethylrhodamine)	Invitrogen/Molecular Probes (A-6397)	1:2000	Goat anti-rabbit IgG, 1:200, Vector Laboratories
Rabbit polyclonal anti-triticum vulgaris lectin	Triticum vulgaris (wheat-germ)	Sigma, St. Louis, MO (T-4144)	1:7500	Goat anti-rabbit IgG, 1:200, Vector Laboratories
Rabbit polyclonal anti-somatostatin	Synthetic human somatostatin (amino acid sequence: Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Ser-Cys)	Immunostar, Hudson, WI (20067)	1:1000	Goat anti-rabbit IgG, 1:200, Vector Laboratories
Rabbit polyclonal anti-met-enkephalin	Synthetic human met-enkephalin (amino acid sequence: Tyr-DAla-Gly-Phe-Met)	Immunostar, Hudson, WI (20065)	1:1000	Goat anti-rabbit IgG, 1:200, Vector Laboratories
Rabbit polyclonal anti-neurotensin	Synthetic human neurotensin (amino acid sequence: Leu-Tyr-Glu-Asn-Lys-Pro-Arg-Arg-Tyr-Ile-Leu)	Immunostar, Hudson, WI (20072)	1:1000	Goat anti-rabbit IgG, 1:200, Vector Laboratories
Mouse monoclonal anti-calbindinD-28k	Purified bovine kidney calbindinD-28k	Sigma, St. Louis, MO (C-9848)	1:10,000	Goat anti-mouse IgG, 1:200, Vector Laboratories

Table 2

Relative placement of retrograde tracer injections into BSTL subdivisions and the overlying striatum.

	BSTLcn	BSTLP	BSTLJ	Striatum
J24FS	++++	+	+	
J28WGA	++++	+	++	
J29WGA	++++	+	+	
J28LY	+	+	+	
J28FS			++++	
J28FR			++++	++
J29FR			++++	++++

Table 3

Relative placement of retrograde tracer injections into the striatum

	Dorsomedial Shell	Ventral shell	Core	Ventromedial caudate nucleus	Central caudate nucleus
J24WGA	+++			++	
J27FS		+++	+		
J26FS		+++			
J27FR		+++	+		
J13WGA			+++		
J24FR				+++	
J25FR				+	+++

Table 4

Summary of amygdala subnuclei inputs to the BSTL and the striatum based on anterograde and retrograde injections (– none, + few, ++ moderate, +++ dense, ++++ heavy)

Amygdala nucleus	Central medial Caudate	Ventromedial Caudate	Core	V. Shell	BSTLJ	DM Shell	BSTLcn
L	–	+	–	–	–	–	–
Bmc	–	+	+	+++	+++	–	–
ABmc	–	++	+	++++	+++	–	–
Bpc	–	–	++	+++	+++	+++	+++
CTA	–	–	–	+	++	+++	+++
ABs	–	–	–	–	+	++++	++++
ABpc	–	–	–	+	–	+	+++
M	–	–	–	–	–	+	+
AHA	–	–	–	–	–	+++	+
CeM	–	–	–	+	+	+	+++
CeLcn	–	–	–	–	–	–	+++