Published in final edited form as:

J Allergy Clin Immunol. 2013 August ; 132(2): 414–425.e6. doi:10.1016/j.jaci.2013.03.025.

TIM-1 deficiency eliminates airway hyperreactivity triggered by the recognition of airway cell death

Hye Young Kim, Ph.D.^{1,6}, Ya-Jen Chang, Ph.D.^{1,6}, Ya-Ting Chuang, Ph.D.¹, Hyun-Hee Lee, Ph.D.¹, David I. Kasahara, Ph.D.², Thomas Martin, M.D, Ph.D.³, Joyce T. Hsu, M.D, Ph.D.¹, Paul B. Savage, Ph.D.⁴, Stephanie A. Shore, Ph.D.², Gordon J. Freeman, Ph.D.⁵, Rosemarie H. DeKruyff, Ph.D.¹, and Dale T. Umetsu, M.D, Ph.D.¹

¹Division of Immunology and Allergy, Children's Hospital, Harvard Medical School, Boston, MA, USA

²Molecular and Integrative Physiological Sciences Program, Dept of Environmental Health, Harvard School of Public Health, Boston, MA

³Division of Respiratory Diseases, Children's Hospital, Harvard Medical School, Boston, MA

⁴Brigham Young University, Provo, Utah, USA

⁵Department of Medical Oncology, Dana-Farber Cancer Institute, Department of Medicine, Harvard Medical School, Boston, Massachusetts, USA

Abstract

Background—Studies of asthma have been limited by a poor understanding of how non-allergic environmental exposures such as air pollution and infection are translated in the lung into inflammation and wheezing.

Objective—Our goal was to understand the mechanism of non-allergic asthma that lead to airway hyperreactivity (AHR), a cardinal feature of asthma independent of adaptive immunity.

Method—We examined mouse models of experimental asthma, in which AHR was induced by Respiratory Syncytial Virus (RSV) infection or ozone exposure, using mice deficient in *TIM1/HAVCR1*, an important asthma susceptibility gene.

Results—*TIM1*^{−/−} mice failed to develop airways disease when infected with RSV or when repeatedly exposed to ozone, a major component of air pollution. On the other hand, the *TIM1*^{−/−} mice developed allergen-induced experimental asthma, as previously shown. The RSV- and ozone-induced pathways were blocked by treatment with caspase inhibitors, indicating an absolute requirement for programmed cell death and apoptosis. TIM-1-expressing, but not TIM-1-deficient, natural killer T (NKT) cells responded to apoptotic airway epithelial cells by secreting cytokines, which mediated the development of airway hyperreactivity.

© 2013 American Academy of Allergy, Asthma and Immunology. Published by Mosby, Inc. All rights reserved.

*To whom correspondence should be addressed: Dale T. Umetsu, MD, PhD, Division of Immunology, Karp Laboratories, Rm 10127, Children's Hospital, Harvard Medical School, One Blackfan Circle, Boston, MA 02115, USA. Phone: 617.919.2439, Fax: 617.730.0384, dale.umetsu@childrens.harvard.edu.

⁶Current address of this author is institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan

⁷These authors contributed equally to the completion of these studies.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of interest: The authors have declared that no conflicts of interest exist.

Conclusion—We defined a novel pathway in which TIM-1, a receptor for phosphatidylserine expressed by apoptotic cells, drives the development of asthma by sensing and responding to injured and apoptotic airway epithelial cells.

Keywords

TIM-1; NKT; apoptosis; asthma

Introduction

Until recently, bronchial asthma has been considered to be a single disease, characterized by allergic inflammation, resulting from the actions of allergen-specific Th2 cells and adaptive immunity. Th2 cells secrete cytokines, such as IL-4, IL-5 and IL-13, which orchestrate many of the classical features of asthma, including eosinophilic inflammation, reversible airway obstruction and airway hyperreactivity (AHR). However, although environmental allergens drive the development of pulmonary Th2 cells that are present in the lungs of many patients with asthma, non-allergic triggers, such as exposure to air pollution or viral infections, are associated with major morbidity and mortality in patients with asthma. These non-allergic environmental triggers for asthma suggest that asthma is far more heterogeneous than previously thought, and that multiple asthma phenotypes exist, some associated with allergy and eosinophils, but others associated with airway neutrophilia, steroid unresponsiveness and innate immunity^{1, 2}. Unfortunately, the precise mechanisms by which non-allergic, innate processes trigger asthma symptoms remain undefined, although a role for airway injury, apoptosis and oxidative stress has been suggested^{3–5}. A consequence of the lack of a satisfactory molecular understanding of non-allergic pathways is that most of the non-allergic forms of asthma lack effective therapies and remain significant clinical challenges.

Genetic analyses support the idea that asthma may involve both allergic and non-allergic processes. For example, the linkage of asthma to polymorphisms in *IL13*, *IL4RA*, and *FCER1*, corroborate a role for Th2 biased allergic inflammation in asthma. However, the linkage of asthma to genes such as *ADAM33*⁶, *ORMDL3*⁷ and *TIM1* (*HAVCR1*)⁸, suggests additional pathogenic pathways that may be dependent on innate immunity or defects in the cells present in the airways. Understanding the role of the *TIM1* gene in particular, might help to link and provide insight into allergic and non-allergic processes in asthma. *TIM1* was identified as a susceptibility gene for AHR using linkage analysis in congenic mice⁹, and was subsequently shown to be a susceptibility gene for atopic diseases, including asthma, in humans^{8, 10–12}.

Previous studies regarding the function of TIM-1 in regulating immunity have shown that TIM-1 can function as a costimulatory molecule for T cells (e.g., Th2 cells), although the precise intracellular signaling events that occur downstream of TIM-1 engagement are not yet understood^{9, 13, 14}. Indeed, blockade of TIM-1 with antibody prevented the development of allergen-induced AHR in mice¹⁵ and in humanized mice (mice containing human lymphoid cells)¹⁶. However, these results have been perplexing, since *TIM1*^{−/−} mice have been reported to normally develop allergen-induced AHR¹⁷. In addition, TIM-1, which is expressed by activated T cells, NKT cells, mucosal epithelial cells, hepatocytes and injured kidney cells, is a receptor for the hepatitis A virus (HAV), infection with which is associated with protection against asthma and allergy^{8, 18, 19}. Further, TIM-1 regulates the severity and progression of HAV infection, HIV infection and cerebral malaria^{20–22}. Moreover, TIM-1 has been shown recently to be a receptor for phosphatidylserine (PtdSer)^{23–25}, a marker of apoptotic cells that serves as an “eat me” signal. Also, the capacity of TIM-1 (as well as TIM-3 and TIM-4) to bind PtdSer appears to allow TIM-1 to enhance entry of enveloped viral particles expressing PtdSer, including Ebola, West Nile virus and dengue virus²⁶.

Finally, regulatory B cells have been shown to express TIM-1^{27, 28}. All of these observations, while supporting an interaction between environmental factors and *TIM1* and an important role for TIM-1 in asthma, have confounded our understanding of how TIM-1 might regulate asthma.

To better understand how TIM-1 affects the development of asthma, we generated *TIM1*^{-/-} mice, which we examined in several distinct experimental models of asthma. First, the *TIM1*^{-/-} mice developed severe *allergen*-induced AHR similarly to wildtype (WT) mice, but failed to develop AHR when infected with respiratory syncytial virus (RSV). RSV is a very common respiratory infection in young children and adults²⁹⁻³¹, and is the most frequent respiratory cause of hospitalization of infants in the US, associated with significant wheezing through processes that are poorly understood.

Second, in a model of asthma associated with air pollution³², *TIM1*^{-/-} mice exposed repeatedly to ozone failed to develop AHR. Exposure to ozone, a major component of air pollution, is known to increase lung oxidant stress, commonly seen in patients with severe asthma³³, and to increase asthmatic symptoms in children, even at concentrations below the U.S. EPA standard³⁴. Further, on days of high ambient ozone concentrations, hospital admissions for asthma are higher³⁵, presumably because ozone exposure induces AHR associated with airway epithelial cell damage³⁶, and with neutrophils and inflammatory mediators in the airways^{37, 38}.

Importantly, the development of both RSV- and ozone-induced AHR was blocked by treatment with caspase inhibitors, while allergen-induced AHR was unaffected. These results indicate that airway injury and apoptosis play critical roles in non-allergic asthma, and are sensed by TIM-1 expressing cells in the lungs that translate airway cell death into AHR and airway inflammation. These results highlight differences between allergic and non-allergic forms of asthma, and provide specific mechanisms for the development of non-allergic asthma.

Materials and Methods

Mice

Wild-type BALB/c ByJ mice on the BALB/c background were purchased from The Jackson Laboratory. *Ja18*^{-/-} mice were gifts from M. Taniguchi/T. Nakayama (Chiba University). The mice were studied at 6–8 wks of age and were age matched. All the animal experiments were approved by the Animal Care and Use Committee of Children's Hospital Boston.

Antibodies and reagents

Anti-mouse TIM-1 mAb blocking antibody (3D10)³⁹ was injected 24 hr before ozone exposure. Anti-Fas mAb (Jo2) and caspase inhibitor (Q-VD-OPH) were purchased from BD Biosciences and SM Biochemicals respectively. α -GalCer was synthesized by P.B. Savage (Brigham Young University, Provo, Utah), and was used for *in vitro* NKT cell cultures.

Measurement of AHR

Mice were anesthetized with 50 mg/kg pentobarbital and instrumented for the measurement of pulmonary mechanics (BUXCO Electronics). Mice were tracheostomized, intubated, and mechanically ventilated at a tidal volume of 0.2 ml and a frequency of 150 breath/min, as previously described⁴¹. Lung resistance (R_L) was measured in response to increasing doses (0.125 to 40 mg/ml) of aerosolized acetyl- β -methylcholine chloride methacholine (Sigma-Aldrich).

Statistical tests

Unpaired Student's *t*-test or ANOVA were used for comparisons between groups (Prism 4; GraphPad Software Inc., San Diego, CA). Significance for all statistical tests are shown in figures for $P<0.05$ (*), $P<0.01$ (**) and $P<0.001$ (***)

Results

A critical role for TIM-1 in ozone-induced AHR

Using an experimental model of air pollution-associated asthma, in which mice were exposed repeatedly to ozone³², we found that *TIM1*^{-/-} mice, generated by targeted deletion of *TIM1/HAVCR1* in BALB/c embryonic stem cells (Fig. 1A), failed to develop ozone-induced AHR and airway inflammation (Fig. 1B, 1C and 1D). The *TIM1*^{-/-} mice had normal numbers of T cells, and B cells in lymphoid organs (Fig. S1), but failed to express TIM-1, for example, on pDC, activated B cells, and NKT cells, which are known to express TIM-1 (Fig.S2). Because ozone-induced AHR and airway inflammation is dependent on NKT cells (it does not occur in *CD1d*^{-/-} mice)³², we compared the number of NKT cells in the lung after air or ozone exposure. The total number of NKT cells in the lung was increased significantly in WT mice but not in *TIM1*^{-/-} mice (Fig. 1E). To our knowledge this is the first report of a specific phenotypic defect in *TIM1*^{-/-} mice.

Allergen-induced AHR does not require TIM-1

The requirement for TIM-1 was specific for ozone-induced AHR, since both WT and *TIM1*^{-/-} mice sensitized and challenged with ovalbumin (OVA) developed allergen-induced AHR with airway eosinophilia (Fig. 2A and B), as previously reported¹⁷. In this response, *TIM1*^{-/-} mice developed normal Th2 and pulmonary NKT cell responses, similar to that seen in WT mice (Fig. 2C). Therefore, TIM-1 was required for the development of ozone-induced but not OVA-induced AHR.

Ozone exposure induces apoptosis of airway epithelial cells

Since TIM-1 is a receptor for PtdSer and mediates the binding and recognition of apoptotic cells²³⁻²⁵, we asked if apoptosis was involved in the ozone-induced AHR response. Indeed, after repeated ozone exposure, increased numbers of apoptotic airway epithelial cells were found in the lung, which were PI and TUNEL double positive, as previously reported⁴⁴ (Fig. 3A). AnnexinV⁺ apoptotic airway cells developed after ozone exposure to the same degree in both WT and *TIM1*^{-/-} mice (Fig.3B). Furthermore, treating ozone exposed wildtype mice with an anti-TIM-1 mAb that blocks TIM-1 recognition of PtdSer, prevented the development of ozone-induced AHR and airway inflammation (Fig. 3C and D). These results suggest that ozone exposure results in apoptosis of airway epithelial cells, and that TIM-1 expressing cells can translate apoptosis into inflammation.

Prevention of apoptosis blocks ozone-induced AHR

We next showed that treatments that block apoptosis surprisingly prevented the development of ozone-induced AHR. Administration of Q-VD-OPH, which binds caspases thereby preventing caspase-dependent apoptosis, prior to ozone exposure abolished ozone-induced AHR and airway inflammation (Fig 4A, B and C). Treatment with Q-VD-OPH reduced the number of apoptotic airway epithelial cells (Fig. 4D), supporting the idea that apoptosis is required for the development of ozone induced AHR. In contrast, treatment with Q-VD-OPH had no effect on OVA-induced AHR and airway eosinophilia (Fig. S3), indicating that apoptosis was required for the development of ozone-, but not OVA-induced AHR.

Damaged and apoptotic cells stimulate NKT cells

Since ozone-induced AHR does not occur in *CD1d*^{-/-} mice and therefore requires the presence of NKT cells³², we next examined whether TIM-1 expressing NKT cells participated in the response. Apoptotic or healthy mouse airway epithelial cells (MLE12) (Fig 5A) were co-cultured with NKT cells from WT or *TIM1*^{-/-} mice. We found that apoptotic but not healthy airway epithelial cells activated WT NKT cells when co-cultured *in vitro* with a suboptimal dose of α -GalCer (1 ng/ml), as demonstrated by production of IL-4 and IL-13 (Fig 5B and Fig. S4). However, apoptotic airway epithelial cells failed to activate NKT from *TIM1*^{-/-} mice, although both WT and *TIM1*^{-/-} NKT cells could be activated by an optimal dose of α -GalCer (100 ng/ml) to produce both IL-4 and IL-13. This TIM-1 dependent response to apoptotic cells required the recognition of PtdSer, since it was blocked by addition of Annexin V (Fig.5B), which binds PtdSer, thus preventing the recruitment of TIM-1 in the apoptotic cell-NKT cell synapse³⁹. Moreover, human NKT cells were also activated by apoptotic but not healthy human type II epithelial cells (A549), and produced significant levels of IL-4 after co-culture with apoptotic cells *in vitro* (Fig. 5C and D). Therefore, TIM-1 delivers a co-stimulatory signal to NKT cells resulting in cytokine production, in both mice and in humans.

TIM-1 expression by NKT cells is required for ozone-induced AHR

We next examined the *in vivo* role of TIM-1 expressing NKT cells in the development of ozone-induced AHR. *Ja18*^{-/-} mice, which lack NKT cells, failed to develop ozone-induced AHR³², but the response in *Ja18*^{-/-} mice was fully reconstituted by adoptive transfer of NKT cells from WT but not *TIM1*^{-/-} mice (Fig. 6A, B and C). Moreover, in the lungs of WT mice exposed to ozone, the number of NKT producing IL-4 (Fig 6D, **left panel**), both IL-4 and IL-13 (**middle panel**) or IL-17 (**right panel**), increased significantly, and this was greatly reduced in the *TIM1*^{-/-} mice. These observations together indicate that apoptotic cells in the airways activate TIM-1 expressing NKT cells, which then drive the development of AHR and airway inflammation.

Infection with RSV induces AHR in wildtype but not *TIM1*^{-/-} mice

We next examined the role of TIM-1 in another innate model of asthma, induced with RSV infection. RSV is an important cause of serious respiratory illness and wheezing in young children and in adults²⁹⁻³¹, through processes that are poorly understood. Unexpectedly, *TIM1*^{-/-} mice failed to develop RSV-induced AHR, whereas WT mice developed robust AHR (Figs. 7A and 7B). Furthermore, TIM-1 expressing NKT cells were required for the development of RSV-induced AHR, since it did not occur in NKT cell deficient *CD1d*^{-/-} mice (Figs. 7C and 7D). RSV infection in WT mice was associated with a significant increase in the number of NKT cells in the BAL fluid (Fig. 7E). RSV infection also resulted in an increase in the number of apoptotic cells in the lungs (Fig. 7F), as has been previously reported⁴⁵⁻⁴⁷. Moreover, treatment with Q-VD-OPH blocked the induction of AHR by RSV (Figs. 7G and 7H), indicating that apoptosis was also required for RSV-induced AHR. Furthermore, adoptive transfer of NKT cells from WT but not from *TIM1*^{-/-} mice reconstituted RSV-induced AHR and airway inflammation (Fig.7I and 7J). These results together strongly support an important role for TIM-1 expressing NKT cells in recognizing apoptotic airway epithelial cells, and in translating programmed cell death into airway inflammation and AHR (Fig. 8).

Discussion

In these experiments, we demonstrated an important role for *TIM1*, an important asthma susceptibility gene, in which NKT cells responding to apoptotic airway cells combine in a novel pathway that triggers the development of AHR. This innate TIM-1 pathway clarifies

several previously unexplained elements of non-allergic asthma, which do not fit with the prevailing Th2 paradigm of asthma, such as its association with lung injury, oxidative stress and airway neutrophilia. We showed that TIM-1 expressing NKT cells responded to apoptotic, but not healthy airway epithelial cells, translating excessive airway cell death into airway inflammation and AHR, in the absence of adaptive immunity. These studies are important because they explain a specific function for TIM-1, a receptor for PtdSer, in mediating the development of airways disease, and provide a specific molecular mechanism for asthma triggered by lung injury. Together, these studies of a TIM-1-apoptotic cell dependent pathway provide an important new dimension in our understanding of the heterogeneity and complexity of asthma.

In the normal lung, apoptotic cells are rarely observed, presumably because the removal of apoptotic cells by alveolar macrophages, a process called efferocytosis, is extremely efficient. However, after repeated exposure to ozone, hyperoxia or following infection or exposure to particulates, large numbers of airway epithelial cells may apoptose and overwhelm the normal clearance mechanisms. Apoptotic cells appeared in the lungs of both WT and *TIM1*^{-/-} mice after exposure to ozone, but only NKT cells that expressed TIM-1 could recognize and translate apoptosis into AHR. Therefore, we suggest that TIM-1, expressed by airway NKT cells, senses apoptosis, mediates the activation of both mouse and human NKT cells and triggers the development of inflammation and AHR. This process develops in the context of lung injury with apoptosis, but not during allergen-induced AHR, explaining why *TIM1*^{-/-} mice develop allergen-induced AHR but not ozone-induced AHR. Thus, although both allergen-induced and ozone-induced AHR require the presence of NKT cells^{32, 41}, the activation of NKT cells during allergen challenge is independent of TIM-1, and was not affected by treatments that interfered with apoptosis. These observations highlight a fundamental difference between the pathological pathways that lead to allergic versus non-allergic asthma.

Although apoptosis has not been thought previously to play a prominent role in asthma, increased numbers of apoptotic cells have been observed in the lungs of patients with asthma, particularly in patients with severe asthma. In such patients the number of apoptotic cells was 10 fold greater than in subjects with mild or no asthma⁴⁸. In children with asthma, apoptotic epithelial cell numbers were also noted to be greater than in non-asthmatic control subjects, as was basement membrane thickening³³. While impaired removal of apoptotic cells by alveolar macrophages is thought to contribute to chronic airway inflammation in cystic fibrosis and COPD⁴⁹, the precise mechanisms by which apoptosis leads to airway inflammation or AHR has been previously unknown. Since NKT cells are normally present in the lungs but are increased in number in the lungs of many, though not all, patients with asthma, particularly in those with severe disease⁵⁰, the colocalization of, and interaction between, apoptotic cells and NKT cells is likely to occur in the lungs of asthmatic patients following exposure to many different environmental agents that are known to cause epithelial cell apoptosis. Our studies now suggest that apoptosis is a critical trigger for the development of asthma associated with lung injury from viral infection or oxidative stress. This mechanism of asthma might be considered a specific “asthma endotype”, defined as an asthma subtype driven by a specific molecular-biological pathway⁵¹.

In our studies, we found that TIM-1 dependent AHR was observed not only with ozone exposure but also with RSV infection. Until now, the conventional wisdom has been that RSV infection causes wheezing by enhancing Th2 immunity, which then causes “allergic” asthma^{52, 53}. This seems an unlikely pathway, since RSV infection very rapidly causes wheezing regardless of the presence of allergy. On the other hand, RSV infection induces airway epithelial cell apoptosis⁴⁶, which we suggest initiates the activation of TIM-1 expressing NKT cells and leads to AHR. Moreover, it is tempting to speculate that the

TIM-1-apoptosis-NKT cell axis may be generalizable to a number of other situations in which AHR occurs in association with airway injury and epithelial cell apoptosis, for example, during rhinovirus infection⁵⁴ or following oxidative stress or ischemia/reperfusion injury.

Our current study greatly extends our previous observation that treatment of mice with anti-Fas mAb induced airway apoptosis, and resulted in AHR through a process that involved TIM-1³⁹. While the previous anti-Fas mAb study suggested a role for apoptosis in lung disease, we now definitively show an essential and non-redundant role for TIM-1, using *TIM1*^{-/-} mice, in two clinically relevant models of asthma, in which mice were exposed to common environmental agents (e.g., viral infection or ozone) causing airway apoptosis and NKT cell activation. Moreover, we demonstrated that both mouse and human NKT cells respond to apoptotic airway epithelial cells, suggesting that TIM-1-expressing NKT cells may indeed regulate human non-allergic asthma. Together, these studies conclusively demonstrate the importance of an axis involving TIM-1, apoptosis and recognition of PtdSer in non-allergic asthma. Finally, since the *TIM1* gene has been shown to be extremely polymorphic, these results could explain how *TIM1* functions as an important asthma susceptibility gene: TIM-1 variants may differentially bind PtdSer and apoptotic cells, resulting in differential activation of NKT cells in different individuals.

Although apoptosis is considered classically as a “silent death” and tolerogenic, it has become apparent that apoptosis in some situations can be highly immunogenic, eliciting strong immune responses^{55, 56}. The immunogenicity of cell death however, depends on many factors, including the characteristics and activation state of the dying cell, what is released from the cell, how quickly the apoptotic cell is removed and where the apoptotic events occur⁵⁶. We suggest that in the lungs, apoptosis may increase during viral infection, toxic exposures or subsequent to ischemia-reperfusion injury^{57–59}, and may be recognized by TIM-1 expressing NKT cells, when present. We demonstrated that ozone exposure and RSV infection increased epithelial cell apoptosis, which was required along with NKT cell activation for the development of AHR. It is possible nonetheless, that additional molecular mechanisms that develop in stressed and injured epithelial cells during RSV infection and ozone exposure, such as secretion of ATP, exposure of calreticulin at the cell surface, and release of HMGB1 protein may contribute to the pulmonary inflammatory process^{55, 56}.

In summary, we identified a novel innate inflammatory pathway in the lungs triggered by the recognition of apoptotic cells through TIM-1 on NKT cells that results in the development of AHR. These results indicate that lung inflammation and AHR are much more complex than suggested by the Th2 paradigm, and that additional pathways and innate cell types beyond Th2 cells can lead to asthma¹. Although Th2 cells, eosinophils, mast cells and adaptive immunity play important roles in asthma, additional innate pathways, such as this TIM-1-apoptosis-NKT cell axis, can lead to AHR and asthma. While each of these pathways can function independently of the other, it is likely that in a given individual, several or all of these pathways may be functional, either separately or synergistically, depending on environmental exposures at a given time. Moreover, it is possible that by developing methods that target the TIM-1-apoptosis pathway, better and more effective therapies for patients with asthma may result.

Acknowledgments

This work was supported in part by NIH grant PO1 AI054456 and R01-A1-089955. We thank Sheena Monahan and Veronica Shaw for excellent technical support.

Abbreviations

AHR	airway hyperreactivity
NKT	Natural Killer T
WT	wild-type
RSV	Respiratory Syncytial Virus

References

- Kim HY, DeKruyff RH, Umetsu DT. The many paths to asthma: phenotype shaped by innate and adaptive immunity. *Nat Immunol*. 2010; 11:577–584. [PubMed: 20562844]
- Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med*. 2012; 18:716–725. [PubMed: 22561835]
- Moussion C, Ortega N, Girard JP. The IL-1-like cytokine IL-33 is constitutively expressed in the nucleus of endothelial cells and epithelial cells in vivo: a novel 'alarmin'? *PLoS One*. 2008; 3:e3331. [PubMed: 18836528]
- Henson PM, Tudor RM. Apoptosis in the lung: induction, clearance and detection. *Am J Physiol Lung Cell Mol Physiol*. 2008; 294:L601–L611. [PubMed: 18178675]
- Auerbach A, Hernandez ML. The effect of environmental oxidative stress on airway inflammation. *Curr Opin Allergy Clin Immunol*. 2012; 12:133–139. [PubMed: 22306553]
- Van Eerdewegh P, Little RD, Dupuis J, RG DM, Falls K, Simon J, et al. Association of the ADAM33 gene with asthma and bronchial hyperresponsiveness. *Nature*. 2002; 418:426–430. [PubMed: 12110844]
- Moffatt MF, Kabisch M, Liang L, Dixon AL, Strachan D, Heath S, et al. Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature*. 2007; 448:470–473. [PubMed: 17611496]
- McIntire J, Umetsu S, Macaubas C, Hoyte E, Cinnioğlu C, Cavalli-Sforza L, et al. Immunology: hepatitis A virus link to atopic disease. *Nature*. 2003; 425:576. [PubMed: 14534576]
- McIntire JJ, Umetsu SE, Akbari O, Potter M, Kuchroo VK, Barsh GS, et al. Identification of Tapr (an airway hyperreactivity regulatory locus) and the linked Tim gene family. *Nat Immunol*. 2001; 2:1109–1116. [PubMed: 11725301]
- Graves PE, Siroux V, Guerra S, Klimecki WT, Martinez FD. Association of atopy and eczema with polymorphisms in T-cell immunoglobulin domain and mucin domain-IL-2-inducible T-cell kinase gene cluster in chromosome 5 q 33. *J Allergy Clin Immunol*. 2005; 116:650–656. [PubMed: 16159638]
- Gao PS, Mathias RA, Plunkett B, Togias A, Barnes KC, Beaty TH, et al. Genetic variants of the T-cell immunoglobulin mucin 1 but not the T-cell immunoglobulin mucin 3 gene are associated with asthma in an African American population. *J Allergy Clin Immunol*. 2005; 115:982–988. [PubMed: 15867855]
- Wu Q, Hu L, Cai P, Li Y, Chen F, Kong L. Association analysis of TIM-1 232G > A and 5383_5397 insertion/deletion polymorphisms with childhood asthma and total serum immunoglobulin E levels in middle China. *J Investig Allergol Clin Immunol*. 2009; 19:146–153.
- Umetsu S, Lee W, McIntire J, Downey L, Sanjanwala B, Akbari O, et al. TIM-1 induces T cell activation and inhibits the development of peripheral tolerance. *Nature Immunol*. 2005; 6:447–454. [PubMed: 15793575]
- Freeman GJ, Casasnovas JM, Umetsu DT, Dekruyff RH. TIM genes: a family of cell surface phosphatidylserine receptors that regulate innate and adaptive immunity. *Immunol Rev*. 2010; 235:172–189. [PubMed: 20536563]
- Encinas JA, Janssen EM, Weiner DB, Calarota SA, Nieto D, Moll T, et al. Anti-T-cell Ig and mucin domain-containing protein 1 antibody decreases TH2 airway inflammation in a mouse model of asthma. *J Allergy Clin Immunol*. 2005; 116:1343–1349. [PubMed: 16337469]

16. Sonar S, Hsu Y-M, Conrad M, Majeau G, Kilic A, Garber E, et al. Antagonism of TIM-1 blocks the development of disease in a humanized mouse model of allergic asthma. *J Clin Invest*. 2010; 120:2767–2781. [PubMed: 20628202]
17. Barlow JL, Wong SH, Ballantyne SJ, Jolin HE, McKenzie AN. Tim1 and Tim3 are not essential for experimental allergic asthma. *Clin Exp Allergy*. 2011; 41:1012–1021. [PubMed: 21470319]
18. Matricardi PM, Rosmini F, Ferrigno L, Nisini R, Rapicetta M, Chionne P, et al. Cross sectional retrospective study of prevalence of atopy among Italian military students with antibodies against hepatitis A virus. *BMJ*. 1997; 314:999–1003. [PubMed: 9112843]
19. Matricardi PM, Rosmini F, Panetta V, Ferrigno L, Bonini S. Hay fever and asthma in relation to markers of infection in the United States. *J Allergy Clin Immunol*. 2002; 110:381–387. [PubMed: 12209083]
20. Kim H, Eyheramonho M, Pichavant M, Cambaceres C, Matangkasombut P, Cervio G, et al. A polymorphism in TIM1 is associated with susceptibility to severe hepatitis A virus infection in humans. *J Clin Invest*. 2011; 121:1111–1118. [PubMed: 21339644]
21. Wichukchinda N, Nakajima T, Saipradit N, Nakayama EE, Ohtani H, Rojanawiwat A, et al. TIM1 haplotype may control the disease progression to AIDS in a HIV-1-infected female cohort in Thailand. *AIDS*. 2010; 24:1625–1631. [PubMed: 20479636]
22. Drube S, Heink S, Walter S, Lohn T, Grusser M, Gerbaulet A, et al. The receptor tyrosine kinase c-Kit controls IL-33 receptor signaling in mast cells. *Blood*. 2010; 115:3899–3906. [PubMed: 20200353]
23. Kobayashi N, Karisola P, Pena-Cruz V, Dorfman DM, Jinushi M, Umetsu SE, et al. TIM-1 TIM-4 Glycoproteins Bind Phosphatidylserine and Mediate Uptake of Apoptotic Cells. *Immunity*. 2007; 27:927–940. [PubMed: 18082433]
24. Santiago C, Ballesteros A, Martinez-Munoz L, Mellado M, Kaplan GG, Freeman GJ, et al. Structures of T Cell Immunoglobulin Mucin Protein 4 Show a Metal-Ion-Dependent Ligand Binding Site where Phosphatidylserine Binds. *Immunity*. 2007; 27:941–951. [PubMed: 18083575]
25. Miyanishi M, Tada K, Koike M, Uchiyama Y, Kitamura T, Nagata S. Identification of Tim4 as a phosphatidylserine receptor. *Nature*. 2007; 450:435–439. [PubMed: 17960135]
26. Jemielity S, Wang J, Chan Y, Ahmed A, Li WW, Monahan S, et al. TIM-family proteins promote infection of multiple enveloped viruses through virion-associated phosphatidylserine. *PLoS Pathogen*. 2013 in press.
27. Ding Q, Yeung M, Camirand G, Zeng Q, Akiba H, Yagita H, et al. Regulatory B cells are identified by expression of TIM-1 and can be induced through TIM-1 ligation to promote tolerance in mice. *J Clin Invest*. 2011
28. Xiao S, Brooks CR, Zhu C, Wu C, Sweere JM, Petecka S, et al. Defect in regulatory B-cell function and development of systemic autoimmunity in T-cell Ig mucin 1 (Tim-1) mucin domain-mutant mice. *Proc Natl Acad Sci U S A*. 2012; 109:12105–12110. [PubMed: 22773818]
29. Hall CB, Weinberg GA, Iwane MK, Blumkin AK, Edwards KM, Staat MA, et al. The burden of respiratory syncytial virus infection in young children. *N Engl J Med*. 2009; 360:588–598. [PubMed: 19196675]
30. Westerly BD, Peebles RS Jr. Respiratory syncytial virus infections in the adult asthmatic--mechanisms of host susceptibility and viral subversion. *Immunol Allergy Clin North Am*. 2010; 30:523–539. vi–vii. [PubMed: 21029936]
31. Falsey AR, Hennessey PA, Formica MA, Cox C, Walsh EE. Respiratory syncytial virus infection in elderly and high-risk adults. *N Engl J Med*. 2005; 352:1749–1759. [PubMed: 15858184]
32. Pichavant M, Goya S, Meyer EH, Johnston RA, Kim HY, Matangkasombut P, et al. Ozone exposure in a mouse model induces airway hyperreactivity that requires the presence of natural killer T cells and IL-17. *J Exp Med*. 2008; 205:385–393. [PubMed: 18250191]
33. Fitzpatrick AM, Teague WG, Holguin F, Yeh M, Brown LA. Airway glutathione homeostasis is altered in children with severe asthma: evidence for oxidant stress. *J Allergy Clin Immunol*. 2009; 123:146–152. e8. [PubMed: 19130935]
34. Gent JF, Triche EW, Holford TR, Belanger K, Bracken MB, Beckett WS, et al. Association of low-level ozone and fine particles with respiratory symptoms in children with asthma. *Jama*. 2003; 290:1859–1867. [PubMed: 14532314]

35. Tolbert PE, Mulholland JA, MacIntosh DL, Xu F, Daniels D, Devine OJ, et al. Air quality and pediatric emergency room visits for asthma in Atlanta, Georgia, USA. *Am J Epidemiol.* 2000; 151:798–810. [PubMed: 10965977]
36. Li Z, Potts-Kant EN, Garantziotis S, Foster WM, Hollingsworth JW. Hyaluronan signaling during ozone-induced lung injury requires TLR4, MyD88, and TIRAP. *PLoS One.* 2011; 6:e27137. [PubMed: 22073274]
37. Aris RM, Christian D, Hearne PQ, Kerr K, Finkbeiner WE, Balmes JR. Ozone-induced airway inflammation in human subjects as determined by airway lavage and biopsy. *Am Rev Respir Dis.* 1993; 148:1363–1372. [PubMed: 8239177]
38. Devlin RB, McDonnell WF, Mann R, Becker S, House DE, Schreinemachers D, et al. Exposure of humans to ambient levels of ozone for 6.6 hours causes cellular and biochemical changes in the lung. *Am J Respir Cell Mol Biol.* 1991; 4:72–81. [PubMed: 1846079]
39. Lee H-H, Meyer E, Goya S, Pichavant M, Kim H-Y, Bu X, et al. Apoptotic cells activate NKT cells through T cell Ig-like mucin-like-1 resulting in airway hyperreactivity. *J Immunol.* 2010; 185:5225–5235. [PubMed: 20889552]
40. Shore SA, Johnston RA, Schwartzman IN, Chism D, Krishna Murthy GG. Ozone-induced airway hyperresponsiveness is reduced in immature mice. *J Appl Physiol.* 2002; 92:1019–1028. [PubMed: 11842035]
41. Akbari O, Stock P, Meyer E, Kronenberg M, Sidobre S, Nakayama T, et al. Essential role of NKT cells producing IL-4 and IL-13 in the development of allergen-induced airway hyperreactivity. *Nature Medicine.* 2003; 9:582–588.
42. Bera MM, Lu B, Martin TR, Cui S, Rhein LM, Gerard C, et al. Th17 cytokines are critical for respiratory syncytial virus-associated airway hyperresponsiveness through regulation by complement C3a and tachykinins. *J Immunol.* 2011; 187:4245–4255. [PubMed: 21918196]
43. Prince GA, Jenson AB, Horswood RL, Camargo E, Chanock RM. The pathogenesis of respiratory syncytial virus infection in cotton rats. *Am J Pathol.* 1978; 93:771–791. [PubMed: 362946]
44. Triantaphyllopoulos K, Hussain F, Pinart M, Zhang M, Li F, Adcock I, et al. A model of chronic inflammation and pulmonary emphysema after multiple ozone exposures in mice. *Am J Physiol Lung Cell Mol Physiol.* 2011; 300:L691–L700. [PubMed: 21355040]
45. Welliver TP, Garofalo RP, Hosakote Y, Hintz KH, Avendano L, Sanchez K, et al. Severe human lower respiratory tract illness caused by respiratory syncytial virus and influenza virus is characterized by the absence of pulmonary cytotoxic lymphocyte responses. *J Infect Dis.* 2007; 195:1126–1136. [PubMed: 17357048]
46. Villenave R, Thavagnanam S, Sarlang S, Parker J, Douglas I, Skibinski G, et al. In vitro modeling of respiratory syncytial virus infection of pediatric bronchial epithelium, the primary target of infection in vivo. *Proc Natl Acad Sci U S A.* 2012; 109:5040–5045. [PubMed: 22411804]
47. Kotelkin A, Prikhod'ko EA, Cohen JI, Collins PL, Bukreyev A. Respiratory syncytial virus infection sensitizes cells to apoptosis mediated by tumor necrosis factor-related apoptosis-inducing ligand. *J Virol.* 2003; 77:9156–9172. [PubMed: 12915532]
48. Cohen L, E X, Tarsi J, Ramkumar T, Horiuchi TK, Cochran R, et al. Epithelial cell proliferation contributes to airway remodeling in severe asthma. *Am J Respir Crit Care Med.* 2007; 176:138–145. [PubMed: 17463414]
49. Vandivier RW, Henson PM, Douglas IS. Burying the dead: the impact of failed apoptotic cell removal (efferocytosis) on chronic inflammatory lung disease. *Chest.* 2006; 129:1673–1682. [PubMed: 16778289]
50. Matangkasombut P, Marigowda G, Irvine A, Idris I, Pichavant M, Kim H, et al. Natural Killer T Cells in the Lungs of Patients with Asthma. *J Allergy Clin Immunol.* 2009; 123:1181–1185. [PubMed: 19356791]
51. Anderson GP. Endotyping asthma: new insights into key pathogenic mechanisms in a complex, heterogeneous disease. *Lancet.* 2008; 372:1107–1119. [PubMed: 18805339]
52. Culley FJ, Pollott J, Openshaw PJ. Age at first viral infection determines the pattern of T cell-mediated disease during reinfection in adulthood. *J Exp Med.* 2002; 196:1381–1386. [PubMed: 12438429]

53. Krishnamoorthy N, Khare A, Oriss TB, Raundhal M, Morse C, Yarlagadda M, et al. Early infection with respiratory syncytial virus impairs regulatory T cell function and increases susceptibility to allergic asthma. *Nat Med.* 2012; 18:1525–1530. [PubMed: 22961107]
54. Deszcz L, Gaudernak E, Kuechler E, Seipelt J. Apoptotic events induced by human rhinovirus infection. *J Gen Virol.* 2005; 86:1379–1389. [PubMed: 15831950]
55. Kepp O, Tesniere A, Schlemmer F, Michaud M, Senovilla L, Zitvogel L, et al. Immunogenic cell death modalities and their impact on cancer treatment. *Apoptosis.* 2009; 14:364–375. [PubMed: 19145485]
56. Griffith TS, Ferguson TA. Cell death in the maintenance and abrogation of tolerance: the five Ws of dying cells. *Immunity.* 2011; 35:456–466. [PubMed: 22035838]
57. Havasi A, Borkan SC. Apoptosis and acute kidney injury. *Kidney Int.* 2011; 80:29–40. [PubMed: 21562469]
58. Uchida Y, Ke B, Freitas MC, Ji H, Zhao D, Benjamin ER, et al. The emerging role of T cell immunoglobulin mucin-1 in the mechanism of liver ischemia and reperfusion injury in the mouse. *Hepatology.* 2010; 51:1363–1372. [PubMed: 20091883]
59. Wong CH, Jenne CN, Lee WY, Leger C, Kubes P. Functional innervation of hepatic iNKT cells is immunosuppressive following stroke. *Science.* 2011; 334:101–105. [PubMed: 21921158]

Key messages

- This study defines a novel disease mechanism in which TIM-1, a receptor for phosphatidylserine exposed on apoptotic cells and an important atopy susceptibility gene, drives the development of non-allergic asthma.
- This pathway results in airway hyperreactivity (AHR), a cardinal feature of asthma, and is initiated by the activation of NKT cells which sense and respond to lung injury and the death of airway epithelial cells through TIM-1.
- RSV infection or ozone exposure induce apoptosis or damage of lung cells, which then stimulate NKT cells to produce cytokines that are critical for the development of AHR.

Figure 1. TIM-1 is required for the development of ozone-induced AHR

(A) The strategy for disruption of the TIM-1 gene. Deletion of TIM-1 gene was confirmed by PCR.

(B) Results represent changes in lung resistance (R_L) in wild type or $TIM1^{-/-}$ mice after air or ozone exposure. * $P < 0.05$, and ** $P < 0.01$, compared to air exposed group (Two-way ANOVA).

(C) Inflammatory cell numbers in BAL fluid of 1B (cells/lung). * $P < 0.05$, and ** $P < 0.01$, compared to air exposed group (Two-tailed t-test).

(D) Representative lung sections stained with hematoxylin/eosin (magnification; X10).

(E) Percentage of tetramer⁺TCR β ⁺ NKT cells in the lung after exposure to either air or ozone (left) and total number of NKT cells (right). *** $P < 0.001$ (WT, air vs ozone exposed group) or *** $P < 0.001$ (WT vs *TIM1*^{-/-} mice after ozone exposure) (Two-tailed t-test). Data are representative of at least three independent experiments with three to five mice per group.

Figure 2. TIM-1 is not required for the development of OVA-induced AHR

(A) WT and $TIM1^{-/-}$ mice develop AHR equivalently after sensitization and challenge with OVA. Graph represents the changes in lung resistance (R_L). $TIM1^{-/-}$ OVA group was compared to WT OVA group (Two-way ANOVA). ns, not significant.

(B) Inflammatory cell numbers in BAL fluid after OVA challenge (cells/lung). The $TIM1^{-/-}$ OVA group was compared to WT OVA group (not significant, Two-tailed t-test). Mac, macrophage; Neu, neutrophils; Eos, eosinophils; Lymph, lymphocytes.

(C) The number of NKT cells was calculated as Fig.1G. Tetramer⁺TCR β ⁺ NKT cells after saline or OVA challenge (left), and graphs represent the total number of NKT cells in each group. *** $P < 0.001$ (WT), *** $P < 0.001$ ($TIM1^{-/-}$, saline vs OVA group), not significant, (WT vs $TIM1^{-/-}$ mice after OVA challenge) (Two-tailed t-test). Data are representative of at least three independent experiments with three to five mice per each group.

Figure 3. Oxidative stress induces apoptosis in the airway

(A) WT mice were exposed to air (top) or ozone (bottom) as in Fig 1, and lungs were taken 24h after last ozone exposure for Immunofluorescence staining. PI (nuclear staining) (red), and TUNEL staining (green). Arrows indicate the apoptotic cells (yellow). Original magnification; X40. Data are representative of two independent experiments with three to four mice in each group.

(B) WT or *TIM-1^{-/-}* mice were exposed to air (top) or ozone (bottom) and apoptotic epithelial cells (CD45⁻ cells) were stained with AnnexinV-FITC (left panel). Graph

represents the percentage of AnnexinV positive cells in each group. *TIM1*^{-/-} mice were compared to WT mice, not significant (Two-tailed t-test)

(C) WT mice were exposed to air or ozone and anti-TIM1 (3D10) mAb was injected 24 hr before the first ozone exposure. Results represent the changes in lung resistance (R_L).

* $P < 0.05$, and ** $P < 0.01$ compared to air exposed group (Two-way ANOVA).

(D) Inflammatory cell numbers in BAL fluid of Fig. 1A (cells/lung). * $P < 0.05$, and ** $P < 0.01$ compared to air exposed mice (Two-tailed t-test). Mac, macrophage; Neu, neutrophils; Eos, eosinophils; Lymph, lymphocytes.

Figure 4. Apoptotic cells were critical for the development of ozone induced AHR

(A) Q-VD-OPH (2mg/kg) was injected 24 hr before every ozone exposure, and AHR measured 6 days after first exposure. * $P < 0.05$, compared to air exposed group (Two-way ANOVA).

(B) Inflammatory cell numbers in BAL fluid (cells/lung). ** $P < 0.01$, and *** $P < 0.001$, compared to air exposed group (Two-tailed t-test). Macs, macrophage; Neu, neutrophils; Eos, eosinophils; Lymph, lymphocytes.

(C) Lung sections from the mice shown in Fig 4A stained with hematoxylin/eosin (magnification; $\times 10$).

(D) Apoptotic airway epithelial cells (left), and the percentage of Annexin V positive cells in each group (right). Q-VD-OPH treated mice were compared to ozone exposed mice. * $P < 0.05$

(Two-tailed t-test). Data are representative of two independent experiments with three to five mice.

Figure 5. Apoptotic cells activate NKT cells through TIM-1

(A) Apoptosis of mouse airway epithelial cell line (MLE12) after H_2O_2 treatment was examined by AnnexinV and PI staining.

(B) WT or *TIM1*^{-/-} NKT cells were cultured with a suboptimal dose of α -GalCer (1ng/ml) (-); an optimal dose of α -GalCer (100 ng/ml); live MLE12 cells + suboptimal α -GalCer (1ng/ml); apoptotic MLE12 + suboptimal α -GalCer (1ng/ml) or apoptotic MLE12 + suboptimal α -GalCer (1ng/ml) + Annexin V (10ug/ml) for 48hr. Data are representative of at least three independent experiments.

(C) Human NKT cells were cultured with a suboptimal dose of α -GalCer (1ng/ml) (-); an optimal dose of α -GalCer (100 ng/ml); live A549 cells + suboptimal α -GalCer (1ng/ml); apoptotic A549+ suboptimal α -GalCer (1ng/ml) or apoptotic A549 + suboptimal α -GalCer (1ng/ml) + Annexin V (10ug/ml) for 48hr. NKT cells were then stained for intracellular cytokines. Data are representative of three independent experiments.

(D) The percentage of IL-4 producing NKT cells from all three experiments was calculated and graphed.

Figure 6. TIM-1 expression by NKT cells is required for ozone-induced AHR

(A) Schematic diagram of adoptive transfer of WT or *TIM-1*^{-/-} NKT cells.

(B) After repeated exposure to ozone, AHR was measured. ** $P < 0.01$ compared to ozone treated *Jα18*^{-/-} mice group (Two-way ANOVA).

(C) Inflammatory cells in BAL fluid (cells/lung). Macs, macrophage; Neu, neutrophils; Eos, eosinophils; Lymph, lymphocytes. * $P < 0.05$, and ** $P < 0.01$ compared to ozone treated *Jα18*^{-/-} mice group (Two-tailed t-test).

(D) WT or *TIM-1*^{-/-} mice were exposed to ozone as in Fig 1A. Lung cells were prepared and stimulated with PMA and ionomycin for 4hr. Cytokines produced from NKT cells were stained for intracellular IL-4, IL-13 and IL-17. * $P < 0.05$, *TIM-1*^{-/-} mice were compared to ozone treated WT mice (Two-tailed t-test). Data are representative of at least two independent experiments.

Figure 7. RSV infection induces AHR in a TIM-1 dependent manner

(A) WT or $TIM1^{-/-}$ mice were infected with RSV or sham, and lung resistance (R_L) measured on day 6 post infection. *** $P < 0.001$ compared to sham infected group (Two-way ANOVA).

(B) BAL fluid was taken from the mice in (A), and assessed for inflammatory cells (cells/lung). Macs, macrophage; Neu, neutrophils; Eos, eosinophils; Lymph, lymphocytes.

** $P < 0.01$, WT vs $TIM1^{-/-}$ mice infected with RSV (Two-tailed t-test).

(C) WT or *CD1d*^{-/-} mice were infected with RSV or sham infected, and lung resistance (*R_L*) was measured. *** *P*<0.001, WT vs *CD1d*^{-/-} mice infected with RSV (Two-way ANOVA).

(D) BAL fluids were assessed for inflammatory cells. Macs, macrophage; Neu, neutrophils; Eos, eosinophils; Lymph, lymphocytes. ** *P*<0.01, and ****P*<0.001, WT RSV infected group was compared to *CD1d*^{-/-} RSV-infected group (Two-tailed t-test).

(E) Tetramer⁺TCRβ⁺ NKT cells in BAL fluid after sham (left panels) or RSV (right panels) infection.

(F) Graph represents the % of annexinV positive cells. ****P*<0.001, compared to sham infected group (Two-tailed t-test).

(G) Q-VD-OPH (2mg/kg) was injected before RSV infection, and AHR measured. ****P*<0.001, compared to RSV infected group (Two-way ANOVA).

(H) BAL fluid was taken from the mice in (G), and assessed for inflammatory cells. Macs, macrophage; Neu, neutrophils; Eos, eosinophils; Lymph, lymphocytes. * *P*<0.05, and ****P*<0.01, RSV vs RSV + Q-VD-OPH group (Two-tailed t-test).

(I) WT or *TIM1*^{-/-} NKT cells (5×10^5 /mouse) were transferred into *Jα18*^{-/-} recipients before RSV infection, and AHR was measured. ***P*<0.01 *Jα18*^{-/-} mice with WT NKT cells were compared to *Jα18*^{-/-} mice with *TIM1*^{-/-} NKT cells (Two-way ANOVA).

(J) BAL fluids were assessed for inflammatory cells. Macs, macrophage; Neu, neutrophils; Eos, eosinophils; Lymph, lymphocytes. * *P*<0.05, *Jα18*^{-/-} mice transferred with WT NKT cells were compared to *Jα18*^{-/-} mice transferred with *TIM1*^{-/-} NKT cells (Two-tailed t-test). Data are representative of at least two independent experiments with four to six mice in each group.

Figure 8. Schematic diagram of the role of TIM-1 in non-allergic forms of asthma

In some forms of asthma (e.g., associated with air pollution and RSV infection), apoptotic airway epithelial cells activate NKT cells via TIM-1 recognition of dead cells, and these in turn activated NKT cells, which induce airway inflammation and AHR.