Published in final edited form as:

Tetrahedron Lett. 2013 August 7; 54(32): 4245–4246. doi:10.1016/j.tetlet.2013.05.136.

Improved synthesis of N1-substituted orotic acid derivatives

Nicholas A. Senger, Jeannette T. Bowler, Rene S. Mercado, Sidney Lin, and Weiming Wu*

Department of Chemistry and Biochemistry, San Francisco State University, San Francisco, CA 94132 USA

Abstract

An improved method for the synthesis of N1-substituted orotic acid derivatives is reported. The method involves sequential incorporation of nitrogen atoms to the pyrimidine structure from simple starting materials and thus allows the synthesis of N1-substituted orotic acid derivatives with single ¹⁵N label at either N-1 or N-3.

Keywords

Orotic acid; orotidine; 1-substituted orotic acid; 1-substituted uracil; dihydrouracil; 5-bromouracil

The decarboxylation of orotic acid and its analogues has been investigated as a model for the enzymatic decarboxylation catalyzed by orotidine-5'-monophosphate decarboxylase (ODCase).^{1–12} Part of the mechanistic investigation requires N1-substituted orotic acid derivatives (structure **1** in Scheme 1) as substrates. Direct alkylation of orotic acid, unfortunately, produces a mixture of disubstituted and N3-substituted orotic acid derivatives because N3 is the more reactive site.¹³ N1-Substituted orotic acid derivatives are commonly synthesized from N1-substituted uracil **2** via 5-bromouracil **3** and 6-cyanouracil **4** as shown in Scheme 1.^{11,14–16} In this Letter, we report a simplified synthetic method for such orotic acid derivatives from readily available 5,6-dihydrouracil derivatives.

A few methods for the synthesis of N1-substituted uracil derivatives have been reported.^{17–19} One of the methods involves sequential integration of the two nitrogen atoms from simple starting materials (as shown in Scheme 2) and thus will allow incorporation of ¹⁵N at one position only.¹⁷ The reported synthesis starts with the conjugated addition of a primary amine to acrylate that yields adduct **5**.^{17,20} The amino group is converted to the urea group via reaction with cyanate. Cyclization as a result of the reaction of the urea group with the ester group gives the 5,6-dihydrouracil **6**. Bromination of **6** produces **7**, which undergoes thermal elimination to yield uracil **2**.

In the process of synthesizing N1-substituted uracil derivatives following this reported synthetic route, we observed that the bromination of 5,6-dihydrouracil **6** yielded desired product **7** mixed with dibrominated product **8** as shown in Scheme 3. Further experiments of purified dibromide **8** illustrated that it underwent elimination upon heating to produce 5-bromouracil **3**, analogous to the conversion of bromide **7** to **2**. We have thus thought to

© 2013 Elsevier Ltd. All rights reserved.

*Corresponding author. Tel.: 1-415-338-1436; fax: 1-415-338-2384; wuw@sfsu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

shorten the synthetic route employing 5,5-dibromo-5,6-dihydrouacil **8** as a synthetic intermediate.

The dibromination of **6** to **8** was effected by using more than two equivalents of bromine. Interestingly, it was observed that a fraction of dibromide **8** underwent elimination spontaneously to produce bromouracil **3** under the reaction conditions (acetic acid, 100 °C). We thought to further simplify the synthetic route by combining the bromination and elimination steps. The bromination reaction was thus carried out in boiling acetic acid. Under the new reaction condition, dihydrouacil **6** was converted to 5-bromouracil **3** in one step, in contrast to the three-step sequence in the original synthesis. The yields for the onestep reactions were excellent for various primary and secondary substituting groups (tertiary groups were found to be labile under the highly acidic reaction conditions as expected) tested as shown in Table 1. In the case when cyclohexyl was the substituting group, the yields for the two synthetic routes were directly compared. The direct conversion of **6a** to **3a** (R = cyclohexyl) was found to be quantitative (the yield is 89% if purified by recrystallization), whereas the reported yields for the three synthetic steps were 69% (**6a** to **7a**), 85% (**7a** to **2a**), and 75% (**2a** to **3a**), respectively, giving an overall yield of 44% from **6a** to **3a**.^{14,17} Therefore, the modified one-step procedure reported here represents a significantly simplified and improved method to synthesize N1-substituted orotic acid derivatives.

It should be pointed out again that this synthetic route allows the incorporation of a single ¹⁵N label at either N1 or N3. Such labeled compounds may aid in the mechanistic investigation of the model decarboxylation of orotic acid derivatives or the enzymatic decarboxylation catalyzed by ODCase.

Experimental Details

All reagents were obtained from commercial sources and used without further purification. Only the experimental procedures for the conversion of dihydrouacil **6** to 5-bromouracil **3** are described here using 1-cyclohexyl-5,6-dihydrouacil as an example. The products were purified through recrystallization or column chromatography. Procedures for other synthetic steps were described in literature and were employed without modification.

Typical experimental procedure

1-Cyclohexyl-5,6-dihydrouacil (1.00 g, 5.1 mmol) was dissolved in 12 mL acetic acid in a round-bottomed flask. A solution of bromine (3.24 g, 20.2 mmol) in 10 mL acetic acid was added to the reaction flask and the reaction mixture was refluxed. The reaction usually took five to six hours and was followed by thin layer chromatography (TLC). Upon completion of the reaction, the reaction mixture was evaporated to dryness under reduced pressure. The solid residue was purified by either column chromatography (10% ethyl acetate/methylene chloride as solvent) or recrystallization from ethanol to produce 5-bromo-1-cyclohexyluracil as colorless flakes; mp 220–222 °C, lit. mp 225 °C.¹⁷

Acknowledgments

This investigation was supported by the National Institutes of Health, Grant SC1 GM095419. N.A.S. and J.T.B. were supported by the Beckman Scholarship. R.S.M. and S.L. were summer students supported by the BRIDGE Program (funded by NIH) and Project SEED (the American Chemical Society), respectively. The NMR facility was funded by the National Science Foundation (DUE-9451624 and DBI 0521342). We thank Professor Ihsan Erden (SFSU) for helpful discussions.

References

1. Beak P, Siegel B. *J Am Chem Soc.* 1976; 98:3601–3606. [PubMed: 1270703]
2. Radzicka A, Wolfenden R. *Science.* 1995; 267:90–93. [PubMed: 7809611]
3. Lee JK, Houk KN. *Science.* 1997; 276:942–945. [PubMed: 9139656]
4. Kakanishi MP, Wu W. *Tetrahedron Lett.* 1998; 39:6271–6272.
5. Feng WY, Austin TJ, Chew F, Gronert S, Wu W. *Biochemistry.* 2000; 39:1778–1783. [PubMed: 10677227]
6. Singleton DA, Merrigan SR, Kim BJ, Beak P, Phillips LM, Lee JK. *J Am Chem Soc.* 2000; 122:3296–3300.
7. Wong FM, Wu W. *Bioorg Chem.* 2006; 34:99–104. [PubMed: 16499947]
8. Yeoh FY, Cuasito RR, Capule CC, Wong FM, Wu W. *Biorg Chem.* 2007; 35:338–343.
9. Wong FW, Capule CC, Wu W. *Org Lett.* 2006; 8:6019–6022. [PubMed: 17165919]
10. Wong FM, Capule CC, Chen DX, Gronert S, Wu W. *Org Lett.* 2008; 10:2757–2760. [PubMed: 18537244]
11. Lewis CA Jr, Wolfenden R. *Biochemistry.* 2009; 48:8738–8745. [PubMed: 19678695]
12. Senger NA, Bliss CE, Keeffe JR, Gronert S, Wu W. *Tetrahedron.* 2013; 69:5287–5292.
13. Curran WV, Angier RB. *J Org Chem.* 1966; 31:201–205.
14. Landesman, PW. PhD Dissertation. State University of New York; Buffalo: 1982.
15. Inoue H, Ueda T. *Chem Pharm Bull.* 1971; 19:1743–1744.
16. Inoue H, Ueda T. *Chem Pharm Bull.* 1978; 26:2657–2663.
17. Okano T, Goya S, Takahashi T. *Yakugaku Zasshi.* 1968; 88:1112–1117. [PubMed: 5751278]
18. Rejman D, Kova ková S, Pohl R, Dra ínský M, Fiedler P, Rosenberg I. *Tetrahedron.* 2009; 65:8513–8523.
19. Winckelmann I, Larsen EH. *Synthesis.* 1986:1041–1044.
20. Padwa A, Hasegawa T, Liu B, Zhang Z. *J Org Chem.* 2000; 65:7124–7133. [PubMed: 11031039]
21. ¹H NMR data of **6** and **3** in CDCl₃: 1-*n*-Butyl-5,6-dihydrouacil **6b**. δ 7.7 (bs, 1H, NH); 3.4 (t, 4H, CH₂NCH₂); 2.7 (t, 2H, CH₂CO); 1.6 (quin, 2H, NCH₂CH₂), 1.3 (sext, 2H, CH₂CH₃); 0.9 (t, 3H, CH₃). 1-*n*-Propyl-5,6-dihydrouacil **6c**. δ 7.5 (bs, 1H, NH); δ 3.4 (m, 4H, CH₂NCH₂); 2.7 (t, 2H, CH₂CO); 1.6 (sext, 2H, CH₂CH₃); 0.9 (t, 3H, CH₃). 1-Isobutyl-5,6-dihydrouacil **6d**. δ 7.6 (s, 1H, NH); 3.4 (t, 2H, NCH₂ ring); 3.2 (d, 2H, NCH₂ side chain); 2.7 (t, 2H, CH₂CO); 2.0 (sept, 1H, CH); 0.9 (6H, CH₃ x 2). 1-*sec*-Butyl-5,6-dihydrouacil **6e**. δ 8.1 (s, 1H, NH); 4.4 (1H, sext, NCH); 3.3 (2H, m, NCH₂); 2.6 (t, 2H, CH₂CO); 1.5 (quint, 2H, CH₃CH₂); 1.2 (d, 3H, CHCH₃); 0.9 (t, 3H, CH₂CH₃). 1-*n*-Butyl-5-bromouracil **3b**. δ 8.7 (bs, 1H, NH); 7.5 (s, 1H, CH); 3.8 (t, 2H, NCH₂); 1.7 (quin, 2H, NCH₂CH₂), 1.4 (sext, 2H, CH₂CH₃); 1.0 (t, 3H, CH₃). 1-*n*-Propyl-5-bromouracil **3c**. δ 8.5 (bs, 1H, NH); 7.5 (s, 1H, CH); 3.7 (t, 2H, NCH₂); 1.7 (sext, 2H, CH₂CH₃); 1.0 (t, 3H, CH₃). 1-Isobutyl-5-bromodouracil **3d**. δ 8.5 (bs, 1H, NH); 7.5 (s, 1H, H-6); 3.6 (d, 2H, NCH₂); 2.1 (sept, 1H, CH); 1.0 (6H, CH₃ x 2). 1-*sec*-Butyl-5-bromodouracil **3e**. δ 9.0 (bs, 1H, NH); 7.5 (s, 1H, H-6); 4.6 (1H, sext, NCH); 1.7 (quint, 2H, CH₃CH₂); 1.3 (d, 3H, CHCH₃); 1.0 (t, 3H, CH₂CH₃).

Scheme 1.

Scheme 2.

Scheme 3.

Table 1Yields for the one-step conversion of 5,6-dihydrouracil **6** to 5-bromouracil **3**²¹

Entry	1-Substituted 5,6-dihydrouracils (6)	Yield (isolated) (%)
1	6a , R = cyclohexyl	100 (89) ^a
2	6b , R = <i>n</i> -butyl	85
3	6c , R = <i>n</i> -propyl	88
4	6d , R = isobutyl	86
5	6e , R = <i>sec</i> -butyl	100

^aYield in parenthesis when purified by recrystallization.