CASE REPORT

Familial ectodermal dysplasia: a peers' agony

Karthik Hegde,¹ Roopashri Rajesh Kashyap,² Gopakumar Nair,³ Preeti P Nair¹

¹Department of Oral Medicine and Radiology, People's College of Dental Sciences and Research Centre, Bhopal, Madhya Pradesh, India

²Department of Oral Medicine and Radiology, A.J. Institute of Dental Sciences, Mangalore, Karnataka, India

³Department of Oral Medicine and Radiology, New Horizon Dental College and Research Institute, Bilaspur, Chhattisgarh, India

Correspondence to
Professor Preeti P Nair,
shajihoss@gmail.com

SUMMARY

Ectodermal dysplasias include a various group of inherited disorders which share primary defect in the development of two or more tissues of embryonic ectodermal origin. Though there are many subtypes, ectodermal dysplasias are mainly hidrotic ectodermal dysplasia and hypohidrotic ectodermal dysplasia, among which the most common variety is X linked hypohidrotic ectodermal dysplasia. We report a rare case of X linked hypohidrotic ectodermal dysplasia occurring in a family with various skin, hair and oral abnormalities.

BACKGROUND

Ectodermal dysplasias are a large group of syndromes that are heterogeneous under clinical and genetic aspects, and are characterised by anomalies in the structures of ectodermal origin.¹ There are more than 192 distinct disorders being described, among them the most common ectodermal dysplasias are X linked recessive hypohidrotic and hidrotic type.¹⁻² They are rare and incidence has been estimated at 1 in 10 000 and 1 in 100 000 births.³ The disorder is determined by the X linked recessive gene, so the disease is more observed in men than in women.¹

The phenotype includes sparse scalp hair, absence of body hair, deficiency of the sweat glands and anodontia or oligodontia with conical teeth.¹⁻⁴ There is often a distinctive facies, with prominence of the forehead, depressed nasal bridge, prominent lips, periorbital wrinkling and pigmentation. Often subcutaneous fat, mucous glands in the respiratory tree and the gastrointestinal tract are diminished.⁴

Thurman (1848) first reported two male first cousins and their maternal grandmother with a hereditary syndrome associated with sparse hair, missing teeth and dry skin¹ and later in the 19th century by Darwin. Thadani in 1921 assigned it to the X chromosome, later reported that the carrier females could manifest signs of the condition.⁴ The full significance of this was not appreciated until Mary Lyon hypothesised the random inactivation of one X chromosome in each female cell early during embryogenesis.⁴

Here we report a case of X linked hypohidrotic ectodermal dysplasia (HED) with complete expression in a male patient and partial expression in a male and two female patients (of the same family) in same generations of a family.

CASE PRESENTATION**Case 1**

A 24-year-old man reported with multiple unerupted teeth. He gave a history of delay in the eruption of the maxillary and mandibular anterior teeth

which were few in number. His deciduous dentition was incomplete with only few primary teeth and later the permanent teeth which erupted were altered in size and shape.

Medical history and general physical examination revealed that palms were dry, sparse hair on the scalp, body, hands and legs and absence of eyebrows, beard and moustache. Eye lashes were sparse. The patient had an average intelligence level. Skin was dry with generalised xerosis and rise of temperature; however, the nails were normal (figure 1A). The patient also had recurrent fever, ear infections and pharyngitis for which the patient had consulted an ear nose and throat surgeon.

Family history revealed that the patient was the eldest child of the two children. There was no parental history of consanguinity. The younger child did not suffer from similar defect. There was a presence of similar deformity among his three maternal first cousins and grandmother (figure 1B).

Extraoral examination revealed a convex facial profile, loss of eyebrows and scanty eyelashes, broad nose, malar hypoplasia, protuberant lips and lower part of face being narrow with prominent chin and height of the lower third face was decreased. Periorbital skin showed wrinkling and hyper pigmentation. Facial skin over the cheeks and forehead region showed chronic eczematous dermatitis (figure 1C).

Intraoral soft tissue examination revealed that the freni in maxillary and mandibular anterior region are attached to the redundant alveolar ridges. The residual ridges were thin and ill developed, covered by a thin mucosa (figure 1D,E).

Hard tissue examination revealed that 12, 14, 16, 17, 21, 22, 25, 27, 31, 32, 34, 35, 36, 37, 41, 42, 44, 45 and 47 were missing. The crown of 11, 13, 23, 24, 33 and 34 were conical in shape and 24 was microdont and over retained 61 and root stumps of 55 and 65 were present (figure 1F).

Case 2, 3 and 4

The three maternal first cousins were interviewed and examined at a later date. The grandmother could not be interviewed as she had deceased.

All the three siblings had multiple unerupted maxillary and mandibular teeth, raised body temperature, intolerance to heat and less sweat production.

Medical history revealed that skin was dry, sparse hair in the body, hands and legs and sparse eyelashes. The scalp hair and nails were normal. In all the three siblings, the general examination, extraoral, intraoral soft tissue and hard tissue examination findings were similar (table 1 and figures 2–5).

To cite: Hegde K, Kashyap RR, Nair G, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-200175

Figure 1 (A) Sparse body hair with generalised xerosis. (B) Pedigree chart representing the family. (C) Loss of eyebrows, eyelashes, protuberant lips, wrinkling and hyper pigmentation of skin and decreased height of lower third of face. (D) Frenum in maxillary anterior region attached to the alveolar ridges. (E) Frenum in mandibular anterior region attached to the alveolar ridges. (F) Orthopantomograph showing conical-shaped crown of 11, 13 23, 24, 33 and 34, over retained 61, microdont 24 and multiple missing teeth.

INVESTIGATIONS

Investigation included radiographs and dermatological consultation in all the cases. Radiographic findings of cases 2, 3 and 4 are described in table 2. Dermatology consultation included starch-iodine test for sweat pore count on the palmar surface which was positive among all the siblings. The radiographic findings in cases 2, 3 and 4 were correlated and confirmed with clinical findings as described in table 1. Based on these findings a final diagnosis of X linked HED (Christ-Siemens-Touraine syndrome) was arrived at.

DIFFERENTIAL DIAGNOSIS

The differential diagnosis familial simple anhidrosis, Werner syndrome and Rothmund Thompson syndrome, were considered.

TREATMENT

For case 1, the root stumps of 55 and 65 and over retained 61 were extracted. The missing teeth were replaced with a removable partial denture for cases 2, 3 and 4.

OUTCOME AND FOLLOW-UP

Healing of the extraction site was uneventful and subsequently managed by partial denture prosthesis for case 1.

DISCUSSION

Anodontia is a rare condition, which represents the congenital absence of all teeth in the primary and permanent dentition. A senile facial appearance due to lack of all teeth and under-development of the alveolar ridges is the characteristic feature

Table 1 General examination, extraoral and intraoral findings of cases 2, 3 and 4

Case	Age/ sex	General examination	Extraoral examination	Intraoral soft tissue examination	Hard tissue examination
Case 2	21/m	Dry skin, eye lashes, hair hands and legs –sparse	Malar hypoplasia, decreased height of the face, protuberant lips, prominent chin, wrinkled periorbital skin (figure 2A)	Moist mucosa, pigmented gingival	Permanent teeth were microdont. Missing teeth: 13, 14, 15, 16, 17, 22, 23, 24, 25, 26, 27, 31, 34, 35, 37, 41, 44, 45 and 47 (figure 2B)
Case 3	17/f	Dry skin, sparse eyebrows and eye lashes	Protuberant lips (figure 3A)	Thin lower residual ridge, high frenal attachment	Narrow incisors conical-shaped canines Spacing between teeth Missing teeth: 12, 14, 22, 24, 31, 32, 33, 34, 41, 42, 43 and 44 (figure 3B,C)
Case 4	14/f	Dry skin, sparse eyebrows and eye lashes	Protuberant lips (figure 4A)	Thin lower residual ridge, high frenal attachment	Reduced mesiodistal width of the incisors conical-shaped 43. Spacing between teeth Missing teeth: 12, 14, 22, 31, 32 and 41 (figure 4B,C)

Figure 2 (A) Extraoral photograph showing malar hypoplasia, protuberant lips and wrinkled periorbital skin. (B) Intraoral image showing retained deciduous teeth, multiple microdont permanent and missing teeth.

of ectodermal dysplasia, which may affect normal social and psychological development in young patients.⁵

Ectodermal dysplasia (ED) is an inherited disorder in which there is a failure in development of two or more ectodermally derived structures, or are abnormal in development. The more common variety of ED is hypohidrotic type. It is inherited as X linked recessive trait, hence, more common among men. Each person with ED has different combination of ectodermal defects. Pure EDs are manifested by defects in ectodermal structures alone, while ED syndromes are characterised by the combination of ectodermal defects in association with other anomalies.⁶ When at least two types of abnormal ectodermal features occur, the patient is diagnosed with ED syndrome.⁷

Freire-Maia redefined this as a developmental defect which, at embryonic level affects the ectoderm and therefore the tissues and structures derived from it. Thus it affects the development

of keratinocytes and cause aberrations in the hair, sebaceous glands, eccrine and apocrine glands, nails, teeth, lenses and conjunctiva of the eyes, anterior pituitary gland, nipples and the ears.²

Christ-Siemens-Touraine syndrome, with X linked recessive inheritance, is the most frequently reported manifestation of ectodermal dysplasia. Depending on the severity of clinical manifestations, Christ-Siemens-Touraine syndrome can be classified as either hypohidrotic or anhidrotic ectodermal dysplasia.⁷ It has full expression only in men. Female carriers show little or no signs of the condition.^{2, 8, 9} The incidence of this condition is 1:100 000, with a mortality rate of 28% in men up to 3 years of age.⁷

Mono *et al* suggested that heterozygous women of unaffected mother having variable clinical features of ED occur because of mosaic of functionally normal and abnormal cells and random X-inactivation (Lyon hypothesis).¹⁰ The diagnosis of patients with ED is based fundamentally on the clinical history which includes ungula dystrophy, hypotrichosis, anodontia, oligodontia, hypodontia; malformed teeth, delayed eruption and dental impaction.¹¹ Oligodontia in ED is most frequently associated with dominant type, hereditary factors, polygenic factors or X linked.¹²

The sex-linked pattern of inheritance for hereditary ED was first described by Darwin in 1844.¹ The gene locus is now known to be on the proximal part of the X chromosome long arm.¹³ Genetic studies of more than 300 cases have revealed X linked mode of inheritance with its gene locus being Xq11–21.1, the gene is carried by women but manifested in the men. However, there are reports of multiple siblings being affected and females suffering with this condition.² Currently, HED is related with a mutation of the protein ectodysplasin-A, related with the EDA gene in the q12–q13 locus of the X chromosome.¹¹

Figure 3 (A) Extraoral photograph showing sparse eyebrows and eye lashes and dry skin. (B) Intraoral image showing thin lower residual ridge and high frenal attachment. (C) Intraoral image showing narrow incisors-shaped and conical-shaped canines.

Figure 4 (A) Extraoral photograph showing malar hypoplasia, protuberant lips and wrinkled periorbital skin. (B) Intraoral image showing high-frenal attachment and multiple missing permanent teeth. (C) Intraoral image showing reduced mesiodistal width of the incisors-shaped and conical-shaped 43.

Figure 5 (A) Orthopantomograph (OPG) showing microdont 11, 12, 31, 32, 41 and 42, over retained deciduous teeth and multiple missing permanent teeth. (B) OPG showing multiple missing permanent teeth, reduced mesiodistal width of 11 and 21, conical-shaped 13 and 23 and under developed mandibular anterior alveolar ridge. (C) Orthopantomograph showing multiple missing permanent teeth, reduced mesiodistal width of 11 and 21 and conical-shaped 42 and 43.

Most of the EDs manifest variable defects in the morphogenesis of ectodermal structures such as the hair, skin, nails, teeth and sweat glands which produce numerous general findings and symptoms, thus the clinical features associated with ED may vary in its presentation.⁸ Characteristic skull deformities include square aspect of the forehead, frontal bossing, prominent supra-orbital ridges, saddle nose, the alae nasi appears hypoplastic, the midface is depressed and hypoplastic, malar hypoplasia and appear flat and depressed with dished-in appearance, the chin may be pointed and the lips are thick and everted, cleft lip or palate. The patient may resemble an edentulous old person. The craniofacial deviations increase with advancing age, in non-

treated patients, with a tendency towards a class III pattern, decreased lower facial height, the maxilla is more retruded as the mandible and the mouth width is smaller than normal.⁸ In our case most of these characteristic features were present.

Most female carriers of HED are recognised by dental anomalies, and in most of them where the teeth are normal, may be recognised by their abnormal pattern of sweating.¹³ The dental anomalies may be number anomalies like anodontia or severe hypodontia, shape anomalies like peg-shaped conical incisors and canines in deciduous as well as permanent dentitions, talon cusp, taurodontia, size anomalies like microdontia in affected males as well as carrier females, structural anomalies like hypoplasia.⁸ In our case all the participants had abnormal pattern of sweating and dental anomalies.

Dystrophic, hypoplastic or absent nails, and sparse hair is also present in ED. Some individuals may lack eyebrows or eyelashes, and present hyperkeratosis of the knees, elbows, knuckles and under the free border of the nails.¹¹ which was seen in our cases.

Nakata *et al* determined tooth size in 49 participants in 15 families with X linked hereditary ED. Affected males and carrier females showed consistently smaller mesiodistal tooth crown diameters than the unaffected individuals.⁹ In our case all the siblings showed this presentation.

Airenne studied 11 men from nine families together with their relatives. He confirmed the reduction in mesiodistal diameter and showed that this was also true for the buccolingual dimension in affected males as well as heterozygous females. Radiographs were used only to assess the degree of hypodontia.⁹ These findings were also reflected in our cases.

According to Crawford *et al*, the first molars, upper central incisors and upper and lower canines were the most 'stable' tooth types, and they were most often present.⁹ Our cases represent the similar pattern of hypodontia.

In studies that utilised serial cephalometric measurements have reported that in general facial growth proportion and pattern of jaw growth appear to be normal in these children despite the absence of tooth development. Therefore, the protuberant lips observed in these children may be attributed to the reduction in the height of the alveolar process.¹⁴

The prognosis in ED is usually good, except in cases with hypo or anhidrosis due fundamentally to respiratory infections resulting from the absence of mucosal glands in the respiratory system where mortality rates can reach 30% in first infancy.⁷

CONCLUSION

The clinical manifestations of ED cause considerable social problems in individuals affected by the condition. There is no general cure for ED, but preventive treatment consists of genetic counselling and prenatal analysis. Many treatments addressing the symptoms are available; however, general treatment may require the coordinated efforts of a multidisciplinary team of specialists who need to systematically and comprehensively plan an individualised treatment. The general treatment is symptomatic and depends on which ectodermal structures are involved. Medical care also improves aesthetics and functions. The dental and sweating signs are useful in early identification of individuals with this condition and female carriers in the family. Hence relatives of carrier female should be counselled so as to understand the problems in the subsequent births occurring in the family, which may minimise the complications in the next generation. When an affected infant is expected it may be that his early management is improved so that he is less likely to succumb to illness or severe fever.

Table 2 Radiographic findings of cases 2, 3 and 4

Case 2	Case 3	Case 4
Permanent teeth were smaller in size Missing 13, 14, 15, 16, 17, 22, 23, 24, 25, 26, 27, 31, 34, 35, 37, 41, 44, 45 and 47 Microdont 11, 12, 31, 32, 41 and 42 Over retained deciduous 53, 55, 63, 65, 75, 84 and 85 Over retained deciduous roots showed no resorption (figure 5A)	Missing 12, 14, 22, 24, 31, 32, 33, 34, 41, 42, 43 and 44 Reduced mesio-distal width of 11 and 21 at the incisal third of the crown-shaped and conical-shaped 13 and 23 Under developed mandibular anterior alveolar ridge (figure 5B)	Missing 12, 14, 22, 31, 32 and 41 Reduced mesio-distal width of 11 and 21 at the incisal third of the crown Conical-shaped 42 and 43 (figure 5C)

Dental treatment is generally essential and requires early and continuous dental intervention during many years, multiple phases of treatment and the involvement of a multidisciplinary dental team.

Learning points

- ▶ Ectodermal dysplasias are associated with disturbances in the development of two or more tissues of embryonic ectodermal derivation.
- ▶ The dental and hypohidrosis are useful in early recognition of individuals with this state and female carriers.
- ▶ Counselling may prove to be cautious in family planning and help reduce the incidence of these kinds of cases.
- ▶ Early oral intervention and rehabilitation in such patients help in normal social and psychological development.

Contributors KH performed the follow-up of cases; RRK recorded patients' details; GN was involved in manuscript writing and PPN edited the manuscript.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Ramesh K, Vinola D, John JB. Hypohidrotic ectodermal dysplasia—diagnostic aids and a report of 5 cases. *J Indian Soc Pedod Prevent Dent* 2010;28:47–54.
- 2 Varghese G, Sathyan P. Hypohidrotic ectodermal dysplasia—a case study. *Oral Maxillofac Pathol J* 2011;2:123–6.
- 3 Gupta S, Tyagi P. Prosthodontic management of anhidrotic ectodermal dysplasia. *Indian J Dent Res* 2011;22:348–51.
- 4 Clarke A. Hypohidrotic ectodermal dysplasia. *J of Med Gen* 1987;24:659–63.
- 5 Ohno K, Ohmori I. Anodontia with hypohidrotic ectodermal dysplasia in a young female: a case report. *Pediatr Dent* 2000;22:49–52.
- 6 Kamran T, Asif T. A 4-year-old male child with hypohidrotic ectodermal dysplasia. *J Pakistan Assoc of Dermatol* 2011;21:69–70.
- 7 Bania M, Tezkirecioglu AM, Akalc N, *et al.* Ectodermal dysplasia with anodontia: a report of two cases. *Eur J Dent* 2010;4:215–22.
- 8 Răducanu AM, Păuna M, Feraru IV. A simple prosthetic restorative solution of a single peg-shaped upper central primary incisor in a case of ectodermal dysplasia. *Rom J Morpholand Embryol* 2010;51:371–4.
- 9 Crawford PJM, Aldred MJ, Clarke A. Clinical and radiographic dental findings in X linked hypohidrotic ectodermal dysplasia. *Med Genet* 1991;28:181–5.
- 10 Ghosh S, Das MK. Hypohidrotic ectodermal dysplasia of sisters in a family. *Indian Pediatr* 2004;41:1176–7.
- 11 Pipa-Vallejo A, López-Arranz-Monje E, González -García M, *et al.* Treatment with removable prosthesis in hypohidrotic ectodermal dysplasia. A clinical case. *Med Oral Patol Oral Cir Bucal* 2008;13:E119–23.
- 12 Tsai PF, Chiou HR, Tseng CC. Oligodontia—a case report. *Quintessence Int* 1998;29:191–3.
- 13 Clarke A, Phillips DIM, Brown TR, *et al.* Clinical aspects of X-linked hypohidrotic ectodermal dysplasia. *Arch Dis Child* 1987;62:989–96.
- 14 Mohammed K, Tony E, Feteih RM, *et al.* Hereditary hypohidrotic ectodermal dysplasia with anodontia: a case report. *Saudi Dent J* 1994;6:31–4.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow