Published in final edited form as:

Curr Opin Immunol. 2009 June ; 21(3): 258–265. doi:10.1016/j.coi.2009.05.006.

The microanatomy of B cell activation

Tri Giang Phan¹, Elizabeth E. Gray², and Jason G. Cyster²

¹Garvan Institute of Medical Research, 384 Victoria St Darlinghurst, Sydney NSW 2010, Australia

²Howard Hughes Medical Institute and Department of Microbiology and Immunology, University of California at San Francisco, 513 Parnassus Ave San Francisco CA 94143, United States of America

Abstract

The logistic problem of B cell antigen encounter in the lymph node has recently been studied by dynamic imaging using two-photon microscopy. These studies combined with early studies of antigen transport have yielded a more complete picture of the orchestration of B cell activation in vivo. Here we summarize the recent advances and focus on the specialized macrophages that are critical to this process and the role of B cells themselves as antigen transporting cells.

Introduction

The immune system comprises a complex system of molecular, cellular and structural components that protects the host against infectious agents and transformed cancer cells by means of exclusion (via mucocutaneous barriers), eradication (via innate and adaptive effector mechanisms) and containment (via granuloma formation). Secondary lymphoid organs (SLOs) such as lymph nodes (LNs) and the spleen are critical for efficient orchestration of the response as they provide a supporting framework within which antigens and rare antigen-specific T and B cells can be efficiently brought together to initiate the adaptive response. As such, understanding the dynamics of cellular migration and antigen trafficking in SLOs and the requirements for antigen encounter in the LN are important steps towards developing a global picture of adaptive immunity. In this review we will discuss recent advances in our understanding of B cell activation with the advent of intravital two-photon microscopy, a technology that has allowed direct visualization of the spatial and temporal organization of the B cell response in intact LNs in real-time. We will examine the logistic problems of bringing antigen into contact with rare antigen-specific B cells and how specialized macrophages that line the subcapsular sinus (SCS) sample the afferent lymph for antigen to present to B cells. We will discuss some biological differences between these SCS macrophages and those that line the medullary sinuses. Finally, we will review the data for B cells as antigen transporting cells and the potential significance of this novel B cell function in FDC antigen deposition and the antibody response.

© 2009 Elsevier Ltd. All rights reserved.

Correspondence should be addressed to: Tri Giang Phan (t.phan@garvan.org.au) and Jason G. Cyster (jason.cyster@ucsf.edu).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Barriers to B cell antigen encounter

The primary role of the LN is to filter the lymph and trap any pathogens that have breached the initial mucocutaneous barriers. The pattern of lymph flow has been extensively studied in the past in small and large animals using colloidal particles, radiotracers and electron-dense particles (reviewed in [1]). Afferent lymphatic vessels drain lymph into the SCS, a sinus that forms an anatomical and functional barrier to the free diffusion of lymph-borne particles into the parenchyma [2-4]. Lymph then slowly percolates through cortical and medullary sinuses that, like the SCS, are lined by LyVE-1-expressing lymphatic vascular endothelial cells [5]. The medullary sinuses comprise a dense labyrinth of branching vessels and are richly populated with macrophages [6] where the bulk of lymph-borne soluble and particulate antigen is trapped and catabolized [3,7,8]. SCS macrophages and the majority of medullary macrophages express the sialic acid-binding immunoglobulin-like C-type lectin sialoadhesin (CD169) recognised by the MOMA-1 mAb [9], a property they share with the metallophilic macrophages that line the marginal sinus of the spleen [10]. Despite some confusion in the literature, CD169⁺ macrophages lining the SCS can clearly be distinguished from those lining the medullary sinuses by the expression of F4/80 [11,12], SIGNR1 [13,14] and MARCO [14,15] by medullary but not SCS macrophages.

The secondary role of the LN is to stage the adaptive immune response and this creates a paradox given that T and B cells are located in distinct anatomical compartments beneath the impermeable barrier formed by the floor of the SCS. For T cells this is partly overcome by the conduit system comprising an interconnected network of collagen fibres ensheathed by fibroblastic reticular cells (FRCs) that express ERTR-7 [16]. Many of these conduits begin in the interfollicular region below the SCS and traverse the T cell zone in the parenchyma to anchor near high endothelial venules (HEVs) [17] (Fig. 1). They provide a mechanism for rapid delivery of small molecules <70 kDa with a hydrodynamic radius <5.5 nm to the T zone [18] where resident dendritic cells (DCs) associated with conduits may sample antigen for presentation to T cells [19,20]. In addition, antigen may be transported from the periphery by DCs travelling in the afferent lymph that must also cross the SCS to reach the T zone. In this case, the network of stromal cells with underlying conduits provide a scaffold for T cells during their random migration as they survey the parenchyma for processed antigen presented on the surface of these DCs [21]•.

Follicular stromal cells that express CD157 recognised by the BP-3 mAb [22,23] form the stromal cell network and likely substratum for B cell migration in the follicle. Although some studies have emphasized that follicles have few conduits [1] others have noted that the conduits present have similar properties to those in the T zone [24-27]. In addition, some EM studies have detected occasional gaps 0.1-1 µm wide in the floor of the SCS through which small particulate antigen may freely diffuse [28,29], but these may be transient migration pores created by the transendothelial migration of lymphocytes [29,30]. Moreover, there is a large discrepancy between their diameter and the physical limits imposed by the barrier to follicular diffusion. An intriguing recent study reported that a bolus of fluorescently-labelled hen egg lysozyme (HEL) injected subcutaneously into the ear was detectable in the follicle by epifluorescence microscopy if the cervical LNs was carefully prepared to avoid washing away the small (14.3 kDa) highly cationic protein [31]•. This study raised the possibility that small antigens may rapidly diffuse across the floor of the SCS into the follicle. In more recent work, Roozendaal et al. tracked the distribution of fluorescently labelled turkey egg lysozyme (TEL) in real-time by two-photon microscopy and found that it first travelled into the follicle via conduits before reaching follicular B cells [27]••. Further comparisons and perturbations of these systems are needed before we can know the exact extent to which small antigens access the follicle via SCS pores versus conduits. The fraction of pathogen-derived antigens that would have the size and

physiochemical properties to access follicles via these routes is not clear but seems likely to be small. Moreover, the inability of antigens opsonized by antibody (>150kD) or C3b (~180kD) to enter conduits suggests that the antigens travelling through these structures will only be transiently available for encounter with rare cognate B cell. So how do most antigen types, including viral particles and opsonized antigens, get from outside in? Here a prominent role for the specialized SCS macrophages has emerged.

Locations of B cell encounter with particulate antigen

Mature recirculating B cells enter the LN via HEVs situated in the outer T zone and interfollicular regions and migrate to the follicle where they are retained for up to 24 hours before they exit the LN and re-enter the circulation [32,33]. This “inside out” B cell migration pattern runs counter-current to the “outside in” trafficking of antigen and provides multiple opportunities for B cell antigen encounter (Fig. 1). Indeed, B cell activation in the vicinity of HEVs has been observed in a study which found CD11c⁺ splenic DCs pulsed with very high concentrations of HEL migrated to the popliteal LN following footpad injection and were able to activate newly arrived B cells [34]•. This study is remarkable for its elegant use of intravital two-photon microscopy to show a recirculating B cell entering the T zone via the HEV fluxing calcium upon immediate contact with an antigen-pulsed DC. Antigen-pulsed DCs have occasionally been shown to induce humoral immune responses after adoptive transfer [35,36], and it would appear that certain DCs do not degrade all the antigen they internalise but recycle some to the cell surface. In the case of immune complexes (ICs), this can occur in an FcγRIIB-dependent manner [37]. Notwithstanding these reports, the chances of antigen encounter are probably highest inside the follicle itself where B cells spend most of their transit time [38]. Consistent with this, early antigen capture studies showed that while most of the antigen is trapped and degraded in the medullary region, small amounts did manage to infiltrate the follicle and can be retained there on FDCs for prolonged periods of time [7,8,39]. This retention is most prominent when the antigen can form ICs [3,40] and may last up to 7 weeks [41]. Thus, a common view held that B cells migrate to follicles to “survey” for antigen displayed on the surface of FDCs.

Early antigen capture studies also noted that antigen-antibody complexes were initially trapped in the floor of the SCS but rarely persisted there for more than a day [7,8,39] (Fig. 2). Therefore, B cells may also sample antigens in the subcapsular region of the follicle, but this possibility was largely overlooked for several decades because of the relatively short-lived nature of their subcapsular retention compared to their retention on FDCs. In fact, the focus then shifted to finding the mechanism of FDC antigen deposition and, subsequently, a non-phagocytic antigen transport cell with dendritic morphology was proposed to shuttle antigen away from the SCS to the follicle for deposition on FDCs [8,42]. This oversight has recently been redressed by three dynamic imaging studies using two-photon microscopy which independently identified SCS macrophages as important sites for particulate antigen encounter with B cells within the first few hours of the immune response [43]••[44]••[45]••.

SCS macrophages capture and present intact antigen to follicular B cells

Two-photon microscopy allows optical sectioning of intact LNs to depths of up to 200μm [46] and therefore is ideal for high-resolution dynamic imaging of the subcapsular region and superficial follicle. Although the three studies utilised different antigens in different experimental systems, the collective data are consistent with a model whereby SCS macrophages capture and present intact antigen (in the form of ICs [44]••, viruses [45]•• and virus-like particles [43]••) to activate follicular B cells. Intravital imaging of the SCS in live animals with intact lymphatic flow and blood circulation demonstrated macrophage capture

of particulate antigen in the form of ICs [44]•• and viruses [45]••. Follicular B cells routinely migrated to the subcapsular region where they interacted closely with SCS macrophages as measured by their slower median velocities and increased turning angles in the macrophage vicinity [44]••. Antigen-specific B cells accumulated in the subcapsular region [43]••[45]••, acquired antigen from the SCS macrophages and subsequently migrated with antigen capped in their uropod with a reduced median velocity [43]•• [44]•• consistent with antigen engagement of their BCR [47]. B cell activation was also evidenced by their downmodulation of the BCR [43]•• [45]••, upregulation of the activation marker CD86 [43]•• [44]••, and relocation to the T-B border [43]••[44]••[45]••. Indeed, when macrophages were depleted by subcutaneous injection of liposomal clodronate, B cell activation was impaired [45]••. As expected, generalized ablation of macrophages in these experiments also led to a failure to contain viruses within the LN and their systemic dissemination. Taking these real-time imaging studies and the antigen catabolism studies together, there appears to be compartmentalization of LN function with the medullary macrophages being indispensable for the primary function of antigen catabolism and SCS macrophages playing a central role in the secondary function of generating an adaptive humoral response.

Not surprisingly, SCS macrophages may themselves become infected directly following capture of lymph-borne pathogens such as viruses [48]•. Consequently, SCS macrophages in the interfollicular zone have been suggested to be a source of antigen presented by nearby DCs to activate CD8⁺ anti-viral T cells [48]•. Neutrophils, macrophages and DCs infected with agents such as *Toxoplasma gondii* draining in the afferent lymphatics may also be trapped by SCS macrophages, possibly via adhesion to CD169 [49], and thereby act as Trojan horses to indirectly infect them [50]•. Of note, early electron microscopy studies of horse-radish peroxidase-containing ICs showed they were deposited in the SCS floor where they were sequestered on the cell surface or in plasma membrane infoldings suggesting they were non-phagocytic [42]. Consistent with the poorly phagocytic nature of these cells, subcutaneous administration of liposomal clodronate results in widespread ablation of medullary macrophages within 1 day but SCS macrophages ablation takes much longer [51]. In this regard, the propensity of SCS macrophages to retain rather than degrade antigens may make them a safe haven for further pathogen replication.

Mechanisms of SCS macrophage antigen capture and presentation

Ultrastructural studies had earlier shown that SCS macrophages often send cytoplasmic protrusions into the sinus to sample the lymph [3,30,42]. Indeed, it was observed by intravital two-photon microscopy that macrophages with 'heads' that protruded more into the lumen of the SCS were well positioned in the lymph flow to capture ICs as they drained from the afferent lymphatic vessel [44]••. The array of macrophages interspersed between lymphatic vascular endothelial cells in the floor of the SCS resembles the network of epidermal dendritic Langerhans cells in the skin [52] and contrasts against the dense filtration mesh created by macrophages in the medulla [6]•. Furthermore, the rate of SCS lymphatic flow is much faster than the sluggish flow in the cortical sinuses and medulla [44]••[6]•. Thus, SCS macrophages appear to sample newly arriving lymph and become coated with antigenic complexes as evidenced by intravital two-photon microscopy [44]•• whereas medullary macrophages serve to stringently filter and trap all and sundry particulates (Fig. 2). This is consistent with the restricted expression pattern of phagocytic receptors by SCS macrophages and in particular the absence of certain C-type lectins and scavenger receptors.

Despite all these studies, the mechanism of antigen capture is still unknown. A potential mechanism for capture of opsonized antigen involves the use of complement receptors CR3

(CD11b/CD18 or Mac1) [53] and CR4 (CD11c/CD18) [54]. Like other macrophage types, SCS macrophages are thought to express CR3 [45] although this may not uniformly be the case [12]. Classical complement activation occurs when pathogens are opsonized with pre-existing 'natural antibodies' [55] or antibodies generated in a primary antibody response. Antigen may also be opsonized by recognition of carbohydrate moieties on bacterial surfaces by C1q or mannose binding lectin to activate the complement cascade independent of antibody binding [56]. Micro-organisms that lack complement inhibitory molecules can become tagged by spontaneously generated C3b, allowing activation of the alternative pathway. Macrophages also typically express Fc receptors and it can be anticipated that this class of receptors will be involved in IC capture. The mannose receptor is shed into the circulation and lymph and may act as a circulating innate pathogen recognition receptor [57]. SCS macrophages bind to the cysteine-rich domain of the mannose receptor, possibly via CD169 [58,59], providing another potential mechanism for antigen capture. In fact, CD169 itself has been proposed to be involved in pathogen recognition and capture [60] although the capacity of CD169-deficient SCS macrophages [61] to capture and present antigen has yet to be tested. Interestingly, capture of viruses by SCS macrophages appeared to be independent of complement C3 [45]•• raising the possibility that other as yet undefined pathogen recognition receptors may be involved.

Interestingly, similar to FDCs that are also able to present intact antigen to B cells, SCS macrophages express high levels of the adhesion molecules VCAM-1 and ICAM-1 [45]••. It is tempting to speculate that this improves cell-cell interactions and facilitates formation of an immunological synapse between SCS macrophages and B cells [62]. There is considerable evidence for a role of these adhesion molecules in vitro but in vivo data is still lacking. Nevertheless, by two-photon microscopy B cells were observed to gather antigen along macrophage processes in a 'contraction-spreading' pattern [43]••[44]••. One implication of this macrophage membrane-bound mode of presentation is that the avidity of soluble antigen will be much higher thereby potentiating any BCR signals that may drive plasmablast differentiation [63].

Follicular B cells as antigen transporting cells

Another insight from two-photon microscopy has been the recognition of follicular B cells themselves as antigen transport cells [44]•• (Fig. 3). In fact, this concept is not new and has been around for several decades [64,65]. As it turns out, the high intrinsic motility of B cells, which was only recognised with the development of two-photon microscopy, and their abundance make them ideal candidates to transport ICs from the SCS to the follicle. Thus, non-cognate B cells were observed to capture ICs from SCS macrophages and transport them at their uropod in a CR2-dependent manner [44]••. It was envisioned that B cell capture of ICs was driven by the higher affinity of CR1/CR2 for complement than the receptor(s) used by SCS macrophages to capture ICs (potentially CR3/CR4 or Fc receptors), and the subsequent IC deposition on FDCs was driven by the higher density of CR1/CR2 on FDCs than on B cells. A similar role of follicular shuttling of ICs has been suggested for marginal zone B cells in the spleen [66][67]•. In another variant on the theme of B cells as antigen transport cells, IgE containing ICs were captured by circulating B cells, most likely via the low affinity IgE receptor (CD23), and delivered rapidly to splenic follicles [68]. Basophils have also recently been implicated in the transport and presentation of IgE-containing ICs during a recall B cell response [69].

So why is follicular B cell antigen transport important? SCS macrophages have only been shown to retain and display ICs for a period of hours. Antigen deposition on primary follicle FDCs is likely to ensure that rare cognate B cells arriving in the LN more than one day after antigen exposure can be activated. Indeed, very recent two-photon microscopy work has

shown cognate B cells capturing opsonized antigen directly from FDC processes one day after immunization [70]••. Even though the antigen was provided as a single bolus, FDCs displayed sufficient antigen to allow on-going activation of B cells newly arriving in the LN over a period of more than a week [70]••. Thus the antigen-specific B cell precursor frequency and dynamics of lymphocyte recirculation under steady state and inflammatory conditions – as well as the properties of the antigen – will impact the site of B cell antigen encounter. Intriguingly, naïve follicular B cells have been observed by two-photon microscopy to routinely enter the germinal center (GC) light zone [71]••[72]••. It has been argued that this allows for recruitment of naïve B cells into an ongoing GC response [72]••. However, another exciting possibility is that these non-cognate B cells transport antigen-containing ICs into the GC to facilitate affinity maturation of the primary antibody response (Fig. 3).

Conclusion

T cell antigen presentation and activation by dendritic cells in the LN has been extensively mapped at the molecular, cellular and structural level over the past decades. In contrast, the logistics of B cell antigen presentation and activation is only beginning to be re-evaluated in the last few years with two-photon microscopy. These studies have ‘re-discovered’ the critical role of SCS macrophages in capturing and presenting antigen to B cells in the earliest phase of the immune response. However, relatively little is known about these specialized macrophages and the mechanisms by which they handle antigen for cognate B cell activation. Another major finding is the role of B cells themselves in the relay of antigen from SCS to follicle and FDCs. The importance of this B cell antigen transporting function in the antibody response remains to be fully elucidated. Clearly these recent advances would not have been possible without two-photon microscopy, and it is expected that further experiments using this technique will shed more light on dynamic aspects of the immune response in vivo.

Acknowledgments

We thank members of the Cyster lab for invaluable discussions. T.G.P. is an National Health and Medical Research Council CJ Martin Fellow and J.G.C. is an Investigator of the Howard Hughes Medical Institute. Work referred to in this review was supported by NIH grants AI45073 and AI40098.

References

1. Gretz JE, Anderson AO, Shaw S. Cords, channels, corridors and conduits: critical architectural elements facilitating cell interactions in the lymph node cortex. *Immunol. Rev.* 1997; 156:11–24. [PubMed: 9176696]
2. Anderson AO, Anderson ND. Studies on the structure and permeability of the microvasculature in normal rat lymph nodes. *Am J Pathol.* 1975; 80:387–418. [PubMed: 1163637]
3. Fossum S. The architecture of rat lymph nodes. IV. Distribution of ferritin and colloidal carbon in the draining lymph nodes after foot-pad injection. *Scand J Immunol.* 1980; 12:433–441. [PubMed: 7466330]
4. Sainte-Marie G, Peng FS, Belisle C. Overall architecture and pattern of lymph flow in the rat lymph node. *Am J Anat.* 1982; 164:275–309. [PubMed: 7137050]
5. Banerji S, Ni J, Wang SX, Clasper S, Su J, Tammi R, Jones M, Jackson DG. LYVE-1, a new homologue of the CD44 glycoprotein, is a lymph-specific receptor for hyaluronan. *J Cell Biol.* 1999; 144:789–801. [PubMed: 10037799]
- 6. Grigorova IL, Schwab SR, Phan TG, Pham TH, Okada T, Cyster JG. Cortical sinus probing, S1P1-dependent entry and flow-based capture of egressing T cells. *Nat Immunol.* 2009; 10:58–65. [PubMed: 19060900] Uses two-photon microscopy to characterize the three-dimensional

organization of cortical sinuses and the macrophage filled medullary sinuses, in the context of defining T cell egress requirements.

7. Nossal GJ, Abbot A, Mitchell J, Lummus Z. Antigens in immunity. XV. Ultrastructural features of antigen capture in primary and secondary lymphoid follicles. *J Exp Med.* 1968; 127:277–290. [PubMed: 4169585]
8. Tew JG, Mandel TE, Phipps RP, Szakal AK. Tissue localization and retention of antigen in relation to the immune response. *Am J Anat.* 1984; 170:407–420. [PubMed: 6475810]
9. Crocker PR, Gordon S. Properties and distribution of a lectin-like hemagglutinin differentially expressed by murine stromal tissue macrophages. *J Exp Med.* 1986; 164:1862–1875. [PubMed: 3783087]
10. Kraal G. Cells in the marginal zone of the spleen. *Int. Rev. Cytology.* 1992; 132:31–73.
11. Hume DA, Robinson AP, MacPherson GG, Gordon S. The mononuclear phagocyte system of the mouse defined by immunohistochemical localization of antigen F4/80. Relationship between macrophages, Langerhans cells, reticular cells, and dendritic cells in lymphoid and hematopoietic organs. *J Exp Med.* 1983; 158:1522–1536. [PubMed: 6355361]
12. Mohr E, Serre K, Manz RA, Cunningham AF, Khan M, Hardie DL, Bird R, MacLennan IC. Dendritic cells and monocyte/macrophages that create the IL-6/APRIL-rich lymph node microenvironments where plasmablasts mature. *J Immunol.* 2009; 182:2113–2123. [PubMed: 19201864]
13. Dijkstra CD, Van Vliet E, Dopp EA, van der Lelij AA, Kraal G. Marginal zone macrophages identified by a monoclonal antibody: characterization of immuno- and enzyme-histochemical properties and functional capacities. *Immunology.* 1985; 55:23–30. [PubMed: 3888828]
14. Kang YS, Yamazaki S, Iyoda T, Pack M, Bruening SA, Kim JY, Takahara K, Inaba K, Steinman RM, Park CG. SIGN-R1, a novel C-type lectin expressed by marginal zone macrophages in spleen, mediates uptake of the polysaccharide dextran. *Int Immunol.* 2003; 15:177–186. [PubMed: 12578847]
15. Elomaa O, Kangas M, Sahlberg C, Tuukkanen J, Sormunen R, Liakka A, Thesleff I, Kraal G, Tryggvason K. Cloning of a novel bacteria-binding receptor structurally related to scavenger receptors and expressed in a subset of macrophages. *Cell.* 1995; 80:603–609. [PubMed: 7867067]
16. Roozendaal R, Mebius RE, Kraal G. The conduit system of the lymph node. *Int Immunol.* 2008; 20:1483–1487. [PubMed: 18824503]
17. Anderson AO, Shaw S. T cell adhesion to endothelium: the FRC conduit system and other anatomic and molecular features which facilitate the adhesion cascade in lymph node. *Semin Immunol.* 1993; 5:271–282. [PubMed: 8219105]
18. Gretz JE, Norbury CC, Anderson AO, Proudfoot AE, Shaw S. Lymph-borne chemokines and other low molecular weight molecules reach high endothelial venules via specialized conduits while a functional barrier limits access to the lymphocyte microenvironments in lymph node cortex. *J Exp Med.* 2000; 192:1425–1440. [PubMed: 11085745]
19. Itano AA, Jenkins MK. Antigen presentation to naive CD4 T cells in the lymph node. *Nat Immunol.* 2003; 4:733–739. [PubMed: 12888794]
20. Sixt M, Kanazawa N, Selg M, Samson T, Roos G, Reinhardt DP, Pabst R, Lutz MB, Sorokin L. The conduit system transports soluble antigens from the afferent lymph to resident dendritic cells in the T cell area of the lymph node. *Immunity.* 2005; 22:19–29. [PubMed: 15664156]
- 21. Bajenoff M, Egen JG, Koo LY, Laugier JP, Brau F, Glaichenhaus N, Germain RN. Stromal cell networks regulate lymphocyte entry, migration and territoriality in lymph nodes. *Immunity.* 2006 doi:10.1016/j.immuni.2006.10.011. Uses real-time imaging to establish that T cells preferentially migrate along the processes of fibroblastic reticular cells.
22. McNagny KM, Bucy RP, Cooper MD. Reticular cells in peripheral lymphoid tissues express the phosphatidylinositol-linked BP-3 antigen. *Eur. J. Immunol.* 1991; 21:509–515. [PubMed: 1847877]
23. Cyster JG, Ansel KM, Reif K, Ekland EH, Hyman PL, Tang HL, Luther SA, Ngo VN. Follicular stromal cells and lymphocyte homing to follicles. *Immunol Rev.* 2000; 176:181–193. [PubMed: 11043777]

24. Ohtani O, Kikuta A, Ohtsuka A, Murakami T. Organization of the reticular network of rabbit Peyer's patches. *Anat Rec.* 1991; 229:251–258. [PubMed: 2012312]
25. Ushiki T, Ohtani O, Abe K. Scanning electron microscopic studies of reticular framework in the rat mesenteric lymph node. *Anat Rec.* 1995; 241:113–122. [PubMed: 7879916]
26. Katakai T, Hara T, Lee JH, Gonda H, Sugai M, Shimizu A. A novel reticular stromal structure in lymph node cortex: an immuno-platform for interactions among dendritic cells, T cells and B cells. *Int Immunol.* 2004; 16:1133–1142. [PubMed: 15237106]
- 27. Roozendaal R, Mempel TR, Pitcher LA, Gonzalez SF, Verschoor A, Mebius RE, von Andrian UH, Carroll MC. Conduits mediate transport of low-molecular-weight antigen to lymph node follicles. *Immunity.* 2009; 30:264–276. [PubMed: 19185517] Demonstrates using real-time imaging that the small antigen HEL travels rapidly from the SCS sinus into follicular conduits where it is quickly accessed by cognate B cells.
28. Clark SL. The reticulum of lymph nodes in mice studied with the electron microscope. *Am. J. Anat.* 1962; 110:217–257. [PubMed: 13879732]
29. Moe RE. Fine structure of the reticulum and sinuses of lymph nodes. *Am. J. Anat.* 1963; 112:311–335.
30. Farr AG, Cho Y, De Bruyn PP. The structure of the sinus wall of the lymph node relative to its endocytic properties and transmural cell passage. *Am J Anat.* 1980; 157:265–284. [PubMed: 7405871]
- 31. Pape KA, Catron DM, Itano AA, Jenkins MK. The humoral immune response is initiated in lymph nodes by B cells that acquire soluble antigen directly in the follicles. *Immunity.* 2007; 26:491–502. [PubMed: 17379546] Using carefully prepared tissue sections, demonstrates that the small antigen HEL gains very rapid access to the follicle and any cognate B cells within.
32. Gowans JL, Knight EJ. The route of re-circulation of lymphocytes in the rat. *Proc. Royal Soc. Lon. Biol.* 1964; 159:257–282.
33. Tomura M, Yoshida N, Tanaka J, Karasawa S, Miwa Y, Miyawaki A, Kanagawa O. Monitoring cellular movement in vivo with photoconvertible fluorescence protein “Kaede” transgenic mice. *Proc Natl Acad Sci U S A.* 2008; 105:10871–10876. [PubMed: 18663225]
- 34. Qi H, Egen JG, Huang AY, Germain RN. Extrafollicular activation of lymph node B cells by antigen-bearing dendritic cells. *Science.* 2006; 312:1672–1676. [PubMed: 16778060] Visualizes calcium flux within B cells encountering cognate HEL antigen on dendritic cells situated near lymph node HEVs.
35. Wykes M, Pombo A, Jenkins C, MacPherson GG. Dendritic cells interact directly with naive B lymphocytes to transfer antigen and initiate class switching in a primary T-dependent response. *J Immunol.* 1998; 161:1313–1319. [PubMed: 9686593]
36. Colino J, Shen Y, Snapper CM. Dendritic cells pulsed with intact *Streptococcus pneumoniae* elicit both protein- and polysaccharide-specific immunoglobulin isotype responses in vivo through distinct mechanisms. *J Exp Med.* 2002; 195:1–13. [PubMed: 11781360]
37. Bergtold A, Desai DD, Gavhane A, Clynes R. Cell surface recycling of internalized antigen permits dendritic cell priming of B cells. *Immunity.* 2005; 23:503–514. [PubMed: 16286018]
38. Cyster JG. Chemokines, sphingosine-1-phosphate, and cell migration in secondary lymphoid organs. *Annu Rev Immunol.* 2005; 23:127–159. [PubMed: 15771568]
39. Szakal AK, Kosco MH, Tew JG. Microanatomy of lymphoid tissue during humoral immune responses: structure function relationships. *Annu Rev Immunol.* 1989; 7:91–109. [PubMed: 2653380]
40. Nossal GJ, Ada GL, Austin CM, Pye J. Antigens in immunity. 8. Localization of 125-I-labelled antigens in the secondary response. *Immunology.* 1965; 9:349–357. [PubMed: 5844810]
41. Tew JG, Mandel T. The maintenance and regulation of serum antibody levels: evidence indicating a role for antigen retained in lymphoid follicles. *J. Immunol.* 1978; 120:1063–1069. [PubMed: 632579]
42. Szakal AK, Holmes KL, Tew JG. Transport of immune complexes from the subcapsular sinus to lymph node follicles on the surface of nonphagocytic cells, including cells with dendritic morphology. *J Immunol.* 1983; 131:1714–1727. [PubMed: 6619542]

- 43. Carrasco YR, Batista FD. B cells acquire particulate antigen in a macrophage-rich area at the boundary between the follicle and the subcapsular sinus of the lymph node. *Immunity*. 2007; 27:160–171. [PubMed: 17658276] References 43, 44 and 45 use real-time imaging by two-photon microscopy to identify SCS macrophages as a site of antigen display and cognate B cell encounter. This study demonstrates SCS macrophage display of viral like particles (100nm diameter fluorescent beads coated with HEL) and capture by cognate B cells that then move to the T zone.
- 44. Phan TG, Grigorova I, Okada T, Cyster JG. Subcapsular encounter and complement-dependent transport of immune complexes by lymph node B cells. *Nat Immunol*. 2007; 8:992–1000. [PubMed: 17660822] Visualizes the capture of phycoerythrin-IgG immune complexes by SCS macrophages and their subsequent capture by cognate B cells. Also demonstrates non-cognate B cells serve as antigen transport cells to deliver immune complexes from SCS macrophages to FDCs.
- 45. Junt T, Moseman EA, Iannacone M, Massberg S, Lang PA, Boes M, Fink K, Henrickson SE, Shayakhmetov DM, Di Paolo NC, et al. Subcapsular sinus macrophages in lymph nodes clear lymph-borne viruses and present them to antiviral B cells. *Nature*. 2007; 450:110–114. [PubMed: 17934446] SCS macrophages are shown to retain and display vesicular stomatitis virus for capture by cognate B cells. Local ablation of macrophages disrupts viral capture in the lymph node and increases systemic spread.
- 46. Germain RN, Miller MJ, Dustin ML, Nussenzweig MC. Dynamic imaging of the immune system: progress, pitfalls and promise. *Nat Rev Immunol*. 2006; 6:497–507. [PubMed: 16799470]
- 47. Okada T, Miller MJ, Parker I, Krummel MF, Neighbors M, Hartley SB, O'Garra A, Cahalan MD, Cyster JG. Antigen-engaged B cells undergo chemotaxis toward the T zone and form motile conjugates with helper T cells. *PLoS Biology*. 2005; 3:e150. [PubMed: 15857154]
- 48. Hickman HD, Takeda K, Skon CN, Murray FR, Hensley SE, Loomis J, Barber GN, Bennink JR, Yewdell JW. Direct priming of antiviral CD8+ T cells in the peripheral interfollicular region of lymph nodes. *Nat Immunol*. 2008; 9:155–165. [PubMed: 18193049] Real-time imaging is used to track vaccinia virus and vesicular stomatitis virus infected cells. SCS macrophages become infected and cognate CD8 T cells are recruited to subcapsular interfollicular regions where they interact with dendritic cells and become activated.
- 49. Kumamoto Y, Higashi N, Denda-Nagai K, Tsuiji M, Sato K, Crocker PR, Irimura T. Identification of sialoadhesin as a dominant lymph node counter-receptor for mouse macrophage galactose-type C-type lectin 1. *J Biol Chem*. 2004; 279:49274–49280. [PubMed: 15364954]
- 50. Chtanova T, Schaeffer M, Han SJ, van Dooren GG, Nollmann M, Herzmark P, Chan SW, Satija H, Camfield K, Aaron H, et al. Dynamics of Neutrophil Migration in Lymph Nodes during Infection. *Immunity*. 2008 Using real-time imaging and fluorescently labeled Toxoplasma, this study shows SCS macrophages become infected by the parasite. Also shows that infected SCS macrophages can be depleted by neutrophils.
- 51. Delemarre FG, Kors N, Kraal G, van Rooijen N. Repopulation of macrophages in popliteal lymph nodes of mice after liposome-mediated depletion. *J Leukoc Biol*. 1990; 47:251–257. [PubMed: 2137849]
- 52. Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature*. 1998; 392:245–252. [PubMed: 9521319]
- 53. Prodingner, WM.; Wurzner, R.; Erdei, A.; Dierich, MP. Complement. In: Paul, WE., editor. *Fundamental Immunology*. edn 4.. Lippincott-Raven: 1999. p. 967-995.
- 54. Probst HC, van den Broek M. Priming of CTLs by lymphocytic choriomeningitis virus depends on dendritic cells. *J Immunol*. 2005; 174:3920–3924. [PubMed: 15778347]
- 55. Ochsenbein AF, Fehr T, Lutz C, Suter M, Brombacher F, Hengartner H, Zinkernagel RM. Control of early viral and bacterial distribution and disease by natural antibodies. *Science*. 1999; 286:2156–2159. [PubMed: 10591647]
- 56. Roozendaal R, Carroll MC. Emerging patterns in complement-mediated pathogen recognition. *Cell*. 2006; 125:29–32. [PubMed: 16615887]
- 57. Martinez-Pomares L, Mahoney JA, Kaposzta R, Linehan SA, Stahl PD, Gordon S. A functional soluble form of the murine mannose receptor is produced by macrophages in vitro and is present in mouse serum. *J Biol Chem*. 1998; 273:23376–23380. [PubMed: 9722572]

58. Martinez-Pomares L, Kosco-Vilbois M, Darley E, Tree P, Herren S, Bonnefoy JY, Gordon S. Fc chimeric protein containing the cysteine-rich domain of the murine mannose receptor binds to macrophages from splenic marginal zone and lymph node subcapsular sinus and to germinal centers. *J Exp Med*. 1996; 184:1927–1937. [PubMed: 8920880]
59. Martinez-Pomares L, Gordon S. Potential role of the mannose receptor in antigen transport. *Immunol Lett*. 1999; 65:9–13. [PubMed: 10065620]
60. Crocker PR, Paulson JC, Varki A. Siglecs and their roles in the immune system. *Nat Rev Immunol*. 2007; 7:255–266. [PubMed: 17380156]
61. Oetke C, Vinson MC, Jones C, Crocker PR. Sialoadhesin-deficient mice exhibit subtle changes in B- and T-cell populations and reduced immunoglobulin M levels. *Mol Cell Biol*. 2006; 26:1549–1557. [PubMed: 16449664]
62. Batista FD, Arana E, Barral P, Carrasco YR, Depoil D, Eckl-Dorna J, Fleire S, Howe K, Vehlou A, Weber M, et al. The role of integrins and coreceptors in refining thresholds for B-cell responses. *Immunol Rev*. 2007; 218:197–213. [PubMed: 17624954]
63. Paus D, Phan TG, Chan TD, Gardam S, Basten A, Brink R. Antigen recognition strength regulates the choice between extrafollicular plasma cell and germinal center B cell differentiation. *J Exp Med*. 2006; 203:1081–1091. [PubMed: 16606676]
64. Brown JC, De Jesus DG, Holborow EJ, Harris G. Lymphocyte-mediated transport of aggregated human gamma-globulin into germinal centre areas of normal mouse spleen. *Nature*. 1970; 228:367–369. [PubMed: 4919774]
65. Heinen E, Braun M, Coulie PG, Van Snick J, Moeremans M, Cormann N, Kinet-Denoel C, Simar LJ. Transfer of immune complexes from lymphocytes to follicular dendritic cells. *Eur J Immunol*. 1986; 16:167–172. [PubMed: 3956570]
66. Ferguson AR, Youd ME, Corley RB. Marginal zone B cells transport and deposit IgM-containing immune complexes onto follicular dendritic cells. *Int Immunol*. 2004; 16:1411–1422. [PubMed: 15326094]
- 67. Cinamon G, Zachariah M, Lam O, Cyster JG. Follicular shuttling of marginal zone B cells facilitates antigen transport. *Nat Immunol*. 2008; 9:54–62. [PubMed: 18037889] Combines an in vivo antibody tagging technique with flow cytometry and immunofluorescence microscopy to provide evidence that marginal zone B cells shuttle between marginal zone and follicle, ferrying opsonized antigen for deposition on FDCs.
68. Hjelm F, Karlsson MC, Heyman B. A novel B cell-mediated transport of IgE-immune complexes to the follicle of the spleen. *J Immunol*. 2008; 180:6604–6610. [PubMed: 18453579]
69. Denzel A, Maus UA, Rodriguez Gomez M, Moll C, Niedermeier M, Winter C, Maus R, Hollingshead S, Briles DE, Kunz-Schughart LA, et al. Basophils enhance immunological memory responses. *Nat Immunol*. 2008; 9:733–742. [PubMed: 18516038]
- 70. Suzuki K, Grigorova I, Phan TG, Kelly L, Cyster JG. Visualizing B cell capture of cognate antigen from follicular dendritic cells. 2009 submitted. By two-photon microscopy, cognate B cells are observed capturing immune complexes from FDCs. Cognate recognition of FDC displayed antigen occurred for at least a week after immunization.
- 71. Allen CD, Okada T, Tang HL, Cyster JG. Imaging of germinal center selection events during affinity maturation. *Science*. 2007; 315:528–531. [PubMed: 17185562] Visualizes B and T cell migration dynamics within germinal center light and dark zones and shows examples of naïve B cells entering light zone.
- 72. Schwickert TA, Lindquist RL, Shakhar G, Livshits G, Skokos D, Kosco-Vilbois MH, Dustin ML, Nussenzweig MC. In vivo imaging of germinal centres reveals a dynamic open structure. *Nature*. 2007; 446:83–87. [PubMed: 17268470] Visualizes germinal center B cell migration dynamics, demonstrating that naïve B cells frequently migrate into the germinal center. Provides evidence that late entering cognate B cells can join an ongoing germinal center response.

Time	Location	Source	Antigen	Reference
Minutes	1	Pores? Follicular conduits	Small proteins Small proteins	Pape et al 2007 Roosendaal et al 2009
Minutes-hours	1	Macrophages	ICs Viruses Virus-like particles	Phan et al 2007 Junt et al 2007 Carrasco et al 2007
Days	2	Migrating DCs	Small proteins	Qi et al 2006
Days-weeks	3	FDCs	ICs	Suzuki et al 2009

Figure 1. Microanatomy of the lymph node

Schematic of a lymph node showing the locations for antigen presentation to B cells. 1) Subcapsular region. 2) Lymphoid follicle. 3) T zone (paracortex) in the vicinity of HEVs. Lymph drains in the afferent lymphatic to the subcapsular sinus (SCS) and flows to the medullary sinuses. The lymphoid follicle (orange) is situated beneath the SCS and is traversed by occasional follicular conduits (white). Follicular dendritic cells (FDCs) are brown. The T zone (blue) contains HEVs (red) through which lymphocytes enter the lymph node. The medulla (light brown) contains numerous macrophages and cords of plasma cells. Three general locations of B cell access to antigen are labelled and the table summarizes the kinetics and properties of antigens arriving at these sites.

Figure 2. Dynamics of antigen capture in the lymph node

Schematic showing the fate of opsonized (irregular shape) and non-opsonized (circle) antigens draining in the afferent lymphatics. Within minutes SCS macrophages have sampled some of the opsonized antigen while the bulk of antigens continues on to the medullary sinus. Within hours SCS macrophages have translocated some of the opsonized antigens from the lumen into the follicle whereas medullary macrophages have phagocytosed the remaining antigens. Opsonized antigens are then relayed to B cells for deposition on FDCs within a day where they can persist for weeks. In contrast, the remaining antigens are catabolized by medullary macrophages.

Figure 3. Model for role of B cell antigen transport in the primary antibody response

Cognate B cells (green) acquire antigen from SCS macrophages (light brown) and migrate to the T-B border. Upon receipt of T cell help they differentiate into early plasmablasts and migrate to the medullary cords where they secrete antibodies; alternatively they differentiate into GC B cells. Early antibodies feedback to opsonize incoming antigen and these ICs are relayed by SCS macrophages to non-cognate B cells (blue) that transport them into the GC light zone. These antigen-containing ICs are then available for deposition on FDCs (dark brown) and affinity-based selection of GC B cells to become high affinity plasma cells. FMZ, follicular mantle zone.