Published in final edited form as:

Dev Psychopathol. 2008 ; 20(1): 139–164. doi:10.1017/S0954579408000072.

Smoking during pregnancy and offspring externalizing problems: An exploration of genetic and environmental confounds

Brian M. D'Onofrio^{1,2}, Carol A. Van Hulle³, Irwin D. Waldman⁴, Joseph Lee Rodgers⁵, K. Paige Harden⁶, Paul J. Rathouz³, and Benjamin B. Lahey³

¹Indiana University

³University of Chicago

⁴Emory University

⁵University of Oklahoma

⁶University of Virginia

Abstract

Previous studies have documented that smoking during pregnancy (SDP) is associated with offspring externalizing problems, even when measured covariates were used to control for possible confounds. However, the association may be due to non-measured environmental and genetic factors that increase risk for offspring externalizing problems. The current project used the National Longitudinal Survey of Youth and their children, ages 4–10 years, to explore the relations between SDP and offspring conduct problems (CP), oppositional-defiant problems (ODP), and attention deficit hyperactivity problems (ADHP) using methodological and statistical controls for confounds. When offspring were compared to their own siblings who differed in their exposure to prenatal nicotine, there was no effect of SDP on offspring CP and ODP. This suggests that SDP does not have a causal effect on offspring CP and ODP. There was a small association between SDP and ADHP, consistent with a causal effect of SDP, but the magnitude of the association was greatly reduced by methodological and statistical controls. Genetically informed analyses suggest that unmeasured environmental variables influencing both SDP and offspring externalizing behaviors account for the previously observed associations. That is, the current analyses imply that important unidentified environmental factors account for the association between SDP and offspring externalizing problems, not teratogenic effects of SDP.

Keywords

smoking during pregnancy; conduct problems; externalizing problems; ADHD; children of twins; causation

Smoking during pregnancy (SDP) has been consistently linked with externalizing problems in offspring, particularly in males (reviews in Cnattingius, 2004; Huizink & Mulder, 2005; Wakschlag & Hans, 2002; Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002). SDP has been associated with parent-reported conduct problems (Ernst, 2001), arrest history from national crime registries (Brennan, Grekin, & Mednick, 1999; Rasanen et al., 1999), contact

²All correspondence should be sent to: Brian D'Onofrio, PhD, Department of Psychological and Brain Sciences, Indiana University, 1101 E. 10th St., Bloomington, IN 47405, Phone: 812-856-0843, Fax: 812-856-4544, bmdonofr@indiana.edu.

with police obtained by city police records (Gibson, Piquero, & Tibbets, 2000), Oppositional Defiant Disorder (Wakschlag & Keenan, 2001), Conduct Disorder (Fergusson, Woodward, & Horwood, 1998; Wakschlag & Hans, 2002; Wakschlag & Keenan, 2001; Wakschlag et al., 1997; Weissman, Warner, Wickramaratne, & Kandel, 1999) and Attention-Deficit/Hyperactivity Disorder (Mick, Biederman, Faraone, Sayer, & Kleinman, 2002; Rodriguez & Bohlin, 2005).

Reviews of the literature note that the association is consistent with a causal connection because the association is specific to externalizing problems, has been found across diverse samples and measures, demonstrates a dose-response relationship, and is consistent with findings from basic research (Cnattingius, 2004; Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002). The interest in SDP as a cause of externalizing problems is driven, at least in part, by the possibility of identifying a risk factor that is amenable to intervention. Many researchers, however, have noted that methodological considerations render causal interpretations impossible at this time because children cannot be randomly assigned to nicotine exposure conditions; the association between SDP and offspring behavior may be due to risk factors associated with SDP rather than the influence of prenatal nicotine exposure (Fergusson, 1999; Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002). Although associations between SDP and offspring characteristics may be due to more than just nicotine (Huizink & Mulder, 2005), we use the phrase “prenatal nicotine exposure” to refer to SDP throughout the article for ease of presentation.

It is important to note that SDP is correlated with many factors that are also correlated with externalizing problems in children. SDP is correlated with low socioeconomic status (Matthews, 2001; Zimmer & Zimmer, 1998), early age of pregnancy (Cnattingius, 2004; Zimmer & Zimmer, 1998), race (Zimmer & Zimmer, 1998), prenatal care (Zimmer & Zimmer, 1998), maternal depression (Breslau, Kilbey, & Andreski, 1993), history of maternal delinquency (D.M. Fergusson, 1999), paternal antisocial behavior (Maughan, Taylor, Caspi, & Moffitt, 2004), and the family environment (Brook, Brook, & Whiteman, 2000), just to name a few possible confounds to the association between SDP and offspring externalizing problems. Most studies have relied on statistical controls of possible confounds by including measured covariates, such as parental antisocial behavior, parenting practices, family cohesion, and socioeconomic status, in the analyses. Researchers have also utilized case-control studies (Mick, Biederman, Faraone, Sayer, & Kleinman, 2002). Overall, associations between SDP and offspring externalizing behaviors have remained significant in most of the extant SDP studies when the measured covariates are included in the analyses (Cnattingius, 2004; Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002). Some parental variables, however, such as ongoing exposure to secondhand smoke (Maughan, Taylor, Taylor, Butler, & Bynner, 2001), maternal report of conduct disorder as a teenager (but not adult externalizing problems) (Silberg et al., 2003), and antisocial behavior of both parents (Maughan, Taylor, Caspi, & Moffitt, 2004), mediate a majority of the relation between SDP and CP.

In addition to the numerous potential environmental risk factors associated with SDP, a number of researchers have noted that genetic confounds may mediate the relationship between SDP and offspring externalizing problems (D'Onofrio et al., 2003; Fergusson, 1999; Moffitt, 2005; Silberg et al., 2003; Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002). Mothers who smoke during pregnancy may pass down genetic risk for externalizing problems to their offspring, a form of passive gene-environment correlation (rGE; Scarr & McCartney, 1983). Passive rGE occurs when genetic factors common to both the parent and the offspring are correlated with one or more measures of the family environment. Genetic risk for antisocial behavior may contribute to a mother's likelihood of SDP and may confer vulnerability to externalizing problems when inherited by offspring. Standard correlational

research approaches, in which environmental and genetic risks are confounded, are ineffective in delineating whether associations between externalizing problems and SDP are due to environmental causation or passive rGE (D'Onofrio et al., 2003; Moffitt, 2005; Rutter, Pickles, Murray, & Eaves, 2001). Moreover, most of the genetically informed studies of SDP and externalizing problems in offspring (Button, Thapar, & McGuffin, 2005; Knopik et al., 2005; Maughan, Taylor, Caspi, & Moffitt, 2004; Thapar et al., 2003) have used an approach, including a measure of SDP in a standard twin study, that assumes the relation is purely environmental (Purcell & Koenen, 2005; Turkheimer, D'Onofrio, Maes, & Eaves, 2005). Therefore, previous studies have been unable to explore the possibility that passive rGE accounts for observed relations between SDP and externalizing.

The standard twin design may be of limited utility in differentiating between causation and passive rGE (Thapar et al., 2003), but alternative behavior genetic designs are effective in examining the processes underlying intergenerational associations (review in Rutter, Pickles, Murray, & Eaves, 2001). In particular, the Children of Twins (CoT) Design is well-suited for studying SDP because the approach can delineate between environmental processes related specifically to a risk factor shared by siblings, genetic transmission from parents to their offspring, and environmental confounds that vary between families (reviews in D'Onofrio et al., 2003; D'Onofrio et al., 2005; Gottesman & Bertelsen, 1989; Heath, Kendler, Eaves, & Markell, 1985; Rutter, Pickles, Murray, & Eaves, 2001). D'Onofrio and colleagues (2003) explored the association between SDP and offspring birth weight as a model system and found that genetic confounds did not mediate the intergenerational association. Recently, Knopik and her colleagues (in press) utilized the CoT Design to explore the association between parental alcoholism and SDP with offspring ADHD. The analyses suggest that the association between parental alcohol dependence and offspring ADHD is genetically mediated and that genetic risk transmitted from parents to their offspring accounts for a significant portion of the SDP-offspring ADHD relation.

The current article explores the association of SDP and offspring externalizing in a sample of women from the National Longitudinal Survey of Youth (NLSY79) and their children (CNLSY), a sample that confers many advantages. The sample is a nationally representative household sample of women and their children, and the survey included measures related to conduct problems (CP), oppositional-defiant problems (ODP), and attention deficit hyperactivity problems (ADHP). Previous analyses in the NLSY79 have documented characteristics of families related to SDP (Zimmer & Zimmer, 1998), and the associations between SDP and offspring health, behavior problems, and academic achievement have been illustrated in offspring of the NLSY79 (Li & Poirier, 2003).

Finally, the sample provides the opportunity to delineate genetic and environmental processes related to SDP and offspring externalizing. There is a history of using the NLSY79 to explore causal mechanisms using differences within and between families, the theoretical basis of the approach used in this article. For example, researchers have used the dataset to explore the relation between birth order and intelligence (Wichman, Rodgers, & MacCallum, 2005), maternal alcohol and illicit drug use and offspring psychopathology (Chatterji & Markowitz, 2001), teenage childbearing and the women's later adjustment (Geronimus & Korenman, 1992), and maternal age at first birth and offspring adjustment (Turley, 2003).

Multiple adult females from the same household and multiple offspring per mother were included in the NLSY79 and CNLSY studies. This clustering of data permits the association between SDP and offspring problems to be decomposed in two ways. First, children exposed to greater SDP can be compared to their own siblings exposed to more or less SDP. This within-mother effect controls for genetic and environmental factors shared by children with

the same mother, whether they are full or half siblings. If SDP causes offspring externalizing problems, the association would be present when siblings who differ in their exposure to prenatal nicotine are compared (e.g. Rodgers, Cleveland, van den Oord, & Rowe, 2000). Second, the externalizing problems of children of adult siblings who differ in their average level of SDP can be compared. The adult NLSY79 sample was originally based on households and included sibling pairs, referred to as the NLSY Household-Level in the analyses. The clustering allowed children exposed to more SDP to be compared to their cousins with less exposure to prenatal nicotine. This comparison is a within-adult-sibling effect. Furthermore, because the NLSY79 has adult sibling pairs who differ in their genetic relatedness, this cousin comparison can be analyzed contingent on the genetic risk associated with SDP. The NLSY is a very powerful design for such purposes because it contains information on genetic relatedness across two generations: for the original NLSY79 youth (Rodgers, Rowe, & Buster, 1999; Rodgers, Buster, & Rowe, 2001) and the children of the mothers of the NLSY (CNLSY; Rodgers, Rowe, & Li, 1994; Rodgers, Rowe, & May, 1994). These analyses are an extension of the CoT Design, taking advantage of multiple genetically-informative groups in the NLSY79 and CNLSY. The methodological controls provided by family relationships in this design were also combined with traditional statistical controls provided by the inclusion of measured family characteristics, thus providing a rigorous, quasi-experimental test of whether associations between SDP and externalizing problems are causal.

Method

Sample

Mother Generation Sample—The NLSY79 survey was funded by the Bureau of Labor Statistics as a comprehensive study of the future U.S. workforce. It included a nationally-representative sample of 6,111 14–21 year old youths who were not in the military, as well as a supplemental over-sample of 3,652 African American and Hispanic youth. The current study used 6,283 females from the combined sample, because there is cross-generational information only for the females and their children. The probability sample for the NLSY79 was selected using a stratified and clustered design. Primary sampling units consisting of standard metropolitan statistical areas and counties were selected randomly, proportionate to population size. Smaller units (census block groups or enumeration districts) within each primary sampling unit were then randomly selected, with households randomly selected in the third or fourth step. In the initial NLSY79 assessment, the response rate was 90%. Participants were re-interviewed annually from 1979 through 1994 and every two years since then. Retention rates during follow-up assessments were 90% or better during the first 16 waves and have since stayed above 80%.

Of the 6,283 women in the total NLSY79 sample, 4,886 had born at least one child by the 2002 report. The sample was racially diverse: 1,002 mothers were Hispanic (16.0%), 1,561 were Black (24.8%), and 3,720 were non-Black, non-Hispanic (59.2%). The characteristics of the sample with children are presented in Table 1. The women reported the highest grade they completed, with 20 years being the maximum. For income, one outlier over \$350,000 was removed from the dataset so that the value would not improperly skew the results. However, the woman was not removed from the models because the analyses used analytical tools that accounted for missing values. No other variables had outliers that were found to skew the results.

Kinship Links—The NLSY79 mother generation is genetically informative because data were collected on all qualified individuals who resided in the sampled households, which included both full-sibling, half-sibling, and other types of kinship pairs. Approximately 60%

of all within household kinship links have been found to be classifiable as twins, full-siblings, half-siblings, and cousins using the data available up to the 1992 survey. Consistent with previous research using the NLSY, the following estimates of the genetic relatedness (R coefficients) were used in the study: $R = .125$ for cousins, $R = .25$ for half-siblings, $R = .375$ for ambiguous-siblings, $R = .50$ for full-siblings, and $R = .75$ for same-sex twins of unknown zygosity (Rodgers, 1996). Most of these values are the standard measures of correlation between different levels of kinship relatedness that can be derived from the quantitative genetic model (e.g. Falconer, 1981). There are two exceptions. The same-sex twins of unknown zygosity contain half MZ and half DZ twins in the population, so that we assigned an R coefficient halfway between those of MZ twins ($R = 1.0$) and DZ twins ($R = .5$). For the siblings that are either full or half siblings (but who could not be classified by the linking algorithm into either category), we also assigned an R coefficient halfway between that of full siblings ($R = .50$) and half siblings ($R = .25$). Although there are more full siblings than half siblings in the population, past research using these kinship links has suggested that there are only minor differences between using the mid-range values and others that account for the unequal numbers of full- and half-siblings.

The kinship links in the NLSY79 sample have been validated using several different mechanisms. First, in the original development of these kinship links, a validity study was run by estimating genetic influences on sex-standardized adult physical height. Consistent with a large literature that indicates heritability for height of about $h^2 = .90$, h^2 for adult height using these kinship links was .88 in the NLSY79 sample; both meta-analytic and the validity study estimates of c^2 were quite small (Rodgers, 1996). Second, a number of published studies using these kinship links contain sensitivity analyses to ascertain the sensitivity of biometrical estimates to the assumptions involved in using these links (e.g., the use of $R = .375$ described above), which informs the use of these kinship links in this and other more recent studies. Third, we have evidence of concurrent validity by using findings based on these kinship links and comparing them to those from other studies. For example, compare results on age-at-first-intercourse from Rodgers, Rowe, and Buster, (1999), to the molecular genetic results from Miller et al, (1999); compare Rodgers, Buster, & Rowe's (2001) delinquency patterns to those in Miles and Carey (1997); or compare the multivariate patterns across education and IQ in Neiss, Rowe, and Rodgers (2002) to those from Tambs et al., (1989).

Offspring Generation Sample—Bi-annual assessments of the children of women in the NLSY79 began in 1986, with an initial response rate of 95% and subsequent average response rate of 90% (Chase-Lansdale, Mott, Brooks-Gunn, & Phillips, 1991). In 1986, 95% of the biological offspring of NLSY79 mothers were assessed. This response rate has stayed high, with an average response rate of 90%. The interview rate in 2000 was 77%, primarily because approximately 40% of the eligible 15–19 year olds in the African American and Hispanic over-sample groups were not assessed because of budgetary limitations. The low response was not a major concern in the current analyses because we explored externalizing problems in 4–10 year old offspring. The full offspring sample was eligible to be reassessed in 2002, however, and the overall response rate returned to 93%.

Three types of interviews were conducted concerning the offspring generation. Mothers were asked to report on their children's characteristics, including behavior, temperament, and home environment. Beginning in 1988, children between the ages of 10 and 14 years were interviewed directly about their interactions with parents, responsibilities, use of leisure time, relationships with peers, expectations, and delinquent activities. Finally, beginning in 1994, adolescents aged 15 years and over were interviewed extensively on family interactions, substance abuse, delinquent activities, and other issues relating to

transitions to adulthood. The current analyses are based on maternal report of offspring ages 4–10 years old.

The analyses in the current paper were drawn from 11,192 children of the mothers from the 2002 NLSY79. The offspring included 4,886 (43.66%) who were first-borns, 3,705 (33.10%) who were second-borns, 1,747 (15.61%) who were third-borns, 637 (5.69%) who were fourth-borns, and 217 (1.94%) who were fifth-borns. Fifty-one percent (N=5703) of the offspring were male, with 49% being female (N=5484). Of the 11,192 children, 8,889 offspring were of appropriate age to have been assessed with the Behavior Problem Index (BPI; see below for description) and thus were used in the current analyses. Nineteen percent of the offspring included in the analyses were assessed once with the BPI, 22% were assessed twice, 38% were assessed three times, and 22% were assessed four times. In general, mothers of children for whom data are available are slightly older, more educated, wealthier, and have higher aptitude scores, as measured by the Armed Services Vocational Aptitude Battery (see below) than mothers of children with no available data. Therefore, mother's age at first birth, highest level of education, income, and intellectual abilities were controlled in the analyses.

It should be noted that a consistent and problematic source of selection bias is rapidly disappearing from research using the children of the NLSY79 mothers. Past research has been plagued by the fact that, until all childbearing by the NLSY79 cohort is completed, the children were necessarily born to the younger mothers. Because well over 95% of all childbearing was completed by the 2002 survey, and because we used relatively young children in our analyses, this form of selection bias was relatively small (though certainly worth noting). Further, the bias reported above in the opposite direction reported above may approximately compensate.

Measures

Maternal Characteristics—In every wave since 1983, the NLSY79 females were asked about their frequency of smoking and alcohol use during 12 months prior to their most recent pregnancy and during pregnancy. Table 2 includes the response categories for smoking and alcohol use during pregnancy, as well as the conversions to packs/day of smoking and days/month of drinking. Of the 8,889 offspring analyzed the current study, the mothers reported their SDP for 6,503 (73%) within approximately one year of the birth. Because few children in the current analyses were born before 1979, mothers reported their SDP for 90% of their children within 4 years of each birth. Unfortunately, we have no measure of offspring's exposure to secondary smoke. When such exposure has been assessed in previous studies, however, maternal smoking during pregnancy almost always been found to account for unique variance in predicting offspring conduct problems (Brook, Brook, & Whiteman, 2000; Day, Richardson, Goldschmidt, & Cornelius, 2000; Griesler, Kandel, & Davies, 1998; Weissman, Warner, Wickramaratne, & Kandel, 1999), with one exception (Maughan, Taylor, Taylor, Butler, & Bynner, 2001).

Because previous studies using the NLSY identified characteristics of families related to SDP (Zimmer & Zimmer, 1998), a number of maternal characteristics were included in the analyses. When they were 15–22 years old, the NLSY79 mothers (and future mothers) were asked about their engagement in 12 delinquent behaviors during the previous year using a version of the Self-reported Delinquency interview (Elliott & Huizinga, 1983). The Self-reported Delinquency measure, which was also administered to the offspring generation, is reliable and valid and is the benchmark measure used in contemporary delinquency research (Loeber, Farrington, Stouthamer-Loeber, & Van Kammen, 1998; Moffitt, 1990; Moffitt, Caspi, Dickson, Silva, & Stanton, 1996). For more details see Rodgers, Buster, and Rowe

(2001). Symptom counts were regressed on the woman's age at which she completed the survey.

Mother's age at first birth was calculated by subtracting the mother's date of birth from her first child's date of birth. Maternal age at each birth was not calculated because previous research has noted that age at first birth, and not maternal age specific to each birth, predicts offspring delinquency in the NLSY (Turley, 2003). Total net family income reported by mothers at age 30, which includes included income from all adults in the household at that time, and highest degree attained by mothers (as of the 2002 survey) were used as measures of families' socioeconomic status. Total net family income was a summary variable calculated from all income received in the household, including government support and food stamps, by the mother and her spouse. Income from cohabitating partners was not included.

During the summer and fall of 1980, NLSY79 respondents completed the Armed Services Vocational Aptitude Battery (ASVAB), which measured knowledge and skill in 10 areas. A composite score derived from select sections of the battery (word knowledge, paragraph comprehension, math knowledge, and arithmetic reasoning) was used to construct an approximate and unofficial Armed Forces Qualifications Test score (AFQT) for each participant, an approximate IQ equivalent. Raw scores were standardized, summed, and converted to a percentile for a measure of maternal intellectual ability. Multiple imputation (Little & Rubin, 1987) was used to account for the maternal characteristics that were missing (see below).

Offspring CP, ODP, and ADHP—At each wave, mothers rated their 4–10 year old children's behavior problems using the Behavior Problem Index (BPI). The BPI was created by selecting items from the Child Behavior Checklist (Achenbach, 1978) that had the strongest correlations with CBCL factor scores (Peterson & Zill, 1986). Mothers rated each of their children in each assessment wave using a 3-point scale for each item: “often true” = 3, “sometimes true” = 2, or “not true” = 1. We used BPI items to create three *a priori* scales based on DSM constructs. However, confirmatory factor analysis of the 13 externalizing items were conducted with MPlus (Muthén & Muthén, 1998–2005) to account for the dichotomous variables, and the results support the three factor solution (results available upon request). The items measuring CP included the following: (a) cheats or lies, (b) breaks things on purpose or deliberately destroys his/her own or another's things, (c) disobedient at home, (d) disobedient at school, (e) has trouble getting along with teachers, (f) does not feel sorry after misbehaving, and (g) bullies other children. ODP included the following: (a) argues too much, (b) is stubborn, sullen or irritable, and (c) has a very strong temper and loses it easily. ADHP included the following: (a) has difficulty concentrating, (b) impulsive or acts without thinking, and (c) restless or overly active, cannot sit still. We obtain standardized Z scores for CP, ODP, and ADHP within each age and, for children assessed repeatedly, calculated the average score. Therefore, each measure represents the average number of problems between the ages of 4 and 10 years.

The child conduct problem items used in the CNLSY overlap substantially with those used in previous population-based longitudinal studies (Fergusson & Horwood, 2002; Moffitt, Caspi, Dickson, Silva, & Stanton, 1996). Moreover, the ADHP and ODP items are very similar to those used in several studies of the development of conduct problems that yielded results very similar to studies that used full DSM measures of ADHD and ODD (Lahey, McBurnett, & Loeber, 2000). The average CP for males ($M = .16$, $SD = 1.09$, $N = 4,531$) was larger than for females ($M = -.17$, $SD = .87$, $N = 4,358$) [$t = 15.63$, $p < .0001$]. Similar results were found for ODP ($M_{males} = .05$, $SD_{males} = 1.02$, $M_{females} = -.06$, $SD_{females} = .97$)

[$t = 5.28, p < .0001$] and ADHP ($M_{males} = .18, SD_{males} = .88, M_{females} = -.18, SD_{females} = .92$)[$t = 17.70, p < .0001$].

Results

Maternal Characteristics Associated with Smoking During Pregnancy

Estimates of the relations between SDP and maternal intellectual abilities, years of education, income, delinquency, age at first birth, and mean alcohol consumption across all pregnancies are in Table 3. Correlations and unstandardized regression weights are presented. The regression weights were calculated based on the dataset with no missing values (using listwise deletion for those with missing values) and with five multiply imputed datasets for the sample of women who had children so that the entire sample, including those with missing values, could be analyzed (Little & Rubin, 1987; Schafer, 1997). The multiply imputed datasets were based on average maternal smoking and alcohol consumption across pregnancies, intellectual abilities, years of education, income at the age of 30, delinquency, and age at first birth.

Mothers who smoked more on average during their pregnancies were more likely to have lower intellectual abilities. For every unit increase in packs/day, maternal intellectual abilities were over seven percentage points lower. Every pack/day was associated with a decrease in over one and a half years of completed education and more than \$10,000 in yearly income at the age of 30. The average packs/day was also associated with maternal delinquency (an increase in .70 reported delinquent activities in the past year), a decrease of 2.32 years in age at first birth, and a .69 increase in days of alcohol consumption per month during pregnancy. Average SDP was also related to maternal race [$F(2,4816) = 78.40, p < .0001$]. The differences among the three groups were all significant, using the Tukey studentized procedure: Hispanic ($M = .10, SD = .25, N = 823$); Black ($M = .18, SD = .35, N = 1,249$), and non-Black, non-Hispanic ($M = .29, SD = .48, N = 2,747$). The findings are consistent with previous reports using the NLSY (Zimmer & Zimmer, 1998).

Association between SDP and Child Behavior Problems

Mean Comparisons—For descriptive purposes, Table 4 presents the mean of CP, ODP, and ADHP by number of packs smoked during the pregnancy and the sample of offspring. The left column shows the mean (partialled for sex of offspring) and sample size for all of the offspring in the entire sample. The results for CP illustrate a dramatic rise in behavior problems as the number of packs/day increased. The second column presents results for a subset of offspring of women for whom there was variation in SDP within their pregnancies (e.g. the mother did not smoke during one pregnancy and smoked during another, or she varied the amount of cigarettes she smoked among her pregnancies). This subset includes 704 mothers and 1,752 offspring. Offspring who were not exposed to prenatal nicotine, but whose mother smoked during other pregnancies, ($M = .12$) had an increase relative to the estimate in the entire sample ($M = -.08$). Furthermore, offspring who were exposed to large amounts of prenatal nicotine (approximately 2.5 packs/day), but whose mother smoked less during other pregnancies, ($M = .37$) had fewer behavior problems than children in the entire sample exposed to the same level of SDP ($M = .55$). The results suggest that at least part of the association between SDP and offspring CP is not due to prenatal nicotine exposure.

The comparison between the subset of mothers with variation in their SDP to the entire sample, however, is largely a between-family comparison. To partially control for these between-family differences, mean offspring externalizing problems are presented for three groups of women in the subset, based on their mean level of smoking across all of their pregnancies: a low level (average Mother SDP $M = 0$), a medium level ($M = .5$), and a high

level ($M \Rightarrow 1.0$). In the first subset (Low Mean Mother SDP), offspring specifically exposed 0 or 0.5 packs/day did not differ greatly in their CP (.08 vs .11, respectively). In the Medium Mean Mother SDP group, exposure to increased individual-level SDP was somewhat associated with more CP ($M_0 = .09$, $M_5 = .16$, $M_{1.5} = .28$), but this trend was not statistically significant. Furthermore, offspring in the High Mean Mother SDP group did not exhibit greater CP if they were exposed to more prenatal nicotine ($M_0 = .43$, $M_5 = .41$, $M_{1.5} = .37$, $M_{2.5} = .37$). The results suggest that offspring CP were strongly associated with a mother's mean SDP across all pregnancies but only minimally, if at all, related to individual-level exposure to SDP.

Similar patterns can be found for associations of SDP with ODP and ADHP. In the entire sample externalizing problems increased with higher levels of SDP; however, the relation was attenuated in the total subset of offspring whose mothers who had variation in SDP among pregnancies. Furthermore, offspring ODP and ADHP appeared to be significantly related to the average level of mother SDP in all pregnancies; yet, within the three subgroups, offspring externalizing problems were not more prevalent as the individual-specific measure of SDP increased. We caution that the comparisons in Table 4 still include between families effects, as the table does not specifically contrast offspring within the same nuclear family. More advanced statistical approaches are required to compare within-mother variation in SDP and behavior problems. The mean analyses, nevertheless, imply that the association between SDP and offspring externalizing problems may largely occur only between mothers. Again, if SDP were to cause offspring externalizing problems, the association would be found at all levels of analysis.

Hierarchical Linear Models—The offspring of cousins/siblings/twins represents a nested three-level design—the offspring level, the mother level, and the NLSY Household level (for more details see D'Onofrio et al., 2005; Lynch et al., 2006; Mendle et al., 2006). Because offspring are nested under mothers, who are nested under households, observations are not independent. The association between SDP and offspring behavior problems can be studied within regression analyses using Hierarchical Linear Models (HLM) to account for the nested nature of the data, as well as provide appropriate standard error estimates and significance testing (Raudenbush & Bryk, 2002).

Six HLMs were fit for each externalizing measure. Model 1 included SDP specific to each child [variable name under the heading SDP – Packs = “Offspring”], child sex, and the interaction of the SDP and offspring sex because previous research has indicated the effect of SDP is larger for male offspring (Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002). The first model compares offspring whose mothers smoked during their pregnancy with unrelated (e.g. not siblings or cousins) offspring whose mothers did not smoke. Because the measures of externalizing were standardized, the coefficients represent the effect size (Cohen's d) associated with each increase in packs smoked/day. Each HLM includes three variance parameters, which represent the variance in externalizing problems attributable to the three levels in the analysis. Model 2, represents the standard approach to control for differences between mothers who differ in their SDP by including measured covariates to help statistically account for confounds. The model included maternal intellectual ability, years of education, income, delinquency, and age at first birth.

Model 3 calculated the average number of packs/day a mother smoked during all of her pregnancies [SDP – Packs = “Mother”] and included it in the model as a second-level variable. The parameter associated with the second-level variable estimated whether mothers who smoke more on average have children who have more externalizing problems on average. The offspring-level SDP variable [SDP – Packs = “Offspring(C)”] was calculated as the difference between the SDP during the specific pregnancy and the average maternal

SDP. If there was no variation in SDP within a mother, the value was zero. Therefore, the offspring-level SDP variable in model 3 compared siblings in the same family in which the mother smoked more during one pregnancy than the others, while holding constant the average SDP for each mother. The parameter associated with the offspring-level SDP variable in Model 3, the within-mother effect, provided a stronger test of the causal connection between SDP and offspring externalizing, because it was not confounded by factors that vary between mothers. Model 4 included the measured covariates to combine the statistical and methodological approaches.

Model 5 included the third level of the data to calculate the average number of packs/day all women from a household smoked during all of their pregnancies [SDP – Packs = “Household”]. The parameter associated with this third-level variable measures whether mothers and aunts, women from the original NLSY households who smoked more on average have children who had more externalizing problems on average. This variable compared children whose mother and aunts smoked more during their pregnancies with unrelated children whose mother and aunts smoked less. In Model 5, the maternal level SDP [SDP – Packs = “Mother(C)”] was the deviation between the third level average household SDP and average maternal SDP. The parameter associated with this variable compared cousins who differed in their exposure to prenatal nicotine, holding constant the average SDP of women in the NLSY Household. The parameter is the within-adult sibling effect and is free of confounds that vary between unrelated adult siblings. The offspring-level SDP variable in model 5 compared siblings who differed in their prenatal nicotine exposure, while holding constant the average maternal and NLSY average household SDP constant. Finally, Model 6 added the statistical covariates to the methodological controls from Model 5. Algebraic representations of the models can be found elsewhere (D'Onofrio et al., 2005).

All analyses were conducted on five multiply imputed datasets, in order to analyze all available data and avoid bias introduced by individuals or families with missing data. The imputed data were based on the maternal characteristics included as covariates and offspring-specific measures of maternal SDP and alcohol consumption. Any lack of precision due to missing values is represented by larger standard errors around the parameters. Unstandardized regression parameters were utilized because of the difficulty comparing standardized coefficients when exploring causal processes (Kim & Ferree, 1981; Kim & Mueller, 1976). A comparison of the between and within parameters can only be made with unstandardized parameters.

The parameters from the HLMs for CP are presented in Table 5. The results of Model 1 suggest that SDP [“Offspring”] was associated with offspring CP, with the relation being larger for male ($b = .29$) offspring than female offspring ($b = .29 - .11 = .18$). Because the interaction between offspring sex and SDP was significant in the first model, the interaction of offspring sex and SDP at each level was included in the models. In Model 2, illustrates how statistically controlling for measured covariates slightly reduced the effect of SDP ($b_{\text{males}} = .21$ and $b_{\text{females}} = .21 - .12 = .09$). It is difficult to interpret the coefficients associated with the maternal covariates because the coefficients were the result of a simultaneous regression analysis. In Model 3, the mean maternal SDP across all pregnancies [“Mother”] was associated with offspring CP ($b_{\text{males}} = .49$ and $b_{\text{females}} = .49 - .24 = .25$). However, when offspring were compared to their siblings who differed in prenatal nicotine exposure [“Offspring(C)”], there was no association ($b_{\text{males}} = .06$ and $b_{\text{females}} = .06 - .07 = -.01$). The same pattern of results occurred in Model 4 when the measured covariates were included. The mean maternal SDP was associated with offspring CP ($b_{\text{males}} = .35$ and $b_{\text{females}} = .35 - .25 = .10$), but there was no effect within mothers ($b_{\text{males}} = .06$ and $b_{\text{females}} = .06 - .07 = -.01$). In Model 5 the average NLSY Household SDP [“Household”] was associated with offspring CP ($b_{\text{males}} = .49$ and $b_{\text{females}} = .49 - .21 = .28$). Furthermore, there

was an association with maternal SDP corrected for average household SDP ["Mother(C)"], ($b_{\text{males}} = .48$ and $b_{\text{females}} = .48-.43 = .05$), but there was no association when siblings were compared ($b_{\text{males}} = .06$ and $b_{\text{females}} = .06-.07 = -.01$). In Model 6, the average SDP at the household-level ($b_{\text{males}} = .33$ and $b_{\text{females}} = .33-.22 = .11$) was significantly associated with offspring CP when statistical covariates were included in the analyses. Average maternal SDP, holding constant average household SDP, ($b_{\text{males}} = .41$ and $b_{\text{females}} = .41-.45 = -.04$.) was significantly associated for males. In contrast, offspring SDP, holding constant the average household and maternal SDP ($b_{\text{males}} = .06$ and $b_{\text{females}} = .06-.07 = -.01$), was not significantly associated for either gender.

Figure 1 presents unstandardized regression parameters for level-one SDP from Models 1 – 4, for each measure of offspring externalizing. Figure 1 illustrates how the effect sizes associated with SDP drop for both males and females in each successive model. The association between SDP and offspring CP was evident when comparing unrelated children (Model 1), was slightly reduced when including statistical controls (Model 2), and was further attenuated for boys and non-existent for girls when comparing siblings who differ in their exposure to prenatal nicotine, with or without statistical controls (Models 3 and 4). In other words, the average amount of SDP in adult siblings was related to the average CP in all their children (cousins), and the average amount of SDP for a mother across all of her pregnancies was associated with the average CP in all her children. However, siblings who were exposed to more prenatal nicotine did not have higher CP than their brothers and sisters exposed to less prenatal nicotine. This pattern of results is not consistent with the hypothesis that SDP causes CP. If SDP truly caused CP, the association would be found at all three levels of the analysis, particularly when comparing siblings (e.g. Rodgers, Cleveland, van den Oord, & Rowe, 2000).

Table 6 presents results for ODP. In Model 1, SDP ["Offspring"] was associated with ODP ($b = .29$), but there is no interaction with offspring sex. Therefore, subsequent models did not include the interaction between SDP and offspring sex. In Model 2, controlling for measured characteristics of the mothers slightly reduced the association ($b = .20$). In Model 3, however, only the mean maternal SDP across all of her pregnancies ["Mother"] was associated with offspring ODP ($b = .41$). There was no relation when siblings were compared ["Offspring (C)"] ($b = -.02$). In Model 4, the measured covariates reduced the association with average maternal SDP. In Model 5, offspring ODP was associated with average household ["Household"] ($b = .43$) and maternal SDP ["Mother (C)"] ($b = .21$) but not with the offspring-level SDP ($b = -.02$). Finally, Model 6 indicated that when covariates are added to the model, offspring ODP was only associated with household-level SDP ($b = .33$), which is a comparison of unrelated offspring. Again, Figure 1 shows how the use of different comparison groups drastically influences the effect sizes associated with SDP.

The HLM results for ADHP are presented in Table 7. The pattern is similar to that of ODP. SDP was associated with offspring ADHP in Model 1 ["Offspring"] ($b = .27$), and offspring gender did not moderate the association. The magnitude of the relation was slightly reduced in Model 2 ($b = .17$). In Models 3 and 4, average maternal SDP ["Mother"] was associated with ADHP, but the sibling comparison [Offspring(C)] suggested a small association that was not statistically significant ($b = .07$ and $b = .07$). In Models 5 and 6, the association between SDP and offspring ADHP was primarily found at the household-level ["Household"] ($b = .41$ and $b = .24$). The comparison of cousins ["Mother(C)"] and siblings ["Offspring(C)"] who differed in exposure to prenatal nicotine found small associations

between ADHP and SDP.¹ Figure 1 notes the small association between SDP and offspring ADHP when siblings who differed in their exposure to prenatal nicotine were compared.

Structural Equation Modeling—The HLM results suggested that the association between SDP and each measure of externalizing is due to unmeasured confounds that are shared by family members. Consequently, we used multilevel Structural Equation Models (SEMs) to examine whether the relevant confounds were genetic or environmental in origin (Harden et al., in press). The two-level SEM is illustrated in Figure 2. The first member of an adult sibship and her respective children are represented on the left side of the figure; the second member and her children are represented on the right side of the figure. The figure is divided into two portions, representing relations between SDP and externalizing problems within children with the same mother and between unrelated children (described in more detail below). This graphical convenience is not meant to imply that these portions of the model are estimated separately; analogous to the estimation of level-specific regressions in HLM, both portions are estimated simultaneously. The reader is referred to Mehta and Neale (2005) for a more complete didactic on multilevel SEM and its equivalence to hierarchical linear models.

The bottom portion of the model (in broken lines) reflects relationships within children with the same mother. The externalizing problems of the j^{th} child in the k^{th} nuclear family of the i^{th} twin family (EXT_{ijk}) is predicted by his or her own exposure to prenatal nicotine (SDP_{ijk}), as represented by the path labeled w . To the extent the child exposed to more SDP than his or her siblings also reports more externalizing problems than his or her siblings, this is reflected in the w path.

The top portion of the model (in unbroken lines) reflects relationships between unrelated mothers and their children. Average maternal SDP across all pregnancies are represented here by the latent variables labeled $\overline{SDP}_{0jk}$. In addition, similar to the standard twin design path model, the variance in average SDP is decomposed into three components: variance due to additive genetic influences (V_a), variance due to other environmental influences that make the adult siblings more similar (V_c), and variance due to environmental influences that make the adult siblings in the same household different (V_e). The covariance parameters between the latent genetic parameters (A_{cov}) were constrained so that the genetic correlations were appropriate for each group. The covariance of the shared environmental factors (C_{cov}) was constrained such that the correlation equaled 1.0, the standard correlation for the shared environmental latent factors (Neale & Cardon, 1992). Readers may be more familiar with a twin model parameterization in which the A , C , and E components are standardized to a variance of 1.0 and the paths from them are freely estimated. The current model is simply a re-parameterization with the paths fixed to one and the variances freely estimated. Because the paths are fixed to one, the scale of each component is defined by the maternal SDP variable, and the total variance in maternal SDP is the sum of the estimated variances of the A , C , and E components. Dividing the estimated variance of each component by the total variance of average maternal SDP yields the familiar proportions of the heritability (h^2), shared environment (c^2), and nonshared environment (e^2). For more detail concerning variations on twin and family models, see Neale & Cardon (1992).

The average number of externalizing problems in children with the same mother ($\overline{EXT}_{0jk}$) is then regressed on the variance components of SDP – A , C , and E . The model is similar to the standard bivariate genetic analysis (Neale & Cardon, 1992), except that the model

¹Fixed Effect Models, a common econometric approach to clustered data (Greene, 2003), were also fit to the data at the nuclear family-level. The parameters associated with SDP were comparable to the HLM results.

estimated the unstandardized variance components in order to compare the intergenerational paths (b_a , b_c , and b_e) on the same scale. The b_a parameter estimates the effect of genetic factors common to both maternal SDP and child externalizing (passive rGE). The b_c parameter estimates the effect of environmental factors that both make adult siblings similar for SDP and influence offspring externalizing. The model, therefore, controls for environmental factors that make adult siblings similar *and* influence offspring externalizing. Finally, the b_e parameter estimates whether mothers who differ in their levels of SDP for non-genetic reasons (as reflected in variance component V_e) have children who differ in their average number of externalizing problems.

If SDP caused offspring externalizing, the within-mother (w) and non-shared environmental intergenerational path (b_e) would be large. These parameters estimate how differences between child siblings in prenatal nicotine exposure are associated with externalizing at the offspring-level (i.e. if a child sibling exposed to more prenatal nicotine had more externalizing problems than his/her sibling exposed to less), as well as how differences between adult siblings in SDP are associated with externalizing at the maternal-level (i.e. if one mother smoked more than her adult sibling and had children who had more externalizing problems on average). If the relation is not causal, a comparison of the genetic (b_a) and common environmental (b_c) parameters would elucidate the source of the confounds. A residual correlation between the cousins in the offspring generation (r_{co}) was also estimated to account for the covariance among cousins not accounted for by the SDP status of their mothers. The correlation was estimated separately in the different groups to account for differing levels of genetic and environmental relatedness.

The SEM were initially fit with data from the cousin, half-sibling, ambiguous-sibling, and full-sibling families.² Thirty-seven sibling pairs and their offspring were dropped from the analyses because the NLSY household included multiple pairs of varying genetic relatedness (e.g. some participants were related as full siblings but others were related as cousins, even though they lived in the same household). The analysis of clustered data in which participants within a single cluster (NLSY household) belong to multiple groups (e.g. full siblings and cousins) was not permitted in Mplus (Muthén & Muthén, 1998–2005). Pairs of individuals in households of unknown genetic relatedness were also not included in the first model because these pairs could not help separate the estimates of genetic and environmental influences. The decomposition of the variance in SDP is based on the correlations among the different adult sibling pairs in the NLSY, found in Table 8. The parent-level correlations were relatively stable as the genetic relatedness of the siblings increase, suggesting that the shared environmental factors account for most of the covariance between siblings. Similarly, SEMs estimated the variance attributable to additive genetic factors (V_a) to be zero. Moreover, the intergenerational parameter associated with genetic variance (b_a) was unstable (i.e., b_a was large but had very large SEs), a result that frequently occurs when one latent variance component is negligible (D'Onofrio et al., 2003).

Therefore, the parameters associated with genetic factors (V_a and b_a) were constrained to be zero in the subsequent models., making the SEMs a between and within-level of SDP at the nuclear family and NLSY household levels. The subsequent SEMs included all adult sibling pairs, including those in the NLSY where the genetic relatedness of the siblings was unknown. All sibling pairs were included in this model because they were informative about environmental factors that make individuals growing up in the same household similar. The models were estimated using MLR, a maximum likelihood estimator with standard errors

²The SEMs were fit using all of the available sibling relationships, including multiple relationships from the same NLSY household. Because the SEMs assumed that each adult sibling pair was independent, all of the models were reanalyzed only using the first sibling pair from each family. The results were consistent with the models based on the complete sample.

that are robust to non-normality (Muthén & Muthén, 1998–2005). The SEM indicated that shared environmental factors accounted for 61% of the variance ($V_c = .083$, $SE = .01$) of mean SDP at the maternal level, and non-shared environmental influences accounted for 39% ($V_e = .054$, $SE = .01$).

The SEM for CP fit the data well ($\chi^2 = 40.72$, $df = 13$, $RMSEA = .015$). The within-mother parameter was minimal ($w = .07$), consistent with HLM results. Environmental factors that vary between families ($b_c = .48$) and within-adult-sibling families ($b_e = .27$) accounted for the association between SDP and CP. Overall, SDP accounted for 0.1% of the variance in offspring CP within mothers and 8% of the variance between mothers. The intergenerational parameters from the SEMs, including unstandardized estimates, standard errors, and standardized estimates, are found in Table 9. The SEM for ODP also fit the data well ($\chi^2 = 42.93$, $df = 13$, $RMSEA = .015$). For ODP, the results indicated a negligible association with SDP within mothers ($w = .03$). Environmental factors that make adult siblings similar ($b_c = .48$) and unique ($b_e = .46$) both contributed to the association between SDP and ODP. SDP accounted for no variance in ODP within-mother but accounted for 11% of the variance in ODP between mothers. Finally, the SEM for ADHP ($\chi^2 = 37.08$, $df = 13$, $RMSEA = .014$) indicated a small association within-mothers ($w = .09$). The parameter associated with non-shared environmental influences was comparable to the within-mother effect but was not statistically significant ($b_e = .10$). Environmental factors that vary between families accounted for most of the intergenerational associations ($b_c = .61$). SDP accounted for only 0.1% of the variance in ADHP within-mothers but 14.4% of the variance between-mothers.

Discussion

The current article used the clustered design in the NLSY sample to explore the association between SDP and three measures of offspring externalizing problems (CP, ODP, and ADHP). The present comparisons of unrelated children were consistent with the results of previous studies (Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002) in several respects: (a) CP, ODP, and ADHP problems were significantly associated with SDP; (b) each association followed a dose-response relationship; (c) the number of CP demonstrated by children exposed to SDP was higher for male children; and (d) each association remained significant after statistically controlling for associated maternal characteristics. In addition to the use of statistical covariates used in previous studies, the current analyses utilized the multiple levels of the NLSY to account for unmeasured confounds. If SDP caused higher conduct and oppositional problems, the relation would have been evident both when comparing related (e.g. within mothers) and unrelated children (e.g. Rodgers, Cleveland, van den Oord, & Rowe, 2000). However, when children were compared with their siblings who differed in their exposure to SDP, the offspring did not differ with respect to conduct and oppositional defiant problems. SDP accounted for no more than 0.1% of the within-nuclear family variance in offspring conduct problems. In contrast, SDP accounted for 8–14% of the between-nuclear family variance in offspring conduct problems. These results suggest that previous studies found a relationship between SDP and offspring conduct problems not because SDP causes increased risk for conduct or oppositional problems, but because environmental influences that vary between families confound associations between SDP and offspring externalizing. This finding is generally consistent with another CoT study of SDP and ADHD (Knopik et al., in press), as well as with studies that have included more precise measurement of adult characteristics that may confound the relation, such as maternal and paternal antisocial characteristics (Maughan, Taylor, Caspi, & Moffitt, 2004) and maternal delinquency during adolescence (Silberg et al., 2003).

The present findings indicate that environmental influences that are risk factors for both maternal SDP and offspring externalizing underlie the previously observed intergenerational

associations between SDP and offspring conduct problems. This has important implications for future research on the causes of conduct problems. The present findings imply that the salient aspects of the familial or social environment that are robustly associated with both SDP and offspring conduct problems have not been the focus of the research on the etiology of externalizing problems. Because SDP is significantly related to offspring conduct problems after controlling for essentially all of the parent and family characteristics that have been examined as risk factors for offspring conduct problems (Wakschlag & Hans, 2002), the present findings suggest that aspects of the family or social environment that are unrelated to maternal antisocial behavior, maternal age, and other well-studied risk factors are responsible for the relations to SDP. Hopefully, the present findings will lead to an expanded search for these environmental risk factors. The important clues offered by these findings is that they are correlated with SDP, but not correlated with the risk factors that have been controlled in this and previous studies of SDP.

The present analyses, however, suggest a minimal role of SDP with offspring ADHP problems, but the magnitude of the association was greatly reduced compared to previous observations. A recent review has documented a small association between SDP and ADHP after controlling for various confounds (Linnet et al., 2003), and the small association, consistent with a direct causal effect of SDP on offspring ADHP, may be related to the influence of SDP on birth weight (Knopik et al., in press).

The current article used two analytical approaches to explore the associations between SDP and offspring externalizing because each approach has its own advantages and limitations. HLMs were used because they enabled the inclusion of every family and offspring. Furthermore, interactions between variables (e.g. SDP and offspring sex) and the effects of measured family characteristics can be more easily and directly included. Although the HLMs indicate that the relation between SDP and offspring externalizing is not causal, it is difficult to identify whether the origin of relevant confounds are genetic or environmental within the HLM approach. Interpreting the between and within-family parameter estimates from the multiple family groups in the NLSY (cousins, half-siblings, ambiguous-siblings, full-siblings, and twins) becomes quite computationally burdensome (results not shown). In contrast, the SEM approach provides a more straightforward approach to identifying the intergenerational effects of genetic, common environmental, and non-shared environmental influences. The SEM approach, however, cannot include adult sibships that differed in genetic relatedness from other pairs within their household. Furthermore, the sibling correlations and SEM suggest no influence of genetic factors on SDP, in contrast to other reports with larger samples (D'Onofrio et al., 2003; Knopik et al., 2005). Accordingly, we conclude that the results *suggest* that the confounds are environmental rather than genetic in origin. Analyses with larger samples will be required to more specifically elucidate the magnitude of genetic and common environmental processes responsible for the intergenerational association. Regardless of the nature of the confounds, the analyses question whether associations between SDP and offspring externalizing in young children are causal.

There are a number of additional limitations of the current study. First, all of the analyses are based on self-report measures completed by the mother. The retrospective report of SDP may have resulted in underreporting or greater measurement error (Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002), but there is a high correlation between self-reported smoking status and serum cotinine measures (McDonald, Perkins, & Walker, 2005). The reliability of retrospective reports is also similar to the recall of other substance use (Petitti, Friedman, & Kahn, 1981). Furthermore, mothers who report SDP may be more likely to report higher externalizing in their children. The magnitude of intergenerational associations in the current paper, however, are consistent with studies that used more sophisticated

measurement strategies (Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002). Second, due to complexity of the analyses we did not analyze the longitudinal responses of the offspring across the age range (4 to 10 years old); rather, we relied on the average across the years. Future analyses that explore offspring characteristics associated with SDP would benefit from using analytical strategies, such as growth curve models, that can account for individual differences in initial level and change across time (e.g. Willett, Singer, & Martin, 1998). Third, our findings are limited to measures of externalizing in young children and may not reflect the underlying mechanisms responsible for the association between SDP and adult criminal activity (Brennan, Grekin, & Mednick, 1999; Rasanen et al., 1999). Fourth, no father information was included in the analyses. Limited information on the fathers is available in the NLSY, and the association between SDP and offspring externalizing could be due to higher rates of paternal delinquency (Maughan, Taylor, Caspi, & Moffitt, 2004). Methodological work on CoT Design has highlighted its strength with characteristics of individuals (such as SDP), but inasmuch as SDP is influenced by the women's spouses, the interpretation of the parameters may be somewhat restricted (Eaves, Silberg, & Maes, 2005). Fifth, we only explored whether offspring gender moderated the influence of SDP, but there may be vulnerability factors (both environmental and genetic in origin) that make individuals more susceptible to the influence of SDP.

Finally, as is the case with all comparisons using different levels of analyses, the design rests on a number of assumptions about women who have multiple offspring and who vary in the levels of SDP among the pregnancies. The design assumes that women who have only one child are similar to women with multiple offspring, because the within-mother parameter is based solely on women with multiple children. Furthermore, the analyses assume that women who have multiple pregnancies and variation in SDP are similar to women who have multiple offspring but smoked at the same level in each pregnancy, including their ability to accurately report SDP. Again, the within-mother parameter can only be estimated based on women with variation in SDP among her pregnancies.³ Ultimately, the limitations of the current article underscore the necessity of replicating the analyses in other samples and exploring the consequences of SDP in animal studies, which have the advantage of random assignment.

The study highlights the importance of using research designs that can pull apart different risk mechanisms and account for non-measured confounds (Rutter, Pickles, Murray, & Eaves, 2001). Furthermore, the results underline the limitations of solely relying on measured covariates to account for confounds. Even though the initial HLMs that compared unrelated children included a measure of maternal delinquency to control for the intergenerational transmission of externalizing problems there was still an association between SDP and offspring externalizing. If we only relied on the measured covariates we would have drawn the incorrect conclusion. Methodologically, these results support the

³Given the seriousness of these assumptions, we conducted a series of additional analyses. First, all of the HLMs were conducted controlling for birth order, number of offspring per mother, and the interaction between birth order and number of offspring. The results of the HLMs did not appreciably change. We also ran the third HLM, which included within and between-mother effects of SDP, using two restricted subsets of the data: (a) a sample that only included families with multiple children and (b) a sample that only included offspring of mothers who varied in their SDP among their multiple pregnancies. The comparison of the between-mothers SDP parameters in the two restricted subsets to the estimated between-mothers parameters in the entire sample provides a test of the assumptions underlying the current analyses (the within-mother parameter is based only on the most restricted subset). The results are presented in the Appendix (Table 10). The between-mothers SDP parameters for each measure of externalizing were only slightly reduced, if at all, when the analyses were conducted on the two restricted subsets. These results are consistent with our assumptions of equality across the groups (e.g. women with variation in SDP versus women without variation in SDP). Table 11 and 12 also present the means for the offspring externalizing problems in more specific subsets of the data. Finally, we explored the relation between SDP and offspring birth weight using the sibling comparison approach to see if our results replicated previous findings that controlled for genetic and environmental confounds (e.g. D'Onofrio et al., 2003). The results indicated that smoking during pregnancy within mothers was associated with decreased birth weight, consistent with a causal connection and previous genetically informed research on the topic.

importance of using design innovations as well as statistical controls to study risk mechanisms rather than just identifying risk factors (Moffitt, 2005; Rutter, Pickles, Murray, & Eaves, 2001).

Similar to other genetically informed studies of familial characteristics, such as parental divorce (D'Onofrio et al., 2006), the underlying processes associated with a risk factor depend on the specific outcome measure being studied. SDP appears to have a specific environmental association with offspring birth weight, in as much as genetic factors, common environmental, and measured covariates do not confound the relation (D'Onofrio et al., 2003).³ However, CP and ODP in young children associated with SDP are completely due to characteristics that lead to both SDP and offspring externalizing behaviors, not the consequences of SDP. As a result, each risk factor and specific measure of adjustment needs to be studied independently with designs that can separate the risk mechanisms.

Finally, we must state that we were surprised by the results of the analyses, given the existent research on SDP (though no previous research has allowed for control over unobserved heterogeneity to the extent that ours has). We want to stress the need to replicate the current findings. Furthermore, we certainly concur with other researchers (e.g. Maughan, Taylor, Caspi, & Moffitt, 2004) in stressing the importance of reducing SDP because of its effects on a wide range of range of developmental characteristics, particularly neurobehavioral functioning (Cnattingius, 2004; Huizink & Mulder, 2005). Animal research has shown that prenatal nicotine exposure has neurotoxic effects (review in Wakschlag, Pickett, Cook, Benowitz, & Leventhal, 2002). Thus, the findings of the current analyses are limited to externalizing problems during childhood. The current article, though, emphasizes that women who smoke during pregnancy are different than mothers who do not. Although the research literature has noted a number of characteristics of mothers and families associated with SDP, the analyses in the current article suggest that unmeasured characteristics accounted for the associations between SDP and offspring externalizing problems. Perhaps there are more differences than previously imagined between mothers who smoke during pregnancy than those who do not. Therefore, addressing the myriad of risk factors associated with SDP may be a more effective approach of minimizing externalizing problems in offspring than specifically focusing on reducing SDP.

Acknowledgments

This research was supported by grant R01-MH070025 to Benjamin B. Lahey. We would like to thank Valerie Knopik, PhD for her helpful comments on an earlier version of the manuscript.

References

- Achenbach TM. The child behavior profile: I. Boys aged 6–11. *Journal of Consulting and Clinical Psychology*. 1978; 46:478–488. [PubMed: 670491]
- Brennan PA, Grekin ER, Mednick SA. Maternal smoking during pregnancy and adult male criminal outcomes. *Archives of General Psychiatry*. 1999; 56:215–219. [PubMed: 10078497]
- Breslau N, Kilbey M, Andreski P. Vulnerability to psychopathology in nicotine-dependent smokers: An epidemiologic study of young adults. *American Journal of Psychiatry*. 1993; 150:941–946. [PubMed: 8494073]
- Brook JS, Brook DW, Whiteman M. The influence of maternal smoking during pregnancy on the toddler's negativity. *Archives of Pediatric and Adolescent Medicine*. 2000; 154:381–385.
- Button TMM, Thapar A, McGuffin P. Relationship between antisocial behaviour, attention-deficit hyperactivity disorder and maternal prenatal smoking. *British Journal of Psychiatry*. 2005; 187:155–160. [PubMed: 16055827]
- Chase-Lansdale PL, Mott FL, Brooks-Gunn J, Phillips DA. Children of the National Longitudinal Survey of Youth: A unique research opportunity. *Developmental Psychology*. 1991; 27:918–931.

- Chatterji P, Markowitz S. The impact of maternal alcohol and illicit drug use on children's behavior problems: Evidence from the children of the National Longitudinal Survey of Youth. *Journal of Health Economics*. 2001; 20:703–731. [PubMed: 11558645]
- Cnattingius S. The epidemiology of smoking during pregnancy: Smoking prevalence, maternal characteristics, and pregnancy outcomes. *Nicotine and Tobacco Research*. 2004; 6:S125–S140. [PubMed: 15203816]
- D'Onofrio BM, Turkheimer E, Eaves LJ, Corey LA, Berg K, Solaas MH, et al. The role of the Children of Twins design in elucidating causal relations between parent characteristics and child outcomes. *Journal of Child Psychology and Psychiatry*. 2003; 44:1130–1144. [PubMed: 14626455]
- D'Onofrio BM, Turkheimer EN, Emery RE, Slutske W, Heath A, Madden PAF, et al. A genetically informed study of marital instability and its association with offspring psychopathology. *Journal of Abnormal Psychology*. 2005; 114:570–586. [PubMed: 16351381]
- D'Onofrio BM, Turkheimer EN, Emery RE, Slutske W, Heath A, Madden PAF, et al. A genetically informed study of the processes underlying the association between parental marital instability and offspring adjustment. *Developmental Psychology*. 2006; 42:486–499. [PubMed: 16756440]
- Day NL, Richardson GA, Goldschmidt L, Cornelius MD. Effects of prenatal tobacco exposure on preschoolers' behavior. *Journal of Developmental and Behavioral Pediatrics*. 2000; 21:180–188. [PubMed: 10883878]
- Eaves LJ, Silberg JL, Maes HH. Revisiting the children of twins: Can they be used to resolve the environmental effects of dyadic parental treatment on child behavior? *Twin Research*. 2005; 8:283–290.
- Elliott DS, Huizinga D. Social class and delinquent behavior in a national youth panel. *Criminology*. 1983; 21:149–177.
- Ernst M. Behavioral and neural consequences of prenatal exposure to nicotine. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001; 40:630–642. [PubMed: 11392340]
- Falconer, DS. Introduction to quantitative genetics. New York: Longman; 1981.
- Fergusson DM. Prenatal smoking and antisocial behavior. *Archives of General Psychiatry*. 1999; 56:223–224. [PubMed: 10078498]
- Fergusson DM, Horwood LJ. Male and female offending trajectories. *Development and Psychopathology*. 2002; 14:159–177. [PubMed: 11893091]
- Fergusson DM, Woodward LJ, Horwood LJ. Maternal smoking during pregnancy and psychiatric adjustment in late adolescence. *Archives of General Psychiatry*. 1998; 55:721–727. [PubMed: 9707383]
- Geronimus AT, Korenman S. The socioeconomic consequences of teen childbearing reconsidered. *The Quarterly Journal of Economics*. 1992; 107:1187–1214.
- Gibson C, Piquero A, Tibbets S. Assessing the relationship between maternal cigarette smoking during pregnancy and age at first police contact. *Justice Quarterly*. 2000; 17:519–542.
- Gottesman II, Bertelsen A. Confirming Unexpressed Genotypes for Schizophrenia. *Archives of General Psychiatry*. 1989; 46:867–872. [PubMed: 2802925]
- Greene, WH. *Econometric Analysis*. 5. Upper Saddle River, NJ: Prentice-Hall; 2003.
- Griesler P, Kandel D, Davies M. Maternal smoking in pregnancy, child behavior problems, and adolescent smoking. *Journal of Research on Adolescence*. 1998; 8:159–185.
- Harden KP, Turkheimer E, Emery RE, D'Onofrio BM, Slutske W, Heath A, et al. Marital conflict and conduct disorder in Children-of-Twins. *Child Development*. in press.
- Heath AC, Kendler KS, Eaves LJ, Markell D. The resolution of cultural and biological inheritance: Informativeness of different relationships. *Behavior Genetics*. 1985; 15:439–465. [PubMed: 4074271]
- Huizink AC, Mulder EJH. Maternal smoking, drinking or cannabis use during pregnancy and neurobehavioral and cognitive functioning in human offspring. *euroscience and Biobehavioral Reviews*. 2005 E-pub ahead of print.
- Kim J, Ferree GD. Standardization in causal analysis. *Sociological Methods and Research*. 1981; 10:187–210.

- Kim J, Mueller CW. Standardized and unstandardized coefficients in causal analysis: An expository note. *Sociological Methods and Research*. 1976; 4:423–438.
- Knopik VS, Heath AC, Jacob T, Slutske WS, Bucholz KK, Madden PAF, et al. Maternal alcoholism and offspring ADHD: Disentangling genetic and environmental effects using a children-of-twins design. *Psychological Medicine*. in press.
- Knopik VS, Sparrow EP, Madden PAF, Bucholz KK, Hudziak JJ, Reich W, et al. Contributions of parental alcoholism, prenatal substance exposure, and genetic transmission to child ADHD risk: A female twin study. *Psychological Medicine*. 2005; 35:625–635. [PubMed: 15918339]
- Lahey, BB.; McBurnett, K.; Loeber, R. Are attention-deficit hyperactivity disorder and oppositional defiant disorder developmental precursors to conduct disorder?. In: Sameroff, A.; Lewis, M.; Miller, S., editors. *Handbook of developmental psychopathology*. 2. New York: Plenum; 2000. p. 431–446.
- Li K, Poirier DJ. The roles of inputs and outputs in predicting health, behaviour and test scores in early childhood. *Statistics in Medicine*. 2003; 22:3489–3514. [PubMed: 14601015]
- Linnet KM, Dalsgaard S, Obel C, Wisborg K, Henriksen TB, Rodriguez A, et al. Maternal lifestyle factors in pregnancy risk of attention deficit hyperactivity disorder and associated behaviors: review of the current evidence. *American Journal of Psychiatry*. 2003; 160:1028–1040. [PubMed: 12777257]
- Little, RJA.; Rubin, DB. *Statistical Analysis with Missing Data*. New York: Wiley; 1987.
- Loeber, R.; Farrington, DP.; Stouthamer-Loeber, M.; Van Kammen, W. *Antisocial behavior and mental health problems: Explanatory factors in childhood and adolescence*. Mahwah, NJ: Lawrence Erlbaum; 1998.
- Lynch SK, Turkheimer E, Emery RE, D'Onofrio BM, Mendle J, Slutske W, et al. A genetically informed study of the association between harsh punishment and offspring behavioral problems. *Journal of Family Psychology*. 2006; 20:190–198. [PubMed: 16756394]
- Matthews TJ. Smoking during pregnancy in the 1990s. *National Vital Statistics Reports*. 2001; 49(7)
- Maughan B, Taylor A, Caspi A, Moffitt TE. Prenatal smoking and early childhood conduct problems. *Archives of General Psychiatry*. 2004; 61:836–843. [PubMed: 15289282]
- Maughan B, Taylor C, Taylor A, Butler N, Bynner J. Pregnancy smoking and childhood conduct problems: A causal association? *Journal of Child Psychology and Psychiatry*. 2001; 42(8):1021–1028. [PubMed: 11806683]
- McDonald SD, Perkins SL, Walker MC. Correlation between self-reported smoking status and serum cotinine during pregnancy. *Addictive Behaviors*. 2005; 30:853–857. [PubMed: 15833588]
- Mehta PD, Neale MC. People are variables too: multilevel structural equations modeling. *Psychological Methods*. 2005; 10:259–284. [PubMed: 16221028]
- Mendle J, Turkheimer E, D'Onofrio BM, Lynch SK, Emery RE, Slutske W, et al. Family structure and age at menarche: A children of twins approach. *Developmental Psychology*. 2006; 42:535–542.
- Mick E, Biederman J, Faraone SV, Sayer J, Kleinman S. Case control study of attention deficit hyperactivity disorder and maternal smoking, alcohol use, and drug use during pregnancy. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2002; 41:378–385. [PubMed: 11931593]
- Miles DR, Carey G. Genetic and environmental architecture of human aggression. *Journal of Personality and Social Psychology*. 1997; 72:207–217. [PubMed: 9008382]
- Miller WB, Pasta DJ, MacMurray J, Chiu C, Wu H, Comings DE. Dopamine receptors are associated with age at first sexual intercourse. *Journal of Biosocial Science*. 1999; 31:43–54. [PubMed: 10081236]
- Moffitt TE. Juvenile delinquency and attention deficit disorder: Boys' developmental trajectories from age 3 to 15. *Child Development*. 1990; 61:893–910. [PubMed: 2364762]
- Moffitt TE. The new look of behavioral genetics in developmental psychopathology: Gene-environment interplay in antisocial behaviors. *Psychological Bulletin*. 2005; 131:533–554. [PubMed: 16060801]

- Moffitt TE, Caspi A, Dickson N, Silva PA, Stanton W. Childhood-onset versus adolescent-onset antisocial conduct in males: Natural history from age 3 to 18. *Development and Psychopathology*. 1996; 8:399–424.
- Muthén, LK.; Muthén, BO. *Mplus user's Guide*. Third Edition. Los Angeles, CA: Muthén and Muthén; 1998–2005.
- Neale, MC.; Cardon, LRD. *Methodology for genetic studies of twins and families*. Dordrecht, The Netherlands: Kluwer Academic Press; 1992.
- Neiss M, Rowe DC, Rodgers JL. Does education mediate the relationship between IQ and age of first birth? A behavioural genetic analysis. *Journal of Biosocial Science*. 2002; 34:259–275. [PubMed: 11926458]
- Peterson JL, Zill N. Marital disruption, parent-child relationships, and behavior problems in children. *Journal of Marriage and the Family*. 1986; 48:295–307.
- Petitti DB, Friedman GD, Kahn W. Accuracy of information on smoking habits provided on self-administered research questionnaires. *American Journal of Public Health*. 1981; 71:308–311. [PubMed: 7468869]
- Purcell S, Koenen KC. Environmental mediation and the twin design. *Behavior Genetics*. 2005; 35:491–498. [PubMed: 15971029]
- Rasanen P, Hakko H, Isohanni M, Hodgins S, Jarvelin J, Tiihonen J. Maternal smoking during pregnancy and risk of criminal behavior among adult male offspring in the Northern Finland 1966 birth cohort. *American Journal of Psychiatry*. 1999; 156:857–862. [PubMed: 10360123]
- Raudenbush, SW.; Bryk, AS. *Hierarchical Linear Models: Applications and Data Analysis Methods*. 2. Thousand Oaks, NJ: Sage Publications; 2002.
- Rodgers J, Cleveland H, van den Oord E, Rowe D. Resolving the debate over birth order, family size, and intelligence. *American Psychologist*. 2000; 55:599–612. [PubMed: 10892201]
- Rodgers JL. NLSY Youth linking algorithm. 1996 Unpublished Manuscript.
- Rodgers JL, Buster M, Rowe DC. Genetic and environmental influences on delinquency: DF analysis of NLSY kinship data. *Journal of Quantitative Criminology*. 2001; 17:145–168.
- Rodgers JL, Rowe DC, Buster M. Nature, nurture and first sexual intercourse in the USA: Fitting behavioural genetic models to NLSY kinship data. *Journal of Biosocial Science*. 1999; 31:29–41. [PubMed: 10081235]
- Rodgers JL, Rowe DC, Li C. Beyond nature vs. nurture: DF analyses of nonshared influences on problem behaviors. *Developmental Psychology*. 1994; 30:374–384.
- Rodgers JL, Rowe DC, May K. DF analysis of NLSY IQ/achievement data: Nonshared environmental influences. *Intelligence*. 1994; 19:157–177.
- Rodriguez A, Bohlin G. Are maternal smoking and stress during pregnancy related to ADHD symptoms in children? *Journal of Child Psychology and Psychiatry*. 2005; 46(3):246–254. [PubMed: 15755301]
- Rutter M, Pickles A, Murray R, Eaves LJ. Testing hypotheses on specific environmental causal effects on behavior. *Psychological Bulletin*. 2001; 127:291–324. [PubMed: 11393298]
- Scarr S, McCartney K. How people make their own environments: A theory of genotype->environment effects. *Child Development*. 1983; 54:424–435. [PubMed: 6683622]
- Schafer, JL. *Analysis of Incomplete Multivariate Data*. London: Chapman Hall; 1997.
- Silberg JL, Parr T, Neale MC, Rutter M, Angold A, Eaves LJ. Maternal smoking during pregnancy and risk to boys' conduct disturbance: An examination of the causal hypothesis. *Biological Psychiatry*. 2003; 53:130–135. [PubMed: 12547468]
- Tambs K, Sundet JM, Magnus P, Berg K. Genetic and environmental contributions to the covariance between occupational status, educational attainment, and IQ: A study of twins. *Behavior Genetics*. 1989; 19:505–522.
- Thapar A, Fowler T, Rice F, Scourfield J, van den Bree M, Thomas H, et al. Maternal smoking during pregnancy and attention deficit hyperactivity disorder symptoms in offspring. *American Journal of Psychiatry*. 2003; 160:1985–1989. [PubMed: 14594745]
- Turkheimer E, D'Onofrio BM, Maes HH, Eaves LJ. Analysis and interpretation of twin studies with measured environments. *Child Development*. 2005; 76:1217–1233. [PubMed: 16274436]

- Turley RNL. Are children of young mothers disadvantaged because of their mother's age or family background? *Child Development*. 2003; 74:465–474. [PubMed: 12705567]
- Wakschlag LS, Hans SL. Maternal smoking during pregnancy and conduct problems in high-risk youth: A developmental framework. *Development and Psychopathology*. 2002; 14:351–369. [PubMed: 12030696]
- Wakschlag LS, Keenan K. Clinical significance and correlates of disruptive behavior symptoms in environmentally at-risk preschoolers. *Journal of Clinical Child Psychology*. 2001; 30:262–275. [PubMed: 11393926]
- Wakschlag LS, Lahey BB, Loeber R, Green SM, Gordon RA, Leventhal BL. Maternal smoking during pregnancy and the risk of conduct disorder in boys. *Archives of General Psychiatry*. 1997; 54:670–676. [PubMed: 9236551]
- Wakschlag LS, Pickett KE, Cook E, Benowitz NL, Leventhal BL. Maternal smoking during pregnancy and severe antisocial behavior in offspring: A review. *American Journal of Public Health*. 2002; 92:966–974. [PubMed: 12036791]
- Weissman MM, Warner V, Wickramaratne PJ, Kandel DB. Maternal smoking during pregnancy and psychopathology in offspring followed to adulthood. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1999; 38:892–899. [PubMed: 10405508]
- Wichman AL, Rodgers JL, MacCallum RC. A multilevel approach to the relationship between birth order and intelligence. *Personality and Social Psychology Bulletin*. 2005
- Willett JB, Singer JD, Martin NC. The design and analysis of longitudinal studies of development and psychopathology in context: statistical models and methodological recommendations. *Development & Psychopathology*. 1998; 10:395–426. [PubMed: 9635230]
- Zimmer MH, Zimmer M. Socioeconomic determinants of smoking behavior during pregnancy. *The Social Science Journal*. 1998; 35:133–142.

Appendix

Table 10

Comparison of SDP Parameter Results from Model Three Using Different Samples.

Parameter	Entire Sample ^a		Only Families with >1 Child ^b		Only >1 Child and Within-Mother SDP Variation ^c	
	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>
Conduct Problems						
Offspring(C)	.06	.05	.07	.05	.06	.06
Offspring*Sex(C)	-.07	.05	-.07	.05	-.06	.06
Mother	.49	.04	.50	.05	.42	.08
Mother*Sex	-.24	.07	-.27	.08	-.20	.11
Oppositional Defiant Problems						
Offspring(C)	-.02	.04	-.02	.05	-.02	.05
Mother	.41	.03	.41	.03	.35	.05
ADHD Problems						
Offspring(C)	.07	.05	.07	.04	.07	.05
Mother	.37	.03	.38	.03	.35	.05

Note. See Tables 5–7 for more details.

^aParameters are based on complete analysis presented in paper.

^bSample only included offspring from families with more than one child.

^cSample only included offspring from families with multiple children and where there was variation within-mothers in SDP.

Table 11

Comparison of Offspring Externalizing by Pack and Detailed Family Structure

SDP Packs/Day	<u>One Child per Family</u>			<u>Sample Multiple Children and No Within-Mother Variation</u>			<u>Multiple Children and Within-Mother Variation</u>		
	<i>M</i>	<i>SD</i>	<i>N</i>	<i>M</i>	<i>SD</i>	<i>N</i>	<i>M</i>	<i>SD</i>	<i>N</i>
<u>Conduct Problems</u>									
0	-.17	.88	486	-.09	.92	4793	.12	1.03	617
.5	.00	.89	178	.10	1.01	749	.23	1.05	745
1.5	.58	1.38	42	.42	1.19	222	.36	1.12	344
2.5	.63	1.26	7	1.22	1.42	11	.37	1.16	46
<u>Oppositional-Defiant Problems</u>									
0	-.14	.98	486	-.10	.96	4793	.10	1.00	617
.5	.12	1.06	178	.15	.96	749	.17	1.00	745
1.5	.67	1.29	42	.54	1.12	222	.35	1.08	344
2.5	.58	1.26	7	1.25	1.23	11	.24	1.12	46
<u>ADHD Problems</u>									
0	.02	.97	486	-.13	.94	4793	.13	.98	617
.5	.34	.98	178	.17	.95	749	.20	.97	745
1.5	.47	1.12	42	.39	1.14	222	.40	1.07	344
2.5	.40	.26	7	.50	1.26	11	.26	1.01	46

Table 12

Comparison of Offspring Externalizing in Samples with No Variation

SDP Packs/Day	<u>One Child per Family</u>			<u>Sample Two Children and No Within-Mother Variation</u>			<u>Three + Children and No Within-Mother Variation</u>		
	<i>M</i>	<i>SD</i>	<i>N</i>	<i>M</i>	<i>SD</i>	<i>N</i>	<i>M</i>	<i>SD</i>	<i>N</i>
<u>Conduct Problems</u>									
0	-.17	.88	486	-.17	.88	2001	-.04	.95	2792
.5	.00	.89	178	.09	.95	357	.10	1.07	392
1.5	.58	1.38	42	.15	1.14	83	.58	1.19	139
2.5	.63	1.26	7	.45	1.71	5	1.86	.77	6
<u>Oppositional-Defiant Problems</u>									
0	-.14	.98	486	-.09	.95	2001	-.11	.96	2792
.5	.12	1.06	178	.27	.94	357	.05	.97	392
1.5	.67	1.29	42	.44	1.12	83	.60	1.12	139
2.5	.58	1.26	7	.78	1.45	5	1.64	.99	6
<u>ADHD Problems</u>									
0	.02	.97	486	-.12	1.14	2001	-.14	.94	2792
.5	.34	.98	178	.21	.91	357	.14	.98	392
1.5	.47	1.12	42	.20	1.13	83	.51	1.13	139
2.5	.40	.26	7	-.20	.97	5	1.09	1.22	6

Figure 1.

Associations between SDP and offspring externalizing problems using different methodological and statistical controls.

Note. There was no interaction between SDP and offspring sex for ODP and ADHP.

Figure 2.

Structural equation model for the offspring of siblings and twins.

Note. Smoking during pregnancy (SDP) and offspring externalizing (EXT). Unbroken lines represent the maternal and NLSY Household-level of the analysis. The hyphenated lines represent the offspring-level (within-maternal). See Harden et al. (in press) for a comparison to the specific Children of Twins model.

Table 1

Sample Characteristics for NLSY Parents.

Maternal Variables	N	M	SD	Min	Max
Age at First Birth	4886	23.20	5.29	10.76	42.78
Intellectual Ability	4637	37.92	27.21	1	99
Years of Education	4879	12.77	2.57	0	20
Income (at 30 yrs old)	3972	32,070	26,186	18	189,918
Delinquency (1980)	4610	.01	1.48	-1.50	9.00

Note. The units of Intellectual Ability are percentage points, and Delinquency is number of delinquent activities during the previous year regressed on mother's age when she completed the SRD.

Table 2

Smoking and drinking during pregnancy responses for each pregnancy.

Smoking Response	Frequency	Percent	Calculated Packs/Day
None	7177	64.13	0
One pack/day or less	2066	18.46	.5
One to two packs/day	787	7.03	1.5
Two or more packs/day	85	.76	2.5
Missing	1077	9.62	-

Alcohol Response	Frequency	Percent	Calculated Days/Month
Never	6965	62.23	0
Less than 1/month	1538	13.74	.5
Once/month	739	6.60	1
3–4 days/month	434	3.88	3.5
1–2 days/week	344	3.07	6
3–4 days/week	66	.59	14
Nearly every day	19	.17	22
Every day	25	.22	28
Missing	1062	9.49	-

Table 3

Relations with Mean Maternal SDP Across all Pregnancies.

Maternal Variables	<i>r</i>	<i>b</i>	<i>N</i>	<i>b^a</i>
Intellectual Abilities	-.11	-7.37	4581	-7.07
Years of Education	-.27	-1.60	4809	-1.60
Income (at 30 yrs old)	-.17	-10,729	3939	-10,605
Delinquency (1980)	.20	.70 ^b	4552	.70 ^b
Age at First Birth	-.19	-2.33	4816	-2.32
Mean Alcohol Consumption	.15	.67	4817	.67

Note. The regression coefficients represent the mother's average SDP across all of her pregnancies regressed on each maternal characteristic. All parameters are significant $p < .001$.

^aRegression estimate based on 5 multiply imputed datasets for the women who had children.

^bEstimate based on maternal income standardized with $M = 0$ and $SD = 1$.

Table 4

Mean Behavior Problems by SDP and Maternal Sample.

Packs/Day	Entire Sample		Total Subset		Within-Mother Variation ^a							
					Mean Mother SDP = 0				Mean Mother SDP = .5			
	M	N	M	N	M	N	M	N	M	N	M	N
<u>Conduct Problems</u>												
0	-.08	5896	.12	617	.08	248	.09	317	.43	52		
.5	.15	1672	.23	745	.11	113	.16	400	.41	232		
1.5	.40	608	.36	344	-	-	.28	37	.37	307		
2.5	.55	64	.37	46	-	-	-	-	.37	46		
<u>Oppositional-Defiant Problems</u>												
0	-.08	5896	.10	617	.02	248	.11	317	.39	52		
.5	.16	1672	.17	745	-.02	113	.11	400	.38	232		
1.5	.44	608	.35	344	-	-	.16	37	.37	307		
2.5	.45	64	.24	46	-	-	-	-	.24	46		
<u>ADHD Problems</u>												
0	-.09	5896	.13	617	-.01	248	.21	317	.29	52		
.5	.20	1672	.20	745	.03	113	.17	400	.32	232		
1.5	.40	608	.40	344	-	-	.28	37	.41	307		
2.5	.31	64	.26	46	-	-	-	-	.26	46		

Note. All means are presented in Z scores.

^aSample only includes women with multiple pregnancies during which she reported different levels of SDP.

Table 5

Parameter Estimates from HLMs for Conduct Problems.

Variables	Model											
	1	2	3	4	5	6	1	2	3	4	5	6
	b	SE	b	SE	b	SE	b	SE	b	SE	b	SE
SDP - Packs												
Offspring	.29	.03	.21	.03								
Offspring*Sex	-.11	.03	-.12	.04								
Offspring(C)					.06	.05	.06	.05	.06	.05	.06	.05
Offspring*Sex(C)					-.07	.05	-.07	.05	-.07	.05	-.07	.05
Mother					.49	.04	.35	.04				
Mother*Sex					-.24	.07	-.25	.07				
Mother(C)							.48	.14	.41	.13		
Mother*Sex(C)							-.43	.18	-.45	.18		
Household					.49	.05	.33	.05				
Household*Sex					-.21	.08	-.22	.08				
Child Sex	-.29	.02	-.29	.02	-.29	.02	-.29	.02	-.29	.02	-.30	.02
Maternal												
Intellectual Abilities			.00	.00	.00	.00					.00	.00
Years of Education			-.02	.00	-.02	.00					-.02	.00
Income ^a			-.11	.01	-.11	.01					-.11	.01
Delinquency			.06	.01	.06	.01					.06	.01
Age at First Birth			-.02	.00	-.02	.00					-.02	.00
Intercept	.07	.02	.69	.13	.04	.02	.64	.13	.03	.03	.64	.13
Covariances												
Household	.10	.03	.06	.03	.10	.03	.06	.03	.10	.03	.06	.03
Mother	.26	.03	.27	.03	.27	.03	.27	.03	.26	.03	.27	.03
Offspring	.58	.01	.57	.01	.57	.01	.57	.01	.57	.01	.57	.01

Note. All parameters are unstandardized. Parameters in bold are significant at $p < .05$. Child sex is coded male=0 and female=1.

^aIncome was converted to a Z score so that the parameter could be accurately estimated in standard units. Offspring(C) represents the within-mother effect of SDP. The Mother(C) represents the within-adult sibling effect.

Table 6

Parameter Estimates from HLMs for Opposition-Defiant Problems.

Variables	Model											
	1	2	3	4	5	6						
	b	SE	b	SE	b	SE	b	SE	b	SE	b	SE
SDP - Packs												
Offspring	.29	.03	.20	.02								
Offspring*Sex	-.03	.04										
Offspring(C)			-.02	.04	-.01	.04	-.02	.04	-.02	.04	-.02	-.04
Mother			.41	.03	.32	.03						
Mother(C)							.21	.10	.16	.10		
Household							.43	.03	.33	.03		
Child Sex	-.08	.02	-.09	.02	-.09	.02	-.09	.02	-.09	.02	-.09	.02
Maternal												
Intellectual			-.002	.001	-.002	.00			-.001	.000		
Abilities												
Years Education			-.02	.01	-.02	.01			-.02	.01		
Income ^a			-.09	.02	-.08	.02			-.08	.02		
Delinquency			.05	.01	.04	.01			.04	.01		
Age at First Birth			-.01	.00	-.01	.00			-.01	.00		
Intercept	-.02	.02	.49	.13	-.05	.02	.39	.13	-.05	.01	.38	.13
Covariances												
Household	.09	.03	.07	.03	.08	.03	.07	.03	.09	.03	.07	.03
Mother	.25	.03	.25	.03	.26	.03	.25	.03	.25	.03	.25	.03
Offspring	.64	.01	.63	.01	.63	.01	.63	.01	.63	.01	.63	.01

Note. See note on Table 5.

Table 7

Parameter Estimates from HLMs for ADHD Problems.

Variables	Model											
	1		2		3		4		5		6	
	b	SE	b	SE	b	SE	b	SE	b	SE	b	SE
SDP - Packs												
Offspring	.27	.03	.17	.03								
Offspring*Sex	.02	.04										
Offspring(C)					.07	.05	.07	.05	.07	.05	.07	.05
Mother					.37	.03	.22	.03				
Mother(C)									.05	.10	-.04	.10
Household									.41	.03	.24	.03
Child Sex	-.38	.02	-.38	.02	-.38	.02	-.38	.02	-.38	.02	-.38	.02
Maternal												
Intellectual			-.001	.001			-.001	.001			-.001	.001
Abilities												
Years Education			-.02	.01			-.02	.01			-.02	.01
Income ^a			-.10	.01			-.10	.01			-.10	.01
Delinquency			.04	.01			.04	.01			.04	.01
Age at First Birth			-.03	.00			-.03	.00			-.03	.00
Intercept	.13	.02	1.05	.11	.11	.02	1.01	.11	.10	.02	.99	.11
Covariances												
Household	.09	.02	.04	.02	.09	.02	.04	.02	.10	.02	.05	.02
Mother	.19	.03	.19	.03	.19	.03	.19	.03	.19	.03	.18	.03
Offspring	.66	.01	.66	.01	.66	.01	.65	.01	.66	.01	.65	.01

Note. See note on Table 5.

Table 8

Correlations between adult siblings in average SDP across all pregnancies.

Relatedness	<i>r</i>	<i>N</i>
Unknown	.41	297
Cousins	.39	31
Half-sibs	.27	16
Ambiguous-sibs	.54	85
Full-sibs	.34	472
Twins	-	0

Note. Correlations in bold are significant at $p < .05$. The correlations were not corrected for multiple sibling pairs per family. There were no twin pairs where both female twins had reports of SDP.

Table 9

SEM results for SDP-offspring externalizing.

Intergenerational Parameters	<u>CP</u>		<u>ODP</u>		<u>ADHD</u>	
	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>	<i>b</i>	<i>SE</i>
Within mothers (<i>w</i>)	.07	.04	.03	.04	.01	.04
Genetic (<i>b_a</i>)	-	-	-	-	-	-
Common environment (<i>b_c</i>)	.48	.14	.48	.12	.26	.13
Non-shared environment (<i>b_e</i>)	.27	.19	.46	.19	.20	.16

Note. Because the initial SEM indicated that no variance in SDP was due to additive genetic factors (*V_a*) the path from the genetic latent factor (*b_a*) was dropped from the SEMs. Parameters in bold are significant at $p < .05$.