Published in final edited form as:

Org Lett. 2013 July 5; 15(13): 3242–3245. doi:10.1021/ol401226y.

Stereoselective Synthesis of Dienyl-Carboxylate Building Blocks: Formal Synthesis of Inthomycin C

Caroline Souris, Frédéric Frébault, Ashay Patel, Davide Audisio, K. N. Houk, and Nuno Maulide*

Max-Planck-Institut für Kohlenforschung, Kaiser-Wilhelm-Platz 1, 45470 Mülheim an der Ruhr, Germany and University of California, Los Angeles, Department of Chemistry and Biochemistry 607 Charles E. Young Drive East, CA 90095-1569, Los Angeles

Abstract

A direct synthesis of stereodefined halodienes is reported. Those key building blocks enable a concise access to polyenic products, as demonstrated in a modular synthesis of Inthomycin C.

Dienyl carboxylate and carbinol subunits are fundamental structural scaffolds present in various natural products¹ (Figure 1). Through simple modifications in the diene substitution pattern and olefin geometry, Nature is able to access remarkable structural diversity. Such functionalized conjugated dienes are conventionally built from simple mono-olefinic fragments² through, *e.g.*, cross-coupling,^{3,4} metathesis⁵ or olefination reactions.⁶ Controlling the configuration of the double bond arrays generated during such transformations represents a major challenge.

The use of stereodefined mono-olefinic fragments as building blocks for cross-coupling reactions leading to the di- or polyenyl frameworks of interest is a strategy that has gained prominence.⁷ An elegant example of such a tactic is the development by Burke of an assembly of so-called MIDA-boronate building blocks, that allow the deployment of iterative cross-couplings en route to polyenic natural products.⁸ Nevertheless, the syntheses of the basic mono-olefinic fragments are typically multi-step and mandate the introduction of a halide and organometallic residue in each fragment. We report herein a strategy for the direct preparation of dienyl carboxylate building blocks that significantly streamlines the total synthesis of polyene natural products.^{9,10,11}

During our studies on allylic alkylation of lactone **1a**,¹² we have discovered an unexpected halide ring-opening relying on the use of alkali halide salts. As shown in Scheme 1(a-b), clean ring-opening of lactone **1a** with either NaI or LiBr provides almost exclusively *trans*-

*maulide@mpi-muelheim.mpg.de.

Supporting Information **Available** Experimental procedures and spectroscopic data for new compounds, computational details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

halocyclobutenes **2a-b** in quantitative yields. Furthermore, nucleophilic chlorination of **1a** with HCl selectively affords *cis*-chlorocyclobutene **2c** with similar efficiency (Scheme 1c).

These thermally stable halocyclobutenes and their ester or amide derivatives are prone to 4π -electrocyclic conrotatory ring opening^{13,14} upon heating. Surprisingly, the iodocyclobutene **2a** and derivatives undergo productive ring-opening leading to a mixture of diene geometrical isomers.^{15,16} In contrast, **2b** and the brominated carboxylate analogues thereof afford the (*E,E*)-halodienes **4a-d** cleanly upon refluxing in THF.¹⁷ In complementary fashion, the *cis*-4-chlorocyclobut-2-ene carboxylic acid **2c**¹⁸ can be readily derivatized and unravelled to deliver the (*Z,E*)-dienyl carboxylates **5a-d** (Scheme 1c).

In order to determine the factors controlling the reactivity and stereoselectivity of the ring openings of **2a-c**, we modeled these 4π -electrocyclic ring opening reactions computationally.¹⁹ Computations indicate that the electrocyclic ring opening reactions of the disubstituted cyclobutenes **2a-c** are all exergonic and thus irreversible with a ΔG_{rxn} ranging from -11 kcal/mol for **2b** and **2c** to -14 kcal/mol for **2a**. The high temperatures required for the ring openings of **2a-c** are consistent with the computed free energy barriers (~30 kcal/mol). The transition structures for the ring opening of **2a**, **2b**, and **2c** are shown in Figure 2.²⁰

Donors like iodide stabilize the transition state by interacting with the transition state LUMO.²¹ The iodide substituent is a weaker donor than chloride or bromide, explaining a 10-fold difference in the rate of reaction of **2a** and **2b**.

The stereochemical outcomes of the electrocyclic reactions of **2b-c** are in agreement with the model regarding torquoselectivity of 4π -electrocyclic ring opening of cyclobutenes.¹⁶ In the ring opening of halocyclobutenes, strong *n* donors rotate outward in order to maximize orbital overlap between the high energy, nonbonding orbital of the donor and the transition state LUMO (σ^*).²² Indeed, computations show that the conrotatory transition state of ring opening in which the halide rotates outward is 13-14 kcal/mol lower in energy than the alternative where the halide rotates inward (Figure 2). Conversely, strong acceptors, like aldehydes, tend to rotate inward in order to maximize overlap of their relatively low energy acceptor orbital with the transition state HOMO (σ).

The $\Delta\Delta G^\ddagger$ values of *trans*-substituted cyclobutenes **2a-c** (13-15 kcal/mol) demonstrate the clear preference of the halide substituent for outward rotation. In the ring opening of *trans*-disubstituted **2a-b**, both the acid²³ and the halide rotate outward, whereas in the ring opening of *cis*-isomer **2c** the acid is forced to rotate inward in the presence of the strong chloride donor. The similarity in the barriers of the *trans*- and *cis*-disubstituted cyclobutene ring openings are in agreement with prior computational work which indicates that the acid has only a small preference for outward rotation.¹⁸

While the formation of the (*E,E*)-iododiene via electrocyclic reaction of **2a** follows the explanation provided above, the (*E,Z*)-iododiene cannot be formed under thermal conditions via a pericyclic mechanism, as the reaction would have to proceed through a forbidden, disrotatory transition state.^{10a} Such a transition state is inaccessible due to its prohibitively high energy. Instead, this (*E,Z*) isomer is likely formed by subsequent isomerization of the allowed (*E,E*) product.

We next sought to demonstrate the synthetic utility of these dienyl-carboxylates.²⁴ Inthomycin C (Scheme 2), isolated in 1991,²⁵ was shown to reduce prostate cancer cell growth as well as to possess selective *in vitro* antimicrobial activity. Our retrosynthetic analysis is shown in Scheme 2, breaking the natural product down to three simple fragments

4e, **6** and **7**. We envisioned that the bromo diene **4e** would undergo cross coupling with a suitable organometallic partner **6** delivering a triene. Further functional group interconversion and Mukaiyama aldol reaction with silylketene acetal **7** would ultimately allow the preparation of Inthomycin C.

In order to obtain validation for the strategy outlined in Scheme 2, we briefly evaluated the reactivity of the stereodefined diene building blocks in cross-coupling reactions (Scheme 3).²⁶ For instance, Suzuki couplings employing aryl- and vinyl boronic acids proceeded to deliver the corresponding substituted dienolic esters **8a-d**. Sonogashira reactions could be performed in good yields with silyl-, alkyl- and aryl-substituted terminal alkynes and various Stille cross-couplings were also successful.²⁷ Similarly, cross-coupling onto the (*Z,E*)-chloro-dienes **5b-d** took place without loss of the diene geometrical configuration (Scheme 4), an important observation.²⁸ To the best of our knowledge, this is an unprecedented use of doubly vinylous chloride esters in cross-coupling reaction.

Armed with this knowledge, we could then confidently complete the synthesis of Inthomycin C (Scheme 5). As shown, lithium bromide smoothly opened the methyl substituted lactone **1b**. As before, a single *trans*-cyclobutenyl bromide **2d** was obtained. Further amide coupling and 4 π electrocyclic ring opening afforded 2-methyl-5-bromodienoic amide **4e** as a single geometrical isomer. Stille cross-coupling with vinyl stannane **6**,²⁹ followed by reduction to the aldehyde **11** and organocatalytic Mukaiyama aldol reaction with silylketene acetal **7** then led to product **12** in 50% yield.³⁰ The conversion of **12** to Inthomycin C has been previously reported.³¹ This modular assembly of substituents around dienes such as **4e** allows considerable flexibility in the context of total synthesis.

In summary, we have reported herein a direct route to functionalised and stereodefined halodiene carboxylate building blocks, which proceeds through electrocyclic ring opening of readily available halocyclobutene precursors. Appropriate control of stereochemical information at the cyclobutene level ensures access to a suitably configured diene fragment upon ring opening. Cross coupling onto these subunits allows ready access to natural product-like polyene substructures without erosion of the geometrical purity. The utility of this strategy is showcased by a modular short formal synthesis of Inthomycin C. We are currently pursuing the application of these and related approaches to the total syntheses of diverse polyene natural products.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the Max-Planck Society and the Max-Planck Institut für Kohlenforschung for funding. This work was funded by the DFG (Grant MA 4861/3-1). The project "SusChemSys" is co-financed by the European Regional Development Fund (ERDF) and the state of NRW (Germany) under the Operational Programme "Regional Competitiveness and Employment" 2007 - 2013. A.P. and K.N.H. thank the National Institute of General Medical Science, NIH (GM 36700) for funding. A.P. acknowledges the support of CBI training program (T32 GM 008496). Computational studies were performed using the Hoffman2 cluster at UCLA as well as resources provided by Extreme Science and Engineering Discovery Environment (XSEDE) program supported by the NSF (OCI-1053575).

References

- (1). Hopf, H.; Maas, G. The Chemistry of dienes and polyenes. Rappoport, Z., editor. Wiley; Chichester: 1997.

- (2). For a recent review on diene synthesis, see: De Paolis M, Chataigner I, Maddaluno J. *Top Curr. Chem.* 2012; 327:87. [PubMed: 22527407]
- (3). Hartwig, JF. *Handbook of Organopalladium Chemistry for Organic Synthesis*. Negishi, E., editor. Wiley; Chichester: 2002.
- (4). (a) Beletskaya, IP.; Cheprakov, AV. Metal Complexes as Catalysts for C-C Cross Coupling Reactions. In: McCleverty, JA.; Meyer, TJ., editors. *Comprehensive Coordination Chemistry II*. Vol. 9. Elsevier Ltd.; Oxford: 2004. p. 305-368. (b) Miyaura N, Suzuki A. *Chem. Rev.* 1995; 95:2457. (c) Miyaura N. *J. Organomet. Chem.* 2002; 653:54. (d) Suzuki A. *Angew. Chem. Int. Ed.* 2011; 50:6722.
- (5). Grubbs, RH., editor. *Handbook of Metathesis*. Wiley-VCH; Weinheim: 2003.
- (6). Takeda, T., editor. *Modern Carbonyl Olefination: Methods and Applications*. Wiley-VCH; Weinheim: 2004.
- (7). (a) Eto K, Yoshino M, Takahashi K, Ishihara J, Hatakeyama S. *Org. Lett.* 2011; 13:5398. [PubMed: 21888385] (b) O'Neil GW, Phillips AJ. *J. Am. Chem. Soc.* 2006; 128:5340. [PubMed: 16620095] (c) Schmidt B, Kunz O. *Eur. J. Org. Chem.* 2012:1008.
- (8). (a) Woerly EM, Struble JR, Palyam N, O'Hara SP, Burke MD. *Tetrahedron.* 2011; 67:4333. [PubMed: 22753994] (b) Gillis EP, Burke MD. *Aldrichimica Acta.* 2009; 42:17. [PubMed: 22523433] (c) Gillis EP, Burke MD. *J. Am. Chem. Soc.* 2007; 129:6716. [PubMed: 17488084] (d) Lee SJ, Gray KC, Paek JS, Burke MD. *J. Am. Chem. Soc.* 2008; 130:466. [PubMed: 18081295] (e) Woerly EM, Cherney AH, Davis EK, Burke MD. *J. Am. Chem. Soc.* 2010; 132:6941. [PubMed: 20441218] (f) Fujita K, Matsui R, Suzuki T, Kobayashi S. *Angew. Chem. Int. Ed.* 2012; 51:7271.
- (9). For previous syntheses of (2*E*,4*E*)-5-iodo-2,4-dienoic acid and ester derivatives, see: Abarbri M, Parrain JL, Duchene A, Thibonnet M. *Synthesis.* 2006:2951. Batsanov AS, Knowles JP, Samsam B, Whiting A. *Adv. Synth. Catal.* 2008; 350:227.
- (10). For previous syntheses of (2*E*,4*E*)-5-bromo-2,4-dienoic esters, see: Macdonald G, Alcaraz L, Wei XD, Lewis NJ, Taylor RJK. *Tetrahedron.* 1998; 54:9823. Zeng FX, Negishi E. *Org. Lett.* 2002; 4:703. [PubMed: 11869106] Crombie L, Horsham MA, Jarrett SRM. *Tetrahedron Lett.* 1989; 30:4299. Wei XD, Taylor RJK. *Tetrahedron Lett.* 1998; 39:3815. Wei XD, Taylor RJK. *J. Org. Chem.* 2000; 65:616. [PubMed: 10813984]
- (11). For previous syntheses of (2*Z*,4*E*)-5-chloro-2,4-dienoic ester derivative, see: Takeda A, Tsuboi S. *J. Org. Chem.* 1973; 38:1709.
- (12). (a) Frébault F, Luparia M, Oliveira MT, Goddard R, Maulide N. *Angew. Chem. Int. Ed.* 2010; 49:5672. (b) Luparia M, Oliveira MT, Audisio D, Frébault F, Maulide N. *Angew. Chem. Int. Ed.* 2011; 50:12631. (c) Luparia M, Audisio D, Maulide N. *Synlett.* 2011:735. (d) Audisio D, Luparia M, Oliveira MT, Klutt D, Maulide N. *Angew. Chem. Int. Ed.* 2012; 51:7314. (e) Niyomchon S, Audisio D, Luparia M, Maulide N. *Org. Lett.* 2013; 15:2318. [PubMed: 23614852] (f) Souris C, Luparia M, Frébault F, Audisio D, Fares C, Goddard R, Maulide N. *Chem. Eur. J.* 2013; 19:6566. [PubMed: 23576421]
- (13). Woodward RB, Hoffmann R. *Angew. Chem. Int. Ed.* 1969; 8:781. See ref 12c and references cited therein.
- (14). For isolated examples of electrocyclic ring opening of cyclobutenes in the context of natural products synthesis, see: Trost BM, McDougal PG. *J. Org. Chem.* 1984; 49:458. Nicolaou KC, Vega JA, Vassilikogiannakis G. *Angew. Chem. Int. Ed.* 2001; 40:4441. Schreiber SL, Santini C. *J. Am. Chem. Soc.* 1984; 106:4038. Ogawa S, Urabe D, Yokokura Y, Arai H, Arita M, Inoue M. *Org. Lett.* 2009; 11:3602. [PubMed: 19637859]
- (15). Brune HA, Schwub W. *Tetrahedron.* 1969; 25:4375.
- (16). For comparison with (2*E*,4*Z*)-5-iodo-2,4-dienoic ester and (2*Z*,4*E*)-5-iodo-2,4-dienoic ester, see: Wang GW, Mohan S, Negishi EI. *P. Natl. Acad. Sci. USA.* 2011; 108:11344. See Supporting Information for details.
- (17). A larger scale reaction afforded 2.5g of diene **4b** in 95% yield from the lactone **1a**.
- (18). Pirkle WH, Mckendry LH. *J. Am. Chem. Soc.* 1969; 91:1179. See also ref 12d.
- (19). Computational experiments were carried out using the M06-2X/6-31+G(d,p) model chemistry with Gaussian 09. For the reaction of **2c**, the core electrons of iodine were modelled using the

LAN2LDZ pseudopotential. Additional details regarding the computational methods employed as well as the full citation for Gaussian are provided in the Supporting Information.

- (20). Comparing the barriers of **2a** and **2b**, we found that ring opening of **2a** proceeds with a $\Delta G^\ddagger$ of 29.8 kcal/mol, whereas the reaction of **2b** has a $\Delta G^\ddagger$ of 28.7. This 1.1 kcal/mol difference leads to a 10-fold greater reaction rate for **2b** compared to its iodo-analogue **2a**.
- (21). Dolbier W, Koroniak H, Houk K, Sheu C. Acc. Chem. Res. 1996; 29:471.
- (22). Niwayama S, Houk KN. Tetrahedron Lett. 1992; 33:883.
- (23). (a) Niwayama S, Houk KN, Kusumi T. Tetrahedron Lett. 1994; 35:527.(b) Lee PS, Sakai S, Hörstermann P, Roth WR, Kallel EA, Houk KN. J. Am. Chem. Soc. 2003; 125:5839. [PubMed: 12733925]
- (24). (a) For a review of the Stille reaction in polyene natural product synthesis, see: Pattenden G, Sinclair DJ. J. Organomet. Chem. 2002; 653:261. For a selected application of the Suzuki reaction in natural product synthesis, see: ref 8e.
- (25). (a) Mori T, Takahashi K, Kashiwabara M, Uemura D, Katayama C, Iwadare S, Shizuri Y, Mitomo R, Nakano F, Matsuzaki A. Tetrahedron Lett. 1985; 26:1073.(b) Henkel T, Zeeck A. Liebigs Ann. Chem. 1991:367.
- (26). For use of a (2*E*,4*E*)-5-bromo-2,4-dienoic ester in a cross coupling reaction, see : Crombie L, Horsham MA, Jarrett SRM. J. Chem. Soc. Perkin Trans. 1. 1991:1511. See also ref 10b, 10d.
- (27). Hadei N, Kantchev EAB, O'Brien CJ, Organ MG. Org. Lett. 2005; 7:3805. [PubMed: 16092880]
- (28). Giannerini M, Fananas-Mastral M, Feringa BL. J. Am. Chem. Soc. 2012; 134:4108. [PubMed: 22352853]
- (29). Henaff N, Whiting A. Org. Lett. 1999; 1:1137.
- (30). Garcia Garcia P, Lay F, Rabalakos C, List B. Angew. Chem. Int. Ed. 2009; 48:4363.
- (31). (a) Senapati BK, Gao L, Lee SI, Hwang GS, Ryu DH. Org. Lett. 2010; 12:5088. [PubMed: 20958087] (b) Webb MR, Donald C, Taylor RJK. Tetrahedron Lett. 2006; 47:549.(c) Webb MR, Addie MS, Crawford CM, Dale JW, Franci X, Pizzonero M, Donald C, Taylor RJK. Tetrahedron. 2008; 64:4778.(d) Yoshino M, Eto K, Takahashi K, Ishihara J, Hatakeyama S. Org. Biomol. Chem. 2012; 10:8164. [PubMed: 22971928]

Figure 1.
Examples of natural products containing a dienyl-carboxylate or –carbinol moiety.

Figure 2.

Lowest energy conrotatory transition structures for the ring opening of **2a**, **2b**, and **2c**. $\Delta H^\ddagger$ and $\Delta G^\ddagger$ are given in red below the corresponding transition structure. The enthalpy and free energy values in blue are the differences between the disfavored (not shown) and preferred transition states. Energies are given in kcal/mol.

* overall isolated yield from lactone **1a**. ** isolated yield after recrystallisation

Scheme 1.

Direct synthesis of halocyclobutenes and their ring-opening reactions.

Scheme 2.
Retrosynthetic strategy towards Inthomycin C.

Scheme 3.

Scope of cross-coupling reactions of (2*E*,4*E*)-5-bromo-2,4-dienoic derivatives **4c-d**.

Scheme 4. Suzuki cross-coupling of (2Z,4E)-5-chloro-dienoic derivatives **5b-d**.

Scheme 5.
Modular formal synthesis of Inthomycin C.