Published in final edited form as:

Biochem Biophys Res Commun. 2011 June 24; 410(1): 134–139. doi:10.1016/j.bbrc.2011.05.121.

Selective cell-surface labeling of the molecular motor protein prestin

Ryan M. McGuire¹, Jonathan J. Silberg^{1,2}, Fred A. Pereira^{1,3}, and Robert M. Raphael^{1,‡}

¹Department of Bioengineering, Rice University, Houston, TX 77251

²Huffington Center on Aging, Department of Biochemistry and Cell Biology, Rice University, Houston, TX 77251

³Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX 77030

Abstract

Prestin, a multipass transmembrane protein whose N- and C-termini are localized to the cytoplasm, must be trafficked to the plasma membrane to fulfill its cellular function as a molecular motor. One challenge in studying prestin sequence-function relationships within living cells is separating the effects of amino acid substitutions on prestin trafficking, plasma membrane localization and function. To develop an approach for directly assessing prestin levels at the plasma membrane, we have investigated whether fusion of prestin to a single pass transmembrane protein results in a functional fusion protein with a surface-exposed N-terminal tag that can be detected in living cells. We find that fusion of the biotin-acceptor peptide (BAP) and transmembrane domain of the platelet-derived growth factor receptor (PDGFR) to the N-terminus of prestin-GFP yields a membrane protein that can be metabolically-labeled with biotin, trafficked to the plasma membrane, and selectively detected at the plasma membrane using fluorescently-tagged streptavidin. Furthermore, we show that the addition of a surface detectable tag and a single-pass transmembrane domain to prestin does not disrupt its voltage-sensitive activity.

Keywords

biotinylation; charge transfer; electrophysiology; membrane protein; molecular motor; prestin; protein engineering

1. Introduction

In response to changes in transmembrane potential, mammalian sensory outer hair cells (OHCs) change length [1; 2; 3]. This motion enhances sound-induced vibrations within the cochlea and improves hearing sensitivity [4]. The transmembrane protein prestin is critical for this signal amplification process [5]. Prestin is a molecular motor within the plasma membrane that responds to changes in transmembrane potential, and the electrical activity of prestin is tightly coupled to changes in the length of OHCs [6; 7]. The physiological role of

© 2011 Elsevier Inc. All rights reserved.

[‡]Correspondent Footnote: Dr. Robert M. Raphael, Department of Bioengineering, Rice University, 6500 Main Street, Houston, TX, 77030, Fax: 713-348-5877, raphael@rice.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

prestin's electromechanical coupling (molecular piezoelectricity) function in hearing has been clearly delineated [8; 9]. Prestin responds to cellular depolarization by reducing cell length, whereas prestin increases cell length when cells become hyperpolarized. Currently, the molecular mechanism of voltage-to-length conversion within this multipass transmembrane protein is not understood.

In contrast to other widely studied molecular motors (*e.g.*, myosin, kinesin, and dynein), prestin does not require ATP or calcium for motor function[3; 10]. Prestin belongs to a family of anionic membrane transport proteins, solute carrier family 26 (SLC26), for which little structural information is available. Like all membrane proteins, prestin function requires proper insertion into the plasma membrane. This latter property makes prestin challenging to study upon heterologous expression in cultured cells, since transiently transfected cells exhibit heterogeneity in both the level of prestin expressed and the amount of prestin trafficked to the plasma membrane [11; 12]. In addition, mutation of prestin can alter these properties as well as electromechanical coupling, making it difficult to interpret the effects of amino acid substitutions or deletions[13]. Since mutagenesis coupled with functional analysis is widely used to study prestin[11; 14; 15; 16; 17; 18; 19; 20; 21], an ability to selectively identify, and potentially quantify, the membrane-localized fraction would greatly enhance the biochemical and biophysical information gleaned from these studies.

To establish a strategy for studying the function of membrane-localized prestin, we investigated whether prestin could be metabolically tagged in living cells with biotin, a vitamin that binds with high affinity[22; 23] to streptavidin-conjugated fluorophores. The lack of endogenous surface biotinylation in mammalian cells[24; 25], and the impermeability of cell membranes to streptavidin conjugates makes this strategy compatible with selective labeling and purification of surface expressed membrane proteins. We based our reporter construct on the most extensively studied endogenously biotinylated protein, the 1.3 S subunit of *Propionibacterium shermanii* transcarboxylase (commonly called PSTCD) [26]. The biotin acceptor peptide (BAP) sequence from PSTCD has been fused to the C-terminus of multiple cytoplasmic enzymes and demonstrated to undergo *in vivo* metabolic biotinylation in both bacteria and yeast[27]. Parrot and Barry extended this technology to metazoans[28], and they demonstrated that both secreted and simple, single-pass integral membrane proteins can be biotinylated [29]. One of their constructs, BAP fused to the N-terminus of the platelet-derived growth factor receptor (PDGFR) transmembrane domain, was used to introduce streptavidin conjugated contrast agents to the surface of tumor cells and enhance magnetic resonance imaging (MRI)[30]. Herein, we demonstrate the first use of BAP-PDGFR for extracellular biotinylation of a multi-pass transmembrane protein.

2. Materials and methods

2.1. Plasmid Construction

The 736 bp BAP-PDGFR reporter gene was synthesized from a 24 primer set, composed primarily of 50 bp oligonucleotides, designed using DNA Works (Helix Systems, NIH) and the PCR-based synthesis methods of Hoover and Lubkowski[31]. The gene includes a 5' NheI restriction site and a 3' HindIII restriction site, and it encodes the murine Ig κ -chain leader sequence, the hemagglutinin A (HA) epitope tag, a biotin acceptor peptide (BAP), the transmembrane domain from platelet-derived growth factor receptor (PDGFR), and a 19 amino acid cytoplasmic linker region. Figure 1A illustrates the protein fusion expressed by this construct. The BAP coding sequence was derived from base pairs 1-390 of the PinPoint Xa vector (Promega, Madison, WI). The murine Ig κ -chain leader sequence, which targets the fusion to the secretory pathway[32], and the PDGFR transmembrane domain, which ensures exoplasmic display of BAP [33], were isolated from base pairs 737-799 and 907-

1053 (respectively) of the pDisplay vector (Invitrogen, Carlsbad, CA). All nucleotide segments were codon optimized from their original sources using a human codon frequency table[31].

The pEGFP-N1 vector (Clontech, Palo Alto, CA), which served as a backbone for subsequent cloning, was modified through Quikchange mutagenesis to include a point mutation within GFP. This mutation (denoted A206K in the literature) is located at residue 207 of the Clontech vector (due to an insertion that enhances mammalian translation efficiency) and prevents non-specific interaction of the fusion proteins[34; 35]. The synthesized BAP-PDGFR gene fusion was digested and inserted into the NheI and HindIII sites of the pEGFP-N1 plasmid. Gerbil prestin (AF230376) was isolated (EcoRI and BamHI) from a previously described prestin-GFP fusion[36] and inserted downstream and in-frame with the BAP-PDGFR gene fusion. The resulting construct is referred to as BAP-prestin-GFP and the translated fusion protein sequence is shown in Figure 1B. The entire fusion was sequenced verified using an eight-primer set (Baylor Sequencing Core, Houston, TX). Prestin-GFP containing the A206K mutation was used for control activity (nonlinear capacitance) measurements. The pSec-BirA vector[29], which expresses *E. coli* biotin ligase (BirA) needed to biotinylate BAP-prestin-GFP was provided as a generous gift from Dr. Michael Barry.

2.2. Plasmid Expression and Cell Culture

HEK293 cells were grown in T75 flasks and 6-well plates containing DMEM supplemented with 10% BCS (Invitrogen, Carlsbad, CA), 1% penicillin-streptomycin, 14.3 mM HEPES, 16.1 mM NaHCO₃, and 10 mg/L (41 μ M) d-Biotin (Sigma-Aldrich, St. Louis, MO). Cells were transfected with prestin-GFP, or co-transfected with BAP-prestin-GFP and pSec-BirA plasmids, using 2 μ g of each plasmid DNA and 3 μ L/plasmid Fugene 6 transfection reagent (Roche, Indianapolis, IN) according to the manufacturer's directions. For electrophysiology and imaging studies, cells were trypsinized and replated onto #1.5 coverslips or MatTek dishes (MatTek Corp., Ashland, MA) 8–12 hours after transfection and then evaluated 48 to 72 hours post-transfection.

2.3. Confocal Imaging

Biotinylation of BAP and localization of the BAP-prestin-GFP fusion to the membrane were verified using confocal microscopy 72 hours post-transfection. Transfected HEKs were incubated with 5 μ g/mL Alexa-633 conjugated streptavidin (Invitrogen, Carlsbad, CA) for 20 minutes at room temperature. Cells were washed with PBS and imaging was performed on an LSM 510 microscope (Zeiss, Thornwood, New York). GFP and the Alexa-633 conjugated streptavidin (SA-Alexa 633) were excited with 488 nm Argon laser light and 633 nm HeNe laser light, respectively. Using a secondary dichotic and separate band pass filters for each channel, GFP fluorescence was collected between 500–530 nm and SA-Alexa 633 fluorescence was collected between 650–710 nm. The pinhole for both channels was adjusted to produce an optical slice less than 13 μ m (2.9 and 2.1 Airy units for the GFP and SA-Alexa 633 channels, respectively). All imaging was performed using a 63X, 1.4NA objective lens.

2.4. Electrophysiology

Kimax-51 capillary tubes (1.5/0.8 mm OD/ID) were heated on a P-97 puller (Sutter, Novato, CA) to form fine tip micropipettes. The micropipettes were filled with pipette blocking solution (130 mM CsCl, 2 mM MgCl₂, 10 mM EGTA, 10 mM HEPES) and threaded onto a Ag⁺/AgCl electrode. Coverslips seeded at low cell density formed the base of a patch clamp chamber in which the cells were bathed in extracellular blocking solution (99 mM NaCl, 20 mM TEA-Cl, 2 mM CoCl₂, 1.47 mM MgCl₂, 1 mM CaCl₂, 10 mM HEPES), and an agar

bridge provided electrical continuity with the reference bath containing a Ag^+/AgCl reference electrode and pipette blocking solution. Blocking solutions[16] were titrated to pH 7.3 and dextrose added to obtain 300 ± 3 mOsm (Osmette A; Precision Systems, Natick, MA). Patch pipettes produced 2–4 M Ω resistance in open-bath and placement was controlled using a Burleigh PCS-6200 Micromanipulator (EXFO Life Sciences, Rochester, NY). On-cell seals in excess of 1 G Ω were obtained from healthy, single cells showing strong GFP fluorescence 48 hours post-transfection. Pipette capacitance was compensated, the cell membrane ruptured to establish whole-cell mode, and only cells with series resistance less than 10 M Ω and membrane resistance in excess of 1 G Ω retained for analysis. We apply an 800 Hz, 10 mV sine wave and measure the current response as DC holding potential is stepped from -140 to $+140$ mV in 2 mV increments (HEKA EPC 10 Plus Amplifier with 18-bit DAC). Using a software-based phase-sensitive detector, implemented in Patchmaster software (HEKA, Mahone Bay, NS), we determine the real and imaginary components of the admittance (the complex current response scaled by the command voltage) and calculate membrane capacitance (C_m) and membrane resistance (R_m)[37]. At each DC potential four complete sinusoidal voltage cycles occur and discrete capacitance values are calculated from the latter three. Capacitance versus voltage (DC holding potential) traces are acquired for each cell and fit to the first derivative of the two-state Boltzmann function (Eqn. 1) using MATLAB (MathWorks, Natick, MA).

$$C_m = \frac{Q_{\max} \left(\frac{ze}{kT} \right)}{\exp \left[\left(\frac{ze}{kT} \right) (V - V_{1/2}) \right] \left(1 + \exp \left[-(V - V_{1/2}) \left(\frac{ze}{kT} \right) \right] \right)^2} + C_{lin} \quad \text{Eqn. 1}$$

$Q_{\max}$ is the maximum nonlinear charge movement provided by prestin. $V_{1/2}$ is the voltage at the peak of the capacitance versus holding potential curve. z is the valence of charge movement by prestin and inversely proportional to the width of the NLC curve. Boltzmann's constant k , absolute temperature T , and the charge carried by an electron e , are all constant values. Since variation in cell size causes differences in the maximal charge transfer ($Q_{\max}$), the charge movement is normalized to C_{lin} , and computed on a cell-by-cell basis. This quantity, designated as charge density, has units of fC/pF[38]. Representative NLC curves are reconstructed from mean $Q_{\max}$, $V_{1/2}$ and z obtained from BAP-prestin-GFP or prestin-GFP expressing cells, and each result is normalized to maximal nonlinear capacitance ($Q_{\max}ze/4kT$) of the corresponding group.

3. Results

3.1. Rationale for protein design

Established biotinylation techniques[39; 40] are not suitable for detecting prestin localized to the plasma membrane, since the N- and C-termini of prestin are both intracellular[41]. This precludes use of direct BAP fusions to achieve extracellular labeling. Moreover, the predicted 12 transmembrane helices[16] of prestin are thought to be tightly packed[19], with high conservation of the entire transmembrane domain region among the eutherian clade[42]. This suggests that the insertion of BAP into a predicted surface exposed loop could adversely affect prestin folding and function. To provide an externally accessible biotinylation site without altering the primary sequence of the prestin hydrophobic core, we fused the BAP-PDGFR reporter to the N-terminus of prestin. Additionally, GFP was fused to the C-terminus of prestin to assay successful transfection and visualize intracellular protein localization. Figure 1A shows the predicted topology of the resultant BAP-prestin-GFP fusion protein.

3.2. Fluorescence imaging

We found that BAP-prestin-GFP is biotinylated and inserts into the plasma membrane, when co-expressed in human embryonic kidney (HEK) cells with the *Escherichia coli* biotin ligase, BirA. Confocal imaging shows that a streptavidin (SA) conjugated fluorophore (SA-Alexa 633) cannot permeate the cell membrane and therefore only labels the membrane-localized fraction of BAP-prestin-GFP in live cells (Figure 2A). In contrast, BAP-prestin-GFP is not biotinylated in the absence of BirA (*data not shown*). To establish if BAP-prestin-GFP is metabolically labeled with biotin, we removed pixels with low fluorescent intensity values (the lower 5% or 35% for GFP and SA-Alexa 633, respectively) and then visualized the remaining colocalized signal (pixels with non-zero fluorescent values in each channel). We found that nearly all the SA-Alexa 633 fluorescence coincided with GFP fluorescence (Figure 2B), indicating that the biotinylated BAP-prestin-GFP is properly trafficked and membrane localized.

3.3. Patch-clamp analysis of prestin function

We next investigated whether the BAP-prestin-GFP fusion retains prestin function. The ability of prestin to move charge within the membrane is tightly coupled to its motor function and has been established as a benchmark for prestin activity[43; 44]. We used the whole-cell patch clamp technique to control the voltage across the cell membrane and assayed transmembrane charge movement. In the absence of prestin, HEK cells show uniform, linear capacitance that is proportional to the size of the cell due to the dielectric properties of the plasma membrane[45]. When functional prestin is present in the plasma membrane, a nonlinear, bell-shaped capacitance versus voltage curve is detected[46]. This charge movement, analogous to the gating current in ion channels, is commonly described as nonlinear capacitance (NLC) and thought to result in part from a voltage-induced conformational change in prestin. Figure 3 shows NLC plots for BAP-prestin-GFP and prestin-GFP without the extra transmembrane domain and BAP reporter. These NLC plots were constructed from mean fit parameters (see experimental section for details) and therefore only display the average increase in capacitance (prestin-associated charge movement) above the linear capacitance (membrane surface area). BAP-prestin-GFP clearly is functional, since it shows bell-shaped NLC comparable to wild-type prestin lacking a BAP fusion. With BAP-prestin-GFP, the voltage at peak capacitance ($V_{1/2}$) and valence of charge movement (z) are indistinguishable from prestin-GFP. This confirms that the operating range and voltage sensitivity of prestin activity were not impacted by the BAP-PDGFR reporter fusion. There was a reduced charge density (Q_{max}/C_{lin}) with the N-terminal fusion that might reflect a lower trafficking and reduced surface expression compared to the unmodified prestin-GFP. This reduced surface expression will be advantageous for future biophysical studies examining single molecule tracking.

4. Discussion

The findings described herein provide the first evidence that a BAP-PDGFR reporter can be used with a multi-pass transmembrane protein for *in vivo* biotinylation and subsequent detection. This approach is particularly useful for identifying the membrane-localized fraction of a target protein or introducing a fluorescent label when commercially available antibodies are insufficient. For prestin, live cell labeling is not currently possible in the absence of BAP-PDGFR using simple affinity tags (*e.g.*, HA, FLAG, or myc tags) that are recognized by commercial antibodies. Intracellular tags have been used as N- or C-termini fusions to detect prestin within fixed, membrane-permeabilized samples. For cell surface detection of prestin, these tags have not been introduced within a surface-exposed region of prestin, due the intracellular localization of prestin's N- and C-termini and because all our attempted insertions have proven to be detrimental to prestin trafficking and function.

Furthermore, it remains challenging to develop antibodies directed at a surface-exposed region of prestin for live cell imaging, since extracellularly available epitopes of prestin are poorly defined and of low antigenicity.

Our findings provide direct evidence that biotinylated BAP-prestin-GFP localizes to the plasma membrane and is functional. While biotinylation was achieved metabolically by coexpressing BirA and BAP-prestin-GFP, this post-translational modification could be achieved through the addition of BirA and biotin to tissue culture cells expressing BAP-prestin-GFP. This latter approach may prove to be more efficient, since the amount of BirA and biotin can be easily varied in the growth medium to maximize labeling efficiency. The results described herein show that biotinylated BAP-prestin-GFP will be useful for future biophysical studies. This construct will enhance studies examining the relationship between changes in prestin sequence and function. In its current form, this reporter system will be useful for deconvoluting the effects of mutations on protein expression, translocation, and function. Furthermore, this system can be used to label membrane-localized prestin in live cells with fluorescent probes having larger molar absorptivity (extinction coefficient)[47] and higher quantum yield[48] than intrinsically fluorescent proteins. These brighter fluorophores are expected to enable quantitative fluorescent microscopy and single molecule tracking in future studies of prestin. In addition, this general approach is expected to be compatible with cell-surface labeling of other multi-pass transmembrane proteins, such as prestin homologs in the SLC26 protein family.

Research Highlights

- > Trafficking to the plasma membrane is required for prestin function.
- > Biotin acceptor peptide (BAP) was fused to prestin through a transmembrane domain.
- > BAP-prestin can be metabolically labeled with biotin in HEK293 cells.
- > Biotin-BAP-prestin allows for selective imaging of fully trafficked prestin.
- > The Biotin-BAP-prestin displays voltage-sensitive activity.

Acknowledgments

The authors thank Shirley Liu, Peter Nguyen, Crystal Stanworth, and Peng Zhai for technical support, and Dr. Michael Barry of the Mayo Clinic for donating the pSec-BirA vector. This work was supported by a Hamill Foundation grant (R.M.R. and J.J.S.), a Robert A. Welch Foundation grant C-1614 (J.J.S.), and through NIH-NIDCD R21 DC008134 (R.M.R. & F.A.P), R01 DC009622 (R.M.R. & F.A.P) and NSF Career BES 044379 (R.M.R.).

References

1. Brownell WE, Bader CR, Bertrand D, de Ribaupierre Y. Evoked mechanical responses of isolated cochlear outer hair cells. *Science*. 1985; 227:194–196. [PubMed: 3966153]
2. Ashmore JF. A fast motile response in guinea-pig outer hair cells: the cellular basis of the cochlear amplifier. *J Physiol*. 1987; 388:323–347. [PubMed: 3656195]
3. Kachar B, Brownell WE, Altschuler R, Fex J. Electrokinetic shape changes of cochlear outer hair cells. *Nature*. 1986; 322:365–368. [PubMed: 3736662]
4. Robles L, Ruggero MA. Mechanics of the mammalian cochlea. *Physiol Rev*. 2001; 81:1305–1352. [PubMed: 11427697]
5. Zheng J, Shen W, He DZ, Long KB, Madison LD, Dallos P. Prestin is the motor protein of cochlear outer hair cells. *Nature*. 2000; 405:149–155. [PubMed: 10821263]

6. He DZ, Zheng J, Kalinec F, Kakehata S, Santos-Sacchi J. Tuning in to the amazing outer hair cell: membrane wizardry with a twist and shout. *J Membr Biol.* 2006; 209:119–134. [PubMed: 16773497]
7. Spector AA, Deo N, Grosh K, Ratnanather JT, Raphael RM. Electromechanical models of the outer hair cell composite membrane. *J Membr Biol.* 2006; 209:135–152. [PubMed: 16773498]
8. Liberman MC, Gao J, He DZ, Wu X, Jia S, Zuo J. Prestin is required for electromotility of the outer hair cell and for the cochlear amplifier. *Nature.* 2002; 419:300–304. [PubMed: 12239568]
9. Dallos P, Wu X, Cheatham MA, Gao J, Zheng J, Anderson CT, Jia S, Wang X, Cheng WH, Sengupta S, He DZ, Zuo J. Prestin-based outer hair cell motility is necessary for mammalian cochlear amplification. *Neuron.* 2008; 58:333–339. [PubMed: 18466744]
10. Holley MC, Ashmore JF. On the mechanism of a high-frequency force generator in outer hair cells isolated from the guinea pig cochlea. *Proc R Soc Lond B Biol Sci.* 1988; 232:413–429. [PubMed: 2895478]
11. McGuire RM, Liu H, Pereira FA, Raphael RM. Cysteine mutagenesis reveals transmembrane residues associated with charge translocation in prestin. *J Biol Chem.* 2010; 285:3103–3113. [PubMed: 19926791]
12. Rajagopalan L, Organ-Darling LE, Liu H, Davidson AL, Raphael RM, Brownell WE, Pereira FA. Glycosylation regulates prestin cellular activity. *J Assoc Res Otolaryngol.* 2010; 11:39–51. [PubMed: 19898896]
13. Dallos P, Zheng J, Cheatham MA. Prestin and the cochlear amplifier. *J Physiol.* 2006; 576:37–42. [PubMed: 16873410]
14. Oliver D, He DZ, Klocker N, Ludwig J, Schulte U, Waldegger S, Ruppersberg JP, Dallos P, Fakler B. Intracellular anions as the voltage sensor of prestin, the outer hair cell motor protein. *Science.* 2001; 292:2340–2343. [PubMed: 11423665]
15. Matsuda K, Zheng J, Du GG, Klocker N, Madison LD, Dallos P. N-linked glycosylation sites of the motor protein prestin: effects on membrane targeting and electrophysiological function. *J Neurochem.* 2004; 89:928–938. [PubMed: 15140192]
16. Deak L, Zheng J, Orem A, Du GG, Aguinaga S, Matsuda K, Dallos P. Effects of cyclic nucleotides on the function of prestin. *J Physiol.* 2005; 563:483–496. [PubMed: 15649974]
17. Zheng J, Du GG, Matsuda K, Orem A, Aguinaga S, Deak L, Navarrete E, Madison LD, Dallos P. The C-terminus of prestin influences nonlinear capacitance and plasma membrane targeting. *J Cell Sci.* 2005; 118:2987–2996. [PubMed: 15976456]
18. Navaratnam D, Bai JP, Samaranayake H, Santos-Sacchi J. N-terminal-mediated homomultimerization of prestin, the outer hair cell motor protein. *Biophys J.* 2005; 89:3345–3352. [PubMed: 16113116]
19. Rajagopalan L, Patel N, Madabushi S, Goddard JA, Anjan V, Lin F, Shope C, Farrell B, Lichtarge O, Davidson AL, Brownell WE, Pereira FA. Essential helix interactions in the anion transporter domain of prestin revealed by evolutionary trace analysis. *J Neurosci.* 2006; 26:12727–12734. [PubMed: 17151276]
20. Kumano S, Tan X, He DZ, Iida K, Murakoshi M, Wada H. Mutation-induced reinforcement of prestin-expressing cells. *Biochem Biophys Res Commun.* 2009; 389:569–574. [PubMed: 19737539]
21. Toth T, Deak L, Fazakas F, Zheng J, Muszbek L, Sziklai I. A new mutation in the human pres gene and its effect on prestin function. *Int J Mol Med.* 2007; 20:545–550. [PubMed: 17786286]
22. Piran U, Riordan WJ. Dissociation rate constant of the biotin-streptavidin complex. *J Immunol Methods.* 1990; 133:141–143. [PubMed: 2212686]
23. Green NM. Avidin. *Adv Protein Chem.* 1975; 29:85–133. [PubMed: 237414]
24. Robinson BH, Oei J, Saunders M, Gravel R. [3H]biotin-labeled proteins in cultured human skin fibroblasts from patients with pyruvate carboxylase deficiency. *J Biol Chem.* 1983; 258:6660–6664. [PubMed: 6406485]
25. Chandler CS, Ballard FJ. Regulation of the breakdown rates of biotin-containing proteins in Swiss 3T3-L1 cells. *Biochem J.* 1988; 251:749–755. [PubMed: 3046602]
26. Samols D, Thornton CG, Murtif VL, Kumar GK, Haase FC, Wood HG. Evolutionary conservation among biotin enzymes. *J Biol Chem.* 1988; 263:6461–6464. [PubMed: 2896195]

27. Cronan JE Jr. Biotinylation of proteins in vivo. A post-translational modification to label, purify, and study proteins. *J Biol Chem.* 1990; 265:10327–10333. [PubMed: 2113052]
28. Parrott MB, Barry MA. Metabolic biotinylation of recombinant proteins in mammalian cells and in mice. *Mol Ther.* 2000; 1:96–104. [PubMed: 10933917]
29. Parrott MB, Barry MA. Metabolic biotinylation of secreted and cell surface proteins from mammalian cells. *Biochem Biophys Res Commun.* 2001; 281:993–1000. [PubMed: 11237761]
30. Tannous BA, Grimm J, Perry KF, Chen JW, Weissleder R, Breakefield XO. Metabolic biotinylation of cell surface receptors for in vivo imaging. *Nat Methods.* 2006; 3:391–396. [PubMed: 16628210]
31. Hoover DM, Lubkowski J. DNAWorks: an automated method for designing oligonucleotides for PCR-based gene synthesis. *Nucleic Acids Res.* 2002; 30:e43. [PubMed: 12000848]
32. Coloma MJ, Hastings A, Wims LA, Morrison SL. Novel vectors for the expression of antibody molecules using variable regions generated by polymerase chain reaction. *J Immunol Methods.* 1992; 152:89–104. [PubMed: 1640112]
33. Gronwald RG, Grant FJ, Haldeman BA, Hart CE, O'Hara PJ, Hagen FS, Ross R, Bowen-Pope DF, Murray MJ. Cloning and expression of a cDNA coding for the human platelet-derived growth factor receptor: evidence for more than one receptor class. *Proc Natl Acad Sci U S A.* 1988; 85:3435–3439. [PubMed: 2835772]
34. Tsien RY. The green fluorescent protein. *Annu Rev Biochem.* 1998; 67:509–544. [PubMed: 9759496]
35. Zacharias DA. Sticky caveats in an otherwise glowing report: oligomerizing fluorescent proteins and their use in cell biology. *Sci STKE.* 2002; 2002:PE23. [PubMed: 11997581]
36. Greeson JN, Organ LE, Pereira FA, Raphael RM. Assessment of prestin self-association using fluorescence resonance energy transfer. *Brain Res.* 2006; 1091:140–150. [PubMed: 16626645]
37. Lindau M, Neher E. Patch-clamp techniques for time-resolved capacitance measurements in single cells. *Pflugers Arch.* 1988; 411:137–146. [PubMed: 3357753]
38. Oliver D, Fakler B. Expression density and functional characteristics of the outer hair cell motor protein are regulated during postnatal development in rat. *J Physiol.* 1999; 519(Pt 3):791–800. [PubMed: 10457091]
39. Howarth M, Takao K, Hayashi Y, Ting AY. Targeting quantum dots to surface proteins in living cells with biotin ligase. *Proc Natl Acad Sci U S A.* 2005; 102:7583–7588. [PubMed: 15897449]
40. Mize GJ, Harris JE, Takayama TK, Kulman JD. Regulated expression of active biotinylated G-protein coupled receptors in mammalian cells. *Protein Expr Purif.* 2008; 57:280–289. [PubMed: 18042400]
41. Zheng J, Long KB, Shen W, Madison LD, Dallos P. Prestin topology: localization of protein epitopes in relation to the plasma membrane. *Neuroreport.* 2001; 12:1929–1935. [PubMed: 11435925]
42. Okoruwa OE, Weston MD, Sanjeevi DC, Millemon AR, Fritzsche B, Hallworth R, Beisel KW. Evolutionary insights into the unique electromotility motor of mammalian outer hair cells. *Evol Dev.* 2008; 10:300–315. [PubMed: 18460092]
43. Santos-Sacchi J. Reversible inhibition of voltage-dependent outer hair cell motility and capacitance. *J Neurosci.* 1991; 11:3096–3110. [PubMed: 1941076]
44. Ashmore JF. Forward and reverse transduction in the mammalian cochlea. *Neurosci Res Suppl.* 1990; 12:S39–S50. [PubMed: 2243636]
45. Solsona C, Innocenti B, Fernandez JM. Regulation of exocytotic fusion by cell inflation. *Biophys J.* 1998; 74:1061–1073. [PubMed: 9533718]
46. Dallos P, Fakler B. Prestin, a new type of motor protein. *Nat Rev Mol Cell Biol.* 2002; 3:104–111. [PubMed: 11836512]
47. Patterson G, Day RN, Piston D. Fluorescent protein spectra. *J Cell Sci.* 2001; 114:837–838. [PubMed: 11181166]
48. Panchuk-Voloshina N, Haugland RP, Bishop-Stewart J, Bhalgat MK, Millard PJ, Mao F, Leung WY. Alexa dyes, a series of new fluorescent dyes that yield exceptionally bright, photostable conjugates. *J Histochem Cytochem.* 1999; 47:1179–1188. [PubMed: 10449539]

Fig. 1.

Predicted BAP-prestin-GFP fusion protein topology. *A*, The biotin acceptor peptide (BAP) was fused to the exoplasmic face of the platelet-derived growth factor receptor (PDGFR) transmembrane domain and attached to a 20 amino acid cytoplasmic linker. This engineered reporter gene and the 12-pass motor protein prestin were cloned to the pEGFP-N1 plasmid (Clontech) to form a fusion (BAP-prestin-GFP) with a convenient external binding moiety. The protein segment from the BAP-PDGFR reporter gene is shown in dashed outline and the transmembrane helices of the prestin-GFP fusion are numbered. BAP can be cleaved from PDGFR by Factor Xa protease and prestin can be liberated from PDGFR by severing the cytoplasmic linker with Enterokinase. *B*, Translated sequence of BAP-prestin-GFP. The

BAP-PDGFR reporter encompasses amino acids 1–237 (bold), prestin 244–987 (gray), and GFP 995–1233 (italics). The BAP-PDGFR reporter contains a leader sequence (amino acids 1–21), hemagglutinin tag (22–30), BAP (35–164), PDGFR transmembrane domain (169–216), and cytoplasmic linker (218–237). Biotinylation occurs at Lys 122 (box). The proteolytic cleavage sites (underlined) allow cleavage of the fusion following residue 164 or 234 by Factor Xa or Enterokinase, respectively. GFP contains A207K (residue 1201 of overall fusion) modification to prevent formation of non-obligate oligomers.

Fig. 2. BAP-prestin-GFP localizes to the cell membrane. *A*, Brightfield, GFP, SA-Alexa 633, and overlay images (counter-clockwise from upper right) for HEK cells co-transfected with pSec-BirA and BAP-prestin-GFP plasmid DNA. The streptavidin conjugated dye selectively labels the membrane localized fraction of BAP-prestin-GFP and does not non-specifically interact with untransfected HEKs. Scale bar 20 μ m. *B*, After thresholding to remove low intensity pixels, an overlay of the remaining colocalized signal demonstrates efficient biotinylation of BAP-prestin-GFP. 92% of pixels with SA-Alexa 633 fluorescence also have GFP signal, and these are displayed in panel *B*. Precise one-to-one correspondence was not

observed due to slight non-specific background staining and photomultiplier tube (PMT) detector noise.

Fig. 3.

BAP-prestin-GFP retains activity. NLC curves constructed from mean $V_{1/2}$, z , and Q_{max} (see experimental section) are normalized to the peak capacitance of each group. BAP-prestin-GFP shows normal $V_{1/2}$ and z , and maintains an ability to move charge despite reduction in charge density (Q_{max}/C_{lin}) compared to prestin-GFP ($p < 0.05$). BAP-prestin-GFP ($n=8$): $V_{1/2} -84.8 \pm 3.7\text{mV}$, $z 0.66 \pm 0.03$, $Q_{max}/C_{lin} 3.15 \pm 0.29\text{fC/pF}$. Prestin-GFP ($n=12$): $V_{1/2} -75.1 \pm 5.1\text{mV}$, $z 0.68 \pm 0.01$, $Q_{max}/C_{lin} 18.01 \pm 2.87\text{fC/pF}$. All values are mean $\pm$ SE.