Published in final edited form as:

J Phys Chem A. 2013 April 25; 117(16): 3414–3427. doi:10.1021/jp401477m.

An Electron Paramagnetic Resonance Spectroscopic Study of Copper Hopping in Doped Bis(L-histidinato)cadmium Dihydrate

Michael J. Colaneri^{†,*}, Jacqueline Vitali^{‡,*}, and Kristin Kirschbaum[¶]

[†]Department of Chemistry and Physics, State University of New York at Old Westbury, Old Westbury, New York 11568

[‡]Department of Physics, Cleveland State University, Cleveland, Ohio, 44115

[¶]Department of Chemistry, University of Toledo, Toledo, Ohio 43606

Abstract

Electron Paramagnetic Resonance (EPR) spectroscopy was used to study Cu(II) dynamic behavior in a doped biological model crystal; bis(L-histidinato)cadmium dihydrate, in order to gain better insight into copper site stability in metalloproteins. Temperature dependent changes in the low temperature X-band EPR spectra became visible around 100 K and continued up to room temperature. The measured 298 K g-tensor (principal values: 2.17, 2.16, 2.07) and copper hyperfine coupling tensor (principal values: −260, −190, −37 MHz) were similar to the average of the 77 K tensor values pertaining to two neighboring histidine binding sites. The observed temperature dependence was interpreted using Anderson's theory of motional narrowing, where the magnetic parameters for the different states are averaged as the copper rapidly hops between sites. The EPR pattern was also found to undergo a sharp sigmoidal-shaped, temperature dependent conversion between two species with a critical temperature $T_c \approx 160$ K. The species below T_c hops between the two low temperature site patterns, and the one above T_c represents an average of the molecular spin Hamiltonian coupling tensors of the two 77 K sites. In addition, the low and high temperature species hop between one another, contributing to the dynamic averaging. Spectral simulations using this 4-state model determined a hop rate between the two low temperature sites $\nu_{h4} = 4.5 \times 10^8 \text{ s}^{-1}$ and between the low and high temperature states $\nu_{h2} = 1.7 \times 10^8 \text{ s}^{-1}$ at 160 K. An Arrhenius relationship of hop rate and temperature gave energy barriers of $\Delta E_4 = 389 \text{ cm}^{-1}$ and $\Delta E_2 = 656 \text{ cm}^{-1}$ between the two low temperature sites, and between the low and high temperature states, respectively.

Keywords

Anderson; Arrhenius; dynamics; temperature; histidine; crystal; structure; rate; energy

Introduction

Electron Paramagnetic Resonance (EPR) spectroscopy was used to study dynamic behavior in a biological model crystal; bis(L-histidinato)cadmium dihydrate, where doped Cu(II) was found to undergo transitions between conformational states. Anderson's theory of motional

*Authors to whom correspondence should be addressed. colanerim@oldwestbury.edu Mailing Address: Department of Chemistry and Physics, SUNY at Old Westbury, Old Westbury, NY 11568, USA, Tel: (516) 876-2756. j.vitali@csuohio.edu Mailing Address: Department of Physics, Cleveland State University, Cleveland, OH 44115, USA, Tel: (216) 687-2431.

Supplementary Information Available:

(S1) CIFs for the structures at 130 K and 200 K.

This material is available free of charge via the Internet at <http://pubs.acs.org>.

narrowing¹, where the magnetic parameters for the individual sites are averaged as the copper rapidly hops between them, was used to interpret the temperature dependence of the EPR spectra. A 4-state hopping model involving two species was employed; a lower temperature species that averages predominantly over two low temperature spectral patterns and a higher temperature species which represents the averaging of the 77 K molecular **g** and copper hyperfine (**A^{Cu}**) tensors. A sigmoidal, temperature dependent conversion between low and high temperature species was also found to affect the dynamic process. The successful application of Anderson's theory advances our understanding of the characteristics of metal site dynamics, and provides fundamental insight into copper site stability in this system. In addition, this work may be useful in interpreting features of metal ion movement in recently characterized copper metalloproteins^{2,3}.

The crystal structure of the host, bis(L-histidinato)cadmium dihydrate, has been previously determined first by X-ray⁴ and later by neutron diffraction methods⁵ at room temperature. Crystals have space group $P4_32_12$, with cell constants $a=b=7.397$ Å and $c=30.53$ Å. In this space group, the **a** axis and **b** axis are indistinguishable, which are denoted here as **a(b)**. The unit cell consists of four cadmium ions and eight histidine molecules. Figure 1A depicts^{6,7} the nitrogen co-ordination around the cadmium which is described as a slightly flattened tetrahedron, where two histidine molecules bind the metal. The histidine molecules are related by a two-fold rotation axis positioned along the diagonal of **a** and **b** (or **a+b**), upon which the cadmium sits. Each histidine is metal ligated through the amide nitrogen (Cd-N bond length=2.287 Å), the imidazole nitrogen (Cd-N bond length= 2.290 Å) and one carboxyl oxygen (Cd-O bond length=2.480 Å).

Our previous single crystal EPR study on ⁶³Cu²⁺-doped bis(L-histidinato)cadmium dideuterate at 77 K revealed two copper site patterns in the **a(b)c** reference planes that are related by the **a+b** two-fold axis.⁸ Rotational EPR measurements determined the **g**-tensor and **A^{Cu}** tensor at 77 K and Electron Spin Echo Envelope Modulation (ESEEM) experiments at 4.2 K were used to obtain the hyperfine and quadrupole coupling tensors of the distant ¹⁴N of the imidazole of the copper ligated histidine.⁸ Based on the alignment of **g** and **A^{Cu}** tensors, and the quadrupole coupling tensor of the remote ¹⁴N nucleus, the copper binding site shown in Figure 1B was postulated. The Cu(II) binds stronger to the amide nitrogen (N1) and imidazole nitrogen (N2) of one of the two histidines related by the **a+b** symmetry axis, and has a weaker interaction with the other. The O1 oxygen of the close histidine is nearly axial to the N1-Cu-N2 basal plane. The unpaired electron mainly occupies the copper $d_{x^2-y^2}$ orbital oriented by this coordination geometry. At that time, an interpretation of the spectral superhyperfine splittings as originating from the two nitrogen ligands and two near protons was offered even though there was concern for crystal twinning and the possibility of a temperature induced phase change to a lower symmetry space group. This study also reported a dramatic difference in the room temperature EPR and raised the possibility that it arises from the average of the 77 K measured tensors pertaining to the two **a+b** symmetry-related histidine binding sites.

Shortly after this report, a similar system, Cu²⁺-doped Zn²⁺-(D,L-histidine)₂ pentahydrate, was investigated using single crystal EPR, X-ray crystallographic and calorimetric analysis.⁹ The two histidine molecules related by the crystallographic C₂ symmetry were postulated to form bidentate ligands to copper which replaced the host zinc ion. Here too, the EPR spectra exhibited a marked temperature dependence characterized by a transition between separate pairs of low temperature patterns and a collapsed region representing the averaged pattern with a $T_c \approx 268$ K. This was interpreted as a consequence of copper hopping between the four states mediated by vibrational stretching of the four copper-nitrogen ligand bonds. A continuous transition from the 80 K pattern pair to the room temperature pattern was explained using Anderson's theory¹ as a dynamic averaging of the two symmetry-related

low temperature EPR patterns with each other and with two unobserved high temperature patterns. An empirically derived, complex sigmoidal-like order parameter, centered slightly lower than T_c , defined the temperature dependent inclusion of the two unobserved high temperature patterns into the averaging process. By employing this dynamic model and using the copper nuclear spin ($I=3/2$; $m_I = 3/2, 1/2, -1/2, -3/2$) associated line-width dependence in the EPR single crystal spectrum when the external magnetic field $\mathbf{H}$ was directed along the $\mathbf{b}$ -axis ($\mathbf{b} // \mathbf{H}$), a hop rate $\nu_h \approx 2.5 \times 10^{10} \text{ s}^{-1}$ at 268 K was found. The authors employed transition state theory to determine the activation energy between the four averaging states as 1000 cm^{-1} . They further concluded that the dynamics of a water molecule, hydrogen bonded to the coordinating histidine amide group, was cooperatively linked to the copper dynamics, and contributed to the non-Boltzmann temperature dependent changes and sharp transition temperature in the EPR spectra. This was based partly on comparing X-ray diffraction determined crystal structures at 150 K and 293 K, where a significant disorder in this water was found, and on Differential Scanning Calorimetric measurements, which showed a broad signal centered slightly lower than T_c , and which correlated with the temperature dependence of the EPR spectral linewidths.

In view of these earlier results, a temperature dependent EPR and crystallographic investigation of Cu^{2+} -doped bis(L-histidinato)cadmium dihydrate was warranted for three reasons. First, given the larger ionic radius of cadmium versus zinc in these two histidine models and the different freedom of motion this afforded to the doped copper ion between its two histidine binding partners, it was of interest to evaluate how the larger space provided by the displaced cadmium in the present crystal would affect the site stability, transition temperature, hop rate and energy barrier between states. Second, since the bis(L-histidinato)cadmium dihydrate crystal structure also contains a water hydrogen bonded to the amide nitrogen of histidine, a study of this system was well suited to test whether structural disorder is an essential component for the dynamic behavior of copper in these systems. And finally, this study provides an opportunity to characterize copper site dynamics in a metal-histidine model for only the second time. It is possible that results stemming from this and similar work may provide useful information concerning the mechanism of copper transport along protein histidine side chains as was recently found in structural studies of caddie proteins^{3b}.

Experimental Methods

Crystalline samples of bis(L-histidinato)cadmium dihydrate were grown in the absence or presence (doped) of ~1% Cu(II) carbonate in normal or isotopically enriched solutions as described in earlier work⁸.

Diffraction data at 130 K and 200 K were measured on a host crystal using a SMART Platform diffractometer with 1 K CCD area detector and Mo $K\alpha$ graphite-monochromated radiation ($\lambda = 0.71073 \text{ \AA}$). Crystal data are given in Table 1. The cell constants are based on the refinement of the XYZ-centroids of reflections above $20 \sigma(I)$. The intensity data were corrected for absorption using SADABS¹⁰. The structures were refined using the Bruker SHELXTL software package¹¹. The room temperature neutron diffraction structure⁵ was the starting model for both refinements. The final anisotropic full-matrix least-squares refinement on F^2 with 155 variables converged at $R1 = 2.52\%$, $wR2 = 6.24\%$, goodness-of-fit = 1.197 for all data (2039 independent reflections) in the 130 K structure and at $R1 = 2.47\%$, $wR2 = 6.07\%$, goodness-of-fit = 1.174 for all data (2150 independent reflections) in the 200 K structure. Data collection and structure refinement details are given in Table 1. Further details, including atomic parameters, complete distances and angles and hydrogen bonds are found in the CIFs provided as Supporting Information.

For EPR measurements on copper-doped crystals, the crystallographic **a(b)** and **c** axes were selected as the reference axes for the external magnetic field direction. The samples were Duco cemented to the end of glass pipettes, aligned under a Leica StereoZoom microscope and placed into conventional 3 by 4 mm EPR tubes. These tubes were fixed through a Bruker goniometer which was mounted on an Oxford Instruments ESR 900 cryostat system and inserted inside the EPR microwave cavity. An Oxford ITC 503S controlled the temperature. Sample position temperatures were double checked for accuracy using a LakeShore Monitor equipped with a Cernox sensor. EPR spectra were obtained with a Varian E-109 X-band spectrometer, whose operating microwave frequency and magnetic field strength setting were monitored with an Agilent 53150A Frequency Counter and F.W. Bell 7010 Gaussmeter, respectively. The spectrometer was interfaced to a Dell Dimension 2400 PC using a National Instruments 6024E PCI card for analog/digital EPR signal conversion and the Gaussmeter digital output was connected to the PC's serial port. Homewritten code based on the LabView language, including commercial code supplied with the Bell Gaussmeter, was used to acquire the EPR signals as a function of magnetic field strength. For the room and temperature dependent EPR experiments, the acquired spectra were integrated in order to facilitate their analysis as discussed below. This was accomplished by the cwEPR program, which was made available by the pulsed-EPR facility at the Albert Einstein College of Medicine. EasySpin¹² was used to fit and simulate the superhyperfine splittings in the 77K spectra. The theoretical chemistry Gaussian 03W package¹³ was used to perform Density Functional Theory (DFT) level calculations. Peakfit (v4.12) was employed to fit Gaussian and Voigt G/L function peak curves and linewidths to the integrated EPR spectra at certain crystal orientations, and also to convolute lorentzian lineshape peaks that were determined using Anderson's theory of motional narrowing¹. Single crystal **g**-tensor and copper hyperfine tensor (**A**^{Cu}) parameters in the EPR molecular Spin Hamiltonian:

$$\mathcal{H} = \beta S \cdot g \cdot H - I \cdot A^{Cu} \cdot S - g_n \beta_n H \cdot I, \quad \text{Eq. 1}$$

with $S=1/2$, $I=3/2$ and remaining terms defined in the usual way¹⁴, were fit to resonance rotational data using local methods as described in earlier reports⁸.

Crystallographic and EPR Results

Low Temperature Crystal Structures

The X-ray determined crystal structures undertaken at 130 K and 200 K gave unit cell and atomic position parameters similar to the room temperature neutron diffraction structure, along with the expected increase in the temperature factors with an increase in temperature. No atomic disorder in the structure or structural phase change was observed. In particular, the water has similar environment at both temperatures as seen in Figure 2.

Symmetry of the Sites in the 77K EPR Spectra

Figure 3 shows the 77K EPR spectra obtained from ⁶³Cu²⁺ and D₂O isotopic-enriched crystals at **c**//**H**, **a(b)**//**H** and at **a+b**//**H**. The copper spectra observed with the external magnetic field along **a** and **b** are identical. At general crystal orientations, eight magnetically related copper hyperfine split, 4-line patterns are observed. The superhyperfine splittings arising from ligand nuclear couplings are discussed below. At orientations with the external field aligned in the crystal planes of the **a(b)c** reference system, four site patterns designated I, I', II and II' are observed, each representing a pair of overlapping site resonances and related by crystal symmetry operations. Sites I and II are related to each other by the **a+b** two-fold symmetry axis. I' and II' are related by the equivalent, two-fold symmetry axis that runs parallel to **a-b**. I and I' are related by a twofold screw axis running parallel to **a(b)** and

the II and II' patterns likewise arise from sites related by a two-fold screw axis parallel to **a(b)**. I and II are neighboring copper sites, as are I' and II'. The pairs of copper site resonances that remain overlapped with I and II, and I' and II' in the reference planes are related to each other by the two-fold rotation axes along **-(a+b)** and parallel to the **b-a** directions, respectively. In Figure 3, at **a(b)//H**, the I and I' patterns stack together as well as the II and II' patterns, and at **a+b//H**, the I and II patterns stack on the low field side of the spectrum, and the I' and II' patterns stack on the high field side. All four coalesce into one 4-line pattern when the external field is directed along the crystal 4_3 screw axis, **c//H**. As reported earlier⁸, these EPR spectral features at 77 K are consistent with the point symmetry of the histidine in the structure.

Analysis of the 77K EPR SuperHyperfine Splittings

The 77 K EPR spectra obtained from crystals grown in native solution had either very complicated or unresolved ligand splittings depending on the sample orientation. Isotopic enriched (^{63}Cu , ^2D) samples were therefore used to improve the resolution by eliminating both the ^{65}Cu m_I split resonances and the couplings due to exchangeable protons. Hyperfine tensor components were well fit to superhyperfine patterns shown in bubbles in Figure 3 using EasySpin according to a model consisting of two strong and one weak ("2+1") ^{14}N ligand coupling and one non-exchangeable ^1H coupling. These are summarized in Table 2 along with theoretical predictions and proposed ligand assignments. Splittings were evaluated at three specific orientations of the crystal, and four specific copper complex orientations. These are **a(b)//H** for the two separate site patterns I and II, **c//H**, and for site I at **a+b//H**. The tabulated experimental isotopic couplings a_{iso} were determined from $a_{\text{iso}} = \frac{1}{3} \text{Trace} \{ \text{on-axis//H splittings} \}$, which is a valid estimate when off-diagonal tensor elements are small. The hyperfine theoretical predictions were done at two levels using the proposed copper site in Figure 1: a point-dipole calculation which approximates the copper orbital spin density and a quantum mechanical DFT/B3LYP level computation. Previous studies have shown that the DFT produced isotopic parameters for the ^{14}N ligands in copper amino acid complexes are poor models. Therefore, for comparative purposes, the experimental isotropic parameters (a_{iso}) were added to the diagonal elements of both the theoretical DFT and the point-dipole determined ^{14}N and ^1H anisotropic hyperfine tensors. With this caveat, Table 2 shows good agreement between experimental fit and calculated hyperfine splittings, supporting the ligand assignments. Referring to Table 2 and Figure 1, the near copper histidine amide (N1) and imidazole (N2) nitrogen ligand a_{iso} couplings of 29.8 MHz and 37.1 MHz, respectively, are similar to those previously reported by Electron Nuclear Double Resonance (ENDOR) studies for directly coordinated nitrogen in copper-doped amino acid crystal complexes (23.5 – 32.1 MHz)¹⁵. The more distant histidine amide (N1') coupling, 20 MHz, is significantly lower than the coupling to N1, and is at the lowest end of this range. This reduction can be attributed to the long N1'-Cu distance (2.6 Å) and the placement of this nucleus 0.75 Å out of the plane containing the copper $d_{x^2-y^2}$ unpaired orbital. The selection of N1' as the origin of this splitting over imidazole N2' was because its theoretical hyperfine components had a much better correspondence with the measured values. The resolved proton splitting was assigned to the C_β carbon-bound H1, as its relatively large a_{iso} of 10.1 MHz can be predicted using the results from a previous survey of ENDOR measured couplings in similar systems¹⁵. Using the Cu-N1- C_β -H1 dihedral angle ($\theta \approx 175^\circ$) with an empirical cosine-square formula found by Colaneri et al.¹⁵ gave an a_{iso} of 7.1 MHz, which is close to but somewhat lower than 10.1 MHz. However, the DFT calculated value $a_{\text{iso}} = 9.8$ MHz confirms this assignment. The overall good agreement between the observed and theoretical splittings supports the proposition that the copper binds tighter to one histidine through N1 and N2 in the equatorial plane and interacts weaker with the other through N1' and possibly O1' out of this plane. The proposed copper site is

positioned 0.41 Å from the displaced cadmium ion location and is located about 0.80 Å distant from the **a+b** axis symmetry-related binding site.

The Room Temperature EPR

Figure 4 compares the integrated EPR spectra recorded at 80 K and 298 K at sample orientations: **c//H** and **a(b)//H**, respectively, for native grown crystals along with PeakFit simulations. Significant differences were observed between the room temperature (298 K) and 77 K acquired spectra. At **c//H** the room temperature copper hyperfine splitting reduced to 63 G from 101 G at 77 K and the *g*-value shifted from 2.161 at 77 K to a slightly lower value of 2.15 at 298 K. At **a(b)//H** the site I and II copper 4-line patterns at 77 K had respectively, splittings of 100 G and 77 G, and *g*-values of 2.180 and 2.107. These collapse into one 4-line spectrum of two stacked patterns; designated as (I_{rt} , II_{rt}) and (I'_{rt} , II'_{rt}), with a copper hyperfine splitting of 60 G and a *g* = 2.12 at 298 K. Note that the room temperature copper splittings and *g*-value are not the expected average of those observed at 77 K. The reason for this is addressed below. We propose that the associated patterns I_{rt} and II_{rt} , and the patterns I'_{rt} and II'_{rt} represent the molecular tensor averaged species for sites related by the 2-fold axes about the **a+b** and **a-b** directions, respectively. These pairs separate from one another when **H** is directed off-axis in the **a(b)** plane but remain overlapped in the **a(b)c** plane and thus follow the point group symmetry of the host cadmium ion. The **a(b)//H** room temperature pattern clearly exhibits a copper m_I dependent broadening which is attributed to a dynamic process with averaging over states⁹. Rotational EPR measurements were conducted at room temperature and the copper hyperfine resonance fields were recorded as a function of applied field orientation in the crystal reference planes. These are plotted in Figure 5. Least-square fit of **g** and **A^{Cu}** hyperfine tensors in Eq. 1 to this data are listed in Table 3A. The sign of the largest hyperfine principal component was assumed negative in order to be consistent with our previous study⁸. The choice among the alternate signs for the tensor direction cosines was decided by matching the observed room temperature Q-band EPR powder spectrum parameters⁸. The directions of the principal g_{max} , g_{mid} and g_{min} values (and the principal **A^{Cu}** values) are found to be aligned with the **a+b**, **c** and **a-b** directions, respectively. The room temperature **g** and copper hyperfine tensors listed in Table 3A are unusual for $d_{x^2-y^2}$ copper model complexes¹⁶. They are more comparable with the room temperature tensors reported in Cu^{2+} -doped Zn^{2+} -(D,L-histidine)₂ pentahydrate⁹ and in copper-doped tutton salt crystals undergoing dynamic Jahn-Teller distortions^{17,18}.

Included in Table 3A are the average of the 77 K **g** and ⁶³Cu hyperfine tensors reported by Colaneri and Peisach⁸ over the two **a+b** axis neighboring binding sites. Also, reproduced in Table 3B are the room temperature **g** and ⁶³Cu hyperfine tensors previously published for Cu^{2+} -doped Zn^{2+} -(D,L-histidine)₂ pentahydrate⁹ as well as the average of the 80 K measured tensors over the C2 axis which relates the two histidines binding to copper in this system. The close correspondence in Table 3 between the averaged 77 K (80 K) tensor principal values and directions with the room temperature tensors found for two different histidine systems suggest the validity of this relationship.

The Temperature Dependence of the EPR Spectra

Temperature dependencies of the low temperature EPR spectrum begin around 100 K and continue up to room temperature. Figure 6A portrays how the integrated EPR spectrum at **c//H** changes with temperature from near 70 K up to room temperature. In general, the low temperature peaks broaden, slightly shift in resonance field, and lose intensity as the temperature is raised. Experiments performed at **c//H** and at other orientations clearly correlate this loss of intensity with the growth of the high temperature spectral pattern. This is shown for example in Figure 6B where the EPR spectra shows two distinct inter-converting patterns as the temperature varies over a relatively narrow range: 155 K to 180

K. Peakfit simulations of the integrated EPR spectrum at $\mathbf{c}/\mathbf{H}$, as displayed in Figure 7A, were used to determine the relative population of the low temperature copper pattern as it transforms into the high temperature pattern. The solid curve in Figure 7B traces out a simple sigmoid function $n_{LT} = 1 - 1 / \{ 1 + e^{-(T-T_c)/\Delta T} \}$, where n_{LT} is the population of the low temperature pattern. Fit parameters $T_c = 163$ K and $\Delta T = 19$ K explain well how the PeakFit curve amplitude of the lowest field line of the low temperature pattern depends on temperature, even though a small amount ($<15\%$) appears to persist at temperatures higher than 220 K. The 77 K pattern lines shift toward the 298 K resonance positions as their peaks broaden. But how these features systematically vary with temperature could not be uniquely determined at $\mathbf{c}/\mathbf{H}$ due to the considerable spectral overlap and changing populations of the two patterns. The most reliable PeakFit simulation shown in Figure 7A is found at 160 K where a low to high temperature species population ratio is near 1:1. Here the best fit occurs with matching full-width, half-maximum linewidth of ~ 70 G for the two sets of outer lines and of ~ 50 G for the two sets of inner lines of the two species. The application of equal linewidths for all 8 resonant lines in PeakFit simulations results in a poor match to the spectrum. Similar features are observed for the EPR temperature dependence at other sample orientations. Figure 8 displays the temperature dependence at $\mathbf{a}+\mathbf{b}/\mathbf{H}$ (Figure 8A) and when $\mathbf{H}$ is directed 110° from the $\mathbf{c}$ -axis (Figure 8B). In both, the lowest field peaks can be observed to shift to higher field as they broaden and lose intensity concomitant with the growth of the high temperature pattern. The conversion between species also follows the functional dependence of Figure 7B. The resonant magnetic fields of the lowest field lines were followed as a function of temperature at these two sample orientations and are plotted in Figure 9. They both trace out non-linear curves until about 170 K, where, at $\mathbf{a}+\mathbf{b}/\mathbf{H}$, the peak overlaps the lowest field line of the growing high temperature pattern and the peak field dependence then follows that of the overlapped high temperature species. With $\mathbf{H}$ oriented 110° from $\mathbf{c}$ -axis, the peak center could not be detected higher than 180 K because of its reducing intensity and increasing line breadth. The analysis of these curves will be discussed in the theory section below.

Theoretical Analysis and Models

The basic theoretical approach follows that described by Dalosto et al.⁹ The important concepts are the following. The temperature variation observations were interpreted using a dynamic model based upon Anderson's theory of motional narrowing of spectral lines¹ The application of this theory provides important information on the molecular motions, specifically the rates and energy barrier between interacting states. Anderson's theory gives the shape and position of the resonance line when the frequency of a spin system jumps randomly between individual states.¹ The intensity distribution of the spectral pattern $I(\omega)$ is just the Fourier transform of a correlation function $\phi(\omega)$ related to the dynamics of the system:

$$I(\omega) = \int d\tau e^{i\omega\tau} \phi(\tau) \quad \text{Eq. 2}$$

Anderson¹ showed that in the absence of saturation, $\phi(\tau)$ becomes:

$$\phi(\tau) = \mathbf{W}_1 \bullet e^{(\tau(i\Omega + \Pi))} \bullet \mathbf{1}, \quad \text{Eq. 3}$$

where the components W_{Ii} of the vector $\mathbf{W}_1$ give the occupation probabilities of the states in equilibrium, $\mathbf{1}$ is a vector with all components equal to unity, and Ω is a diagonal matrix whose elements ω_i are the resonant absorption frequencies in the absence of dynamics. The matrix Π has elements $\Pi_{jk} = p_{jk}$ and $\Pi_{jj} = -p_{jk}$, with $j \neq k$ and where p_{jk} is the transition rate between the accessible states j and k . Anderson¹ and later Sack¹⁹ solved Eq. 2 incorporating Eq. 3 and obtaining for the spectral intensity distribution:

$$I(\omega) \approx 2\text{Re}\mathbf{W}_1 \bullet (i\omega\mathbf{E} - i\mathbf{\Omega} - \mathbf{\Pi})^{-1} \bullet \mathbf{1}, \quad \text{Eq. 4}$$

where $\omega\mathbf{E}$ is the unit matrix $\mathbf{E}$ times the constant ω . Eq. 4 is used below in the analysis of the EPR data in terms of dynamical models. For Cu(II) ions with nuclear spin $I=3/2$, we follow the assumption of Dalosto et al.⁹ that hopping transitions occur only between states with the same m_I , and the hop rate ν_h is independent of m_I . The transition rate p_{jk} is taken as the product $W_j\nu_h$, where W_j is the population of the departing state j and ν_h is the hopping rate. In a 4-state model with $S=1/2$, $I=3/2$, Eq. 4 becomes a 16×16 matrix, which reduces to 8×8 for a 2-state model.

The general consequences of the dynamic averaging on the EPR pattern are spectral narrowing by the shifting of line positions and changes in the line shape until an eventual collapse of the resonances occurs.^{1,9} This happens when the transition rate p_{jk} becomes comparable to the resonant frequency difference between the exchanging lines. Figure 10 displays a simulated EPR pattern in the presence of dynamic averaging based upon a 2-state model. The spectrum in Figure 10A shows two non-equivalent 4-line copper split patterns. As the hop rate between the two patterns increases, the lines with matching copper m_I states draw towards each other, broaden and finally collapse together as the hop frequency in magnetic field units become equivalent to their separation. It is important to note that Anderson developed Eq. 4 on the assumption that the spectral tensors of the averaging states were diagonal in the same reference frame. Eq. 4 therefore is not valid for the general case of motional averaging of molecular spin Hamiltonian tensors in different frames. This is why the patterns exhibited by the tensor averaged species at room temperature (I_{rt} , II_{rt}) in Figure 4 are not the spectral average of the patterns arising from the individual sites I and II at 77 K. However, since the I_{rt} and II_{rt} (and I_{rt}' and II_{rt}') patterns remain overlapped throughout the observations and their hopping transition $I_{rt} \leftrightarrow II_{rt}$ (and $I_{rt}' \leftrightarrow II_{rt}'$) does not directly impact the observations below ~ 160 K, this limitation in Eq. 4 was overlooked in the dynamic analysis of the I and II states. The application of Eq. 4 to determine the spectral intensity distribution gives Lorentzian line-shapes. These require convolution with a Gaussian function, which represents the line-shape in the absence of dynamics, in order to make a comparison with observed spectral lines. Figure 10B shows the consequence of dynamic averaging between sites with identical site patterns. Here no changes occur.

Dalosto et al.⁹ has derived the following formula based on Eq. 4 using a 2-state model that gives a relationship between the spectral linewidth in the presence of dynamics (ΔH_m) to the static linewidth (ΔH_0), the hop rate ν_h and the field separation between the hopping lines δH_m .

$$\Delta H_m(\theta, \phi) = \Delta H_0 + \{\sqrt{3}g(\theta, \phi)\mu_B/8h\nu_h\}\delta H_m(\theta, \phi)^2 \quad \text{Eq. 5}$$

The angular dependence (θ, ϕ) comes about because of the orientation anisotropy of the spectral patterns. Other terms have been defined by Dalosto et al.⁹.

Two-state Model: Hopping (ν_h) from Low to High Temperature Species

The extensive overlap of spectra from the different site patterns allowed only a limited use of the Eq. 5. Because EPR spectra of sites I, II, I' and II' stack at $c//H$ (Figure 3A), as does I_{rt} , II_{rt} , I_{rt}' and II_{rt}' , the temperature dependence can be analyzed according to an effective 2-state hopping model between the low and high temperature species, that is between $I \leftrightarrow II_{rt}$ and between the equivalent and overlapping $II \leftrightarrow I_{rt}$, $I' \leftrightarrow II_{rt}'$ and $II' \leftrightarrow I_{rt}'$. The reason is that that jumping between identical patterns; $I \leftrightarrow II$, $I' \leftrightarrow II'$, $I_{rt} \leftrightarrow II_{rt}$ and $I_{rt}' \leftrightarrow II_{rt}'$ at this orientation leave the spectrum unchanged (see Figure 10B and Dalosto et al.⁹), reducing the 4-state hopping to an effective 2-state model. Employing Eq. 5 with the PeakFit line

curve fits at 160 K reported in Figure 7A as well as the 80 K and 298 K spectra shown in Figure 4A, and using a departing population W_j of $1/4$ (see below for explanation) in $W_j v_{h2}$ leads to a low to high species hop rate (v_{h2}) of $\sim 1.2 \times 10^8 \text{ s}^{-1}$. However, the best match to the 160 K spectrum was achieved from a dynamic simulation by diagonalizing Eq. 4 using a slightly higher jump rate of $v_{h2} = 1.7 \times 10^8 \text{ s}^{-1}$ as described in Figure 11. Also displayed are the measured integrated EPR and a 1:1 composite spectrum consisting of the measured 77 K pattern and the 298 K EPR pattern. A comparison clearly shows the superiority of the dynamic model. The composite fails to reproduce the observed spectral narrowing and line broadening.

An effective 2-state dynamic model also explains the temperature induced EPR changes observed at **a+b//H**. As indicated in Figure 12A, at this orientation the lower field part of the integrated spectrum at 77 K is due to overlapped sites I and II, and sites I' and II' stack together at the higher field part. These sites hop between the corresponding stacked pairs of room temperature patterns, i.e., on the low field side, I_{rt} , II_{rt} and on the high field side, I_{rt}' , II_{rt}' . Therefore, essentially two separate hopping transitions affect the temperature dependence of the spectrum, one on the low field side; $I \leftrightarrow II_{rt}$ (along with the equivalent and overlapped $II \leftrightarrow I_{rt}$) and one on the high field side; $I' \leftrightarrow II_{rt}'$ (along with the equivalent and overlapped $II' \leftrightarrow I_{rt}'$). Since I and II, and I_{rt} and II_{rt} , are equivalent sites related by crystal symmetry, it is assumed that the hop rates between $I \leftrightarrow II_{rt}$ and $II \leftrightarrow I_{rt}$ are the same. The same goes for the primed site transitions. Diagonalizing Eq. 4 with $W_j = 1/4$ and using the same hop rate $v_{h2} = 1.7 \times 10^8 \text{ s}^{-1}$ that was found above at **c//H** produced a simulation that also best matched the observed integrated 160 K EPR spectrum at **a+b//H**. Shown in Figure 12B, this spectrum is a composition of the two unique dynamic simulations, i.e., due to jumps between $I \leftrightarrow II_{rt}$ and between $I' \leftrightarrow II_{rt}'$. The figure also depicts the measured, integrated EPR spectrum at 160 K and a 1:1 composite of the 77 K and the 298 K spectra. Here again, a simple addition of the low and high temperature patterns does a poor job explaining the observed spectral narrowing and broadening as compared to the dynamic model.

Four-state Model: Evidence for Hopping (v_{h4}) Between Neighboring Sites

At low temperature, with the magnetic field **H** oriented at 110° from **c** in the reference plane, the lowest field m_I line of site I becomes clearly resolved from its **a+b** related site II peaks, as well as those lines from other symmetry related sites. Figure 13A depicts the integrated EPR spectra at this orientation at 80 K and 298 K along with PeakFit simulations which were guided by line field positions determined in earlier work⁸ and from Figure 4. Figure 13B provides the integrated EPR spectrum measured at 160 K. Dynamic simulations performed using a 2-state hopping between $I \leftrightarrow II_{rt}$ with a rate $v_{h2} = 1.7 \times 10^8 \text{ s}^{-1}$ failed to reproduce the pronounced field shift of this low field resonance line as the temperature changes. In a 4-state dynamic model, the hopping states are: $I \leftrightarrow II$, $I \leftrightarrow II_{rt}$, $II \leftrightarrow I_{rt}$ and $I_{rt} \leftrightarrow II_{rt}$, as well as the corresponding primed states. We have assumed that the hopping rates are equivalent for $I \leftrightarrow II$ and $I_{rt} \leftrightarrow II_{rt}$, which is denoted as v_{h4} . The population of the leaving state at 160 K is $W_j = 1/4$ since all four patterns are equally present. Using this model and the hop rate $v_{h2} = 1.7 \times 10^8 \text{ s}^{-1}$ determined above, v_{h4} was adjusted in simulations using Eq. 4 to match the resonance field position of the lowest field m_I line of site I at 160 K. The position was best fit with $v_{h4} = 4.5 \times 10^8 \text{ s}^{-1}$. Figure 13B displays the dynamic simulation of the spectrum at 160 K using these quantities. This is a composite including all the dynamic transitions at this orientation. Also shown is the 1:1 composite spectrum of the integrated EPR measured at 77 K and 298 K. As shown, the dynamic model provides a much better match to the observed resonance position of the low field peak and to the overall spectral features, thus providing evidence for hopping between the neighboring I and II copper sites across the **a+b** symmetry axis.

Resonant Field Shifts as a Function of Hopping Rate (ν_h) for the Low Field Peak

Dynamic simulations performed using Eq. 4 determined the field dependence of the lowest resonance line of species I on the transition rate $p_{jk2} = W_j \nu_{h2}$ at two crystal orientations; $\mathbf{a} + \mathbf{b}/\mathbf{H}$ and when $\mathbf{H}$ is directed 110° from $\mathbf{c}$. Following the discussion above, the hopping at $\mathbf{a} + \mathbf{b}/\mathbf{H}$ can be considered as a 2-state jump. The lowest field peak of site I (LF in Figure 12A) shifts in magnetic field towards the same (lowest field) m_I line of the Π_{rt} pattern. How the simulated resonant field of this peak depends on transition rate p_{jk2} is plotted in Figure 14. The curve is non-linear and can be fit to a simple $H(\text{G}) = a + b(p_{jk2})^c$ function, with $a = 2815.6$, $b = 199$ and $c = 2.35$. Because the hopping initiates from the low temperature pattern, the population of the departing state changes according to the sigmoidal function in Figure 7B. The W_j in Eq. 4 therefore varies as $n_{\text{LT}}/2$ or from $1/2$ at 80 K to $1/4$ at 160 K. Using this dependence and the results displayed in Figures 9A and 14, a correlation between hop rate ν_{h2} and temperature was determined. This correlation will be analyzed below.

A slightly different approach was used to determine the dependence of the low field resonant line of site I on temperature when $\mathbf{H}$ is directed 110° from $\mathbf{c}$ (LF in Figure 13B). Here, 4-state model dynamic simulations between transitioning sites described above were performed incorporating the p_{jk2} transition rate at each specific temperature found above, and then adjusting the p_{jk4} transition rate to match the lowest field line resonant position displayed in Figure 9B. From the deduced p_{jk4} dependence on temperature and the W_j variation in Figure 7B, the hop rate ν_{h4} as a function of temperature was also determined and will be discussed below.

Discussion

At low temperature (77 K) the copper is strongly bound to one of the two histidine molecules related by the $\mathbf{a} + \mathbf{b}$ axis, and weaker to the other. The evidence comes from both the alignment of previously measured 77 K $\mathbf{g}$ and $\mathbf{A}^{\text{Cu}}$ tensors, and the 4.2 K remote nitrogen quadrupole tensor with the molecular site, and from the present analysis of the EPR superhyperfine splittings from isotopically enriched crystals. The splitting features could be explained by couplings to two strong and one weaker ^{14}N ligand, and a nearby non-exchangeable proton (Table 2). To our knowledge, this system represents the first characterized copper complex exhibiting a “2+1” nitrogen coordination. One would therefore anticipate that the stability of this complex is considerably less than the more commonly found, four-coordinate copper complexes. It is notable that the 77 K $\mathbf{g}$ and $\mathbf{A}^{\text{Cu}}$ tensor parallel direction principal values fall mid-range for previously characterized 3N1O copper model systems¹⁴ and thus provides no indication of its unusual coordination or unstable nature.

As the temperature rises above 100 K, the copper begins to jump between the two low temperature symmetry-related states I and II, swapping its major histidine partner and causing the 77 K EPR site patterns to approach their average. The spectral change begins at a relatively low temperature which signifies the instability of the 77 K complex. As the hop rate increases with temperature, the resonant lines reflect this dynamic effect by shifting and broadening towards a collapse of the averaging spectral lines. However, when the rate reaches a certain threshold, the low temperature averaging patterns convert into a high temperature species. This distinct conversion traces out a sigmoidal dependence with a $T_c \approx 160$ K over a narrow temperature range (Figure 7B). It is important to emphasize that, in general, the spectrum of the average of crystal tensors (or site patterns) is not equivalent to the spectrum arising from the average of the local or molecular tensors²⁰. We contend that the high temperature species ($\text{I}_{\text{rt}}, \text{II}_{\text{rt}}$) is the resonant pattern resulting from the average of the 77 K molecular $\mathbf{g}$ and $\mathbf{A}^{\text{Cu}}$ coupling tensors. Support for this comes from the close agreement of the measured room temperature tensors to the average of the corresponding

low temperature tensors in both this system and in Cu^{2+} -doped Zn^{2+} -(D,L-histidine)₂ pentahydrate (see Table 3). Even though the averaged tensors have slightly greater **g** and **A^{Cu}** principal values than those measured at room temperature, their principal directions are almost exactly aligned. We suggest that the small disparities in these quantities are the result of slight temperature-dependent changes in the potential energy surface.

As the low temperature states convert to the high temperature species they also hop between one another. So that at T_c a 4-state dynamic process exists between equally populated states I, II, I_{rt} and II_{rt} (and their primed state counterparts). The conformational sites and hopping pathways are depicted in Figure 15. As mentioned above, the $I_{rt} \leftrightarrow II_{rt}$ transition represents the average over the molecular tensors of the two sites and is therefore not governed by Eq. 4. However, since the I_{rt} and II_{rt} patterns remain overlapped and their hopping transition does not directly affect the analysis of the I and II states, we have assumed that (1) Eq. 4 can be applied to the dynamic analysis below $\sim T_c$, (2) the hopping rates between $I \leftrightarrow II$ and $I_{rt} \leftrightarrow II_{rt}$ are the same and (3) each state hops between two others, specifically; $I \leftrightarrow II$, $I \leftrightarrow II_{rt}$, $II \leftrightarrow I_{rt}$ and $I_{rt} \leftrightarrow II_{rt}$. For temperatures greater than T_c , the high temperature species dominates the spectrum as the intensity of low temperature pattern reduces to a smaller percentage. The remaining amount of low temperature pattern continues to hop between the high temperature species. This, added to the dynamic averaging of the molecular tensors of the two sites, causes the m_I dependent broadening observed in the room temperature EPR spectra.

In Figure 16A, an Arrhenius-type relationship $v_h = v_0 e^{(-\Delta E/kT)}$ characterizes the dependencies of the v_{h2} and v_{h4} hop rates on temperature. In this formula ΔE has been interpreted in past studies as the activation energy needed to overcome the barrier between hopping states.¹⁷ The energy barrier between the low and high temperature species $\Delta E_2 = 656 \text{ cm}^{-1}$ and between the two low temperature sites $\Delta E_4 = 389 \text{ cm}^{-1}$ were determined from the slopes of the lines in Figure 16A. These values are both lower than the energy barrier between conformational states found in copper-doped Zn^{2+} -(D,L-histidine)₂ pentahydrate (1000 cm^{-1})⁹ but higher than that associated with an axial-equatorial ligand bond switch found in copper-doped zinc tutton salt (120 cm^{-1})¹⁵. The potential energy profile displayed in Figure 16B summarizes the dynamic behavior of the copper complex in the current system. The blue curve depicts this profile for the low temperature states, site I and II, and the red depicts that for the high temperature states I_{rt} and II_{rt} . A sharp sigmoidal temperature dependent conversion between the low and high temperature profiles happens with a $T_c \approx 160 \text{ K}$.

Results from copper-doped bis(L-histidinato)cadmium dihydrate, can be compared with those from doped Zn^{2+} -(D,L-histidine)₂ pentahydrate⁹. In the zinc system, the authors proposed that the copper undergoes a dynamic pseudo-Jahn-Teller distortion mediated by the four nitrogens of the two ligated histidines, characterized by a transition from a 2-state to a 4-state hopping model slightly below $T_c \approx 268 \text{ K}$. The two high temperature states, although thermally unoccupied, nevertheless contribute to the model until a collapse of states occurs at T_c to become a 4-state averaged pattern. The hop rate at 268 K between the four states was determined to be $2.5 \times 10^{10} \text{ s}^{-1}$. This is a much higher rate than found in the present system ($v_{h4} = 4.5 \times 10^8 \text{ s}^{-1}$ and $v_{h2} = 1.7 \times 10^8 \text{ s}^{-1}$ at 160 K) and may be related to the cooperative lattice dynamics proposed for this site⁹. The authors further postulated that the copper dynamic behavior is correlated with a fluctuating disorder of a water molecule found in the crystal structure. Their transition temperature (268 K) is significantly higher than found in the copper-doped bis(L-histidinato)cadmium dihydrate (160 K), which means that the present system is a far more unstable complex. This is consistent with the much lower energy barrier found between interacting states 389 cm^{-1} and 656 cm^{-1} , compared to 1000 cm^{-1} found in doped Zn^{2+} -(D,L-histidine)₂ pentahydrate.

The bis(L-histidinato)cadmium dihydrate crystal structure also displays a water (O3 in Figure 1) that makes a hydrogen bond to the coordinated amino nitrogen of histidine.⁵ However, X-ray diffraction experiments at 130 K and 200 K (on the low and high side of the ~160 K transition temperature) displayed similar crystal structures and, as shown in Figure 2, almost identical water environments, which discounts the possibility that the motions of disordered water moderates the copper dynamic behavior. The distinct 1:1 conversion, displayed in Figure 6B, between low and high temperature copper states was not observed in Cu²⁺-doped Zn²⁺-(D,L-histidine)₂ pentahydrate⁹. Instead the authors report a sharp but continuous conversion slightly below T_c from the copper having two states to having four states available for the hopping transitions. In the present system, the conversion between distinct species therefore suggests a different dynamic mechanism. A possible explanation for this difference is a physical jumping of the copper ion between the two histidine binding positions. The larger ionic radius²¹ of cadmium (1.03 Å) versus zinc (0.69 Å) in the two systems provides the doped metal ion a much greater freedom of motion between the histidines. In addition, the copper, in a more unstable “2+1” coordination, may have a greater tendency to hop between the energetically equivalent, neighboring binding sites.

Recently determined protein crystal structures of Tyrosinase complexes with mutant caddie proteins found multiple copper sites depending on crystal-copper solution soak times.^{3b} Based on the structural evidence for the locations of these sites, a mechanism for Cu(II) transportation into the Tyrosinase active site via the caddie histidine side chains was proposed. Here, imidazole-histidine residue motion suggested by disorder in the structures was used to explain how one copper site can be brought into vicinity of another along a pathway to the dicopper active site. It is interesting that the closest approach found for any two coppers along a path to the active site (Protein Data Bank code 3AXO) is much greater (~5 Å) than the hop distance (0.8 Å) found in doped bis(L-histidinato)cadmium dihydrate. It remains to be explored on whether copper can jump rapidly over such large distances to nearby histidine binding sites.

Conclusion

Copper complex dynamic behavior was characterized in doped bis-(L-histidinato)cadmium dihydrate crystals using Anderson's theory of exchange narrowing¹. A distinct 1:1 conversion from a low temperature species that averages the 77 K patterns from neighboring copper sites into a high temperature pattern representing the average of the molecular spin Hamiltonian parameters of these sites was observed with a relatively low transition temperature T_c ≈ 160 K. Hop rates between the low and high temperature species (ν_{h2}) of $1.7 \times 10^8 \text{ s}^{-1}$ and between the two low temperature states (ν_{hd}) of $4.5 \times 10^8 \text{ s}^{-1}$ at 160 K were determined through dynamic spectral simulations using a 4-state (or effective 2-state) model. Arrhenius plots of the jump rate dependences on temperature gave energy barriers $\Delta E_2 \approx 656 \text{ cm}^{-1}$ and $\Delta E_4 \approx 389 \text{ cm}^{-1}$ between the interacting states. These findings are in contrast with previous results from copper-doped Zn²⁺-(D,L-histidine)₂ pentahydrate⁹ where the transition temperature (268 K), hop rate ($2.5 \times 10^{10} \text{ s}^{-1}$) and activation energy between states (1000 cm^{-1}) were all significantly higher. An explanation is the different stabilities and dynamic mechanisms of the copper complexes in the two systems. In the present system, copper was found to have an unusual “2+1” nitrogen coordination making the complex unstable relative to the four nitrogen coordination in the zinc crystal system. There was no evidence for any water disorder in the host bis(L-histidinato)cadmium dihydrate structure, discounting the contribution of water motions to the copper dynamics as was proposed in Zn²⁺-(D,L-histidine)₂ pentahydrate. In addition, the larger space of the replaced cadmium ion compared to zinc appears to plays an important role in regulating the copper dynamic behavior.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank SUNY/Old Westbury undergraduate student Ms. Brenda Marmol for her help in the early stages of the EPR analysis. The following grant is acknowledged for financial support: NIGMS 5S06GM008180 (to MJC). The computations were supported in part by an allocation of computing time from the Ohio Supercomputer Center.

References

1. Anderson PW. A Mathematical Model for the Narrowing of Spectral Lines by Exchange or Motion. *J. Phys. Soc. Jpn.* 1954; 9:316–339.
2. Steiner RA, Kooter IM, Dijkstra BW. Functional Analysis of the Copper-Dependent Quercetin 2,3-Dioxygenase. 1. Ligand-Induced Coordination Changes Probed by X-ray Crystallography: Inhibition, Ordering Effect, and Mechanistic Insights. *Biochemistry*. 2002; 41:7955–7962. [PubMed: 12069585] Kooter IM, Steiner RA, Dijkstra BW, van Noort PI, Egmond MR, Huber M. EPR characterization of the mononuclear Cu-containing *Aspergillus japonicus* quercetin 2,3-dioxygenase reveals dramatic changes upon anaerobic binding of substrates. *Eur. J. Biochem.* 2002; 269:2971–2979. [PubMed: 12071961]
3. (a) Matoba Y, Kumagai T, Yamamoto A, Yoshitsu H, Sugiyama M. Crystallographic Evidence That the Dinuclear Copper Center of Tyrosinase Is Flexible during Catalysis. *J. Biol. Chem.* 2006; 281:8981–8990. [PubMed: 16436386] Sendovski M, Kanteev M, Ben-Yosef VS, Adir N, Fishman A. First Structures of an Active Bacterial Tyrosinase Reveal Copper Plasticity. *J. Mol. Biol.* 2011; 405:227–237. [PubMed: 21040728] (b) Matoba Y, Bando N, Oda K, Noda M, Higashikawa F, Kumagai T, Sugiyama M. A Molecular Mechanism for Copper Transportation to Tyrosinase That Is Assisted by a Metallochaperone, Caddie Protein. *J. Biol. Chem.* 2011; 286:30219–30231. [PubMed: 21730070]
4. Candlin R, Harding MM. The Crystal Structure of Bis(histidino)cadmium Dihydrate. *J. Chem. Soc. (A)*. 1967:421–423.
5. Fuess H, Bartunik H. Neutron Diffraction Structure Determination of Bis-(L-histidinato)cadmium Dihydrate. *Acta Cryst.* 1976; B32:2803–2806.
6. ORTEP: A Fortran Thermal-Ellipsoid Plot Program for Crystal Structure Illustrations. Johnson, C.K. ORNL-3794 (Rev.), Union Carbide Corp., Oak Ridge Natl. Lab. Jun. 1965
7. Persistence of Vision Raytracer Pty. Ltd. 2006. <http://www.povray.org>
8. Colaneri MJ, Peisach J. A Single Crystal EPR and ESEEM Analysis of Cu(II)-Doped Bis(L-histidinato)cadmium Dihydrate. *J. Amer. Chem. Soc.* 1995; 117:6308–6315.
9. Dalosto SD, Calvo R, Pizarro JL, Arriortua MI. Structure, Disorder, and Molecular Dynamics in Zn(D,L-histidine)₂: EPR of Copper Ion Dopants, X-ray Diffraction, and Calorimetric Studies. *J. Phys. Chem. A*. 2001; 105:1074–1085.
10. Sheldrick, GM. SADABS. Göttingen, Germany: University of Göttingen; 1996.
11. Sheldrick GM. A short history of SHELX. *Acta Cryst. A*. 2008; 64:112–122. [PubMed: 18156677]
12. Stoll S, Schweiger A. EasySpin, a comprehensive software package for spectral simulation and analysis in EPR. *J. Magn. Reson.* 2006; 178:42–55. [PubMed: 16188474]
13. Frisch, MJ.; Trucks, GW.; Schlegel, HB.; Scuseria, GE.; Robb, MA.; Cheeseman, JR.; Montgomery, JA., Jr; Vreven, T.; Kudin, KN.; Burant, JC., et al. Gaussian 03, revision E.01. Wallingford, CT: Gaussian, Inc; 2007.
14. Wertz, JE.; Bolton, JR. Electron Spin Resonance: Elementary Theory and Practical Applications. New York: McGraw-Hill, Inc; 1972.
15. Colaneri MJ, Vitali J, Peisach J. Aspects of Structure and Bonding in Copper-Amino Acid Complexes Revealed by Single-Crystal EPR/ENDOR Spectroscopy and Density Functional Calculations. *J. Phys. Chem. A*. 2009; 113:5700–5709. [PubMed: 19378965]

16. Peisach J, Blumberg WE. Structural Implications Derived from the Analysis of Electron Paramagnetic Resonance Spectra of Natural and Artificial Copper Proteins. Arch. Biochem. Biophys. 1974; 165:691–708. [PubMed: 4374138]
17. Silver BL, Getz D. ESR of $\text{Cu}^{2+}(\text{H}_2\text{O})_6$. II. A quantitative study of the dynamic Jahn-Teller effect in copper-doped zinc Tutton's salt. J. Chem. Phys. 1974; 61:638–650.
18. Riley MJ, Hitchman MA, Mohammed AW. Interpretation of the temperature dependent g values of the $\text{Cu}(\text{H}_2\text{O})_6^{2+}$ ion in several host lattices using a dynamic vibrational coupling model. J. Chem. Phys. 1987; 87:3766–3778. Hoffmann SK, Goslar J, Hilczer W, Augustyniak MA, Marciniak M. Vibronic Behavior and Electron Spin Relaxation of Jahn-Teller Complex $\text{Cu}(\text{H}_2\text{O})_6^{2+}$ in $(\text{NH}_4)_2\text{Mg}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ Single Crystal. J. Phys. Chem. A. 1998; 102:1697–1707.
19. Sack RA. A contribution to the theory of the exchange narrowing of spectral lines. Mol. Phys. 1958; 1:163–167.
20. Hoffmann SK, Szczepaniak LS. Dynamical Averaging of Two Symmetry-Related Magnetic Resonance Tensors in Single Crystals. J. Mag. Reson. 1983; 52:182–192.
21. Cotton, FA.; Wilkinson, G. Advanced Inorganic Chemistry. 3rd ed.. NY: John Wiley & Sons; 1972. p. 52

Figure 1. Ortep-III⁶/PovRay⁷ renderings of (A) the atomic spatial arrangement of the host bis(L-histidinato)cadmium dihydrate showing the metal ion coordination and (B) the proposed copper binding site in doped bis(L-histidinato)cadmium dihydrate crystals at 77 K. The copper position was deduced from the $\mathbf{g}$ and $\mathbf{A}^{\text{Cu}}$ tensors and the ESEEM measured ^{14}N hyperfine and quadrupole coupling tensors of the remote nitrogen (N3) of the coordinated imidazole. The Cu^{2+} binds stronger to one of the two neighboring histidine molecules which are related by a crystallographic two-fold rotation axis. This gives rise to two adjacent copper sites related by this symmetry. Coloring scheme in this and subsequent figures: red for O, blue for N, grey for C, tan for H, gold for Cu and scarlet for Cd.

Figure 2.
The similar environment of water at 130 K and 200 K in bis(L-histidinato)cadmium dihydrate. Hydrogen bonds are shown as dashed lines and distances are indicated between the hydrogens and corresponding acceptor atoms. The symmetry relationships for the labeled atoms are: (a) $x - 1, y, z$ (b) $y - 1, x - 1, -z$ and (c) $y - 1/2, -x + 1/2, z + 1/4$.

Figure 3.

The 77 K EPR spectra of crystals of $^{63}\text{Cu(II)}$ -doped bis(L-histidinato)cadmium dideuterate acquired at $c//H$, $a(b)//H$ and $a+b//H$ portraying the superhyperfine splittings. The separate bubble sections illustrate experimental splittings and EasySpin fits (green curves) using a “2+1” ^{14}N and one ^1H nuclear coupling model as described in the text. I, II, I' and II' denote the different copper m_I split patterns in the spectra that arise from sites related by crystal symmetry as described in the text.

Figure 4.

The integrated EPR spectra of Cu(II)-doped bis(L-histidinato)cadmium dihydrate obtained at (A) $c//H$ and (B) $a(b)//H$ and recorded at 80 K (blue) and at 298 K (green). PeakFit curves were used to mimic the spectra. The copper $I=3/2$ hyperfine splittings and spectral g-values are reported in the text.

Resonant Magnetic Field

Figure 5.

Room Temperature (298 K) single crystal EPR measurements of the dependence of the resonance magnetic fields for the copper pattern as a function of the magnetic field orientation relative to the crystal reference planes. The solid curves were calculated variations based upon the least square fit g and A^{Cu} hyperfine tensors. The slightly split lines indicate the quality and symmetry of the fit data as these occur when crystal orientation angles ranged between 90° – 180° in the left plot and 45° – 90° in the right plot.

B**Figure 6A.**

A The temperature dependence of the integrated EPR spectrum at **c//H**. Indicative of the changes with rising temperature from 77 K, the lowest field peak of the 4-line pattern broadens, shifts slightly to higher resonant field and loses intensity as the 4 line pattern of the high temperature species grows.

B Changes in the EPR spectrum at **c//H** as the temperature varies between 155 K and 180 K clearly showing a distinct conversion from a low temperature to the high temperature pattern. The vertical lines indicate the field positions of the lowest field resonance peaks of the low and high temperature species.

Figure 7.

(A) PeakFits of the measured integrated EPR spectra at c//H at four temperatures. n_{LT} was determined as the ratio of the amplitude of the lowest field PeakFit peak of the low temperature species with the sum of the amplitudes of the lowest field peak of the low and high temperature species. (B) Plot of the fractional population of the low temperature species n_{LT} as a function of temperature. The temperature dependence below 200 K fit a sigmoidal function $n_{LT} = 1/(1 + e^{-(T-T_c)/\Delta T})$, with $T_c = 163 \text{ K}$ and $\Delta T = 19 \text{ K}$.

Figure 8.

(A) The temperature dependence of the integrated EPR spectrum at $a+b//H$. The lowest field peak of the low temperature pattern broadens, shifts slightly to higher field and merges with the growing lowest field peak of the high temperature pattern. (B) The temperature dependence of the integrated EPR spectrum when H is directed 110° from c in the $a(b)c$ reference plane. As the temperature increases from 80 K, the lowest field peak of the 4-line pattern broadens, shifts to higher resonant fields and loses intensity as the 4-line pattern of the room temperature species increases in intensity. The vertical dashed lines indicate the field positions of the lowest field peaks of the low and high temperature species.

Figure 9.

Plots of the resonant field of the lowest field peak of the low temperature pattern of site I as a function of temperature at two sample orientations. (A) when $\mathbf{a}+\mathbf{b} \parallel \mathbf{H}$ and (B) when $\mathbf{H}$ is directed 110° from the $\mathbf{c}$ -axis in the $\mathbf{a}(\mathbf{b})\mathbf{c}$ reference plane. The peak in (A) overlaps the lowest field peak of the high temperature pattern of site II_H near 165 K. The peak in (B) loses intensity and broadens to become immeasurable at temperatures >180 K.

Figure 10.

(A) Simulated EPR spectra computed by diagonalizing Eq. 4 for two copper patterns, Site 1 and Site 2. The spectrum shows a narrowing when the hop rate becomes comparable to the separation of lines of the two patterns. As the ratio of hop rate v_h to the line separation v_o (in frequency units) increase, the lines shift in position, broaden and collapse at the midpoint of the hopping m_I line resonances. Note that the resonant shift and broadening become evident even at a very low ratio (also shown in Anderson¹). (B) Dynamic simulations when the two hopping spectral patterns are identical and overlap. Here the spectrum remains unaltered by the hopping rate and process.

Figure 11.

Integrated EPR spectra at 160 K when $c//H$. The blue curve represents the experimental spectrum, the red is a dynamic simulated spectrum by diagonalizing Eq. 4 with a hop rate $\nu_{h2} = 1.7 \times 10^8 \text{ s}^{-1}$ and $W_j = 1/4$. The stick diagrams represent the field positions of the resonant m_I lines of the I, II, I_{rt} and II_{rt} patterns (and their overlapped, primed counterparts). In this effective 2-state model, hopping occurs between $I \leftrightarrow \text{II}_{rt}$ and between the equivalent and overlapping $\text{II} \leftrightarrow \text{I}_{rt}$. It is assumed that jumping occurs to equivalent m_I states. Simulated lines were convoluted with a gaussian function having a breadth of 39 G to mimic the experimental linewidths in Figure 4A. The pink curve is a 1:1 composite of measured spectra at 80 K and 298 K.

Figure 12.

(A) Integrated EPR spectra at $\mathbf{a+b//H}$ measured at 80 K (blue) and 298 K (green). Peak curves for patterns I, II (80 K) and I_{rt} , II_{rt} (298 K) at the lower field section, and I' , II' (80 K) and I_{rt}' , II_{rt}' (298 K) at the higher field section were determined using PeakFit. The lowest field m_I resonance line of I discussed in the text is indicated LF. (B) A comparison of the integrated EPR spectrum at $\mathbf{a+b//H}$ at 160 K. The stick diagrams represent the field positions of the resonant m_I lines of patterns I, II, I_{rt} and II_{rt} and patterns I' , II' , I_{rt}' and II_{rt}' . The blue curve represents the experimental spectrum, the red is a dynamic simulated spectrum using Eq. 4 with a hop rate $\nu_{h2} = 1.7 \times 10^8 \text{ s}^{-1}$ $W_j = 1/4$. In this effective 2-state model, hopping occurs between $I \leftrightarrow II_{rt}$ (and between the equivalent and overlapping $II \leftrightarrow I_{rt}$) at the lower field section of the spectrum and between $I' \leftrightarrow II_{rt}'$ (and between the equivalent and overlapping $II' \leftrightarrow I_{rt}'$) at the higher field section of the spectrum. It is assumed that jumping occurs to the same m_I state. Simulated lines were convoluted with a gaussian function to mimic the experimental linewidth. The pink curve is a 1:1 composite of measured spectra at 80 K and 298 K.

Figure 13.

(A) 80 K and 298 K integrated EPR with H oriented 110° from the c -axis in the $a(b)c$ reference plane showing the individual m_I lines of the different sites. Stick diagrams denote the patterns for sites I and II in the 80 K spectrum and I_{rt} and II_{rt} in the 298 K spectrum. The peak curves were determined using PeakFit. The resonant field positions of all the patterns in these fits were guided by measured values reported in the earlier study (Reference 8, Figure 3, upper plot at 70°) and from the present work. (B) A comparison of the integrated EPR spectrum with H oriented 110° from the c -axis in the $a(b)c$ reference plane at 160 K. The blue curve represents the experimental spectrum, the red is a dynamic simulated spectrum using Eq. 4 with hop rates of $\nu_{h4} = 4.5 \times 10^8 \text{ s}^{-1}$ and $\nu_{h2} = 1.7 \times 10^8 \text{ s}^{-1}$, and with $W_j = 1/4$. In this 4-state model, hopping occurs between $I \leftrightarrow II_{rt}$, $I \leftrightarrow II$, $II \leftrightarrow I_{rt}$ and $I_{rt} \leftrightarrow II_{rt}$, as well as between their primed counterparts. It is assumed that jumping occurs to equivalent m_I states. Simulated lines were convoluted with a gaussian function to mimic the experimental linewidth at 80 K and 298 K. The pink curve is a 1:1 composite of measured spectra at 80 K and 298 K. The lowest field m_F resonance line of I discussed in the text is indicated LF.

Figure 14.

Dynamic simulation plot of resonant field position as a function of transition rate p_{jk2} for the lowest field peak of site I of the low temperature pattern when $\mathbf{a+b/H}$. The hopping transitions are described in Figure 12 and the text. The simulation was performed by diagonalizing Eq. 4 using a 2-state model as described in the text. The curve $H(G) = a + b(p_{jk2})^c$ function, with $a = 2815.6$, $b = 199$ and $c = 2.35$ was fit to the theoretical points up to a rate $p_{jk2} (=W_j v_{h2})$ of $\sim 0.45 \times 10^8 \text{ s}^{-1}$. Simulations with higher rates produced too large an overlap of the low field line of I with the low field line of the I_{rt} to distinguish their resonant positions.

Figure 15.

A conformational view of the low temperature states I and II, and the high temperature states I_{rt} and I_{rt} . The high temperature states are the average of the two molecular spin Hamiltonian tensors at low temperature. In the 4-state model, the hopping transitions are $I \leftrightarrow II$, $I \leftrightarrow I_{rt}$, $II \leftrightarrow I_{rt}$, and $I_{rt} \leftrightarrow I_{rt}$. The hop rate between the low and high temperature states $I \leftrightarrow I_{rt}$, $II \leftrightarrow I_{rt}$ is v_{h2} , and between the low temperature states $I \leftrightarrow II$ and between the high temperature states $I_{rt} \leftrightarrow I_{rt}$ is v_{h4} .

A**B****Figure 16.**

(A) Arrhenius plots of $\ln(v_h)$ versus $(kT)^{-1}$ for the hopping transitions in copper-doped bis(L-histidinato)cadmium dihydrate crystals in the 120 to 180 K range. For the $I \leftrightarrow II_{rt}$ and $II \leftrightarrow I_{rt}$ transitions, the hop rate is v_{h2} and for $I \leftrightarrow II$ and $I_{rt} \leftrightarrow II_{rt}$, the hop rate is v_{h4} . (B) Potential Energy profile for the low temperature (blue) I and II, and high temperature (red) I_{rt} and II_{rt} states. The energy barriers between jumping states; $\Delta E_2 = 656 \text{ cm}^{-1}$ and $\Delta E_4 = 389 \text{ cm}^{-1}$, were determined from the slopes of the lines in (A).

TABLE 1

Crystal data, data collection and structure refinement details

	130 K	200 K
Chemical formula	C ₁₂ H ₂₀ CdN ₆ O ₆	C ₁₂ H ₂₀ CdN ₆ O ₆
Formula weight	456.74	456.74
Wavelength	0.71073 Å	0.71073 Å
Space group	P 4 ₃ 2 ₁ 2	P 4 ₃ 2 ₁ 2
Unit cell dimensions	a = 7.3625(3) Å c = 30.591(1) Å	a = 7.3515(7) Å c = 30.456(3) Å
Volume	1658.23(11) Å ³	1646.0(3) Å ³
Z	4	4
q range for data collection	2.85 to 28.28°	2.85 to 29.56°
Reflections collected	7648	7386
Independent reflections	2039	2150
Data / restraints / parameters	2039 / 0 / 155	2150 / 0 / 155
Goodness-of-fit on F ²	1.197	1.174
	2027 data; I > 2σ(I)	2134 data; I > 2σ(I)
Final R indices	R ₁ = 0.025, wR ₂ = 0.062 all data R ₁ = 0.025, wR ₂ = 0.062	R ₁ = 0.024, wR ₂ = 0.061 all data R ₁ = 0.025, wR ₂ = 0.061
Weighting scheme	w=1/[σ ² (F _o ²)+(0.0212P) ² +2.1391P] where P=(F _o ² +2F _c ²)/3	w=1/[σ ² (F _o ²)+(0.0179P) ² +1.7887P] where P=(F _o ² +2F _c ²)/3
Max and min peaks	0.590 and −0.917 eÅ ^{−3}	0.581 and −0.701 eÅ ^{−3}

$$^a R_1 = \sum ||F_o| - |F_c|| / \sum |F_o|;$$

$$^b wR_2 = [\sum [w (F_o^2 - F_c^2)^2] / \sum [w (F_o^2)^2]]^{1/2}.$$

Ligand assignments, experimental fits and predictions of the nuclear couplings responsible for the superhyperfine splittings in the 77 K EPR spectra for the proposed $^{63}\text{Cu}^{2+}$ binding site in bis(L-histidinato)cadmium dideuterate. Quantities in (black) were determined by experimental fits using EasySpin, in (red) were predicted using a point-dipole model for the electron spin and in (blue) were obtained from quantum mechanical DFT/B3LYP theoretical calculations using TZVP atomic basis functions. The experimental isotropic hyperfine component a_{iso} of each coupling was determined using the average of the three on-axis splittings. The a_{iso} values in parentheses were those determined either from the DFT computations or, for the ^1H coupling (asterisk), from an empirically derived relationship described in the text. All predicted splittings listed in the table incorporated the experimental a_{iso} components.

Table 2

Nuclear	Coupling (MHz)	a_{iso}	$a, b/H$ (A)	$a, b/H$ (B)	c/H	$a+b/H$ (low field)
N2		37.1	35.7	37.4	38.6	36.2
			35.0	39.8	37.2	35.6
		(16.4)	35.5	37.0	38.8	36.3
N1		29.8	37.8	25.9	25.6	28.2
			35.8	25.9	27.2	29.6
		(28.1)	35.8	26.3	27.7	29.6
N1'		20.0	19.2	23.5	18.2	21.1
			18.2	24.9	17.8	19.5
		(15.5)	18.7	21.8	19.5	19.6
H1		10.1	12.6	9.6	8.9	9.9
		(7.1)*	13.3	8.6	9.1	12.0
		(9.80)	12.0	9.1	9.2	9.5

Table 3

(A) Top: The g and Cu(II) hyperfine coupling (A^{Cu}) tensors measured from doped bis(l-histidinato)cadmium dihydrate crystals at room temperature. Bottom: The averages of the 77 K g and Cu(II) hyperfine coupling (A^{Cu}) tensors related by the two-fold axis along the a+b axis. The tensor principal directions (direction cosines) refer to the crystallographic a(b)c axes.

Room Temperature Tensors

	Principal Values	Direction Cosines		
		a	b	c
g	2.168	0.700	0.714	-0.008
	2.157	0.005	0.006	0.999
	2.074	0.714	-0.700	0.001
	Principal Values (MHz)	Direction Cosines		
		a	b	c
A^{Cu}	-256.	0.692	0.722	-0.005
	-188.	0.001	0.006	0.999
	-37.2	0.722	-0.692	0.003

Average of 77 K Tensors

	Principal Values	Direction Cosines		
		a	b	c
g	2.186	0.707	0.707	0.000
	2.163	-0.114	0.114	0.987
	2.089	0.698	-0.698	0.161
	Principal Values (MHz)	Direction Cosines		
		a	b	c
A^{Cu}	-265.	0.707	0.707	0.000
	-191.	-0.125	0.125	0.984
	64.2	0.696	-0.696	0.177

(B) Top: The g and Cu(II) hyperfine coupling (A^{Cu}) tensors reproduced from the EPR study of the room temperature pattern in doped Zn^{2+} (D,L-histidine)₂ pentahydrate crystals⁹. Bottom: The averages of the 80 K g and Cu(II) hyperfine coupling (A^{Cu}) tensors related by the two-fold axis along the b-axis. The tensor principal directions (direction cosines) refer to the crystallographic a'bc axes (with a' = b × c).

Room Temperature Tensors

	Principal Values	Direction Cosines		
		a'	b	c
g	2.204	0.692	-0.022	0.721
	2.124	0.002	0.999	0.033
	2.071	0.721	-0.025	-0.692
	Principal Values (MHz)	Direction Cosines		
		a'	b	c
A^{Cu}	360.	0.69	-0.01	0.72
	144.	0.01	0.99	0.03
	15.0	0.72	-0.02	-0.69

(B) Top: The g and Cu(II) hyperfine coupling (A^{Cu}) tensors reproduced from the EPR study of the room temperature pattern in doped $\text{Zn}^{2+}(\text{D,L-histidine})_2$ pentahydrate crystals⁹. Bottom: The averages of the 80 K g and Cu(II) hyperfine coupling (A^{Cu}) tensors related by the two-fold axis along the b -axis. The tensor principal directions (direction cosines) refer to the crystallographic $a'b'c'$ axes (with $a' = b \times c$).

Room Temperature Tensors

Principal Values		Direction Cosines		
		a'	b	c
<hr/>				
Average of 80 K Tensors				
Principal Values		Direction Cosines		
		a'	b	c
<hr/>				
g	2.241	0.601	0.000	0.799
	2.132	0.000	1.000	0.000
	2.061	0.799	0.000	-0.601
Principal Values (MHz)		Direction Cosines		
		a'	b	c
<hr/>				
A^{Cu}	391.	0.680	0.000	0.734
	189.	0.000	1.000	0.000
	31.5	0.734	0.000	-0.680