Ultradian metabolic rhythm in the diazotrophic cyanobacterium *Cyanothece* sp. ATCC 51142

Jan Červený^{a,1}, Maria A. Sinetova^{a,b,1}, Luis Valledor^{a,c}, Louis A. Sherman^d, and Ladislav Nedbal^{a,e,2}

^aGlobal Change Research Centre–CzechGlobe, Academy of Sciences of the Czech Republic, 664 24 Drásov, Czech Republic; ^bLaboratory of Intracellular Regulation, Institute of Plant Physiology, Russian Academy of Sciences, Moscow 127276, Russia; ^cDepartment of Molecular Systems Biology, University of Vienna, A1090 Vienna, Austria; ^dDepartment of Biological Sciences, Purdue University, West Lafayette, IN 47907; and ^eSystems Biology Laboratory, Faculty of Informatics, Masaryk University, 602 00 Brno, Czech Republic

Edited by Steven L. McKnight, The University of Texas Southwestern Medical Center, Dallas, TX, and approved June 26, 2013 (received for review January 31, 2013)

The unicellular cyanobacterium *Cyanothece* sp. American Type Culture Collection (ATCC) 51142 is capable of performing oxygenic photosynthesis during the day and microoxic nitrogen fixation at night. These mutually exclusive processes are possible only by temporal separation by circadian clock or another cellular program. We report identification of a temperature-dependent ultradian metabolic rhythm that controls the alternating oxygenic and microoxic processes of *Cyanothece* sp. ATCC 51142 under continuous high irradiance and in high CO₂ concentration. During the oxygenic photosynthesis phase, nitrate deficiency limited protein synthesis and CO₂ assimilation was directed toward glycogen synthesis. The carbohydrate accumulation reduced overexcitation of the photosynthetic reactions until a respiration burst initiated a transition to microoxic N₂ fixation. In contrast to the circadian clock, this ultradian period is strongly temperature-dependent: 17 h at 27 °C, which continuously decreased to 10 h at 39 °C. The cycle was expressed by an oscillatory modulation of net O₂ evolution, CO₂ uptake, pH, fluorescence emission, glycogen content, cell division, and culture optical density. The corresponding ultradian modulation was also observed in the transcription of nitrogenase-related *nifB* and *nifH* genes and in nitrogenase activities. We propose that the control by the newly identified metabolic cycle adds another rhythmic component to the circadian clock that reflects the true metabolic state depending on the actual temperature, irradiance, and CO₂ availability.

cyanobacteria | diurnal | metabolism | oscillation

Cyanobacteria have helped form the Earth's biosphere since they first evolved some 3 billion years ago. These microorganisms made our atmosphere oxygenic and their fixation of CO₂ and N₂ has made important contributions to the elemental cycles. In past decades, cyanobacteria of the genus *Cyanothece* have attracted strong attention for their contributions to the nitrogen cycle (1); as unique models to study the relationship between N₂ fixation, photosynthesis, and respiration (2); and as promising candidates for bioenergy production (3–5). The strain *Cyanothece* sp. American Type Culture Collection (ATCC) 51142 (hereafter *Cyanothece*) has particularly robust properties (2, 4). *Cyanothece* is a unicellular cyanobacterium in which spatial compartmentalization of the mutually exclusive oxygenic photosynthesis and microoxic nitrogen fixation is impossible (6). The strategy used by this organism is to temporally separate the molecular oxygen released by photosynthesis from the nitrogenase that would otherwise be irreversibly O₂-inactivated. The capacity to separate the antagonistic metabolic processes in time is usually attributed to circadian control. The circadian clock in cyanobacteria relies on cyclic (de-)phosphorylation involving complexes of the KaiA, KaiB, and KaiC proteins (7–9). The clock mechanism has been studied in great detail in *Synechococcus elongatus* PCC 7942 and also in vitro (10). The clock period in this organism has been shown to be temperature-compensated—a feature essential for controlling the daily rhythm,

particularly in organisms that do not sustain a stable temperature for their metabolism (11).

In *Cyanothece* the *kai* genes exist in multiple copies (12), and the Kai proteins have not been studied in the detail achieved for *S. elongatus*. The daily modulation of the metabolic activity and gene transcription, namely alternation of photosynthetic and N₂-fixation phases, has been attributed to the control by the circadian clock largely because the observed period was ~24 h, even in continuous light following a 12-h light/12-h dark entrainment (13). Certain genes, especially those associated with nitrogenase assembly and function, demonstrated a circadian-type of regulation when the light-dark pattern was modified for growth under continuous light or short day–night periods (14).

However, there are no published data on *Cyanothece* to show that the period of the oscillatory modulation in continuous light is temperature-compensated as expected for circadian control. One can expect an involvement of metabolic processes (see ref. 2) that may result in distinct ultradian rhythms, such as those demonstrated in yeast, which occur with temperature-dependent periods that are significantly shorter than 24 h. The hypothesis suggesting involvement of an ultradian metabolic cycle in *Cyanothece* is supported by oscillations with *ca.* 12-h periods that occur in continuous light following an initial 12-h light/12-h dark entrainment (15) as well as in continuous light in a batch culture (16). Components with an approximate 12-h period have also been detected by a Fourier transform analysis in transcript data in continuous light following a 12-h light/12-h dark entrainment (17).

In searching for a potential ultradian metabolic rhythm in *Cyanothece* we have performed experiments similar to those in yeast in which ultradian oscillations were induced by a starvation period (18). *Cyanothece* was grown to late exponential or linear phase in regular medium containing nitrate and supplied with saturating CO₂ and light. Strong ultradian oscillations occurred after the cells were moved to a minus-nitrate medium. This newly identified ultradian metabolic rhythm is strongly temperature-dependent. We also show that the circadian cycle is well temperature-compensated. The contrasting temperature dependencies document that the ultradian and circadian cycles are independent. The ultradian rhythm dominates in saturating CO₂ and light, whereas the circadian rhythm prevails when irradiance and/or CO₂ concentration are lowered.

Author contributions: J.Č., M.A.S., L.A.S., and L.N. designed research; J.Č., M.A.S., and L.V. performed research; J.Č., M.A.S., L.V., L.A.S., and L.N. analyzed data; and L.A.S. and L.N. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹J.Č. and M.A.S. contributed equally to this work.

²To whom correspondence should be addressed. E-mail: nedbal.lad@gmail.com.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1301171110/-DCSupplemental.

Ultradian and Circadian Rhythms in a Diurnally Entrained Culture.

Cyanotheca was grown in flat-panel photobioreactors with highly time-resolved, automated sampling to follow cyclic processes over days and weeks (19). Before the experiment shown in Fig. 1, the culture was entrained in 12-h light/12-h dark cycles in a turbidostat mode in which the culture optical density was kept in a narrow range by a photobioreactor-controlled feedback dilution. The experiment started after the last period of the diurnal entrainment (interval 0–24 h in Fig. 1) by switching off the dilution that allows culture batch growth and by keeping the culture in continuous light for the subsequent 10 d (interval 24–264 h in Fig. 1). In response to the pretreatment, the culture exhibited a complex oscillatory pattern in which both the shorter ultradian and the longer circadian rhythms combined. The periods in which the ultradian rhythm dominated (see the 72–96 h interval in Fig. 1, *Lower Left*) were followed by periods dominated by the circadian rhythm (see the 216–264 h interval in Fig. 1, *Lower Right*).

The concentration of dissolved oxygen oscillated with a minimum in the first half of subjective night in the circadian phase (216–264 h, see open circles in Fig. 1, *Lower Right*). The minimum in the concentration of the dissolved oxygen during the ultradian phase (~48–144 h, open circles in Fig. 1, *Lower Left*) was not firmly anchored to the subjective day–night cycle,

Fig. 1. Transition from ultradian to circadian rhythms in diurnally entrained batch culture. *Cyanothece* culture was diurnally entrained in a turbidostat without nitrate. The dark–light entrainment was followed by constant light 136 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ of 627 nm and 10 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ of cool white light with turbidostat switched off. (*Upper*) The experiment period starting with the last 24-h entrainment period and ending after 10 d of continuous light. The black rectangle at the bottom of the graph between 12 and 24 h shows the last night dark period; the subsequent striped rectangles show subjective (anticipated) night periods. (*Lower*) Typical ultradian (72–96 h) and circadian (216–264 h) rhythms. The solid line shows the culture growth dynamics as measured by OD_{680} . The thick line with open circles shows relative changes in the dissolved O_2 (dO_2). The other measurements are shown only in the lower two panels: The thin full line represents the relative change in the dissolved CO_2 , the dashed line represents the relative change of pH, and the long-dashed line shows the relative change of the steady-state pigment fluorescence emission Ft.

Fig. 2. Batch culture transition from ultradian to circadian rhythms after an exchange to nitrate-less medium. The culture growing rapidly in nitrate supplemented ASP2 medium was harvested and resuspended in nitrate-less medium (time 0). The transition introduced oscillations in dO_2 , in OD_{680} optical density, steady-state fluorescence emission F_t , and in pH. The dissolved CO_2 is represented here by the antiparallel pH dynamics. The symbols are the same as in Fig. 1.

because its period was neither equal to 12 h nor 24 h. The minima of the dissolved oxygen preceded the maxima of the dissolved CO₂ concentration by 1–2 h (thin full line in Fig. 1). The oscillations of the medium pH (short-dashed line in Fig. 1, all *Lower* panels) were nearly antiparallel to the dissolved CO₂ concentration, as anticipated from the water chemistry of inorganic carbon forms. During the rapid growth (Fig. 1, *Lower Left*), OD₆₈₀ of the suspension was weakly modulated with a period of the ultradian rhythm so that the maxima of the dissolved oxygen concentration matched the steepest increase of the optical density. This suggests that high photosynthetic activity indicated by the high concentration of dissolved O₂ correlated with an increase in chlorophyll concentration represented by the surrogate OD₆₈₀. The maxima of the stationary pigment fluorescence emission (long-dashed line in Fig. 1, all *Lower* panels) approximately coincided with the O₂ minima and with the CO₂ maxima, a pattern that was not always reproduced in parallel experiments. This dynamic pattern can be tentatively interpreted as high fluorescence resulting from a reduction of the plastoquinone pool by high rates of respiration and low rates of photosynthesis and vice versa.

Metabolic Rhythms in a Diurnally Nonentrained Culture. A complex oscillatory pattern similar to the diurnally entrained culture (Fig. 1) was also observed in continuous light, i.e., without the diurnal entrainment, when cyanobacteria growing rapidly in a nitrate-rich medium were transferred to a medium without nitrate (time 0 in Fig. 2). The dynamic pattern during the first 3 d after the media replacement exhibited a dominant ultradian character (period ~ 10.5 h). As the culture grew ($OD_{680} \sim 2.2$), the oscillations became dominated by the circadian rhythm (Fig. 2, *Lower Right*). Note that the residual involvement of the ultradian dynamics remained also in the dense culture, as demonstrated by the two high amplitudes at hours 160.5 and 234 of the experiment. The period separating the two high amplitudes corresponded to seven ultradian periods (7×10.5 h = 73.5 h) and to, roughly, three circadian periods (3×24 h = 72 h). Thus, at every

seventh ultradian and every third circadian period, the respective amplitudes combined rather than canceled each other.

We argue that the transition from the dominant ultradian to circadian rhythm in Fig. 2 was induced by decreasing the mean effective irradiance in the dense culture. This hypothesis is supported by the results in Fig. 3 in which the culture density was stabilized by the turbidostat that was diluting the suspension to the same optical density and, thus, the mean irradiance was fully controlled by the photobioreactor to high irradiance in the initial days and to low irradiance toward the end of the experiment. The data demonstrate that the ultradian rhythm can be converted to a circadian cycle by lowering of the incident irradiance in the photobioreactor. This result raises the question as to what is the decisive factor controlling the transition of the ultradian phenomena to the circadian rhythm: Is it light as the result in Fig. 3 suggests, CO_2 availability, or is it fast growth that requires both high light and high CO_2 ? Data shown in Figs. S1 and S2 suggest that high light and high CO_2 , rather than fast growth, lead to the dominating ultradian rhythm. Another condition necessary for the ultradian rhythm to occur is lack of nitrate in the medium. An addition of nitrate quickly damped the ultradian oscillations to zero (Fig. S3). Also, the ultradian oscillations in Fig. 2 were induced by transferring the cells to a nitrate-free medium.

Temperature is another important factor affecting the ultradian metabolic rhythm. The oscillations shown in Figs. 1–3 occurred at 36 °C with an ultradian period of 10.55 ± 0.05 h (mean and SD of five biological replicates). We found that the period of ultradian oscillations was strongly temperature-dependent. It increased to 17 h at 27 °C and decreased to 10 h at 39 °C (Fig. 4). The temperature coefficient (Q_{10}) was 1.55, documenting the ultradian temperature dependence. We also measured the periods of the 24-h rhythms in different temperatures and found Q_{10} close to 1, typical for the temperature-compensated circadian rhythm. The different response to temperature change shows that the ultradian rhythm is truly an independent phenomenon that is not derived from the circadian clock.

Data in Fig. 5A demonstrate the ultradian oscillations of dissolved O_2 concentration, pH, and partial synchronization of cell division (bars). The increasing dissolved oxygen and rising pH

Fig. 3. Transition from ultradian to circadian rhythms after decrease of light in a turbidostat. The oscillations of dissolved O_2 , CO_2 , fluorescence F_t , and pH were dominated by the ultradian rhythm during the first 64 h when exposed to high irradiance ($160 \mu\text{mol photons m}^{-2}\text{s}^{-1}$). Switching to low irradiance ($50 \mu\text{mol photons m}^{-2}\text{s}^{-1}$) led to a decrease in the ultradian rhythm amplitude and to prevalence of the circadian oscillation. Noteworthy is the saw-like OD_{680} curve that shows that the culture was automatically diluted whenever its density became >0.7 . The dilutions were less frequent around minima in the concentration of dissolved O_2 , indicating slow growth when photosynthesis is suppressed.

Fig. 4. Temperature dependence of the ultradian (open circles) and circadian (closed circles) cycles. The data were obtained as averages of at least two independent experiments. At 33 °C and 36 °C, more independent experiments were performed to determine the SD of ± 0.05 h. The Q_{10} rates were calculated to quantify the temperature dependence of a rate proxy of 1 per period.

(i.e., decreasing dissolved CO_2) reflected active photosynthesis leading to an accumulation of biomass carbon between *ca.* the seventh and 13th hours (closed squares in Fig. 5B). A large part of the organic carbon was stored in the form of semiamylopectin granules (20), frequently referred to as glycogen (Fig. 5B, open squares). The proportion of glycogen carbon to total biomass carbon increased during the photosynthesis phase, likely because of the lack of nitrogen that limited protein synthesis. The imbalance between the surplus of fixed carbon and the lack of nitrate in the medium grew to a point where photosynthesis was down-regulated; the dissolved oxygen and, *ca.* 2 h later, the pH started decreasing; and the accumulation of organic carbon was suspended. The glycogen content started decreasing and oxygen dropped to a level that allowed nitrogen fixation. Nitrogen fixation dominated between *ca.* the third and seventh hours and between hours 13.5 and 18 of the experiment—periods indicated by the dashed lines surrounding the respective segments in Fig. 5. The rising nitrogen content of the biomass is indicated by the dashed line and closed circles in Fig. 5C. The genes *nifH* and *nifB* necessary for nitrogen fixation are expressed to higher mRNA levels in the early phase of the nitrogen fixation periods.

Discussion

Linking the Ultradian Metabolic Rhythm to Other Cyclic Processes.

This paper has demonstrated that *Cyanothece* displays ultradian rhythms of alternating photosynthesis and N_2 fixation when grown without nitrate in continuous culture conditions. Whether the cultures were entrained (Fig. 1) or not (Fig. 2), the exponential growth phase showed ultradian rhythms that eventually degenerated into more circadian-like rhythms. This finding ties together many disparate results for *Cyanothece* over the years and may provide a basis for the interrelationship of the circadian and ultradian types of temporal regulation.

The earlier experiments typically used low light/low CO_2 and showed circadian patterns without an obvious ultradian modulation (3, 21). Here, we demonstrated that the ultradian rhythm occurred when photosynthesis was saturated by high light and high CO_2 concentration. In respect to high carbon availability, it is important to note that an ultradian component was also observed over a circadian pattern when *Cyanothece* was grown heterotrophically with glycerol (22).

Both the ultradian and circadian cycles exhibit many similar features. During the photosynthesis phase, cells accumulated large

Fig. 5. The processes occurring during the ultradian oscillations relative to the photosynthesis and nitrogen fixation phases. (A) Oscillation of the dissolved O_2 (open circles), pH (closed circles), and percentage of dividing cells as counted through a microscope (bars). (B) Total carbon in cells (closed rectangles) increased during the photosynthesis phase, whereas it remained stagnant during the nitrogen fixation phase. Glycogen content (open rectangles) decreased during the nitrogen-fixing phase. (C) Dynamics of the total nitrogen content of photobioreactor biomass together with expression levels of nitrogenase genes. The dashed line shows the increase of N content during the nitrogen-fixation phase and stagnant level during the oxygenic photosynthesis phase. The solid lines show expression levels (NRQ) of the *nifB* (x) and *nifH* (+) genes together with measured nitrogenase activity (open triangles).

amounts of glycogen. The transition from photosynthetic to nitrogen fixation phase correlated with degradation of glycogen and a respiration burst. When enough ATP and NADH/NADPH accumulated and intracellular oxygen was lowered by respiration, the cells start fixing nitrogen.

Earlier studies of *Cyanothece* were performed mostly on 12-h light/12-h dark grown cells and also on cells grown under continuous light and short light–dark periods (21). Invariably, the nitrogenase genes in the 35-gene cluster demonstrated the most consistent circadian behavior (13, 14, 21). Interestingly, the nitrogenase genes were among the large majority of genes displaying a circadian rhythm even in experiments in which some other genes oscillated with a 12-h period. The observation of gene oscillation with a 12-h period led to an earlier proposal of an ultradian rhythm (17). Here, we probed expression of the *nifB* and *nifH* genes; both were clearly modulated by the ultradian rhythm, a feature consistent with the ultradian modulation of the nitrogenase activity (Fig. 5). The ultradian modulation of the nitrogenase genes supports the notion that the ultradian rhythm is, under favorable conditions, dominant in all aspects attributed earlier exclusively to circadian control. In this sense, it is highly relevant to ask if the ultradian and circadian processes are not reflecting a single control mechanism (23).

To clarify the interrelationship between the ultradian and circadian rhythms, we measured temperature dependence of the respective periods (Fig. 4). The circadian period was shown to be well temperature-compensated ($Q_{10} \sim 1$), whereas the ultradian period was found to be strongly temperature-dependent ($Q_{10} =$

1.55). We conclude that the ultradian metabolic oscillation represents a process that is truly independent from the circadian cycle.

The ultradian rhythms were replaced or dominated by circadian rhythms when the cell irradiance was lowered either by self-shading in high cell density (Figs. 1 and 2) or by dimming the irradiance incident at the photobioreactor cuvette (Fig. 3). In *SI Materials and Methods*, we show an experiment that supports the conclusion that the ultradian rhythm is largely suppressed when photosynthesis and growth are not saturated by light (Fig. S14).

The ultradian rhythm may also depend on fast growth that requires both high light and high CO_2 . However, we show in Fig. S1B that the growth rate was maximal at ca. 33 °C and declined toward 36 °C and 39 °C. This is a qualitatively contrasting temperature dependence compared with that of the ultradian period that declined monotonously between 27 °C and 39 °C. We conclude that the growth rate is not a driver of the ultradian phenomena. This conclusion was further supported by the demonstration (Fig. S2) that the change of the specific growth rate by increasing the sparging gas CO_2 concentration in the 700–1,000 ppm range was small compared with the much steeper decrease of the ultradian period.

It appears that two independent mechanisms of temporal separation of photosynthesis and nitrogen fixation are possible in *Cyanothece*: the ultradian rhythm that prevails when photosynthesis is saturated by light and CO_2 and the circadian rhythm that dominates when the rate of photosynthesis is limited.

Interestingly, the experiments reported here also shed light on an earlier observation that the circadian patterns of photosynthesis and nitrogen fixation often appeared even during the first entrainment period, a feature that was hard to understand without the context shown in Fig. 1. The experimental cultures used to be inoculated with batch cultures that were 10–14 d old and, thus, closely resembled ca. >200-h cells in Fig. 1 that were probably already poised for the circadian regimen before inoculation. It was the photobioreactor that permitted us to work under a variety of environmental conditions; we can now assimilate the information from these different experiments.

On the Homology Between the Metabolic Rhythms in *Cyanothece* and in *Saccharomyces cerevisiae*.

Aerobically growing, dense continuous cultures of *S. cerevisiae* exhibit autonomous energy-metabolite ultradian oscillations of ca. 2–5 h periods that are observed under low nitrogen and low carbon conditions (18, 24–26). Although yeast are eukaryotic organisms that separate incompatible processes in different compartments, they also use temporal separation to carry out incompatible metabolic reaction and further alleviate futile cycles (27). A difference between the ultradian oscillations in yeast and in *Cyanothece* relates to the culture density. In contrast to yeast, the *Cyanothece* ultradian oscillations do not occur in high cell density that leads to self-shading of the penetrating light and to a reduction of the ultradian oscillatory amplitudes. This difference is only due to the role of light penetration through the culture that is critical for cyanobacteria and irrelevant to yeast.

However, several principal homologous features of metabolic oscillations of *Cyanothece* and *S. cerevisiae* can be found despite their largely different prokaryotic and eukaryotic character. The yeast oscillations, similar to the oscillations reported here for *Cyanothece*, consisted of two distinct phases: the anabolic, carbohydrate-storing phase (reductive in the yeast and photosynthetic in the cyanobacterium) and the catabolic (oxidative in the yeast and N_2 fixing in the cyanobacterium). The ultradian oscillations of dissolved O_2 and pH (CO_2) were observed in both organisms, accompanied by accumulation of carbohydrates in the anabolic/photosynthetic phase that was concluded by a burst of respiration, (see refs. 18 and 24 for yeast). Also, cell division occurred only in the anabolic/photosynthetic phase (see ref. 18 for yeast). In yeast, many genes participating in the regulation of metabolic processes were oscillating with the same ultradian period as the

energetic-metabolic oscillations. In *Cyanothece*, the nitrogenase-related genes *nifB* and *nifH* were also clearly modulated by the ultradian rhythm, a feature consistent with the ultradian modulation of the nitrogenase activity. The homology between the metabolic ultradian oscillations in yeast and in cyanobacteria suggests that both organisms may have the same strategy of benefiting from the ultradian cycling in separating the mutually exclusive metabolic processes, anabolic or photosynthetic and catabolic or nitrogen-fixing reactions, and avoiding futile cycles (18, 27).

The finding of the ultradian metabolic cycle of *Cyanothece* also resonates with the recent report (2) that highlighted significant differences among six *Cyanothece* strains subjected to identical external cues. The differences were interpreted as indicating importance of metabolic signals that are specific for each strain rather than pure regulation by the circadian clock. An important question to ask in future research relates to the role of such metabolic signals and of the ultradian cycle in natural conditions. *Cyanothece* is a benthic organism that lives in mats or clusters on rocks in intertidal areas (28). Ultradian synchronization of alternating oxygenic and microoxic conditions in the *Cyanothece*-containing biofilms can be considered as a potentially important supplement to the circadian control in adjusting to actual temperature, light, and carbon availability in the environment.

Materials and Methods

Growth Conditions, Real-Time Measurements, and Sampling. The *Cyanothece* sp. strain ATCC 51142 (28) was obtained from Jana Stöckel (Washington University, St. Louis), and maintained in complete ASP2 medium (29, 30) and buffered with TAPS and TAPSO as described earlier (16).

For growing and real-time monitoring of cultures, we used flat panel photobioreactors (FMT-150, Photon Systems Instruments), which were described in detail previously (19). Unless specified otherwise, the culture temperature was stabilized at 36.0 ± 0.3 °C, irradiance was $130\text{--}160$ $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ of red light (~ 627 nm) supplemented with 25 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ of cool white or blue light (~ 455 nm). A high concentration of dissolved CO_2 was generated by sparging with 0.5% CO_2 .

For diurnal entrainment, the culture grown previously on a shaker in an Erlenmeyer flask was inoculated in a nitrate-free ASP2 medium in the

photobioreactor with alternating 12-h light and 12-h dark periods. After the entrainment lasting at least 5 d, the lights were switched on continuously.

For ultradian metabolic cycles induced by removal of nitrate from the medium, the culture was first grown in photobioreactors in ASP2 medium with nitrate until the late exponential or early linear phase. The cells were harvested by centrifugation and resuspended in nitrate-free ASP2 medium. Cells continued to grow in continuous light without nitrate in batch or in turbidostat mode ($\text{OD}_{680} = 0.7$). Aliquots for cell counting; quantification of chlorophyll, carbohydrate storage, total carbon, and total nitrogen content; and measurement of nitrogenase activity were collected every 2–2.5 h for 24–54 h. Concentrations of dissolved oxygen and CO_2 were measured by the photobioreactor in situ every minute as described in ref. 31.

Analytical Measurements. Estimation of chlorophyll, carbohydrate, carbon, and nitrogen content and cell counts were done as described previously (16).

The nitrogenase activity was determined using the standard acetylene reduction method: Culture aliquots of 15 mL were incubated in gas-tight 40-mL volume glass vials with 8 mL of acetylene for 2 h in the light and temperature parameters that were used for growing. Then, 4 mL of head-space gas were assayed by gas chromatography. Potential ethylene contamination of acetylene was checked in 4 mL of gas measured immediately after the addition of the acetylene.

Quantitative RT-PCR scans were performed in a Mini-Opticon System (Bio-Rad). Normalized relative quantities (NRQ) and SEs of relative quantities were determined according to previous work (32). Expression levels of 16S and *rpnA* genes were used as endogenous controls for normalizing the abundance of *nifB* and *nifH* transcripts. Detailed information about RNA extraction, RT-PCR conditions is available in (SI Materials and Methods and Table S1).

ACKNOWLEDGMENTS. We thank Kristina Felcmanová for contributions during nitrogenase activity samples analysis. This publication is an output of the CzechGlobe Centre (Reg. CZ.1.05/1.1.00/02.0073) and the Education for Competitiveness Operational Programme (EC OP) Project financed by the European Union with the support of Czech Ministry of Education, Youth and Sports (MEYS CR) Reg. CZ.1.07/2.3.00/20.0256 (to J.C., M.A.S., L.V., and L.N.). This work was supported by research project Russian Foundation for Basic Research (RFRB) 12-04-32148 (to M.A.S.), Grant GACR 206/09/1284 (to L.N.), and by a grant from the US Department of Energy Genomics: GTL Program (to L.A.S.).

- Zehr JP (2011) Nitrogen fixation by marine cyanobacteria. *Trends Microbiol* 19(4):162–173.
- Bandyopadhyay A, Elvitigala T, Liberton M, Pakrasi HB (2013) Variations in the rhythms of respiration and nitrogen fixation in members of the unicellular diazotrophic cyanobacterial genus *Cyanothece*. *Plant Physiol* 161(3):1334–1346.
- Sherman LA, Min H, Toepel J, Pakrasi HB (2010) Recent Advances in Phototrophic Prokaryotes, ed Hallenbeck PC (Springer, New York), pp 275–290.
- Bandyopadhyay A, Stöckel J, Min H, Sherman LA, Pakrasi HB (2010) High rates of photobiological H_2 production by a cyanobacterium under aerobic conditions. *Nat Commun* 1(9):139.
- Aryal UK, et al. (2013) Proteome analyses of strains ATCC 51142 and PCC 7822 of the diazotrophic cyanobacterium *Cyanothece* sp. under culture conditions resulting in enhanced H_2 production. *Appl Environ Microbiol* 79(4):1070–1077.
- Schneegurt MA, Sherman DM, Nayar S, Sherman LA (1994) Oscillating behavior of carbohydrate granule formation and dinitrogen fixation in the cyanobacterium *Cyanothece* sp. strain ATCC 51142. *J Bacteriol* 176(6):1586–1597.
- Dong G, Golden SS (2008) How a cyanobacterium tells time. *Curr Opin Microbiol* 11(6):541–546.
- Iwasaki H, Kondo T (2004) Circadian timing mechanism in the prokaryotic clock system of cyanobacteria. *J Biol Rhythms* 19(5):436–444.
- Johnson CH, Mori T, Xu Y (2008) A cyanobacterial circadian clockwork. *Curr Biol* 18(17):R816–R825.
- Nakajima M, et al. (2005) Reconstitution of circadian oscillation of cyanobacterial KaiC phosphorylation in vitro. *Science* 308(5720):414–415.
- Johnson CH, Elliott J, Foster R, Honma K-I, Kronauer R (2004) *Chronobiology: Biological Timekeeping*, eds Dunlap JC, Loros JJ, DeCoursey PJ (Sinauer Associates, Sunderland, UK), pp 66–105.
- Welsh EA, et al. (2008) The genome of *Cyanothece* 51142, a unicellular diazotrophic cyanobacterium important in the marine nitrogen cycle. *Proc Natl Acad Sci USA* 105(39):15094–15099.
- Toepel J, Welsh E, Summerfield TC, Pakrasi HB, Sherman LA (2008) Differential transcriptional analysis of the cyanobacterium *Cyanothece* sp. strain ATCC 51142 during light-dark and continuous-light growth. *J Bacteriol* 190(11):3904–3913.
- Toepel J, McDermott JE, Summerfield TC, Sherman LA (2009) Transcriptional analysis of the unicellular, diazotrophic cyanobacterium *Cyanothece* sp. ATCC 51142 grown under short day/night cycles. *J Phycol* 45(3):610–620.
- Cervený J, Nedbal L (2009) Metabolic rhythms of the cyanobacterium *Cyanothece* sp. ATCC 51142 correlate with modeled dynamics of circadian clock. *J Biol Rhythms* 24(4):295–303.
- Sinetova MA, Cervený J, Zavřel T, Nedbal L (2012) On the dynamics and constraints of batch culture growth of the cyanobacterium *Cyanothece* sp. ATCC 51142. *J Biotechnol* 162(1):148–155.
- Elvitigala T, Stöckel J, Ghosh BK, Pakrasi HB (2009) Effect of continuous light on diurnal rhythms in *Cyanothece* sp. ATCC 51142. *BMC Genomics* 10(1):226.
- Tu BP, Kudlicki A, Rowicka M, McKnight SL (2005) Logic of the yeast metabolic cycle: Temporal compartmentalization of cellular processes. *Science* 310(5751):1152–1158.
- Nedbal L, Trtílek M, Cervený J, Komárek O, Pakrasi HB (2008) A photobioreactor system for precision cultivation of photoautotrophic microorganisms and for high-content analysis of suspension dynamics. *Biotechnol Bioeng* 100(5):902–910.
- Suzuki E, et al. (2013) Physicochemical variation of cyanobacterial starch, the insoluble α -Glucans in cyanobacteria. *Plant Cell Physiol* 54(4):465–473.
- McDermott JE, et al. (2011) A model of cyclic transcriptomic behavior in the cyanobacterium *Cyanothece* sp. ATCC 51142. *Mol Biosyst* 7(8):2407–2418.
- Schneegurt MA, Tucker DL, Ondr JK, Sherman DM, Sherman LA (2000) Metabolic rhythms of a diazotrophic cyanobacterium, *Cyanothece* sp. strain atcc 51142, heterotrophically grown in continuous dark. *J Phycol* 36(1):107–117.
- Elvitigala T, Pakrasi HB, Ghosh BK (2009) Dynamic network modeling of diurnal genes in cyanobacteria. *Emergent Problems in Nonlinear Systems and Control*, eds Ghosh B, Martin CF, Yishao Z (Springer, Heidelberg), pp 21–41.
- Chen C-I, McDonald KA (1990) Oscillatory behavior of *Saccharomyces cerevisiae* in continuous culture: II. Analysis of cell synchronization and metabolism. *Biotechnol Bioeng* 36(1):28–38.
- Müller D, Exler S, Aguilera-Vázquez L, Guerrero-Martín E, Reuss M (2003) Cyclic AMP mediates the cell cycle dynamics of energy metabolism in *Saccharomyces cerevisiae*. *Yeast* 20(4):351–367.
- Saito T, Mitsui K, Hamada Y, Tsurugi K (2002) Regulation of the Gts1p level by the ubiquitination system to maintain metabolic oscillations in the continuous culture of yeast. *J Biol Chem* 277(37):33624–33631.
- Tu BP, McKnight SL (2006) Metabolic cycles as an underlying basis of biological oscillations. *Nat Rev Mol Cell Biol* 7(9):696–701.
- Reddy KJ, Haskell JB, Sherman DM, Sherman LA (1993) Unicellular, aerobic nitrogen-fixing cyanobacteria of the genus *Cyanothece*. *J Bacteriol* 175(5):1284–1292.

29. Provasoli L, McLaughlin JJA, Droop MR (1957) The development of artificial media for marine algae. *Arch Mikrobiol* 25(4):392–428.
30. van Baalen C (1962) Studies on marine blue-green algae. *Bot Mar* 4:129–139.
31. Červený J, Šetlík I, Trtílek M, Nedbal L (2009) Photobioreactor for cultivation and real-time, *in-situ* measurement of O₂ and CO₂ exchange rates, growth dynamics, and of chlorophyll fluorescence emission of photoautotrophic microorganisms. *Eng Life Sci* 9(3):247–253.
32. Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J (2007) qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. *Genome Biol* 8(2):R19.