What Does the Cochrane Collaboration Say about Stretching Exercises?

Bartels EM, Lund H, Hagen KB, et al. Aquatic exercise for the treatment of knee and hip osteoarthritis. *Cochrane Database Syst Rev.* 2007;(4):CD005523. <http://dx.doi.org/10.1002/14651858.CD005523.pub2>. Medline:17943863

Blyton F, Chuter V, Walter KEL, et al. Non-drug therapies for lower limb muscle cramps. *Cochrane Database Syst Rev.* 2012;1(1):CD008496. <http://dx.doi.org/10.1002/14651858.CD008496.pub2>. Medline:22258986

Busch AJ, Barber KA, Overend TJ, et al. Exercise for treating fibromyalgia syndrome. *Cochrane Database Syst Rev.* 2007;(4):CD003786. <http://dx.doi.org/10.1002/14651858.CD003786.pub2>. Medline:17943797

Carvalho APV, Vital FMR, Soares BGO. Exercise interventions for shoulder dysfunction in patients treated for head and neck cancer. *Cochrane Database Syst Rev.* 2012;4(4):CD008693. <http://dx.doi.org/10.1002/14651858.CD008693.pub2>. Medline:22513964

Goldman EF, Jones DE. Interventions for preventing hamstring injuries. *Cochrane Database Syst Rev.* 2010;(1):CD006782. <http://dx.doi.org/10.1002/14651858.CD006782.pub2>. Medline:20091605

Harvey LA, Brosseau L, Herbert RD. Continuous passive motion following total knee arthroplasty in people with arthritis. *Cochrane Database Syst Rev.* 2010;(3):CD004260. <http://dx.doi.org/10.1002/14651858.CD004260.pub2>. Medline:20238330

He ML, Xiao ZM, Lei M, et al. Continuous passive motion for preventing venous thromboembolism after total knee arthroplasty. *Cochrane Database Syst Rev.* 2012;1(1):CD008207. <http://dx.doi.org/10.1002/14651858.CD008207.pub2>. Medline:22258981

Herbert RD, de Noronha M, Kamper SJ. Stretching to prevent or reduce muscle soreness after exercise. *Cochrane Database Syst Rev.* 2011;(7):CD004577. <http://dx.doi.org/10.1002/14651858.CD004577.pub3>. Medline:21735398

Katalinic OM, Harvey LA, Herbert RD, et al. Stretch for the treatment and prevention of contractures. *Cochrane Database Syst Rev.* 2010;(9):CD007455. <http://dx.doi.org/10.1002/14651858.CD007455.pub2>. Medline:20824861

Kay TM, Gross A, Goldsmith CH, et al. Exercises for mechanical neck disorders. *Cochrane Database Syst Rev.* 2012;8(8):CD004250. <http://dx.doi.org/10.1002/14651858.CD004250.pub4>. Medline:22895940

Mason DL, Dickens VA, Vail A. Rehabilitation for hamstring injuries. *Cochrane Database Syst Rev.* 2012;12(12):CD004575. <http://dx.doi.org/10.1002/14651858.CD004575.pub3>. Medline:23235611

McNeely ML, Campbell K, Ospina M, et al. Exercise interventions for upper-limb dysfunction due to breast cancer treatment. *Cochrane Database Syst Rev.* 2010;(6):CD005211. <http://dx.doi.org/10.1002/14651858.CD005211.pub2>. Medline:20556760

Romano M, Minozzi S, Bettany-Saltikov J, et al. Exercises for adolescent idiopathic scoliosis. *Cochrane Database Syst Rev.* 2012;8(8):CD007837. <http://dx.doi.org/10.1002/14651858.CD007837.pub2>. Medline:22895967

Yeung SS, Yeung EW, Gillespie LD. Interventions for preventing lower limb soft-tissue running injuries. *Cochrane Database Syst Rev.* 2011;(7):CD001256. <http://dx.doi.org/10.1002/14651858.CD001256.pub2>. Medline:21735382

The Cochrane Collaboration is an international not-for-profit and independent organization dedicated to making up-to-date, accurate information about the effects of health care readily available worldwide. It produces and disseminates systematic reviews of health care interventions and promotes the search for evidence in the form of clinical trials and other studies of interventions. For more information, visit www.cochranecollaboration.org.

DOI:10.3138/ptc.65.3.cochrane