Published in final edited form as:

Biochemistry. 2013 July 2; 52(26): 4553–4562. doi:10.1021/bi4004556.

Kinetics of Reversible Reductive Carbonylation of Heme in Human Cystathionine β -synthase†

Sebastián Carballal^{‡,§,||}, Ernesto Cuevasanta^{‡,||}, Inés Marmisol^{‡,||}, Omer Kabil[¶], Carmen Gherasim[¶], David P. Ballou[¶], Ruma Banerjee^{¶,*}, and Beatriz Alvarez^{‡,||,*}

[‡]Laboratorio de Enzimología, Facultad de Ciencias

[§]Departamento de Bioquímica, Universidad de la República, Montevideo, Uruguay

^{||}Center for Free Radical and Biomedical Research, Facultad de Medicina, Universidad de la República, Montevideo, Uruguay

[¶]Department of Biological Chemistry, Medical Center, University of Michigan, Ann Arbor, MI 48109-0600, USA

Abstract

Cystathionine β -synthase (CBS) catalyzes the condensation of homocysteine with serine or cysteine to form cystathionine and water or hydrogen sulfide (H_2S), respectively. In addition to pyridoxal phosphate, human CBS has a heme cofactor with cysteine and histidine as ligands. While Fe(III)-CBS is inert to exogenous ligands, Fe(II)-CBS can be reversibly inhibited by carbon monoxide (CO) and reoxidized by O_2 to yield superoxide radical. In this study, we have examined the kinetics of Fe(II)CO-CBS formation and reoxidation. Reduction of Fe(III)-CBS by dithionite showed square root dependence on concentration, indicating that the reductant species was the sulfur dioxide radical anion ($\text{SO}_2^{\bullet-}$) that exists in rapid equilibrium with $\text{S}_2\text{O}_4^{2-}$. Formation of Fe(II)CO-CBS from Fe(II)-CBS and 1 mM CO occurred at $(3.1 \pm 0.4) \times 10^{-3} \text{ s}^{-1}$ (pH 7.4, 25 °C). The reaction of Fe(III)-CBS with the reduced form of the flavoprotein methionine synthase reductase in the presence of CO and NADPH resulted in its reduction and carbonylation to form Fe(II)CO-CBS. Fe(II)-CBS was formed as intermediate with a rate constant of $(9.3 \pm 2.5) \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$. Reoxidation of Fe(II)CO-CBS by O_2 was multiphasic. The major phase showed hyperbolic dependence on O_2 concentration. Although H_2S is a product of the CBS reaction and a potential heme ligand, we did not find evidence for exogenous H_2S effect on activity or heme binding. Reversible reduction of CBS by a physiologically relevant oxidoreductase is consistent with a regulatory role for the heme and could constitute a mechanism for crosstalk between the CO, H_2S and superoxide signaling pathways.

Keywords

Cystathionine β synthase; heme; carbon monoxide; hydrogen sulfide; heme autooxidation; methionine synthase reductase; kinetics; dithionite

[†]This work was supported by grants and fellowships from CSIC, Universidad de la República, Uruguay (to B. A.), ANII (Agencia Nacional de Investigación e Innovación, Uruguay) (to S. C, I. M. and E. C) and the National Institutes of Health (HL58984, to R. B.)

*Correspondence should be addressed to Beatriz Alvarez, Laboratorio de Enzimología, Facultad de Ciencias, Iguá 4225, 11400 Montevideo, Uruguay; Phone/Fax: +598-2-5250749; Beatriz.Alvarez@fcien.edu.uy or Ruma Banerjee, Biological Chemistry Department, University of Michigan, Ann Arbor, MI 48109-0600, USA; Phone: +1-734-615-5238; rbanerje@umich.edu.

SUPPORTING INFORMATION AVAILABLE

Figures with control experiments of heme reduction by MSR, UV-visible absorption spectra of Fe(III)- and Fe(II)-CBS in the presence of CO or H_2S and enzymatic activity of CBS in the presence of H_2S are available free of charge via the Internet at <http://pubs.acs.org>.

Cystathionine β -synthase (CBS¹) is a pyridoxal phosphate (PLP)-dependent enzyme that plays a central role in mammalian sulfur metabolism. It catalyzes the condensation of serine with homocysteine to give cystathionine and water in the first step of the canonical transsulfuration pathway that leads to cysteine, the limiting substrate for glutathione synthesis. CBS also catalyzes the condensation of homocysteine and cysteine to form cystathionine and hydrogen sulfide (H₂S), that modulates several physiological functions in the cardiovascular, gastrointestinal and nervous systems (1–3). Mutations in CBS are the single most common cause of severe hyperhomocysteinemia (4). Increased levels of homocysteine in plasma are associated with a range of pathologies including cardiovascular diseases and neural tube defects (5, 6).

In addition to PLP, human CBS harbors a *b*-type heme cofactor that is low spin and six-coordinate in both the ferric and ferrous states (7). The axial heme ligands in human CBS are cysteine (Cys52) and histidine (His65) (8–11). The UV-visible absorption spectrum of Fe(III)-CBS shows a Soret peak at 428 nm and broad α/β bands centered at 550 nm. Upon reduction to Fe(II)-CBS, the Soret peak shifts to 449 nm and the α/β bands are resolved into two peaks at 571 and 540 nm, respectively (12).

The function of the heme in CBS remains unknown. A direct catalytic role in the PLP-mediated enzymatic mechanism was initially excluded by spectroscopic studies (13) and is consistent with its absence from the highly homologous yeast and *Trypanosoma cruzi* enzymes, which catalyze the same overall reaction (14, 15). Although crystal structures reveal a considerable distance (~ 20 Å) between the heme and PLP cofactors (10, 11, 16), a regulatory role has been suggested on the basis of the observation that perturbations in the heme coordination affect the equilibrium between PLP tautomers and modulate CBS activity (17). Bidirectional communication between the heme and PLP sites occurs through an α -helix that interacts at one end with the Cys52 heme ligand *via* Arg266 and at the other end with PLP *via* two conserved threonine residues (Thr 257 and Thr 260) (17–22). Ferric CBS is relatively stable, as reflected by the fairly low reduction potential of -0.350 V (20), and is inert to typical ferric heme ligands (23, 24). It is sensitive only to the thiol chelator, mercuric chloride, and to the strong oxidant, peroxynitrite (25, 26). In contrast, ferrous heme can bind ligands such as isonitriles, cyanide, nitric oxide ($\bullet$ NO), and carbon monoxide (CO), although with different affinities (23, 24, 27). CO binds to ferrous CBS by displacing the axial cysteine ligand and forming Fe(II)CO-CBS (Scheme 1), a species with a Soret maximum at 420 nm and decreased catalytic activity. The K_i value for CO is 5.6 μ M, which might be in a physiologically relevant range for endogenous CO concentrations generated by heme oxygenases in different tissues (23, 28, 29). In addition, Fe(II)-CBS can slowly decay to an inactive form that absorbs at 424 nm (Fe(II)424-CBS), in which the cysteine is replaced by a neutral ligand of unknown identity (30). Finally, the reaction of Fe(II)-CBS with O₂ occurs rapidly with a second-order rate constant of $(1.13 \pm 0.05) \times 10^5$ M⁻¹ s⁻¹ (pH 7.4, 25 °C), leading to formation of Fe(III)-CBS (which is fully active) and superoxide radical anion (O₂^{•-}). No intermediates are detected during the oxidation reaction, which exhibits a linear dependence on O₂ concentration, suggesting that it occurs *via* an outer sphere electron transfer (31).

Knowing whether the heme can be reduced by biological systems is central to evaluating its possible roles in sensing CO or producing O₂^{•-} in the cellular milieu. The low heme reduction potential in CBS makes it a challenging reduction. On the other hand, the heme

¹Abbreviations: CBS, cystathionine β -synthase; Fe(III)-CBS, CBS with ferric heme; Fe(II)-CBS, CBS with ferrous heme; CO, carbon monoxide; Fe(II)CO-CBS, CBS with ferrous heme and CO as ligand; Fe(II)424-CBS, CBS with ferrous heme and a neutral ligand replacing cysteine; Fe(II)5c-CBS, pentacoordinate ferrous CBS; PLP, pyridoxal 5'-phosphate; SOD, superoxide dismutase; DTPA, diethylenetriaminepentaacetic acid; MSR, methionine synthase reductase.

appears to be fairly surface exposed in the structures of both truncated and full-length CBS, indicating that it should be accessible to cellular reducing systems (10, 11, 16). We have recently demonstrated that human methionine synthase reductase (MSR) and NADPH can reduce CBS (Scheme 1). In the presence of CO, reduction is coupled to carbonylation, leading to formation of Fe(II)CO-CBS with concomitant loss of enzyme activity (32). MSR is a cytosolic flavoprotein oxidoreductase that harbors both FAD and FMN (33). The reduction potentials for the semiquinone/hydroquinone couples are -0.291 and -0.227 V for FAD and FMN, respectively (34). MSR has been shown to mediate electron transfer from NADPH to the cob(II)alamin cofactor of methionine synthase. The cob(II)alamin/cob(I)alamin reduction potential in methionine synthase is estimated to be -526 mV (35), and kinetic coupling of reduction with methylation by S-adenosylmethionine results in reductive methylation and reactivation of methionine synthase (33, 34).

In this study, we have investigated the kinetics of the reduction of the heme in CBS by dithionite and MSR/NADPH in the presence of CO. We have also examined the kinetics of Fe(II)CO-CBS reoxidation by O_2 and assessed the effect of H_2S on CBS activity. Our results provide insights into the kinetics of heme-mediated regulation of human CBS.

EXPERIMENTAL PROCEDURES

Enzyme Purification and Assays

Recombinant full-length human CBS, the truncated $\Delta C143$ variant and MSR were purified as described previously (12, 33, 36). CBS activity was measured using either the ninhydrin method (37) or the radiolabel assay (12) as described. Specific activities of the CBS preparations used in this study were 320 ± 26 (for full-length CBS) and $878 \pm 52 \mu\text{moles h}^{-1} \text{mg}^{-1}$ (for the truncated variant). Protein concentration was determined by the Bradford method using bovine serum albumin as standard (38). The absorption spectrum of MSR exhibited maxima at 380 nm and 454 nm and a shoulder at 480 nm. The ratio of A_{280}/A_{454} , was ~ 5.5 , similar to that reported previously (36). Reduction of MSR by NADPH under aerobic conditions led to a decrease in absorbance at 454 nm and the appearance of a small shoulder in the vicinity of 585 nm and 640 nm, consistent with the formation of an air stable semiquinone. The specific activity of MSR, determined by the NADPH-dependent cytochrome *c* reduction, was $3.8 \pm 0.2 \mu\text{mol min}^{-1} \text{mg}^{-1}$ at 37°C , similar to that reported previously (36).

Kinetics of Fe(III)-CBS reduction by dithionite

Sodium dithionite stock solutions were prepared in degassed 0.1 M NaOH and quantified by ferricyanide reduction ($\epsilon_{420} = 1020 \text{ M}^{-1} \text{cm}^{-1}$) assuming a 2:1 (ferricyanide:dithionite) stoichiometry (39, 40). An anaerobic solution of Fe(III)-CBS ($2 \mu\text{M}$) was mixed with dithionite (0.1–4 mM) in phosphate buffer (0.2 M, pH 7.4, 0.2 mM DTPA) at 25°C . Formation of Fe(II)-CBS was followed at 449 nm in a Varian Cary 50 spectrophotometer coupled to an Applied Photophysics RX2000 stopped flow machine. No changes in pH were detected after mixing.

Kinetics of Fe(II)-CBS reaction with CO

A solution of Fe(III)-CBS in CO-saturated phosphate buffer (0.1 M, pH 7.4) was mixed with a CO-saturated water solution to which dithionite had been added in low concentrations, using a stopped-flow spectrophotometer (SX.MV18, Applied Photophysics) with a diode array detector, placed inside an anaerobic chamber (Vacuum Atmospheres Co., Hawthorne, CA) that had an atmosphere of 95/5 N_2/H_2 and <1 ppm O_2 .

Kinetics of Fe(III)-CBS reduction by MSR/NADPH in the presence of CO

Quartz cuvettes containing CBS, MSR and NADPH were prepared in the anaerobic chamber, sealed with a rubber septum and the headspace was purged with CO for 15 min outside the anaerobic chamber. Then, UV-visible absorption spectra were recorded every minute up to ~50 min. The ionic strength dependence of the reduction of CBS by MSR was determined in 25, 50 or 100 mM potassium phosphate buffer, pH 7.4, to which variable amounts of KCl were added to obtain final ionic strengths ranging from 75 to 760 mM. The ionic strength (μ) was calculated according to $\mu = \frac{1}{2} \sum (M_i Z_i^2)$, where M_i and Z_i are the molarity and net charge of the ion. Rapid reaction kinetic studies were performed using the stopped-flow spectrophotometer placed inside the anaerobic chamber or a Hi-Tech Scientific SF-61DX stopped-flow spectrophotometer in the photodiode array mode placed outside the anaerobic chamber. In the latter case, to minimize O₂ contamination, the stopped-flow instrument was flushed with an anaerobic solution of protocatechuate (200 μ M) and protocatechuate 3,4-dioxygenase (~ 1 μ M) in phosphate buffer (0.1 M, pH 7.4) (41). This solution was replaced with O₂-free buffer before introduction of the anaerobic reaction mixtures. CBS and MSR solutions were made anaerobic in glass tonometers by repeated cycles of vacuum and equilibration with argon and left under a positive pressure of argon. Anaerobic 1 μ M CO-saturated Fe(III)-CBS was mixed with increasing concentrations (2, 5 or 10 μ M) of MSR pre-reduced with 1 mM NADPH and spectra were recorded from 0.02 to 1425 s. The experiments were performed at 25.0 ± 0.1 °C.

Kinetics of Fe(II)CO-CBS reoxidation by O₂

Fe(II)CO-CBS was obtained by first placing anaerobic Fe(III)-CBS (10 μ M) in phosphate buffer (0.2 M, pH 7.4) in a tonometer equipped with an L-shaped side arm fused to a spectrophotometer quartz cuvette. Next, CBS was stoichiometrically reduced with dithionite delivered *via* a gastight Hamilton syringe, monitoring the heme redox status spectrophotometrically by tipping the solution from the tonometer to the cuvette. Since O₂^{•-} radical can be formed in the presence of O₂, care was taken to avoid high concentrations of dithionite, and 0.25 μ M superoxide dismutase (SOD) was included. Following reduction, CBS was saturated with CO by consecutive cycles of vacuum and CO flow. Fe(II)CO-CBS formation was confirmed by the appearance of the 420 nm peak. The resulting Fe(II)CO-CBS solution was then mixed with solutions containing a series of O₂ concentrations obtained by equilibrating distilled water for 20 min with air (21% O₂) or with certified O₂/N₂ gas mixtures (10, 50, and 100% O₂) at 25.0 ± 0.1 °C.

Exposure of CBS to H₂S

Solutions of H₂S were freshly prepared by dissolving NaHS in ultrapure water and kept in sealed vials. The concentration of stocks was standardized by iodometric titration (42). The final solutions contained a mix of H₂S and HS⁻, governed by the value of $pK_{a1} = 7.0$ (43). At pH 7.4, sulfide distributes into 72% HS⁻ and 28% H₂S. In some experiments, dithionite (280 μ M) was added to the NaHS stock solution (0.13 M). When required, reduction of Fe(III)-CBS was achieved by addition of dithionite (60 μ M) in nitrogen-purged sealed vials.

Simulations and data analysis

Computer-assisted kinetic simulations were performed with Gepasi (44). Kinetic traces from the stopped-flow experiments were analyzed and fitted using the Pro-Data Viewer Software (Applied Photophysics), KinetAsyst3 software (Hi-Tech Scientific, Salisbury, UK), or Origin 6.1. All experiments were repeated at least two times and representative data are shown.

RESULTS

Kinetics of Fe(III)-CBS reduction by dithionite

When Fe(III)-CBS was mixed with excess dithionite, the heme was reduced rapidly. The formation of Fe(II)-CBS followed exponential kinetics, in agreement with the reaction being first-order in Fe(III)-CBS (Fig. 1A). The observed rate constant (k_{obs}) did not increase linearly with the concentration of dithionite. Instead, as with other hemeproteins (45–47), the k_{obs} showed a dependence on the square root of dithionite concentration (Fig. 1B) and plots of k_{obs} versus $[dithionite]^{1/2}$ were linear (Fig. 1B, inset). These results are consistent with the reductant being the sulfur dioxide radical anion ($SO_2^{\bullet-}$), formed by rapid homolysis of dithionite as described by equations 1–2.

The forward and reverse rate constants for equation 1 are: $k_1 = 2.5 \text{ s}^{-1}$ and $k_{-1} = 1.8 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ (25 °C) (45, 48). This mechanism yields k_{obs} as described by equation 3, where k_2 is the second-order rate constant for Fe(III)-CBS reduction by $SO_2^{\bullet-}$,

$$k_{obs} = k_2(k_1/k_{-1})^{1/2} [S_2O_4^{2-}]^{1/2} \quad (3)$$

From the fit of the values of k_{obs} to equation 3 (Fig. 1B), the dependence on $[S_2O_4^{2-}]^{1/2}$ was found to be $k_2(k_1/k_{-1})^{1/2} = 353 \pm 7 \text{ M}^{-1/2} \text{ s}^{-1}$ and k_2 was calculated to be $(9.5 \pm 0.2) \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ (pH 7.4, 25 °C).

Kinetics of Fe(II)-CBS reaction with CO

Fe(II)CO-CBS formation was first analyzed by mixing anaerobic Fe(III)-CBS (1 μM) with dithionite (26 μM) in a stopped-flow spectrophotometer, in the presence of CO (1 mM). Complete heme reduction was achieved in <10 s, which is consistent with the fast kinetics of CBS reduction by dithionite described above. A half-time of 0.4 s for heme reduction can be estimated under these conditions. Thus, the first spectrum shown is of the reduced CBS heme. Once reduced, Fe(II)-CBS reacted with CO and the heme Soret band shifted from 449 nm to a new maximum at 420 nm, consistent with previously reported spectra (Fig. 2) (23). The kinetics of CO binding to Fe(II)-CBS, monitored by the decrease in absorbance at 449 nm for 500 s, was fitted to a single exponential function (Fig. 2, inset) and yielded a value for k_{obs} of $(3.1 \pm 0.4) \times 10^{-3} \text{ s}^{-1}$ at 1 mM CO (pH 7.4, 25 °C). After 500 s, the absorbance maximum shifted slowly to 414 nm. The origin of this spectral shift is presently not understood and was not studied further. Of note, the carbonylated species obtained for *Drosophila melanogaster* CBS has a peak maximum at 414 nm (49).

Kinetics of Fe(III)-CBS reduction by MSR/NADPH in the presence of CO

In the absence of CO (Fig. S1), no formation of Fe(II)-CBS was detected, even at ratios of MSR:CBS 1:1, 2:1 or 4:1. In the presence of CO, conversion to Fe(II)CO-CBS with a Soret maximum near 420 nm was observed (Fig. 3A). This suggests that the heme in CBS is first reduced by MSR in the presence of NADPH and then reacts with CO forming Fe(II)CO-CBS. The apparent rate for Fe(II)CO-CBS formation was estimated to be $\sim 2 \times 10^{-10} \text{ M s}^{-1}$ during the first minutes of spectral registration (0.1 M phosphate buffer, pH 7.4, ionic strength $\sim 260 \text{ mM}$). Dithionite was used to obtain the difference extinction coefficient between Fe(II)CO-CBS and Fe(III)-CBS at 420 nm ($57,360 \text{ M}^{-1} \text{ cm}^{-1}$, Fig. S2). Dividing

the rate by the concentrations of MSR and CBS, an apparent second order rate constant of the order of $10^2 \text{ M}^{-1} \text{ s}^{-1}$ was estimated for Fe(II)CO-CBS formation. Exponential fits of the time courses yielded similar estimations (not shown). Next, the dependence of the apparent rate of Fe(II)CO-CBS formation on ionic strength was examined and found to be bell-shaped, with maximal rates observed at $\sim 200 \text{ mM}$ (Fig. 3B).

Since the initial reaction time course was not captured in the above experiments, stopped-flow experiments were performed. As shown in Fig. 4, when anaerobic CO-saturated Fe(III)-CBS was mixed with NADPH-reduced MSR, Fe(II)-CBS was formed in $< 200 \text{ s}$, followed by a time-dependent formation of Fe(II)CO-CBS as described by equations 4 and 5

To determine the rate constant (k_4) for reduction of the heme by MSR to form Fe(II)-CBS, the absorbance traces at 449 nm (Fig. 5A) were analyzed using an initial rate approach, assuming that the concentration of the reagents remained constant and equaled the initial concentration. The difference absorption coefficient at 449 nm was determined to be $63,900 \pm 5,020 \text{ M}^{-1} \text{ cm}^{-1}$ ($n = 11$) using dithionite as a reductant (not shown). The initial rate of Fe(II)-CBS formation, determined from the first 15 s of the kinetic traces, increased linearly with MSR concentration (Fig. 5B). Dividing the slope of the plot by the concentration of CBS, the rate constant k_4 for the reduction of Fe(III)-CBS was determined to be $9.3 \pm 2.5 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$ (pH 7.4, 25°C).

Although conversion of Fe(II)-CBS to Fe(II)CO-CBS (equation 5) is slow relative to the initial reduction step (equation 4), only $\sim 15\%$ of the initial CBS concentration formed the Fe(II)-CBS intermediate. This low yield is likely due to a relatively rapid reverse reaction, i.e. oxidation of Fe(II)-CBS to Fe(III)-CBS, consistent with the low reduction potential of CBS (-0.350 V (20)) vs. that of MSR (-0.227 V (34)). Based on the overall ΔE° for reduction of CBS by MSR, a ΔG° value of $+2.8 \text{ kcal mol}^{-1}$ and an equilibrium constant of 7.7×10^{-3} was calculated for equation 4, representing $[\text{Fe(II)-CBS}][\text{MSR}_{\text{ox}}]/([\text{Fe(III)-CBS}][\text{MSR}_{\text{red}}])$, which correlates with the concentration quotient, $\sim 8 \times 10^{-3}$, that can be estimated from the amount of Fe(II)-CBS formed (Fig. 5A). From the equilibrium constant and the forward reaction, $k_4 = 9.3 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$, the reverse rate constant k_{-4} , was estimated to be $1.2 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$.

Analysis of the formation of Fe(II)CO-CBS at 420 nm yielded an average k_{obs} value of $9.7 \pm 3.0 \times 10^{-4} \text{ s}^{-1}$ (Fig. 6A). For a mechanism involving the initial formation of Fe(II)-CBS in a relatively rapid equilibrium step followed by reaction with CO (equations 4–5), the net k_{obs} for Fe(II)CO-CBS formation is given by k_5 multiplied by the fraction of CBS that is in the reduced state. This fraction depends on the concentrations of reduced and oxidized MSR and CBS and on the rate constants k_4 and k_{-4} , changes during the time course of the reaction and is difficult to evaluate, because it involves second-order steps both in the forward and reverse directions. In addition, regeneration of reduced MSR is likely to occur according to the relatively high rates reported for MSR reduction by NADPH (36, 50). Assuming that the fraction of reduced CBS was $\sim 15\%$, k_5 can be roughly estimated to be $\sim 6 \times 10^{-3} \text{ s}^{-1}$ ($\sim 500 \mu\text{M CO}$), in reasonable agreement with the value of $3.1 \times 10^{-3} \text{ s}^{-1}$ obtained for Fe(II)CO-CBS formation when dithionite was used as reductant (Fig. 2). To complement the data analysis, the rates of Fe(II)CO-CBS formation were measured from the slopes of the linear portions of kinetic traces at 420 nm (40–80 or 100–250 s, depending on the trace) divided by the difference extinction coefficient of $57,360 \text{ M}^{-1} \text{ cm}^{-1}$ (Fig. 6B). The rates increased with

MSR concentration. Kinetic simulations performed with Gepasi using the initial concentrations of CBS and MSR and values of k_4 and k_{-4} of 9.3×10^2 and $1.2 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$, respectively, showed good agreement with experimental values when k_5 was assigned a value between 5 and $8 \times 10^{-3} \text{ s}^{-1}$ (Fig. 6B).

Kinetics of Fe(II)CO-CBS reaction with O₂

As observed previously, exposure of Fe(II)CO-CBS to air resulted in its rapid reoxidation to Fe(III)-CBS concomitant with recovery of enzyme activity (32). The kinetics of the O₂-dependent Fe(II)CO-CBS oxidation were characterized by stopped-flow spectrophotometry. For this, Fe(III)-CBS was reduced with dithionite in the presence of SOD (31) and saturated with CO. Formation of Fe(II)CO-CBS was confirmed spectrophotometrically (Fig. S2). As shown in Fig. 7, when Fe(II)CO-CBS was mixed with excess O₂, it was converted to Fe(III)-CBS. To estimate the rate constants associated with the reoxidation process, both multi-wavelength global fits and single wavelength kinetic traces were analyzed, with similar results. The decrease in absorbance at 420 nm was multiphasic and was studied in different time frames (Fig. 8). The first phase occurred with a $k_{\text{obs}} > 0.2 \text{ s}^{-1}$ (half-time $< 3 \text{ s}$) and its amplitude represented $< 25\%$ of the absorbance change. The second phase had a k_{obs} of $\sim 10^{-2} \text{ s}^{-1}$ and an amplitude of $\sim 50\%$. The third phase had a k_{obs} of $\sim 10^{-3} \text{ s}^{-1}$ and an amplitude of $< 25\%$. The first and third phases did not show a clear pattern of O₂ dependence and were not studied further. The second phase was associated with the largest amplitude change and its k_{obs} showed evidence of saturation behavior (Fig. 9). From the hyperbolic fit, the maximal k_{obs} was determined to be $0.033 \pm 0.013 \text{ s}^{-1}$ and the O₂ concentration at the half maximal k_{obs} was $0.10 \pm 0.07 \text{ mM}$.

Lack of interaction of CBS with exogenous H₂S

Since CBS is a source of H₂S in mammals (51–53), we tested the hypothesis that its heme might have a role as sulfide sensor. However, the presence of up to 13 mM exogenous H₂S had no effect on the enzymatic activity of Fe(III)-CBS or Fe(II)-CBS (Fig. S3). In addition, exposure to 0.5–3.2 mM H₂S had no effect on the absorption spectra of Fe(III)-CBS (Fig. S4). With Fe(II)-CBS, the presence of either oxidizing contaminants or decomposition products in the H₂S solution resulted in oxidation to Fe(III)-CBS (data not shown), but when dithionite (280 μM) was added to the H₂S stock solution (0.13 M), no changes in the absorption spectra of Fe(II)-CBS were observed in the presence of 0.1–4.1 mM H₂S (Fig. S5). Thus, no evidence was found for potential feedback regulation of CBS by H₂S.

DISCUSSION

In the present study, we have characterized the kinetics of reductive carbonylation of the heme in CBS using a chemical (dithionite), and a biochemical (MSR and NADPH) reductant (Table 1, Scheme 1). The kinetics of CO binding to Fe(II)-CBS, obtained using very low dithionite concentrations at pH 7.4, were slow and monophasic, with a k_{obs} of $3.1 \times 10^{-3} \text{ s}^{-1}$, probably representing cysteine dissociation. This result is comparable to the timescale of PLP tautomerization from the active ketoenamine to the inactive enolimine that occurs when CO binds to Fe(II)-CBS (17). It is also consistent with the kinetics of decay of Fe(II)-CBS to Fe(II)424-CBS, where the Cys52 ligand is replaced by a neutral ligand (54). A previous study on CO binding kinetics to CBS was conducted at pH 8.6 and reported biphasic kinetics. The fast phase increased hyperbolically with CO concentration, with a limiting k_{obs} of 0.0166 s^{-1} and a half-maximal value at 0.109 mM CO, while the slower phase was independent of CO concentration with a $k_{\text{obs}} \sim 1.1 \times 10^{-3} \text{ s}^{-1}$ (55). Another more recent study (49) reported that the kinetics of CO binding are affected by pH and by the presence of Fe(II)424-CBS. The amount and purity of dithionite used are probably also critical considerations (56).

Reductive carbonylation of CBS in the presence of MSR exhibited a dependence on ionic strength, suggesting that electrostatic contacts are involved in protein-protein interactions. The environment around the heme pocket in CBS is electropositive, while that around the FMN cofactor in MSR is negative (31, 57). A similar ionic strength maximum at 220 mM has been reported for the interaction of MSR and methionine synthase (33), with acidic residues in MSR being critical (36, 57). The unfavorable redox potentials between MSR and the CBS heme lead to a rather small rate constant for CBS heme reduction by MSR, $9.3 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$, coupled with a large reverse rate constant, $1.2 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$. An alternative reductase with a lower redox potential than MSR is likely to be more effective at reducing CBS. For example, the cytosolic human novel reductase 1, also in the same reductase family but of unknown biological function, has an FMN semiquinone to hydroquinone reduction potential of -0.305 V (58) and is expected to be a better reductant of the heme in CBS.

In contrast to the monophasic reaction of Fe(II)-CBS with O_2 to Fe(III)-CBS and superoxide, which shows a linear dependence on O_2 concentration ($1.13 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$), consistent with an outer sphere process (31), reoxidation of Fe(II)CO-CBS was multiphasic and involved a step with hyperbolic dependence on O_2 concentration. This multiphasic behavior is compatible with several parallel or consecutive reactions and suggests the coexistence of several heme species. In principle, reoxidation of Fe(II)CO-CBS to Fe(III)-CBS may occur directly or may involve the intermediate formation of Fe(II)-CBS. In addition, pentacoordinate Fe(II)5c-CBS may also be involved as an intermediate during the reoxidation process, and, at relatively long time frames, Fe(II)424-CBS. The kinetics of reactions of these latter two species with O_2 are not known. We hypothesize that the phase exhibiting saturation behavior reflects a step prior to reaction with O_2 being limiting, e.g., dissociation of CO (0.033 s^{-1}).

Studies with cell culture and tissues suggest that the reductive carbonylation of CBS occurs *in vivo*, since an increase in CO concentration decreases the concentration of cystathionine, presumably due to inhibition of CBS (59, 60). The interaction of CBS with CO is proposed to be involved in the regulation of cerebral vasodilation triggered by hypoxia (61). According to this model, diminished O_2 availability leads to decreased production of CO by heme oxygenase, which in turn prevents down-regulation of CBS by CO, resulting in greater H_2S production and consequently increased vasodilation. This hypothesis implies that the heme must be in the reduced ferrous state under normoxic conditions for CBS to be inhibited by CO; therefore, the relatively high rate of reoxidation by O_2 must be countered by some efficient reductive process(es) (31).

Assigning a sensor role to the heme in CBS heme depends critically on the feasibility of its reduction to the ferrous state, since the ferric form is stable and does not appear to interact with ligands that could result in inhibiting enzymatic activity (23, 24). Indeed, as shown herein, the heme does not interact with H_2S , itself a product of the enzymatic activity and despite the fact that a cysteine thiolate serves as a heme ligand in the ferric and ferrous states. Considerations on the possible roles of the CBS heme need to take into account both reduction to the ferrous state and its subsequent reaction with CO to form the inactive Fe(II)CO-CBS species or with O_2 to form Fe(III)-CBS and $\text{O}_2^{\bullet-}$. The finding that the heme in CBS can be reversibly reduced by a biochemical system reported previously (32) and characterized in more detail in this study, supports the possibility of a sensor role for this cofactor and provides a mechanism for crosstalk between the CO, H_2S and $\text{O}_2^{\bullet-}$ signaling pathways.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Drs Gerardo Ferrer (Facultad de Ciencias, Universidad de la República, Uruguay) and Tatyana Spolitak (Department of Biological Chemistry, University of Michigan, Ann Arbor) for helpful discussions and technical assistance.

REFERENCES

1. Abe K, Kimura H. The possible role of hydrogen sulfide as an endogenous neuromodulator. *J Neurosci.* 1996; 16:1066–1071. [PubMed: 8558235]
2. Zhao W, Zhang J, Lu Y, Wang R. The vasorelaxant effect of H₂S as a novel endogenous gaseous K(ATP) channel opener. *Embo J.* 2001; 20:6008–6016. [PubMed: 11689441]
3. Kabil O, Banerjee R. Redox biochemistry of hydrogen sulfide. *J Biol Chem.* 2010; 285:21903–21907. [PubMed: 20448039]
4. Kraus JP, Janosik M, Kozich V, Mandell R, Shih V, Sperandio MP, Sebastio G, de Franchis R, Andria G, Kluijtmans LA, Blom H, Boers GH, Gordon RB, Kamoun P, Tsai MY, Kruger WD, Koch HG, Ohura T, Gaustadnes M. Cystathionine beta-synthase mutations in homocystinuria. *Hum Mutat.* 1999; 13:362–375. [PubMed: 10338090]
5. Mills JL, McPartlin JM, Kirke PN, Lee YJ, Conley MR, Weir DG, Scott JM. Homocysteine metabolism in pregnancies complicated by neural-tube defects. *Lancet.* 1995; 345:149–151. [PubMed: 7741859]
6. Refsum H, Ueland PM, Nygard O, Vollset SE. Homocysteine and cardiovascular disease. *Annu Rev Med.* 1998; 49:31–62. [PubMed: 9509248]
7. Kery V, Bukovska G, Kraus JP. Transsulfuration depends on heme in addition to pyridoxal 5'-phosphate. Cystathionine beta-synthase is a heme protein. *J Biol Chem.* 1994; 269:25283–25288. [PubMed: 7929220]
8. Kery V, Poneleit L, Kraus JP. Trypsin cleavage of human cystathionine beta-synthase into an evolutionarily conserved active core: structural and functional consequences. *Arch Biochem Biophys.* 1998; 355:222–232. [PubMed: 9675031]
9. Taoka S, Widjaja L, Banerjee R. Assignment of enzymatic functions to specific regions of the PLP-dependent heme protein cystathionine beta-synthase. *Biochemistry.* 1999; 38:13155–13161. [PubMed: 10529187]
10. Meier M, Janosik M, Kery V, Kraus JP, Burkhard P. Structure of human cystathionine beta-synthase: a unique pyridoxal 5'-phosphate-dependent heme protein. *Embo J.* 2001; 20:3910–3916. [PubMed: 11483494]
11. Taoka S, Lepore BW, Kabil O, Ojha S, Ringe D, Banerjee R. Human cystathionine beta-synthase is a heme sensor protein. Evidence that the redox sensor is heme and not the vicinal cysteines in the CXXC motif seen in the crystal structure of the truncated enzyme. *Biochemistry.* 2002; 41:10454–10461. [PubMed: 12173932]
12. Taoka S, Ohja S, Shan X, Kruger WD, Banerjee R. Evidence for heme-mediated redox regulation of human cystathionine beta-synthase activity. *J Biol Chem.* 1998; 273:25179–25184. [PubMed: 9737978]
13. Kabil O, Toaka S, LoBrutto R, Shoemaker R, Banerjee R. Pyridoxal phosphate binding sites are similar in human heme-dependent and yeast heme-independent cystathionine beta-synthases. Evidence from ³¹P NMR and pulsed EPR spectroscopy that heme and PLP cofactors are not proximal in the human enzyme. *J Biol Chem.* 2001; 276:19350–19355. [PubMed: 11278994]
14. Jhee KH, McPhie P, Miles EW. Yeast cystathionine beta-synthase is a pyridoxal phosphate enzyme but, unlike the human enzyme, is not a heme protein. *J Biol Chem.* 2000; 275:11541–11544. [PubMed: 10766767]
15. Nozaki T, Shigeta Y, Saito-Nakano Y, Imada M, Kruger WD. Characterization of transsulfuration and cysteine biosynthetic pathways in the protozoan hemoflagellate, *Trypanosoma cruzi*. Isolation and molecular characterization of cystathionine beta-synthase and serine acetyltransferase from *Trypanosoma*. *J Biol Chem.* 2001; 276:6516–6523. [PubMed: 11106665]

16. Koutmos M, Kabil O, Smith JL, Banerjee R. Structural basis for substrate activation and regulation by cystathionine beta-synthase (CBS) domains in cystathionine (beta)-synthase. *Proc Natl Acad Sci U S A*. 2010; 107:20958–20963. [PubMed: 21081698]
17. Weeks CL, Singh S, Madzelan P, Banerjee R, Spiro TG. Heme regulation of human cystathionine beta-synthase activity: insights from fluorescence and Raman spectroscopy. *J Am Chem Soc*. 2009; 131:12809–12816. [PubMed: 19722721]
18. Evande R, Ojha S, Banerjee R. Visualization of PLP-bound intermediates in hemeless variants of human cystathionine beta-synthase: evidence that lysine 119 is a general base. *Arch Biochem Biophys*. 2004; 427:188–196. [PubMed: 15196993]
19. Singh S, Madzelan P, Banerjee R. Properties of an unusual heme cofactor in PLP-dependent cystathionine beta-synthase. *Nat Prod Rep*. 2007; 24:631–639. [PubMed: 17534535]
20. Singh S, Madzelan P, Stasser J, Weeks CL, Becker D, Spiro TG, Penner-Hahn J, Banerjee R. Modulation of the heme electronic structure and cystathionine beta-synthase activity by second coordination sphere ligands: The role of heme ligand switching in redox regulation. *J Inorg Biochem*. 2009; 103:689–697. [PubMed: 19232736]
21. Smith AT, Su Y, Stevens DJ, Majtan T, Kraus JP, Burstyn JN. Effect of the Disease-Causing R266K Mutation on the Heme and PLP Environments of Human Cystathionine beta-Synthase. *Biochemistry*. 2012; 51:6360–6370. [PubMed: 22738154]
22. Yadav PK, Xie P, Banerjee R. Allosteric communication between the PLP and heme sites in the H2S-generator human cystathionine beta-synthase. *J Biol Chem*. 2012; 287:37611–37620. [PubMed: 22977242]
23. Taoka S, West M, Banerjee R. Characterization of the heme and pyridoxal phosphate cofactors of human cystathionine beta-synthase reveals nonequivalent active sites. *Biochemistry*. 1999; 38:2738. [PubMed: 10052944]
24. Vadon-Le Goff S, Delaforge M, Boucher JL, Janosik M, Kraus JP, Mansuy D. Coordination chemistry of the heme in cystathionine beta-synthase: formation of iron(II)-isonitrile complexes. *Biochem Biophys Res Commun*. 2001; 283:487–492. [PubMed: 11327727]
25. Taoka S, Green EL, Loehr TM, Banerjee R. Mercuric chloride-induced spin or ligation state changes in ferric or ferrous human cystathionine beta-synthase inhibit enzyme activity. *J Inorg Biochem*. 2001; 87:253–259. [PubMed: 11744063]
26. Celano L, Gil M, Carballal S, Duran R, Denicola A, Banerjee R, Alvarez B. Inactivation of cystathionine beta-synthase with peroxynitrite. *Arch Biochem Biophys*. 2009; 491:96–105. [PubMed: 19733148]
27. Taoka S, Banerjee R. Characterization of NO binding to human cystathionine beta-synthase: possible implications of the effects of CO and NO binding to the human enzyme. *J Inorg Biochem*. 2001; 87:245–251. [PubMed: 11744062]
28. Ingi T, Chiang G, Ronnett GV. The regulation of heme turnover and carbon monoxide biosynthesis in cultured primary rat olfactory receptor neurons. *J Neurosci*. 1996; 16:5621–5628. [PubMed: 8795618]
29. Wu L, Wang R. Carbon monoxide: endogenous production, physiological functions, and pharmacological applications. *Pharmacol Rev*. 2005; 57:585–630. [PubMed: 16382109]
30. Pazicni S, Cherney MM, Lukat-Rodgers GS, Oliveriusova J, Rodgers KR, Kraus JP, Burstyn JN. The heme of cystathionine beta-synthase likely undergoes a thermally induced redox-mediated ligand switch. *Biochemistry*. 2005; 44:16785–16795. [PubMed: 16363792]
31. Carballal S, Madzelan P, Zinola CF, Grana M, Radi R, Banerjee R, Alvarez B. Dioxxygen reactivity and heme redox potential of truncated human cystathionine beta-synthase. *Biochemistry*. 2008; 47:3194–3201. [PubMed: 18278872]
32. Kabil O, Weeks CL, Carballal S, Gherasim C, Alvarez B, Spiro TG, Banerjee R. Reversible heme-dependent regulation of human cystathionine beta-synthase by a flavoprotein oxidoreductase. *Biochemistry*. 2011; 50:8261–8263. [PubMed: 21875066]
33. Olteanu H, Banerjee R. Human methionine synthase reductase, a soluble P-450 reductase-like dual flavoprotein, is sufficient for NADPH-dependent methionine synthase activation. *J Biol Chem*. 2001; 276:35558–35563. [PubMed: 11466310]

34. Wolthers KR, Basran J, Munro AW, Scrutton NS. Molecular dissection of human methionine synthase reductase: determination of the flavin redox potentials in full-length enzyme and isolated flavin-binding domains. *Biochemistry*. 2003; 42:3911–3920. [PubMed: 12667082]
35. Banerjee RV, Harder SR, Ragsdale SW, Matthews RG. Mechanism of reductive activation of cobalamin-dependent methionine synthase: an electron paramagnetic resonance spectroelectrochemical study. *Biochemistry*. 1990; 29:1129–1135. [PubMed: 2157485]
36. Gherasim CG, Zaman U, Raza A, Banerjee R. Impeded electron transfer from a pathogenic FMN domain mutant of methionine synthase reductase and its responsiveness to flavin supplementation. *Biochemistry*. 2008; 47:12515–12522. [PubMed: 18980384]
37. Kashiwamata S, Greenberg DM. Studies on cystathionine synthase of rat liver. Properties of the highly purified enzyme. *Biochim Biophys Acta*. 1970; 212:488–500. [PubMed: 5456996]
38. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*. 1976; 72:248–254. [PubMed: 942051]
39. Schellenberg KA, Hellerman L. Oxidation of reduced diphosphopyridine nucleotide. *J Biol Chem*. 1958; 231:547–556. [PubMed: 13538990]
40. Balahura RJ, Johnson MD. Outer-sphere dithionite reductions of metal complexes. *Inorganic Chemistry*. 1987; 26:3860–3863.
41. Patil PV, Ballou DP. The use of protocatechuate dioxygenase for maintaining anaerobic conditions in biochemical experiments. *Anal Biochem*. 2000; 286:187–192. [PubMed: 11067739]
42. Carballal S, Trujillo M, Cuevasanta E, Bartsaghi S, Moller MN, Folkes LK, Garcia-Bereguian MA, Gutierrez-Merino C, Wardman P, Denicola A, Radi R, Alvarez B. Reactivity of hydrogen sulfide with peroxynitrite and other oxidants of biological interest. *Free Radic Biol Med*. 2011; 50:196–205. [PubMed: 21034811]
43. Cotton, F.; Wilkinson, G. *Advanced Inorganic Chemistry*. New York: 1988.
44. Mendes P. GEPASI: a software package for modelling the dynamics, steady states and control of biochemical and other systems. *Comput Appl Biosci*. 1993; 9:563–571. [PubMed: 8293329]
45. Lambeth DO, Palmer G. The kinetics and mechanism of reduction of electron transfer proteins and other compounds of biological interest by dithionite. *J Biol Chem*. 1973; 248:6095–6103. [PubMed: 4353631]
46. Aviram I, Sharabani M. Kinetic studies of the reduction of neutrophil cytochrome b-558 by dithionite. *Biochem J*. 1986; 237:567–572. [PubMed: 3026324]
47. Weiland TR, Kundu S, Trent JT, Hoy JA 3rd, Hargrove MS. Bis-histidyl hexacoordination in hemoglobins facilitates heme reduction kinetics. *J Am Chem Soc*. 2004; 126:11930–11935. [PubMed: 15382928]
48. Creutz C, Sutin N. Kinetics of the reactions of sodium dithionite with dioxygen and hydrogen peroxide. *Inorganic Chemistry*. 1974; 13:2041–2043.
49. Su Y, Majtan T, Freeman KM, Linck RC, Ponter S, Kraus JP, Burstyn JN. Comparative study of enzyme activity and heme reactivity in *Drosophila melanogaster* and *Homo sapiens* cystathionine beta-synthases. *Biochemistry*. 2013; 52:741–751. [PubMed: 23002992]
50. Wolthers KR, Scrutton NS. Electron transfer in human methionine synthase reductase studied by stopped-flow spectrophotometry. *Biochemistry*. 2004; 43:490–500. [PubMed: 14717604]
51. Chen X, Jhee KH, Kruger WD. Production of the neuromodulator H₂S by cystathionine beta-synthase via the condensation of cysteine and homocysteine. *J Biol Chem*. 2004; 279:52082–52086. [PubMed: 15520012]
52. Chiku T, Padovani D, Zhu W, Singh S, Vitvitsky V, Banerjee R. H₂S biogenesis by human cystathionine gamma-lyase leads to the novel sulfur metabolites lanthionine and homolanthionine and is responsive to the grade of hyperhomocysteinemia. *J Biol Chem*. 2009; 284:11601–11612. [PubMed: 19261609]
53. Singh S, Padovani D, Leslie RA, Chiku T, Banerjee R. Relative contributions of cystathionine beta-synthase and gamma-cystathionase to H₂S biogenesis via alternative trans-sulfuration reactions. *J Biol Chem*. 2009; 284:22457–22466. [PubMed: 19531479]

54. Cherney MM, Pazicni S, Frank N, Marvin KA, Kraus JP, Burstyn JN. Ferrous human cystathionine beta-synthase loses activity during enzyme assay due to a ligand switch process. *Biochemistry*. 2007; 46:13199–13210. [PubMed: 17956124]
55. Puranik M, Weeks CL, Lahaye D, Kabil O, Taoka S, Nielsen SB, Groves JT, Banerjee R, Spiro TG. Dynamics of carbon monoxide binding to cystathionine beta-synthase. *J Biol Chem*. 2006; 281:13433–13438. [PubMed: 16505479]
56. Mayhew SG. The redox potential of dithionite and SO₂ from equilibrium reactions with flavodoxins, methyl viologen and hydrogen plus hydrogenase. *Eur J Biochem*. 1978; 85:535–547. [PubMed: 648533]
57. Wolthers KR, Scrutton NS. Protein interactions in the human methionine synthase-methionine synthase reductase complex and implications for the mechanism of enzyme reactivation. *Biochemistry*. 2007; 46:6696–6709. [PubMed: 17477549]
58. Finn RD, Basran J, Roitel O, Wolf CR, Munro AW, Paine MJ, Scrutton NS. Determination of the redox potentials and electron transfer properties of the FAD- and FMN-binding domains of the human oxidoreductase NR1. *Eur J Biochem*. 2003; 270:1164–1175. [PubMed: 12631275]
59. Shintani T, Iwabuchi T, Soga T, Kato Y, Yamamoto T, Takano N, Hishiki T, Ueno Y, Ikeda S, Sakuragawa T, Ishikawa K, Goda N, Kitagawa Y, Kajimura M, Matsumoto K, Suematsu M. Cystathionine beta-synthase as a carbon monoxide-sensitive regulator of bile excretion. *Hepatology*. 2009; 49:141–150. [PubMed: 19085910]
60. Yamamoto T, Takano N, Ishiwata K, Suematsu M. Carbon monoxide stimulates global protein methylation via its inhibitory action on cystathionine beta-synthase. *J Clin Biochem Nutr*. 2011; 48:96–100. [PubMed: 21297920]
61. Morikawa T, Kajimura M, Nakamura T, Hishiki T, Nakanishi T, Yukutake Y, Nagahata Y, Ishikawa M, Hattori K, Takenouchi T, Takahashi T, Ishii I, Matsubara K, Kabe Y, Uchiyama S, Nagata E, Gadalla MM, Snyder SH, Suematsu M. Hypoxic regulation of the cerebral microcirculation is mediated by a carbon monoxide-sensitive hydrogen sulfide pathway. *Proc Natl Acad Sci U S A*. 2012; 109:1293–1298. [PubMed: 22232681]

Figure 1. Kinetics of Fe(III)-CBS reduction by dithionite

(A) Fe(III)-CBS (2 μ M) was mixed with sodium dithionite (0.1, 0.5, 1, 3 and 4 mM, gray traces) in sodium phosphate buffer (0.2 M, pH 7.4, 0.2 mM DTPA) under anaerobic conditions at 25 °C. All concentrations given are for after mixing. The absorbance at 449 nm was fit to a single exponential function to determine the k_{obs} . (B) The observed rate constants (k_{obs}) were plotted against dithionite concentration and against the square root of dithionite concentration (*inset*).

Figure 2. Kinetics of Fe(II)-CBS reaction with CO

Stopped-flow UV-visible absorption spectra after mixing Fe(III)-CBS in CO-saturated phosphate buffer (0.1 M, pH 7.4) with dithionite in CO-saturated water at 25 °C. The concentrations of Fe(III)-CBS, dithionite and CO after mixing were 1 μ M, ~26 μ M and ~1 mM, respectively. Spectra were collected every 25 s for 500 s. The arrows indicate the direction of the absorbance change over time. *Inset.* Absorbance at 449 nm (circles) was plotted against time. The solid line represents the fit to an exponential function.

Figure 3. CBS reduction by MSR in the presence of CO

(A) Fe(III)-CBS (1 μM) was mixed with MSR (2 μM) and NADPH (1 mM) in phosphate buffer (0.1 M, pH 7.4) at 25 $^{\circ}\text{C}$ under anaerobic conditions. Then, the reaction mixture was saturated with CO ($\sim 1 \text{ mM}$) for 15 min and changes in the UV-visible spectrum were monitored. The dotted line represents the spectra of the reaction mixture before saturation with CO. *Inset*. Absorbance at 420 nm was plotted against time. (B) Ionic strength dependence of the reductive carbonylation of CBS by MSR. The reaction mixtures were prepared in phosphate buffer with increasing KCl concentrations as described under Experimental Procedures. The rates of the reaction (v_o) were determined from the linear increment of absorbance at 420 nm in the first 10 min.

Figure 4. Stopped-flow kinetics of CBS reduction by MSR in the presence of CO. A solution containing Fe(III)-CBS (1 μ M) and CO ($\sim$ 500 μ M) was mixed with MSR (2 μ M) pre-reduced by NADPH (500 μ M) in phosphate buffer (0.1 M, pH 7.4 with DTPA 0.1 mM) at 25 $^{\circ}$ C under anaerobic conditions (concentrations after mixing). Spectra were collected from 0.75 to 1425 s. For clarity, only selected spectra are shown. *Inset.* Expanded view of the UV-visible spectra between 400 and 475 nm. The arrows indicate the direction of the spectral evolution as a function of time.

Figure 5. Stopped-flow analysis of the formation of Fe(II)-CBS in the presence of MSR and CO
 (A) Kinetic traces at 449 nm were obtained from the UV-visible spectra in Fig. 4, with Fe(III)-CBS (1 μM) and CO ($\sim 500 \mu\text{M}$) in phosphate buffer (0.1 M, pH 7.4 with DTPA 0.1 mM) at 25 $^{\circ}\text{C}$ under anaerobic conditions, in the absence (dotted line) or presence of various concentrations of NADPH-reduced MSR (as indicated). Traces shown are the mean of duplicates for a representative experiment. (B) Initial rates were obtained from linear fits in the first 15 s. Each value is the mean $\pm$ standard deviation ($n = 5$) of data obtained on two different days.

Figure 6. Stopped-flow analysis of the formation of Fe(II)CO-CBS in the presence of MSR and CO

(A) Kinetic traces at 420 nm were obtained from the UV-visible spectra in Fig. 4, with Fe(III)-CBS (1 μM) and CO (~500 μM) in phosphate buffer (0.1 M, pH 7.4 with DTPA 0.1 mM) at 25 °C under anaerobic conditions, in the presence of various concentrations of NADPH-reduced MSR (as indicated). Representative traces from two independent experiments are shown. (B) Rates were obtained from linear fits of data at 40–80 or 100–250 s, depending on the trace. Each experimental value (closed circles) is the mean $\pm$ standard deviation ($n = 3$) of data from two independent experiments. Open symbols represent computer simulations performed for the reaction sequence shown in equations 4 and 5 using the initial concentrations of CBS and MSR and values for k_4 and k_{-4} of 9.3×10^2 and $1.2 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$, respectively. k_5 was assumed to be $5 \times 10^{-3} \text{ s}^{-1}$ (triangles) or $8 \times 10^{-3} \text{ s}^{-1}$ (squares).

Figure 7. Oxidation of Fe(II)CO-CBS by O₂

An anaerobic solution of Fe(II)CO-CBS (5 μ M) was mixed with O₂ (0.625 mM) in the presence of CO (500 μ M) and SOD (0.25 μ M monomer), in phosphate buffer (0.1 M, pH 7.4, DTPA 0.1 mM) at 25 °C. UV-visible absorption spectra were registered from 0.75 to 1425 s. Selected spectra (from top to bottom at 420 nm) recorded at 2.25, 3.75, 5.25, 8.25, 12.75, 20.25, 30.75, 52.5, 100.5, 150, 300, 603, 801, 1017, 1209, and 1425 s after mixing are shown.

Figure 8. Kinetic traces of the reaction between Fe(II)CO-CBS and O₂

The absorbance traces at 420 nm as a function of time are shown up to (A) 28 s, (B) 142 s and (C) 1400 s. The data were obtained from the UV-visible spectra in Fig. 7. The gray lines represent the best fits to a double exponential (A and C) and double exponential plus straight line (B) functions.

Figure 9. Dependence of the k_{obs} for Fe(II)CO-CBS reaction with O_2
Kinetic spectra were obtained as in Fig. 7 at increasing O_2 concentrations (0.0625–0.625 mM). Representative k_{obs} for the second phase of the reaction obtained from the global fit to a double exponential function of two independent experiments recorded for 148 s are shown. The black line represents the best fit to a hyperbolic function.

Scheme 1. Reversible reduction of CBS by MSR in the presence of CO

Table 1

Reactions involved in the reduction and reoxidation of the heme in CBS

Reaction	Rate constant ^a	Reference
$S_2O_4^{2-} \rightleftharpoons 2 SO_2^{\bullet -}$	$2.5 s^{-1}$ (f) ^b $1.8 \times 10^9 M^{-1} s^{-1}$ (r)	(45, 48)
$Fe(III)\text{-CBS} + SO_2^{\bullet -} \rightarrow Fe(II)\text{-CBS} + SO_2$	$(9.5 \pm 0.2) \times 10^6 M^{-1} s^{-1}$	This study
$Fe(III)\text{-CBS} + MSR_{red} \rightleftharpoons Fe(II)\text{-CBS} + MSR_{ox}$	$(93 \pm 2.5) \times 10^{-2} M^{-1} s^{-1}$ (f) ^b $1.2 \times 10^5 M^{-1} s^{-1}$ (r)	This study
$Fe(II)\text{-CBS} + O_2 \rightarrow Fe(III)\text{-CBS} + O_2^{\bullet -}$	$(1.13 \pm 0.05) \times 10^5 M^{-1} s^{-1}$	(31)
$Fe(II)\text{-CBS} + CO \rightarrow Fe(II)CO\text{-CBS}$	$(3.1 \pm 0.4) \times 10^{-3} s^{-1}$ at 1 mM CO ^c	This study
$Fe(II)CO\text{-CBS} + O_2 \rightarrow Fe(III)\text{-CBS} + \text{Products}$	$0.033 \pm 0.013 s^{-1}$ at infinite O_2 ^d	This study
$Fe(III)\text{-CBS} + H_2S \rightarrow \text{Products}$	Not detected	This study
$Fe(II)\text{-CBS} + H_2S \rightarrow \text{Products}$	Not detected	This study

^aRate constants at pH 7.4 and 25 °C except for the homolysis of dithionite, reported at pH 6.5(45, 48).

^b(f), Forward; (r), reverse.

^cPseudo-first order rate constant (k_{OBS}) obtained at 1 mM CO, see Discussion.

^dMultiphasic process, limiting pseudo-first order rate constant (k_{OBS}) of the major phase, that has hyperbolic dependence on oxygen concentration, see Results and Discussion.