Published in final edited form as:

Org Lett. 2013 July 5; 15(13): 3480–3483. doi:10.1021/ol401514s.

Enantioselective Synthesis of the 5-6-7 Carbocyclic Core of the Gagunin Diterpenoids

Grant M. Shibuya, John A. Enquist Jr., and Brian M. Stoltz

The Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology, 1200 East California Boulevard, MC 101-20, Pasadena, CA 91125, USA

Brian M. Stoltz: stoltz@caltech.edu

Abstract

A catalytic enantioselective double allylic alkylation reaction has been employed in the synthesis of the core of the gagunin diterpenoids. Enantioenriched material was advanced in eleven steps to afford the core of the highly-oxygenated target, which includes two all-carbon quaternary stereocenters.

The gagunin family of diterpenoids (**1–7**, Figure 1) were isolated from the sponge *Phorbas* sp. off the coast of South Korea by Shin and coworkers.¹ A wide range of cytotoxicity (LC_{50} = 0.03–50 $\mu\text{g/mL}$) was reported for gagunins A–G against the human leukemia cell line K562. The cytotoxicity of the gagunins is modulated by differential hydroxylation or esterification level present in a specific gagunin, as Shin and coworkers found gagunins A (**1**) and B (**2**) to be less active than other family members, likely due to the presence of a butyrate group at C(11) of these molecules.¹ Additionally, a perhydroxylated derivative of gagunin A (**8**) exhibited no cytotoxicity, further suggesting the importance of substitution patterns. Gagunin E (**5**) displayed an LC_{50} value of 0.03 $\mu\text{g/mL}$ (50 nM), the most potent cytotoxicity within this family of natural products. In the course of a reisolation of the gagunins from *Phorbas* sp., gagunin E was not found despite the reisolation of all other known gagunins and the discovery of ten new gagunins.^{1b} Given the scarcity of gagunin E in nature and broad range of biological activity displayed by the gagunin family, we sought to pursue a general synthetic route toward the gagunins to allow for additional biological

Correspondence to: Brian M. Stoltz, stoltz@caltech.edu.

The authors declare no competing financial interest.

Supporting Information Available: Experimental details and NMR spectra of all intermediates. These materials are available free of charge via the Internet at <http://pubs.acs.org>.

profiling and structure-activity relationship studies. Herein we report our progress toward a synthesis of the carbocyclic core of the gagunin family.

The carbon skeleton of the gagunins resembles the carbon skeleton of the cyathane family of diterpenoids.² A general route toward the cyathanes was previously reported by our laboratory using our enantioselective Tsuji allylation methodology.³ Bis- β -ketoester **9**, comprised of a mixture of racemic and *meso* diastereomers, was treated with a catalytic amount of Pd₂(pmdba)₃ and the ligand (*S*)-*t*-Bu-PHOX (**10**)⁴ to forge two all-carbon quaternary stereocenters in a single operation, affording diketone **11** in 99% ee as a 4.4:1 mixture of (*R,R*) to *meso*-diastereomers (Scheme 1). Diketone **11** was converted to enol triflate **12**, a common intermediate used in the syntheses of cyanthiwiggins F (**13**),⁵ B (**14**),^{5b} and G (**15**).^{5b}

Our retrosynthetic analysis of gagunin E (**5**) is shown in Scheme 2. We planned to employ a similar overall strategy in our approach to the gagunins, as was used in the cyanthiwiggins. Retrosynthetically, we envisioned a late-stage installation of the 5-membered ring, preceeded by establishing the 7-membered ring through a ring-closing methathesis (RCM) reaction. Such an approach would again allow for the early and essentially simultaneous introduction of both quaternary stereocenters.

In the forward direction, enol triflate **12** was subjected to conditions reported by Mulzer and coworkers to effect an intermolecular Heck reaction at the hindered neopentyl enol triflate.⁶ The addition of silyl ketene acetal **17**,⁷ Pd(PPh₃)₄, and LiOAc afforded methyl ester **18** in 77% yield (Scheme 3).⁸ Next, ketone **18** was converted to the corresponding ketal (**19**) in 47% yield by treatment with ethylene glycol and catalytic *p*-toluenesulfonic acid in benzene at reflux. Methyl ester **19** was then exposed to *N,O*-dimethylhydroxylamine hydrochloride and isopropylmagnesium chloride to afford Weinreb amide **20** in 93% yield. The addition of vinylmagnesium bromide to amide **20** at –20 °C furnished enone **21** in 74% yield.

Enone **21** was treated with Hoveyda–Grubbs generation 2 catalyst **22**⁹ in benzene at 40 °C to afford RCM adduct **23** in 85% yield (Scheme 4). The enone of **23** was next functionalized to the corresponding enol carbonate (**24**) in 93% yield after treatment with LHMDs followed by the addition of methyl chloroformate. A selective Wacker oxidation was performed on the terminal allyl group of tetraene **24**, giving methyl ketone **25** in 61% yield.¹⁰ Further purification by HPLC allowed **25** to be isolated as a single diastereomer. Previously, the separation of such diastereomers was accomplished only after closure of the final ring.⁵ Ketone **25** was next diazotized using Danheiser's diazo transfer conditions¹¹ to afford **26** in 47% yield over two steps.

With diazoketone **26** in hand, we proceeded to form the final carbocycle of **5**. Treatment of **26** with catalytic Rh₂(OAc)₄ afforded cyclopropane product **27** in 71% yield.^{12,13} The structure of **27** was confirmed by single crystal X-ray analysis (Figure 2).¹⁴ Finally, enol carbonate cleavage and cyclopropane opening was accomplished by exposing ketone **27** to K₂CO₃ in methanol at 0 °C, giving the desired ring-opening product **28** and an unexpected rearranged product (**29**) in a 1:1.8 ratio as determined by crude ¹H NMR analysis. The structure of **29** was again unambiguously confirmed by single crystal X-ray diffraction following chromatographic purification of **28** and **29**.¹⁵ Ultimately, desired dienone **28** was isolated in 31% yield and cyclopropane **29** in 27% yield; the change in ratio between the crude analysis and isolated yields suggest cyclopropane **29** may be unstable to chromatography conditions. Cyclopropane **29** is presumably formed from desired product **28** by a retro-norcaradiene rearrangement after formation of an enolate following deprotonation at C(6).¹⁶

In summary, an expedient route toward the tricyclic core of the gagunins has been established, giving sixteen of twenty carbons present in the core of these diterpenoids and the full tricyclic skeleton. The 7-membered ring was synthesized via an intermolecular Heck reaction at a hindered neopentyl carbon followed by a ruthenium-catalyzed RCM reaction of an enone and an allyl group. The 5-membered ring was prepared from an allyl group via a Wacker oxidation and ring-forming cyclopropanation from a diazoketone. Efforts to optimize this sequence and carry **28** and **29** forward to the gagunins and their analogs are ongoing.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors wish to thank NIH-NIGMS (R01GM080269-01), Amgen, Abbott, Boehringer Ingelheim, and Caltech for financial support. Mr. Austin Moehle (Caltech) is thanked for the preparation of early compounds. Dr. David Vander Velde (Caltech) is gratefully acknowledged for assistance with the characterization of compounds by NMR spectroscopy. Mr. Lawrence Henling (Caltech) is gratefully acknowledged for X-ray crystallographic structural determination and Dr. Alexander Goldberg (Caltech) is acknowledged for assisting in the preparation of crystallographic images for publication. Ms. Kelly Kim (Caltech) is acknowledged for assistance with the IR data of a late-stage compound. Materia, Inc. and Dr. Keith Keitz (Caltech) are thanked for the generous donation of ruthenium catalysts. The Bruker KAPPA APEXII X-ray diffractometer was purchased via an NSF CRIF:MU award to the California Institute of Technology, CHE-0639094. We also wish to acknowledge helpful discussions with Christina White (UIUC), Huw Davies (Emory), and other members of the CCI Center for Selective C–H Functionalization supported by NSF (CHE-1205646).

References

1. (a) Rho JR, Lee HS, Sim CJ, Shin J. *Tetrahedron*. 2002; 58:9585–9591. (b) Jang KH, Jeon J, Ryu S, Lee HS, Oh KB, Shin J. *J Nat Prod*. 2008; 71:1701–1707. [PubMed: 18811205]
2. (a) Green D, Goldberg I, Stein Z, Ilan M, Kashman Y. *Nat Prod Lett*. 1992; 1:193–199. (b) Peng J, Walsh K, Weedman V, Bergthold JD, Lynch J, Lieu KL, Braude IA, Kelly M, Hamann MT. *Tetrahedron*. 2002; 58:7809–7819. (c) Peng J, Avery MA, Hamann MT. *Org Lett*. 2003; 5:4575–4578. [PubMed: 14627387] (d) Peng J, Kasanah N, Stanley CE, Chadwick J, Fronczek FR, Hamann MT. *J Nat Prod*. 2006; 69:727–730. [PubMed: 16724830] (e) Enquist JA Jr, Stoltz BM. *Nat Prod Rev*. 2009; 26:661–680.
3. (a) Behenna DC, Stoltz BM. *J Am Chem Soc*. 2004; 126:15044–15045. [PubMed: 15547998] (b) Mohr JT, Behenna DC, Harned AM, Stoltz BM. *Angew Chem Int Ed*. 2005; 44:6924–6927. (c) Behenna DC, Mohr JT, Sherden NH, Marinescu SC, Harned AM, Tani K, Seto M, Ma S, Novák Z, Krout MR, McFadden RM, Roizen JL, Enquist JA Jr, White DE, Levine SR, Petrova KV, Iwashita A, Virgil SC, Stoltz BM. *Chem–Eur J*. 2011; 17:14199–14223. [PubMed: 22083969]
4. Helmchen G, Pfaltz A. *Acc Chem Res*. 2000; 33:336–345. [PubMed: 10891051]
5. (a) Enquist JA Jr, Stoltz BM. *Nature*. 2008; 453:1228–1231. [PubMed: 18580947] (b) Enquist JA Jr, Virgil SA, Stoltz BM. *Chem–Eur J*. 2011; 17:9957–9969. [PubMed: 21769952]
6. Magauer T, Mulzer J, Tiefenbacher K. *Org Lett*. 2009; 11:5306–5309. [PubMed: 19824621]
7. Oisaki K, Suto Y, Kanai M, Shibasaki M. *J Am Chem Soc*. 2003; 125:5644–5645. [PubMed: 12733898]
8. Our initial investigations centered on a direct RCM approach to a cycloheptadiene via pentaene **ii**, available by cross-coupling of **i** with enol triflate **12**. Unfortunately, RCM of **ii** produced a mixture of **iii** and **iv** when employing a range of ruthenium catalysts. Therefore, the approach outlined above was pursued.

9. (a) Garber SB, Kingsbury JS, Gray BL, Hoveyda AH. *J Am Chem Soc.* 2000; 122:8168–8179. (b) Gessler S, Randl S, Blechert S. *Tetrahedron Lett.* 2000; 41:9973–9976.
10. Using a Parr shaker was necessary to obtain product in good yield.
11. Danheiser R, Miller RF, Brisbois RG, Park SZ. *J Org Chem.* 1990; 55:1959–1964.
12. (a) Fernández-Mateos A, Pérez Alonso JJ, Rubio González R. *Tetrahedron.* 1999; 55:847–860. (b) Srikrishna A, Jagadreswar Reddy J. *J Chem Soc, Perkin Trans 1.* 2001:2040–2046.
13. Concurrent to the preparation of diazo ketone **26**, we prepared diazodiketone **v** using a similar approach. Treatment of **v** with $\text{Rh}_2(\text{OAc})_4$ did not yield a cyclopropanation, but instead a C–O insertion product tentatively assigned as **vi** was isolated in 38% yield.

14. Two conformations of the ketal of **27** were observed in the X-ray structure. Only one has been shown here.
15. For similar structures containing cyclopropane rings, please see: Nani RR, Reisman SE. *J Am Chem Soc.* 2013; 135:7304–7311. [PubMed: 23642221] Levin S, Nani RR, Reisman SE. *J Am Chem Soc.* 2011; 133:774–776. [PubMed: 21174417] Tseng CC, Ding H, Li A, Guan Y, Chen DYK. *Org Lett.* 2011; 13:4410–4413. [PubMed: 21770370] Ng W, Wege D. *Tetrahedron Lett.* 1996; 37:6797–6798. Ruppert JF, White JD. *J Am Chem Soc.* 1981; 103:1808–1813.
16. Alternatively, enol carbonate cleavage of **27** without cyclopropane fragmentation would give an enone that could be deprotonated at the C(6) position and allow for the formation of **29** directly from **27** after cyclopropane opening.

Figure 1. Gagunins A–G (1–7) and perhydroxylated gagunin A (8) and their reported biological activities.

Figure 2.
Single-crystal X-ray structures of **27** and **29**.

Scheme 1.

Enantioselective Tsuji Allylic Alkylation Applied to the Synthesis of Cyanthiwigins F (**13**), B (**14**), and G (**15**).

Scheme 2.
Retrosynthesis of gagunin E (5).

Scheme 3.
Synthesis of Enone **21**.

Scheme 4.
Synthesis of Diazoketone **26**.

Scheme 5. Completion of the Tricyclic Core of **5** via Cyclopropanation and Ring Opening.