Published in final edited form as:

J Pharm Sci. 2013 September ; 102(9): 2930–2940. doi:10.1002/jps.23590.

Moving from Basic towards Systems Pharmacodynamic Models

William J Jusko¹

¹Department of Pharmaceutical Sciences, School of Pharmacy and Pharmaceutical Sciences, State University of New York at Buffalo, Buffalo, NY 14214

Abstract

Building upon many classical foundations of pharmacology, a diverse array of mechanistic pharmacokinetic-pharmacodynamic (PK/PD) models have emerged based on mechanisms of drug action and primary rate-limiting or turnover processes in physiology. An array of basic models can be extended to handle various complexities including tolerance and can readily be employed as building blocks in assembling enhanced PK/PD or small systems models. Our corticosteroid models demonstrate these concepts as well as elements of horizontal and vertical integration of molecular to whole-body processes. The potential advantages and challenges in moving PK/PD towards systems models are described.

Keywords

Pharmacokinetics; pharmacodynamics; pharmacometrics; systems pharmacology; indirect response models

INTRODUCTION

The areas of pharmacokinetics and pharmacodynamics (PK/PD) have evolved from a long history of appreciation of basic pharmacological principles mostly applied to static or *in vitro* systems. An array of simple PK/PD models for the time course of *in vivo* drug effects have evolved to the present era of use of extended or enhanced PK/PD models and small-to-large systems models to capture drug actions at various levels of biological organization. This overview will describe the various arenas that have embraced PK/PD and pharmacometrics, highlight major concepts and features of commonly used PK/PD models, demonstrate model-building approaches leading to enhanced PK/PD and small systems models, and indicate the complications faced in evolving better quantitative methods for larger systems models.

Evolution of PK/PD and Pharmacometrics

The recognition that new mathematical relationships were needed to extend basic pharmacologic equations from static systems to *in vivo* time courses of drug effects began in 1965 (1) with the Levy “ $k \cdot m$ ” equation which provided the connection between pharmacokinetics (the k reflecting the monoexponential elimination rate constant) and pharmacology (the m being the mid-range slope of the Effect versus log drug concentration function). Gerhard Levy has been viewed as the “Father of Pharmacodynamics” for this and his many later contributions to PK/PD. Subsequently, with simulation studies, Wagner (2)

This article is partly based on presentations given at the American College of Clinical Pharmacology, Sept 2012, upon my receipt of the Distinguished Investigator Award and at the American Society of Clinical Pharmacology and Therapeutics, March 2013, upon my receipt of the Sheiner-Beal Pharmacometrics Award.

popularized the use of the Hill Function and demonstrated the value of ‘signature profiles’ [my term] to portray basic expectations of simple PK/PD functions.

These early contributions have blossomed into wide acceptance with many advancements in theory, and numerous applications of PK/PD and pharmacometrics in the pharmaceutical industry, government regulation, research institutes, and academia. The recent review by Lalonde et al (3) describes the utilization of modeling and simulation in the pharmaceutical industry pointing out how quantitative pharmacology can be implemented in each phase of drug development. The Food & Drug Administration embraced PK/PD in the early 1990’s and both early and recent reviews by the leadership of Peck, Lesko, and Gobburu (4–7) provide perspectives on how pharmacometrics has impacted the search for safer and more efficacious drugs in more efficient and timely fashion. The National Institutes of Health held a meeting in 2002 to assess training needs in the pharmacologic sciences. They concluded “There was a remarkable consensus that the core subject matter of pharmacology remains the principles of pharmacokinetics and pharmacodynamics” (8). This area is now widely taught, especially in Schools of Pharmacy. The NIH sponsored two recent symposia bringing together PK/PD modelers and system biologists and pharmacologists to consider the state-of-the-art and future of quantitative and systems pharmacology (QSP). This resulted in an extensive ‘white paper’ (9) which will hopefully lead to improved funding and new research (10). Accompanying these avenues of advancement of PK/PD has been the evolution of computational power and software programs such as WinNonlin, NonMem, Adapt, and many others. Numerous small companies provide consultation, data analysis, simulations, and pharmacometric reports for both the Pharma and generic companies. Led by the 1973 appearance of the *Journal of Pharmacokinetics and Biopharmaceutics* (now *Pharmacodynamics*) edited by Sidney Riegelman, Leslie Benet, and Malcom Rowland [I am currently Editor-in-Chief], the number of journals and published articles in quantitative pharmacology has exploded over the past 50 years. Journal articles which include ‘pharmacodynamics’ in the title in recent decades number in MEDLINE (11) as follows: (1963–72) **153**, (1973–82) **255**, (1983–92) **970**, (1993–2012) **1564**, and (2003–12) **1772**. Numerous scientific meetings include PK/PD and QSP topics and symposia. Several series of PK/PD specialty meetings have promoted the field including those hosted by David D’Argenio (Biomedical Simulation Resource) in Los Angeles, Meindert Danhof in the Netherlands, the American Conference on Pharmacometrics, and the Population Approach Group Europe (PAGE) meetings. Recently established was the International Society of Pharmacometrics (www.go-isop.org).

Basic Mechanism-Based PK/PD Models

Most “mechanism-based” PK/PD models include some recognition that one or more critical steps in drug action are controlled by the pharmacokinetics, receptor or target binding mechanism, and/or physiology including homeostatic and disease mechanisms yielding resolvable parameters for the major rate-limiting process(es). These three main components are depicted in Figure 1 (12). The applications of such models can be described as a “top-down” approach where the modeler should gain appreciation of the underlying pharmacokinetics, pharmacology, and physiology in ascribing a general model and designing studies and analysis of experimental data. The models typically seek parsimony. Model parameters are expected to have strong statistical reliability and relate to underlying processes (e.g. production or clearance of endogenous substances). Systems models, on the other hand, are “bottom up” in configuration with a wealth of equations and assumed parameters and are used for simulation purposes seeking to match expectations with profiles of experimental data and for exploratory purposes.

Pharmacokinetics

The pharmacokinetics serves as the driving force for drug access to target sites and the input rates and elimination processes control both the duration of drug exposure and often the duration of drug action. As portrayed in Figure 2, the major types of pharmacokinetic models are noncompartmental, compartmental, and physiological. Noncompartmental Analysis (NCA) is often a useful ‘black-box’ starting point to assess Clearance (CL) and steady-state volume of distribution (V_{ss}) for diagnostic purposes and preliminary assessment of properties such as linearity and stationarity. Compartmental models can be semi-mechanistic adding improved insights into distribution properties of drugs and organisms. Physiologically-based PK (PBPK) models are systems models where parameters are assigned using physiological measurements (blood flow, organ sizes) and resolved by direct analysis of plasma concentrations and tissue transport, binding, and metabolic properties. There is a long history of evolution of highly useful PBPK models for small molecules (13) and more recent success for biologics (14). They have great value in translating preclinical data to man, anticipating changes of physicochemical properties of drugs, and assessing the impact of altered physiology such as occurs in infants and children. We have recently demonstrated how highly ‘lumped’ or minimal PBPK models provide more useful assessment of PK properties of drugs than do classical compartment models (15).

One major PK goal is gaining understanding and resolving the time-course and factors determining drug access to target sites. This is known as ‘biophase distribution’ (16). The PBPK models may provide such insights if target sites are known and if local binding, metabolism, and transporters do not complicate predictions. Difficult to measure or verify are free or unbound drug concentrations in interstitial fluids and tissue cells. Two key PK/PD principles often followed in early drug development are demonstration of target exposure and target activity. Assessment of drug at target sites should be part of preclinical development of PK/PD models. Possible biophases other than plasma need consideration in modeling human data. Such alternatives may include arterial rather than venous blood or plasma, free drug at various sites, cerebrospinal fluid, drug in urine (for diuretics), and -of-course- specific organs or tissues as either measured, evaluated with PBPK models, or assessed by employing imaging methods.

Distribution of drugs to peripheral sites of action is often accompanied by a delay or slow onset of observed effects. Sheiner et al (17) in 1979 popularized the use of a diffusion-like equation to describe drug access via rate constant k_{eo} to a hypothetical biophase compartment (concentration C_e):

$$dC_e/dt = k_{eo} \cdot (C_p - C_e) \quad (1)$$

This equation is helpful for assessing clinical PD data if the biophase is unknown and is not measurable and if the rate-limiting step for a delayed drug effect is truly the rate of peripheral access. Values of k_{eo} should be examined to assure compatibility with PBPK principles (usually relatively rapid distribution rates). As will be described, numerous alternative mechanisms can account for slow onset of drug effects. This biophase model has often been misapplied for systems where other explanations for delayed effects are more plausible (18).

Receptor Binding

The nature and time-course of drug effects on the body have been observed and reported for many centuries, but the underlying quantitative principles emerged with the recognition by Ehrlich that “Corpora non agunt nisi fixata” [Substances do not act unless bound.] which laid

the foundation of receptor theory (19). Pharmacologists have long embraced the concept of “Receptor Occupancy” with equations predicated on the law of mass action as formulated by Clark (20) in terms of rate of association (k_{on}) and dissociation (k_{off}) of drug (A) and Receptor (R) concentrations:

This can be viewed as a rate equation:

$$dAR/dt = k_{on} \cdot A \cdot R - k_{off} \cdot AR \quad (3)$$

or equilibrium binding:

$$F_{occ} = AR/B_{max} = A/(K_D + A) \quad (4)$$

where F_{occ} is Fractional Occupancy, B_{max} is the total receptor concentration, and K_D is the Equilibrium Dissociation Constant (k_{off}/k_{on}). This relationship underlies the Hill Function (21) which is commonly employed for quantitating the nonlinear relationship between drug effects (E) and plasma (C_p) or biophase drug concentrations:

$$E = (E_{max} \cdot C_p^\gamma) / (EC_{50}^\gamma + C_p^\gamma) \quad (5)$$

where E_{max} is the maximum achievable effect, EC_{50} is the drug concentration associated with $1/2$ of E_{max} and γ is the Hill coefficient. An important later advance was the recognition by Black and Leff (22) that receptor binding may be an interface for signaling cascades which can be represented by a connected second nonlinear function (the operational model of agonism). The complex array of concepts and relationships needed to describe various drug and receptor or target interactions has grown to require textbooks for more complete consideration (23). One of the basic goals of PK/PD modeling is a translational one (12), from in vitro and preclinical to clinical, by seeking to identify the connections between Eq. (4) and (5) by relating B_{max} to E_{max} , K_D to EC_{50} , and in vitro (A) to plasma or biophase drug concentrations (C_p). This is where quantitative pharmacology can be employed in using an array of basic to systems PK/PD models.

In the most elemental PD situations, biophase distribution and receptor binding are extremely rapid and drug effects can be related to drug concentrations in plasma with the Hill Function. This was essentially the situation with the muscle relaxant effects of succinylcholine modeled by Levy (1). Such simple direct effects are most commonly encountered in situations where enzyme activity is measured in blood. One example is with the ACE inhibitory effects of benazeprilat where the target and drug are present together in blood (24). This is the one situation where time delays do not exist to produce hysteresis in the relationship between effects and plasma drug concentrations. Equations other than the Hill Function may be required for seemingly simple drug effects. Mandema et al (25) provided an early application of Black and Leff concepts to demonstrate that EEG effects of several benzodiazepines were similar in rats if plasma concentrations and K_D values were employed to calculate receptor occupancy (Eq. 4) as controlling drug effects.

The extent (capacity) of receptor binding and/or physiological signaling mechanisms often determine the efficacy of drug effects (viz. B_{max} may control E_{max}). Sometimes the binding k_{on} and k_{off} rates determine the time-course of drug effects. Shimada et al (26) used rate equations analogous to Eq. (3) to model the relationship between plasma concentrations and time-course of blood pressure effects of 8 calcium channel blockers, including nifedipine, in

hypertensive subjects. Of translational importance, they found a good correlation between the clinical K_D values calculated as k_{off}/k_{on} and in vitro receptor measurements of K_D for the 8 compounds. The PK phenomenon of 'target-mediated drug disposition' (TMDD) uses similar principles of target binding to account for certain types of nonlinear drug disposition (27). Such models were extended to describe the complexities in PD of interferon-beta by employing the drug-receptor complex as the controlling factor for the delayed effects of this compound (28).

Turnover and Homeostasis

The two basic tenets of pharmacodynamics are the expectations of capacity-limitation in target binding or activation of the drug and the presence of turnover and homeostasis in the functioning of physiological and disease processes (Figure 1). Figure 3 provides a range of biological structures and functions with an indication of their approximate turnover rates or durations. An essential feature of most organisms is that their components are continuously being produced and degraded, viz. undergoing turnover. The homeostatic element is that these systems are at relative steady-state, albeit subject to circadian, developmental, aging, and disease changes or other perturbations. Tolerance or functional adaptation mechanisms exist to maintain homeostasis. Measurements of any of the listed components could comprise the pharmacodynamics of a drug. The turnover rates help determine the time frame to carry out PD measurements. Rapid rates such as EEGs and EKGs can be captured in a few seconds while very slow rates such as changes in bone structure requires years of monitoring. The upper-mid part of the list tends to represent biomarkers which may signal alterations in more substantive body functions. The lower part of the list reflects the clinical effects (usually major body symptoms) including patient survival.

In conjunction with the mechanism of action, recognition of turnover rates also helps in identifying the type of mechanism-based PKPD model which may be necessary. Simple direct effect and biophase models are sufficient for rapid turnover processes as the time-course of drug concentrations (either in plasma or a biophase) will be rate-controlling. Indirect Response Models (IRM) will capture processes with intermediate turnover rates where production (k_{in}) or loss (k_{out}) may control observed responses. The slower turnover components of the body may have multiple steps contributing to production or loss which can often be captured with transduction or more complex models. These may suffice in the absence of a more complete array of measurements where enhanced PD or small systems models could be applied.

Diversity of Basic Turnover Models

Figure 4 demonstrates the seven primary mechanism-based PKPD models which are often employed as a starting point in the top-down analysis of the time-course of drug responses. The figure includes the direct effect (center right), biophase (top right), and receptor-binding (top left) models already described. The remaining models all include a turnover process along with a pharmacologic function for the mechanism of action of the drug and, of course, are controlled by the PK. Various complexities can be added to each of these models where deemed appropriate.

The center-left diagram depicts a basic model with self-replication of cells (k_s) and a drug-induced loss process (k_L). Originally proposed to account for the actions of cytotoxic anticancer drugs (29), this type of model has also been applied to some antibiotics (30) and antimalarial drugs (31). The center diagram shows a basic model with zero-order production (k_{in}), first-order loss (k_{out}), and an irreversible (second-order, $k \cdot C_p$) inactivation of the response parameter. This type of model has been used to account for the anti-platelet

inactivation of COX-1 by aspirin (32) as well as the effects of pantoprazole and other proton pump inhibitors causing reduction in gastric acidity (33).

The bottom-left diagram symbolizes the four basic indirect response models (IRM) which capture drug effects on either inhibition or stimulation of k_{in} , or inhibition or stimulation of k_{out} . These models have been widely applied to a variety of drug responses (34–35), including diverse antidiabetic drugs (18). An early example was capturing the antinociceptive (pain relief) effects of 6 dose levels of tolmetin in an inflammation model in rats using population PKPD modeling (36). The bottom-right model depicts a transduction model as compartments or steps with turnover of each with a common time constant τ (37). The drug is assumed to act in the first compartment (e.g. receptor binding) with the observed response evident at a later time as the last compartment. Many useful applications of this model have been published (38) including capturing chemotherapeutic effects on tumor xenografts (39).

Some extensions of these models exist which are useful for describing tolerance or functional adaptation in pharmacodynamics. For example, a biophase-like compartment can reflect receptor desensitization (40), a precursor step to an IRM can capture tolerance and rebound (41), a transit step can be added to an IRM to provide feedback alteration of production or loss (42), and counter-regulation can be represented as serial compartments with an intervening transduction step (43). The homeostatic and signaling mechanisms of the body feature a diverse array of feedback loops and control mechanisms (44) which will present both challenges and future opportunities for evolving ever more complex and mechanistic systems pharmacology models.

Enhanced Models

The types of basic mechanisms and models which have been described can be melded into more complex PKPD models if more comprehensive experiments, measurements, and data assemblies permit. The receptor-gene mediated effects of corticosteroids have been assessed in our “giant rat” studies where a group of animals were given various doses of methylprednisolone and blood and multiple tissues sampled at sacrifice to assess PK, tissue receptor content, tissue gene expression (both specific and with gene arrays), tissue enzymes or proteins, various circulating biomarkers, and sometimes measurable structures or functions. Our 5th-generation model had several simple components: PK, receptor-binding, transduction, and two types of IRM: one for stimulation of biomarker gene production and one for feedback inhibition of receptor gene production (45). This enhanced PKPD model was subsequently incorporated into a larger small systems model to describe disease progression in a rat model of rheumatoid arthritis (RA) (paw edema) and the effects of dexamethasone (Dex) on inhibition of pro-inflammatory cytokine genes in inflamed rat paws (46, 47). Figure 5 shows part of our experimental data including the PK of Dex, the disease progression of paw edema captured with transit compartments and reduction in cytokine expression by Dex producing decreased paw edema as an inhibitory IRM. The PK/PD/DIS model used to capture these data is shown in Figure 6. This small systems model utilizes our 5th-generation corticosteroid model (45) for receptor dynamics for binding of Dex and corticosterone (top left section) and largely employs transit compartments for the slower unmeasured immune responses producing inflammation and bone loss (right side and bottom). In principle, transit compartments serve as place-holders for natural or pathological phenomena which might be measured with additional methodologies or experiments. These efforts represent a mix of both horizontal and vertical integration as described in systems modeling (9). Our tissue measurements are horizontal in assessing hundreds of genes (and now proteins) at the molecular level (48). The modeling is vertical in connecting measurements of drug molecules, genes, proteins, presumed cellular events, and end-organ

(paw and bone) responses. Some RA animal studies also utilize a symptom score to reflect the degree of whole body dysfunction, much as employed in clinical studies in patients (49). These types of models have offered us a basis for general understanding of the major determinants of corticosteroid action, a means of assessing conceptual limitations when the models do not fit, and simulations allow for the optimal design of new experiments for seeking the next-generation advanced PKPD model.

Iyengar et al (50) provide other examples of enhanced PD (ePD) models to describe this middle ground of small systems or targeted PD models. Their report lists major features and utility of ePD models. It provides a process diagram for building ePD models beginning with available genomic/epigenomic data, incorporates drug-centered regulatory networks, and includes drug-target information to be used in enacting and assessing results from suitable PK/PD studies.

Systems Models

Systems biology and pharmacology models seek to identify all known biological processes controlling body events typically utilizing a vast array of differential equations with parameters assigned as best identified from diverse experiments found in the literature and/or by direct studies (9). With such extensive networks, the models operate in simulation mode with somewhat subjective adjustments made when they can be tested with available experimental data. This is traditionally how PBPK models have been operated, although our current computers and software have allowed for complete and population-type modeling of multiple-animal group PBPK studies (51). Complex systems models exist for describing some aspects of inflammation (52, 53) and for control of calcium homeostasis and bone remodeling (54). Commercial system model assemblies have been implemented by at least 3 companies (Entelos, Archimedes, Rosa) to model diabetes and allow theoretical assessment of various effects of possible antidiabetic agents.

Vodovitz et al (53) and Iyengar et al (50) have indicated the considerable potential of systems models as a repository for biologic, pharmacologic, and clinical knowledge. Figure 7 generalizes similar systems model perspectives as might be implemented for specific areas of disease and pharmacology. Building such models and data depositories would require a team approach utilizing the input of experts from various disciplines. They offer a platform for common understanding, enlightened discussion, efficient experimentation, and continual advancement of improved quantitative models.

Transitioning from Basic toward Systems Models

The NIH QSP Workshops (9) identified a basic tension as well as opportunities that exist in seeking targeted or enhanced PKPD models. As depicted in Figure 8, the bioinformatic, statistical, and computational needs vary depending on the complexity of the modeling task. Fitting basic and enhanced PKPD models, even with mixed-effect population assessments, can be readily managed with current software. The modeling metrics, while steadily evolving, are reasonably well established. There exist well-accepted diagnostic plots to determine how well model functions fit the data, procedures such as Visual Predictive Checks have become popular, standardization is evolving, and the displayed metrics are growing overly elaborate. Numerous software platforms exist offering both comparable and alternative fitting algorithms. For systems models, the array of equations are considerable, parameter sources are not always available, assumptions are sometimes unclear or approximate, and the models operate by simulation with some tweaking when comparing with experimental data. Basic PKPD modeling is moving towards a middle ground dealing with models of ever-increasing complexity where many components must be assigned based on best experimental or theoretical evidence, missing elements might need approximation

with empirical components (e.g. transit compartments), while major rate-limiting steps can be captured by more traditional fitting procedures. This was largely the approach needed for our PK/PD/DIS model in RA (Figure 6). Finding the best approaches between robust model fitting and simulation is our next major computational challenge in moving from basic towards systems models.

Acknowledgments

This work was supported by NIH Grants GM 24211 and 57980. The author appreciates the many contributions of his esteemed colleagues, collaborators, fellows, students, and technicians for those aspects of this work which is cited herein as well as forming the basis for his PKPD modeling efforts over the course of his career. The assistance of Katherine Majewski from the National Library of Medicine is also appreciated. Our efforts were part of the activities of the UB Center of Excellence in Pharmacokinetics and Pharmacodynamics. I also cherish the long-term friendship and interactions with Leslie Z. Benet in whose honor this article was constructed.

References

1. Levy G. Kinetics of pharmacologic effects. *Clin Pharmacol Ther.* 1966; 7(3):362–372. [PubMed: 5937003]
2. Wagner JG. Kinetics of pharmacologic response. I. Proposed relationships between response and drug concentration in the intact animal and man. *J Theor Biol.* 1968; 20(2):173–201. [PubMed: 5727238]
3. Lalonde RL, Kowalski KG, Hutmacher MM, Ewy W, Nichols DJ, Milligan PA, Corrigan BW, Lockwood PA, Marshall SA, Benincosa LJ, Tensfeldt TG, Parivar K, Amantea M, Glue P, Koide H, Miller R. Model-based drug development. *Clin Pharmacol Ther.* 2007; 82(1):21–32. [PubMed: 17522597]
4. Peck CC, Barr WH, Benet LZ, Collins J, Desjardins RE, Furst DE, Harter JG, Levy G, Ludden T, Rodman JH, Sanathanan L, Schentag JJ, Shah VP, Sheiner LB, Skelly JP, Stanski DR, Temple RJ, Viswanathan CT, Weissinger J, Yacobi A. Opportunities for integration of pharmacokinetics, pharmacodynamics, and toxicokinetics in rational drug development. *Pharm Res.* 1992; 9(6):826–833. [PubMed: 1409369]
5. Lesko LJ, Rowland M, Peck CC, Blaschke TF. Optimizing the science of drug development: opportunities for better candidate selection and accelerated evaluation in humans. *J Clin Pharmacol.* 2000; 40(8):803–814. [PubMed: 10934664]
6. Bhattaram VA, Bonapace C, Chilukuri DM, Duan JZ, Garnett C, Gobburu JV, Jang SH, Kenna L, Lesko LJ, Madabushi R, Men Y, Powell JR, Qiu W, Ramchandani RP, Tornoe CW, Wang Y, Zheng JJ. Impact of pharmacometric reviews on new drug approval and labeling decisions--a survey of 31 new drug applications submitted between 2005 and 2006. *Clin Pharmacol Ther.* 2007; 81(2): 213–221. [PubMed: 17259946]
7. Gobburu JV, Lesko LJ. Quantitative disease, drug, and trial models. *Annu Rev Pharmacol Toxicol.* 2009; 49:291–301. [PubMed: 18851702]
8. Preusch PC. News from the National Institute of General Medical Sciences: What is training in the pharmacological sciences? *Pharm Res.* 2002; 19(12):1771–1772. [PubMed: 12523653]
9. Sorger, PK.; Allerheiligen, SRB.; Abernethy, DR.; Altman, RB.; Brouwer, KLR.; Califano, A.; D'Argenio, DZ.; Iyenger, R.; Jusko, WJ.; Lalonde, R.; Lauffenburger, DA.; Shoichet, B.; Stevens, JL.; Subramaniam, S.; Van der Graaf, P.; Vicini, P. An NIH White Paper from the QSP Workshop Group – July, 2011. Quantitative and systems pharmacology in the postgenomic era: New approaches to discovering and understanding therapeutic drugs and mechanisms. <http://www.nigms.nih.gov/News/Reports/201110-syspharma.htm>
10. Rogers M, Lyster P, Okita R. NIH support of the emergence of quantitative and systems pharmacology. *CPT: Pharmacometrics & Systems Pharmacology.* 2013; 2:e37–e39. [PubMed: 23887687]
11. Search: pharmacodynamics[ti] AND medline[sb] AND yyyy[PDAT]: yyyy[PDAT]. Bethesda (MD): National Library of Medicine (US); 1946. PubMed [Internet]. [cited 2013 Apr 9]. Available from: <http://pubmed.gov/>

12. Mager DE, Jusko WJ. Development of translational pharmacokinetic-pharmacodynamic models. *Clin Pharmacol Ther.* 2008; 83:909–912. [PubMed: 18388873]
13. Rowland M, Peck C, Tucker G. Physiologically-based pharmacokinetics in drug development and regulatory science. *Annu Rev Pharmacol Toxicol.* 2011; 51:45–73. [PubMed: 20854171]
14. Garg A, Balthasar JP. Physiologically-based pharmacokinetic (PBPK) model to predict IgG tissue kinetics in wild-type and FcRn-knockout mice. *J Pharmacokinet Pharmacodyn.* 2007; 34(5):687–709. [PubMed: 17636457]
15. Cao Y, Jusko WJ. Applications of minimal physiologically-based pharmacokinetic models. *J Pharmacokin Pharmacodyn.* 2012; 39:711–723.
16. Furchgott RF. The pharmacology of vascular smooth muscle. *Pharmacol Rev.* 1955; 7(2):183–265. [PubMed: 13245382]
17. Sheiner LB, Stanski DR, Vozeh S, Miller RD, Ham J. Simultaneous modeling of pharmacokinetics and pharmacodynamics: application to d-tubocurarine. *Clin Pharmacol Ther.* 1979; 25(3):358–371. [PubMed: 761446]
18. Landersdorfer CB, Jusko WJ. Pharmacokinetic/pharmacodynamic modelling in diabetes. *Clin Pharmacokin.* 2008; 47:417–448.
19. Ehrlich P. Chemotherapeutics: Scientific principles, methods, and results. *Lancet.* 1913; 2:445–451.
20. Clark, AJ. The Mode of Action of Drugs on Cells. London: Edward Arnold; 1933.
21. Hill AV. The possible effects of the aggregation of the molecules of haemoglobin on its dissociation curves. *J Physiol.* 1910; 40:iv–vii.
22. Black JW, Leff P. Operational models of pharmacological agonism. *Proc R Soc Lond B Biol Sci.* 1983; 220:141–162. [PubMed: 6141562]
23. Kenakin, TP. Pharmacologic Analysis of Drug-Receptor Interaction. 3rd ed. New York: Raven Press; 1997.
24. Toutain PL, Lefebvre HP, King JN. Benazeprilat disposition and effect in dogs revisited with a pharmacokinetic/pharmacodynamic modeling approach. *J Pharmacol Exp Ther.* 2000; 292(3):1087–1093. [PubMed: 10688627]
25. Mandema JW, Sansom LN, Dios-Vièitez MC, Hollander-Jansen M, Danhof M. Pharmacokinetic-pharmacodynamic modeling of the electroencephalographic effects of benzodiazepines. Correlation with receptor binding and anticonvulsant activity. *J Pharmacol Exp Ther.* 1991; 257(1):472–478. [PubMed: 1850477]
26. Shimada S, Nakajima Y, Yamamoto K, Sawada Y, Iga T. Comparative pharmacodynamics of eight calcium channel blocking agents in Japanese essential hypertensive patients. *Biol Pharm Bull.* 1996; 19(3):430–437. [PubMed: 8924914]
27. Mager DE, Jusko WJ. General pharmacokinetic model for drugs exhibiting target-mediated drug disposition. *J Pharmacokin Biopharm.* 2001; 28:507–532.
28. Mager DE, Neuteboom B, Jusko WJ. Pharmacokinetics and pharmacodynamics of pegylated IFN- α 1a following subcutaneous administration in monkeys. *Pharm Res.* 2005; 22:58–61. [PubMed: 15771230]
29. Jusko WJ. Pharmacodynamics of chemotherapeutic effects: Dose-time-response relationships for phase-nonspecific agents. *J Pharm Sci.* 1971; 60:892–895. [PubMed: 5166939]
30. Zhi JG, Nightingale CH, Quintiliani R. Microbial pharmacodynamics of piperacillin in neutropenic mice of systematic infection due to *Pseudomonas aeruginosa*. *J Pharmacokinet Biopharm.* 1988; 16(4):355–375. [PubMed: 3193364]
31. Gordi T, Xie R, Jusko WJ. Semi-mechanistic pharmacokinetic/pharmacodynamic modelling of the antimalarial effect of artemisinin. *Brit J Clin Pharmacol.* 2005; 60:594–604. [PubMed: 16305583]
32. Yamamoto K, Abe M, Katashima M, Yamada Y, Sawada Y, Iga T. Pharmacodynamic analysis of antiplatelet effect of aspirin in the literature - modeling based on inhibition of cyclooxygenase in the platelet and the vessel wall endothelium. *Jpn J Hosp Pharm.* 1996; 22:133–141.
33. Ferron GM, McKeand W, Mayer PR. Pharmacodynamic modeling of pantoprazole's irreversible effect on gastric acid secretion in humans and rats. *J Clin Pharmacol.* 2001; 41(2):149–156. [PubMed: 11210394]

34. Dayneka N, Garg V, Jusko WJ. Comparison of four basic models of indirect pharmacodynamic responses. *J Pharmacokin Biopharm.* 1993; 21:457–478.
35. Jusko WJ, Ko HC. Physiologic indirect response models characterize diverse types of pharmacodynamic effects. *Clin Pharmacol Ther.* 1994; 56:406–419. [PubMed: 7955802]
36. Flores-Murrieta FJ, Ko HC, Flores-Acevedo DM, López-Muñoz FJ, Jusko WJ, Sale ME, Castañeda-Hernández G. Pharmacokinetic-pharmacodynamic modeling of tolmetin antinociceptive effect in the rat using an indirect response model: a population approach. *J Pharmacokin Biopharm.* 1998; 26(5):547–557. [PubMed: 10205770]
37. Sun Y-N, Jusko WJ. Transit compartments versus gamma distribution function to model signal transduction processes in pharmacodynamics. *J Pharm.Sci.* 1998; 87:732–737. [PubMed: 9607951]
38. Mager DE, Jusko WJ. Pharmacodynamic modeling of time-dependent transduction systems. *Clin Pharmacol Ther.* 2001; 70:210–216. [PubMed: 11557908]
39. Yang J, Mager DE, Straubinger RM. Comparison of two pharmacodynamic transduction models for the analysis of tumor therapeutic responses in model systems. *AAPS J.* 2010; 12(1):1–10. [PubMed: 19902363]
40. Mandema JW, Wada DR. Pharmacodynamic model for acute tolerance development to the electroencephalographic effects of alfentanil in the rat. *J Pharmacol Exp Ther.* 1995; 275(3):1185–1194. [PubMed: 8531080]
41. Sharma A, Ebling WF, Jusko WJ. Precursor-dependent indirect pharmacodynamic response model for tolerance and rebound phenomena. *J Pharm Sci.* 1998; 87:1577–1584. [PubMed: 10189270]
42. Karlsson MO, Anehall T, Friberg LE, Henningsson A, Kloft C, Sandström M, Xie R. Pharmacokinetic/pharmacodynamic modelling in oncological drug development. *Basic Clin Pharmacol Toxicol.* 2005; 96(3):206–11. [PubMed: 15733216]
43. Bauer JA, Fung HL. Pharmacodynamic models of nitroglycerin-induced hemodynamic tolerance in experimental heart failure. *Pharm Res.* 1994; 11(6):816–823. [PubMed: 7937519]
44. Brandman O, Meyer T. Feedback loops shape cellular signals in space and time. *Science.* 2008; 322:390–395. [PubMed: 18927383]
45. Ramakrishnan R, DuBois D, Almon RA, Pyszczynski NA, Jusko WJ. Fifth-generation model for corticosteroid pharmacodynamics: Steady-state receptor down-regulation and enzyme induction patterns during seven-day continuous infusion of methylprednisolone in rats. *J Pharmacokin Pharmacodyn.* 2002; 29:1–24.
46. Earp JC, DuBois DC, Molano DS, Pyszczynski NA, Keller CE, Almon RR, Jusko WJ. Modeling corticosteroid effects in a rat model of rheumatoid arthritis I: Mechanistic disease progression model for the time course of collagen-induced arthritis in Lewis rats. *J Pharmacol Exp Ther.* 2008; 326:532–545. [PubMed: 18448865]
47. Earp JC, DuBois DC, Molano DS, Pyszczynski NA, Almon RR, Jusko WJ. Modeling corticosteroid effects in a rat model of rheumatoid arthritis II: Mechanistic pharmacodynamic model for dexamethasone effects in Lewis rats with collagen-induced arthritis. *J Pharmacol Exp Ther.* 2008; 326:546–554. [PubMed: 18448864]
48. Yang E, Almon RR, DuBois DC, Jusko WJ, Androulakis IP. Extracting global system dynamics of corticosteroid genomic effects in rat liver. *J Pharmacol Exp Ther.* 2008; 324:1243–1254. [PubMed: 18086872]
49. Lon H-K, Liu D, Jusko WJ. Pharmacokinetic/pharmacodynamic modeling in inflammation. *Crit Rev Biomed Eng.* 2012; 40:295–312. [PubMed: 23140121]
50. Iyengar R, Zhao S, Chung SW, Mager DE, Gallo JM. Merging systems biology with pharmacodynamics. *Sci Transl Med.* 2012; 4(126):126ps7.
51. Pawaskar DK, Straubinger RM, Fetterly GJ, Hylander BH, Repasky EA, Ma WW, Jusko WJ. Physiologically based pharmacokinetic models for everolimus and sorafenib in mice. *Cancer Chemother Pharmacol.* 2013 (online).
52. An G. In silico experiments of existing and hypothetical cytokine-directed clinical trials using agent-based modeling. *Crit Care Med.* 2004; 32(10):2050–2060. [PubMed: 15483414]
53. Vodovotz Y, Csete M, Bartels J, Chang S, An G. Translational systems biology of inflammation. *PLoS Comput Biol.* 2008; 4(4):e1000014. [PubMed: 18437239]

54. Peterson MC, Riggs MM. A physiologically based mathematical model of integrated calcium homeostasis and bone remodeling. *Bone*. 2010; 46:49–63. [PubMed: 19732857]

Figure 1. Mechanism-based pharmacokinetic – pharmacodynamics (PK/PD) models typically integrate the time-course of drug concentrations (PK) including biophase distribution, the nature of drug-target interaction (Pharmacology), and turnover processes reflecting the relevant Physiology and Disease (12).

Array of PK Models

Figure 2. Various modeling approaches to assessment of pharmacokinetic data including Noncompartmental (NCA), Compartmental, and Physiologically-Based PK.

Biological Turnover Rates of Structures or Functions

Figure 3.
Selected physiological structures and functions and their approximate turnover rates.

Major PK/PD Models

Figure 4. Seven major types of basic PK/PD models commonly utilized based on mechanisms of drug action and primary rate-limiting steps in the kinetic, target-binding, or physiological process.

Figure 5.

Multi-level assessment of the pharmacokinetics of dexamethasone (Dex), production of a pro-inflammatory gene (IL-1 β) in the rat paw and its inhibition by Dex, and the time-course of Paw Edema after collagen injection in rats showing natural RA Disease Progression and inhibitory effects of various dosing regimens of Dex. Curves with the data are PK/PD model fittings using the model shown in Figure 6. Experimental data are from studies of Earp et al (46, 47).

Figure 6. Pharmacokinetic/Pharmacodynamic/Disease (PK/PD/DIS) model of Earp et al (46, 47) used to characterize the PK, receptor binding, disease progression, and Dex alteration of gene expression resulting in amelioration of Paw Edema in rats with RA. Measured entities are marked in yellow.

Figure 7.
Conceptualization of the information flow, integration of knowledge, and potential applications of systems pharmacology data bases and models (50, 53).

Levels of Modeling Complexity

Figure 8.
Range and types of modeling complexity at three modeling levels of Quantitative and Systems Pharmacology (QSP).