Published in final edited form as:

ACS Chem Biol. 2013 June 21; 8(6): 1110–1116. doi:10.1021/cb4001052.

Persulfide Reactivity in the Detection of Protein S-Sulfhydration¹

Jia Pan and Kate S. Carroll*

Department of Chemistry, The Scripps Research Institute, Jupiter, FL 33456, USA

Abstract

Hydrogen sulfide (H₂S) has emerged as new member of the gaseous transmitter family of signaling molecules and appears to play a regulatory role in the cardiovascular and nervous systems. Recent studies suggest that protein cysteine S-sulfhydration may function as a mechanism for transforming the H₂S signal into a biological response. However, selective detection of S-sulfhydryl modifications is challenging since the persulfide group (RSSH) exhibits reactivity akin to other sulfur species, especially thiols. A modification of the biotin switch technique, using S-methyl methanethiosulfonate (MMTS) as an alkylating reagent, was recently used to identify a large number of proteins that may undergo S-sulfhydration, but the underlying mechanism of chemical detection was not fully explored. To address this key issue, we have developed a protein persulfide model and analog of MMTS, S-4-bromobenzyl methanethiosulfonate (BBMTS). Using these new reagents, we investigated the chemistry in the modified biotin switch method and examined the reactivity of protein persulfides toward different electrophile/nucleophile species. Together, our data affirm the nucleophilic properties of the persulfide sulfane sulfur and afford new insights into protein S-sulfhydryl chemistry, which may be exploited in future detection strategies.

Introduction

Oxidative post-translational modification (oxPTM) of protein cysteines plays an important regulatory role in many physiological systems.^{1–3} Of these, protein S-sulfhydration has recently received attention in the context of hydrogen sulfide (H₂S) mediated signaling pathways^{4–7} involved in cardiovascular and cerebral vasodilatory responses.^{8,9} To date, the precise mechanisms of H₂S action are still the subject of active investigation. A significant challenge in protein S-sulfhydration research is that the persulfide group (RSSH) exhibits reactivity akin to other sulfur species, in particular thiols,^{10–12} which compounds difficulties in developing selective detection methods.

In regards to detection, our interest was piqued when a commonly used alkylating agent, known as S-methyl methanethiosulfonate (MMTS),¹³ was reported to differentiate thiols and persulfides using a modified biotin switch technique (BST),⁴ originally designed to study protein S-nitrosylation.¹⁴ As shown in Figure 1a, in the modified BST thiols are first alkylated with MMTS and persulfides were postulated to remain unreacted and be available for subsequent conjugation to N-[6-(biotinamido) hexyl]-3'-(2'-pyridyldithio) propionamide (biotin-HPDP). Using this modified version of the BST, a large number of proteins were reported as targets of H₂S in the mouse liver and the basal sulfhydration level of some

¹This work was supported by the National Institutes of Health (Grant No. GM102187).

*Correspondence should be addressed to: Kate S. Carroll, The Scripps Research Institute, Scripps Florida, 130 Scripps Way, Jupiter, FL 33458, kcarroll@scripps.edu, Phone: (561) 228-2460, Fax: (561) 228-2919.

Supporting Information

This material is available free of charge via the Internet at <http://pubs.acs.org>.

proteins was estimated to be as high as 25%.⁴ Despite these intriguing results, the structure of reaction products in each step of the modified BST and the underlying mechanism of thiol vs. persulfide MMTS selectivity was not investigated.

Herein, we report several persulfide models and new MMTS analog, *S*-4-bromobenzyl methanethiosulfonate (BBMTS). Using these reagents, we investigated the chemistry in the modified BST and examined the reactivity of protein persulfides toward different electrophile/nucleophile species. These data reaffirm the nucleophilic properties of the persulfide sulfane sulfur and afford new insights into protein *S*-sulfhydryl chemistry, which may be exploited in future detection strategies.

Results

Although selective (albeit slow) methods for H₂S detection have rapidly accumulated,^{15–18} methods for detecting protein persulfide modifications have been slower to advance. In this case, the challenges are two fold: i) the chemical reactivity of the terminal sulfane sulfur in the protein persulfide is similar to the thiol, and ii) few examples of robust *S*-sulfhydryl protein models have been reported. To this end, we initiated studies to explore the generation and reactivity of protein persulfides. Our second objective was to understand the chemical mechanisms underlying current chemoproteomic methods of persulfide detection.

In an effort to identify suitable models, we first investigated the reported method¹⁹ for preparing persulfide derivatives of the tripeptide glutathione. As illustrated in Figure 2a, the GSH persulfide was obtained as an equilibrium mixture of glutathione thiol (GSH, **2**) and persulfide (GSSH, **3**) after the reaction of glutathione disulfide (GSSG, **1**) with Na₂S. Though GSH persulfide **3** could not be directly observed by liquid chromatography - mass spectrometry (LC-MS), its formation was confirmed through conjugation with alkylating agents such as iodoacetamide (IAM) and *N*-ethylmaleimide (NEM), which yielded detectable products in LC-MS analyses (Figure 2b and Figure S1). The inability to directly observe the persulfide species likely stems from the instability of the underivatized modification in acidic conditions and/or poor ionization properties.

In parallel investigations, single protein persulfide modifications were generated *via* a stepwise procedure in the cysteine protease, papain and the C64,82S thiol peroxidase, Gpx3. In this method, reduced thiol **6** was first reacted with 5,5'-dithiobis-2-nitrobenzoic acid (DTNB or Ellman's reagent) to form activated disulfide **7**, which was then cleaved by Na₂S to generate persulfide **8** (Figure 2c). After conditions such as the equivalents of DTNB and Na₂S were optimized, papain and Gpx3 persulfides were successfully prepared and confirmed by intact MS analysis post-trapping with IAM or NEM (Figure 2d). In addition, to the desired papain persulfide derivative **9**, we observed signals consistent with papain sulfinic acid **10** (as-isolated from the original commercial sample) as well as persulfide oxidation. Of note, these species were also observed in the reaction with NEM (Figure S2), indicating these mass signals were not unique to a particular alkylating reagent. Interestingly, in the case of Gpx3, the alkylation reaction revealed the presence of both the persulfide (major derivative) and the trisulfide (minor derivative; Figures 2e and S3).

Next, we tested the reaction of the persulfide models with MMTS. As shown in Figure 3a, the reaction of glutathione persulfide with MMTS gave a small amount of GSSH derivative **14**, with major product **15** derived from GSH, and other byproducts GSSG (**1**) and GSSSG (**13**). The desired papain persulfide **16** and Gpx3 persulfide **18** products were clearly observed as the major species, albeit with some overlap between protein sulfonates and thiosulfates (*i.e.*, papain **17** in Figure 3b; Gpx3 **20** and **21** in Figure 3c). These analyses also highlight the small mass difference (~2 Da; Figure 3d) between byproducts from the

oxidation of thiols and persulfides, relative to the MMTS-modified sulfur, which makes it difficult to distinguish between these chemically distinct species by intact protein MS analysis.

To address the above issue and clarify whether the ~80 Da mass increase corresponds to –SSMe or –SSO₃[–] forms of GSH, papain, and Gpx3, we designed and synthesized an analog of MMTS, *S*-4-bromobenzyl methanethiosulfonate (BBMTS; Figure 4a). Compared to MMTS, the advantage of this reagent is two-fold: i) a larger mass change results from the thio-BBMTS adduct (+200 Da), and ii) BBMTS exhibits the telltale bromine mass pattern (*i.e.*, strong peaks at *M*–1 and *M*+1), which can be discerned in low-molecular-weight peptides, including GSH. Initially, we verified the reactivity of BBMTS, which proved comparable to MMTS (Tables S1 and S2). We then tested the BBMTS reagent in reactions with the GSH and protein persulfide models. As shown in Figure 4b, treatment of glutathione persulfide with BBMTS, gave a small amount of the desired persulfide derivative **25** and a large peak of the thiol derivative **26** (both confirmed using the bromine signature afforded by the new reagent), as well as the thiosulfate byproduct **27** (also detailed in Figures S4–7 and Table S3). On the other hand, the reaction between protein persulfides and BBMTS clearly showed the formation of the desired persulfide derivatives, as well as alkyl polysulfides (Figures 4c and 4d). Taken together, these data unequivocally establish that the nucleophilic sulfane sulfur of protein persulfides reacts with electrophilic MMTS and BBMTS reagents.

Since the MMTS and BBMTS electrophiles react with protein persulfides, an important question arises: what does the biotin signal in the modified BST stem from? To address this issue, we treated papain and Gpx3 products of the MMTS/BBMTS alkylation step with the biotin-HPDP analog, *N*-acetylcysteine pyridyldisulfide (NACP). Ensuing intact MS analysis of these reactions indicated the following: i) no product corresponding to a NACP-labeled persulfide was detected in papain or Gpx3, ii) a small amount of NACP-labeled thiol was observed in papain (**32**; Figures 5a and 5b), and iii) no NACP-labeled thiol was observed in Gpx3 (Figures 5c and 5d). These studies suggest that the protein-NACP labeling product is derived from thiol alkylation.

Finally, since little has been reported about the chemical reactivity of protein persulfides, we investigated the behavior of the Gpx3-SSH sulfane sulfur with various nucleophilic and electrophilic reagents (Table 1 and Figure S8). The nucleophilicity of the sulfane sulfur was reaffirmed in reactions with electrophilic alkylating reagents such as IAM, NEM and MMTS (**Entry 1–3**), as well as disulfide-containing compounds, like DTNB and NACP (**Entry 4–5**). As an electrophile, the sulfane sulfur could be reduced to the free thiol form by DTT (**Entry 6**), GSH (note: free thiol and mixed disulfide forms; **Entry 7**), and NAC (**Entry 8**). By contrast, weaker sulfur-based nucleophiles like sulfite or methyl sulfinatate did not reduce the Gpx3 persulfide (**Entry 9–10**). We also tested the reaction of Gpx3 persulfide with carbon nucleophiles known to react with sulfenic acid (RSOH), but not disulfides or nitrosothiols (RSNO). These studies show that dimedone, malononitrile, methyl acetoacetate and barbituric acid carbon nucleophiles exhibited no reactivity toward the Gpx3 persulfide (**Entries 11–14**). As before, thiosulfate formation was also noted in the persulfide samples, most likely resulting from oxidation by trace metal ions in the presence of oxygen.²⁰ The thiosulfate species was stable and also did not react with the carbon nucleophiles (Figures S9, S13F-K and Table S4).

Discussion

We have investigated GSH and protein persulfide models for chemical reactivity studies. Among these prototypes, GSH and papain persulfides have been previously described;¹⁹

however, the protein persulfides were not evaluated or confirmed by mass analysis. Using this early study as inspiration, in this work we have examined the utility of a new persulfide protein model, based on a single-cysteine mutant of thiol peroxidase, Gpx3. This model proved quite facile for the generation of protein persulfides and intact MS analysis was not complicated by the presence of a large fraction sulfinic acid-modified protein, in contrast to that observed for commercial sources of papain. With respect to the GSH persulfide model, we observed only a small amount of persulfide IAM and NEM alkylation products. The most likely explanation for these data is that the reaction of sterically unhindered persulfide derivatives **14** and **25** with low molecular weight thiols (*i.e.*, GSH, GSSH) is rapid, relative to chemical alkylation. Thus, compared to protein persulfide models, the GSH persulfide may have less utility in reactivity studies.

In addition, we have examined the chemistry underlying the derivitization steps in the modified BST using the GSH and protein persulfide model systems. In contrast to earlier supposition,⁴ we demonstrate that the terminal persulfide sulfur undergoes facile reaction with both MMTS and BBMTS, and that these alkylated forms do not react with the NACP pyridyldisulfide. These findings thus reaffirm the nucleophilic reactivity of the sulfane sulfur atom in protein persulfides and are consistent with physical organic studies^{10, 11} and the IAM/NEM labeling observed in this work. In fact, the only NACP-labeled species that we observed in this study corresponded to the reaction product derived from thiol, as opposed to the persulfide. As shown in Figure 6, two possible models could account for this finding. In one scenario, free thiols may be incompletely blocked in the first MMTS alkylation step and subsequently react with the pyridyldisulfide biotin reagent (Figure 6a). Alternatively, or in addition, biotin labeling may be achieved *via* stepwise thiol-disulfide exchange in a reaction catalyzed by trace free thiols (Figure 6b). A common element in both models is that not all free thiols are blocked during the MMTS labeling step.

Recently, it was also reported that the persulfide-modified active site cysteine of protein tyrosine phosphatase 1B (PTP1B) can be alkylated by iodoacetic acid (IAA). Subsequent reduction of the persulfide adducts and labeling with iodoacetamide-linked biotin (IAP) has also been proposed as an indirect approach for detecting protein *S*-sulfhydration (Figure 1b).⁶ However, it is not apparent how this method distinguishes persulfide modifications from other DTT-reducible modifications, such as disulfide bonds, sulfenic acids, and nitrosothiols (Figure 1c). Nevertheless, IAA derivitization of the PTP1B persulfide provides further evidence that the persulfide is nucleophilic with similar reactivity to thiols. Likewise, the γ -sulfur of persulfide is a weak electrophile that shares some reactivity with disulfides (also alluded to by sulfur transfer mechanisms in Fe-S biosynthetic pathways¹²), but is chemically distinct from the sulfur atom in sulfenic acid, which is significantly more electrophilic in nature. Collectively these data reaffirm the nucleophilic properties of the persulfide sulfane sulfur and afford new insights into protein *S*-sulfhydryl chemistry, which can serve as a resource in developing future detection strategies.

Materials and Methods

Sodium sulfide nonahydrate ($\text{Na}_2\text{S}\cdot 9\text{H}_2\text{O}$), extra pure, was obtained from ACROS. DTNB was obtained from Sigma and the purity was 98%. Papain 2x crystallized was purchased from Sigma. All other chemicals were purchased from Sigma-Aldrich or Fisher at the highest available purity.

Preparation of persulfides

(1) Glutathione persulfide¹⁹—To a solution of freshly prepared glutathione (oxidized form, 1 mM final concentration) in 1 mL Tris-HCl buffer (100 mM, pH 7.4), was added freshly prepared sodium sulfide (1 mM final concentration), and the solution was stirred at

37 °C for 15 min. The resulting glutathione persulfide (1 mM final concentration) was directly used without further purification.

(2) Papain persulfide¹⁹—A solution of papain (10 mg) in 1 mL Tris-HCl (pH 7.4, 100 mM, degassed with N₂ for 10 min) was incubated with cysteine (1.7 mg) at rt for 10 min, and the protein was purified through a PD10 column (GE Healthcare) containing G-25 Sephadex. Protein-containing fractions were pooled from this column (500 PL, 270 μM) and incubated with DTNB (50 μL, 4 mM) at rt for 20 min, and purified by passing through a PD-10 column. A fraction of the pooled protein (200 μL, 120 μM) was then incubated in a solution of Na₂S (4 μL, 30 mM) at rt for 10 min, and purified by PD-10 column to give the papain persulfide, which was immediately used in further reactions.

(3) Gpx3 persulfide—C64,82S Gpx3 was prepared as previously described.^{21,22} A solution of C64,82S Gpx3 (100 μM, 100 μL) was buffer exchanged through a P-30 spin filtration column (BioRad) that had been equilibrated in Tris-HCl buffer (pH 7.4, 100 mM, degassed with N₂ for 10 min). After determination of protein concentration (80 μM), the solution was incubated with DTNB (4 μL, 4 mM) at rt for 20 min, and purified again spin filtration. The resulting protein (100 μL, 62 μM) was then incubated with a Na₂S solution (1 μL, 30 mM) at rt for 10 min, and purified by spin column to give the Gpx3 persulfide, which was immediately used in downstream reactions.

Verification of persulfide formation and analysis of chemical reactivity—

Freshly prepared GSH or protein persulfide (final concentration: 1 mM for glutathione; 20 μM for protein) was incubated with IAM or other nucleophilic or electrophilic reagents (final concentration: 4 mM for glutathione; 2 mM for protein) at rt for 1 h. After purification, aliquots of the samples were analyzed by electrospray LC/MS on Poroshell 120 (Agilent) or ZIC[®]-pHILIC HPLC column (the Nest group).

Reaction of GSH with MMTS or BBMTS—To a solution of freshly prepared glutathione persulfide (1 mM final concentration) in 1 mL Tris-HCl buffer (100 mM, pH 7.4), was added MMTS or BBMTS (4 mM final concentration), and the solution was stirred at rt for 30 min. Sample aliquots were generated at each step and analyzed by LC-MS.

Reaction of protein persulfides with MMTS or BBMTS followed by NACP—A freshly prepared protein persulfide solution (20 μM final concentration) was incubated with the methanethiosulfonate (2 mM final concentration) at rt for 30 min, and then purified through a P-30 spin column. The sample solution was then incubated with NACP (2 mM final concentration) at rt for 3 h, and purified as before. Sample aliquots were generated at each step and analyzed by LC-MS.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Walsh CT, Garneau-Tsodikova S, Gatto GJ. Protein post translational modifications: the chemistry of proteome diversifications. *Angew. Chem. Int. Ed.* 2005; 44:7342–7372.
2. Reddie KG, Carroll KS. Expanding the functional diversity of proteins through cysteine oxidation. *Curr. Opin. Chem. Biol.* 2008; 12:746–754. [PubMed: 18804173]
3. Mieyal JJ, Chock PB. Posttranslational modification of cysteine in redox signaling and oxidative stress: focus on S-glutathionylation. *Antioxidants & Redox Signaling.* 2012; 16:471–475. [PubMed: 22136616]

4. Mustafa AK, Gadalla MM, Sen N, Kim S, Mu W, Gazi SK, Barrow RK, Yang G, Wang R, Snyder SH. H₂S signals through protein S-sulfhydration. *Sci. Signal.* 2009; 2:ra72. [PubMed: 19903941]
5. Kabil O, Banerjee R. Redox biochemistry of hydrogen sulfide. *J. Biol. Chem.* 2010; 285:21903–21907. [PubMed: 20448039]
6. Krishnan N, Fu C, Pappin DJ, Tonks NK. H₂S-induced sulfhydration of the phosphatase PTP1B and its role in the endoplasmic reticulum stress response. *Sci. Signal.* 2011; 4:ra86. [PubMed: 22169477]
7. Sen N, Paul BD, Gadalla MM, Mustafa AK, Sen T, Xu R, Kim S, Snyder SH. Hydrogen sulfide-linked sulfhydration of NF- κ B mediates its antiapoptotic actions. *Mol. Cell.* 2012; 45:13–24. [PubMed: 22244329]
8. Zhao W, Zhang J, Lu Y, Wang R. The vasorelaxant effect of H₂S as novel endogenous gaseous gaseous KATP channel opener. *EMBO J.* 2001; 20:6008–6016. [PubMed: 11689441]
9. Olson KR, Dombkowski RA, Russell MJ, Doellman MM, Head SK, Whitfield NL, Madden JA. Hydrogen sulfide as an oxygen sensor/transducer in vertebrate hypoxic vasoconstriction and hypoxic vasodilation. *J. Exp. Biol.* 2006; 209:4011–4023. [PubMed: 17023595]
10. Flavin M. Microbial transsulfuration: the mechanism of an enzymatic disulfide elimination reaction. *J. Biol. Chem.* 1962; 237:768–777. [PubMed: 13893449]
11. Heimer NE. Biologically oriented organic sulfur chemistry. 21-Hydrogendisulfide of a penicillamine derivative and related compounds. *J. Org. Chem.* 1981; 46:1374–1377.
12. Mueller EG. Trafficking in persulfides: delivering sulfur in biosynthetic pathways. *Nat. Chem. Biol.* 2006; 2:185–194. [PubMed: 16547481]
13. Smith DJ, Maggio ET, Kenyon GL. Simple alkanethiol groups for temporary blocking of sulfhydryl groups of enzymes. *Biochemistry.* 1975; 14:766–771. [PubMed: 163643]
14. Jaffrey SR, Snyder SH. The biotin switch method for the detection of S-nitrosylated proteins. *Sci. STKE.* 2001; 86:pl1. [PubMed: 11752655]
15. Lippert AR, New EJ, Chang CJ. Reaction-based fluorescent probes for selective imaging of hydrogen sulfide in living cells. *J. Am. Chem. Soc.* 2011; 133:10078–10080. [PubMed: 21671682]
16. Liu C, Pan J, Li S, Zhao Y, Wu LY, Berkman CE, Whorton AR, Xian M. Capture and visualization of hydrogen sulfide by a fluorescent probe. *Angew. Chem. Int. Ed.* 2011; 50:10327–10329.
17. Qian Y, Karpus J, Kabil O, Zhang S, Zhu H, Banerjee R, Zhao J, He C. Selective fluorescent probes for live-cell monitoring of sulphide. *Nat. Commun.* 2011; 2:495. [PubMed: 21988911]
18. Lin VS, Chang CJ. Fluorescent probes for sensing and imaging biological hydrogen sulfide. *Curr. Opin. Chem. Biol.* 2012; 16:595–601. [PubMed: 22921406]
19. Francoleon NE, Carrington SJ, Fukuto JM. The reactivity of H₂S with oxidized thiols: Generation of persulfides and implications to H₂S biology. *Arch. Biochem. Biophys.* 2011; 516:146–153. [PubMed: 22001739]
20. Everett SA, Wardman P. Perthiols as antioxidants: radical-scavenging and prooxidative mechanisms. *Methods in Enzymology.* 1995; 251:55–69. [PubMed: 7651231]
21. Mason JT, Kim SK, Knaff DB, Wood MJ. Thermodynamic basis for redox regulation of the Yap1 signal transduction pathway. *Biochem.* 2006; 45:13409–13417. [PubMed: 17087494]
22. Paulsen CE, Carroll KS. Chemical dissection of an essential redox switch in yeast. *Chem. Biol.* 2009; 16:217–225. [PubMed: 19230722]

Figure 1.

Reported methods for detecting protein persulfide modifications. **a)** Modified biotin switch technique (BST): protein samples are alkylated by MMTS to block free thiols, while sparing persulfides. In the next step, biotin-HPDP is applied to alkylate persulfides and form protein-biotin conjugates. **b)** Thiols and persulfides are reacted with iodoacetic acid (IAA). The persulfide is then reduced with dithiothreitol (DTT) and conjugated to biotinylated IAP. **c)** Other cysteine oxPTMs that may be identified using the method in **b**.

Figure 2.

Preparation and verification of persulfide formation in GSH and protein models. **a)** Formation of glutathione (GSH) persulfide. **b)** HPLC UV trace at 254 nm and corresponding masses from the reaction in **a**. **c)** Formation of protein persulfides. Mass analysis of the reaction between iodoacetamide (IAM) and: **d)** papain persulfide, and **e)** Gpx3 persulfide [*indicates the product derived from alkylation of Gpx3-S-S-SH (i.e., trisulfide)].

Figure 3.

Reactivity of GSH and protein persulfide models with MMTS. **a**) HPLC UV trace at 254 nm and corresponding masses from the reaction of GSH persulfide and MMTS (*the mass in this peak could not be assigned to a chemical structure). Mass analysis of the reaction between: **b**) papain persulfide and MMTS, and **c**) Gpx3 persulfide and MMTS. **d**) Increase in molecular mass that results from MMTS modification (–SSMe) or oxidation to the thiosulfate (–SO₃[–]).

Figure 4. Reactivity of the GSH and protein persulfide models with BBMTS. **a)** Synthesis of *S*-4-bromobenzyl methanethiosulfonate (BBMTS). Mass analyses of the reaction between: **b)** GSH persulfide and BBMTS. **c)** papain persulfide and BBMTS [⁺papain-SO₃⁻ (sulfonic acid); *papain-S-SO₃⁻ (thiosulfate)], and **d)** Gpx3 persulfide and BBMTS.

Figure 5. Sequential labeling reactions of protein persulfides with MMTS or BBMTS and biotin-HPDP analog, *N*-acetylcysteine pyridyldisulfide (NACP). Mass analyses of the reaction between: **a)** papain persulfide, MMTS and then NACP, **b)** papain persulfide, BBMTS, and then NACP [⁺papain-SO₃⁻ (sulfonic acid); *papain-S-SO₃⁻ (thiosulfate)], **c)** Gpx3 persulfide, MMTS, and then NACP [*Gpx3-S-S-S-SMe (tetrasulfide)], and **d)** Gpx3 persulfide, BBMTS, and then NACP as described in the Methods [*Gpx3-S-S-S-SCH₂C₆H₄Br (tetrasulfide)].

Figure 6.

Proposed labeling mechanism of protein persulfide, methanthiosulfonate (MMTS or BBMTS), and pyridyldisulfide NACP. Two possible models are possible. **a**) Free thiols may be incompletely blocked in the first MMTS alkylation step and subsequently react with the pyridyldisulfide biotin reagent. **b**) Alternatively, or in addition, labeling may be achieved *via* stepwise thiol-disulfide exchange in a reaction catalyzed by trace free thiols (RSH).

Table 1

Reactivity of Gpx3 persulfide towards electrophilic and nucleophilic reagents.

Entry	Reagent	Product	Entry	Reagent	Product
1	IAM		8	NAC
2	NEM		9	Sulfite	No reaction
3	MMTS		10	Methylsulfinic acid	No reaction
4	DTNB		11	Dimedone	No reaction
5	NACP		12	Malononitrile	No reaction
6	DTT		13	Methyl acetoacetate	No reaction
7	GSH		14	N,N'-Dimethylbarbituric acid	No reaction