Sleep Apnea and Cardiometabolic Risk in South Asians

Commentary on Leong et al. The prevalence and severity of obstructive sleep apnea in severe obesity: the impact of ethnicity. *J Clin Sleep Med* 2013;9:853-858.

Namratha R. Kandula, M.D., M.P.H.¹; Sanjay R. Patel, M.D.²

¹Division of General Internal Medicine, Northwestern University, Chicago, IL;

²Division of Sleep Medicine, Brigham and Women's Hospital, Boston, MA

People of South Asian descent, including those from India, Pakistan, Bangladesh, Sri Lanka, and Nepal, make up a quarter of the world's population. For over a century, the South Asian diaspora has also led to a substantial number of South Asians living in Western countries.¹ South Asians represent 7% and 3% of the population in England and Canada, respectively. In the US, South Asians have been one of the fastest growing population segments and now represent the second largest Asian American group. South Asians throughout the world have been shown to have a disproportionate risk of cardiovascular disease (CVD) and type 2 diabetes mellitus (DM).²⁻⁴ There are clearly lifestyle factors contributing to this increased cardiometabolic risk⁵; however, surprisingly little is known about sleep in South Asians and whether it may also be contributing to the increased risk. This is of particular interest given that obstructive sleep apnea (OSA) and other causes of poor sleep are increasingly linked to CVD and DM.^{6,7}

Epidemiologic studies from India suggest the prevalence of OSA is similar to US whites, despite a lower level of obesity.^{8,9} However, few studies have directly compared OSA risk in South Asians to other groups. Questionnaire-based surveys suggest people of South Asian heritage are at higher risk than those of European or Chinese heritage, but the relative performance of these questionnaires across races has never been assessed.¹⁰⁻¹²

In this issue of the *Journal*, Leong et al. report on differences in the prevalence of OSA in severely obese South Asians and whites in a UK bariatric clinic where all patients underwent limited-channel home sleep testing.¹³ South Asians had a higher prevalence of OSA (43%) than whites (22%) and a substantially greater apnea hypopnea index. Differences remained significant even after adjustment for potential confounders such as body mass index (BMI). These data confirm that severely obese South Asians are at elevated risk for OSA compared to whites. The authors are to be commended for performing the first cross-ethnic assessment including South Asians utilizing an objective assessment of OSA severity. While the possibility of differential biases for referral to a bariatric program needs to be considered, these findings support the need for additional studies in more representative South Asian populations in order to understand the reasons for the difference in OSA risk and to examine OSA as a potential driver of CVD and DM in South Asians.

It is well known that South Asians manifest greater visceral adiposity, insulin resistance, and DM at a lower BMI compared to whites.^{14,15} Because South Asians commonly develop adverse metabolic profiles at a lower BMI,¹⁶ the International Diabetes Federation has recommended race-specific thresholds to define central obesity.¹⁷ A similar approach might be needed to define the thresholds for OSA risk in South Asians and other high risk racial/ethnic groups. In the meantime, clinicians need to recognize that the use of "one size fits all" BMI thresholds in OSA screening instruments may not accurately identify South Asians who are at increased risk for OSA.

In terms of obesity independent risk factors, a small cross-ethnic study identified bony craniofacial predictors of OSA in some ethnic groups but not Asian Indians.¹⁸ There is a notable paucity of work evaluating ethnic differences in physiological risk factors for OSA such as arousal threshold or loop gain. Further work is needed to identify both anatomic and non-anatomic risk factors for OSA in South Asians and other non-white groups.

An even more interesting question is whether the consequences of OSA may differ by race. The Sleep Heart Health Study suggested common OSA symptoms such as snoring, witnessed apnea, and sleepiness differ substantially by race/ethnicity.¹⁹ However, the difference in sleepiness between African Americans and whites was largely explained by differing responses to two questions in the Epworth Sleepiness Scale, suggesting that differences in the performance characteristics of this instrument may underlie racial/ethnic differences.²⁰ No studies have used objective assessments to assess sleepiness across diverse OSA populations. Nevertheless, reports of genetic polymorphisms that modify the impact of OSA provide a mechanism for racial heterogeneity in OSA consequences.²¹ A prior study found South Asians with OSA had higher hemoglobin A1c levels and lower high density lipoprotein concentrations than whites with OSA.¹⁰ Interestingly, Leong et al found the prevalence of diabetes was twice as high in South Asians compared to whites in their cohort. These cross-sectional data provide important observations that need further investigation. Specifically, longitudinal multiethnic cohorts that include South Asians are needed to understand the causal pathways and strength of the association between OSA and cardiometabolic outcomes, and to answer questions about whether these pathways and consequences differ by race/ethnicity.

A final issue that should be considered is to what extent does race/ethnicity influence treatment options, adherence, and outcomes? There is increasing evidence that African Americans have disproportionate failure rates following adenotonsillectomy for OSA in children and reduced rates of CPAP adherence for OSA in adults.^{22,23} Other vulnerable groups have also been found to have lower CPAP adherence.²⁴ Little information is available regarding the success of specific OSA treatment options in South Asians. While the heterogeneous results of CPAP studies on metabolic syndrome may be due to differences in study design,²⁵⁻²⁷ another possible explanation is that the metabolic impact of OSA therapy in South Asians may differ from whites.

Reducing the cardiometabolic risk of South Asians is a vital public health priority. The data from Leong et al. support the notion that sleep disorders may be contributing to this increased risk. Further well-designed studies are needed to determine if and how sleep disorders such as OSA are contributing to the elevated CVD and DM risk observed in South Asian populations and the potential of sleep interventions to minimize this disparity.

CITATION

Kandula NR; Patel SR. Sleep apnea and cardiometabolic risk in South Asians. *J Clin Sleep Med* 2013;9(9):859-860.

REFERENCES

1. A demographic snapshot of South Asians in the United States. South Asian Americans leading together. 2012 [accessed July 25, 2013]; Available from: <http://saalt.org/resources/reports-and-publications/#factsheets>.
2. Joshi P, Islam S, Pais P, et al. Risk factors for early myocardial infarction in South Asians compared with individuals in other countries. *JAMA* 2007;297:286-94.
3. Lee JW, Brancati FL, Yeh HC. Trends in the prevalence of type 2 diabetes in Asians versus whites: results from the United States National Health Interview Survey, 1997-2008. *Diabetes Care* 2011;34:353-7.
4. McKeigue PM, Shah B, Marmot MG. Relation of central obesity and insulin resistance with high diabetes prevalence and cardiovascular risk in South Asians. *Lancet* 1991;337:382-6.
5. Yusuf S, Hawken S, Ounpuu S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004;364:937-52.
6. Shaw JE, Punjabi NM, Wilding JP, Alberti KG, Zimmet PZ. Sleep-disordered breathing and type 2 diabetes: a report from the International Diabetes Federation Taskforce on Epidemiology and Prevention. *Diabetes Res Clin Pract* 2008;81:2-12.
7. Somers VK, White DP, Amin R, et al. Sleep apnea and cardiovascular disease: an American Heart Association/American College of Cardiology Foundation Scientific Statement from the American Heart Association Council for High Blood Pressure Research Professional Education Committee, Council on Clinical Cardiology, Stroke Council, and Council on Cardiovascular Nursing. In collaboration with the National Heart, Lung, and Blood Institute National Center on Sleep Disorders Research (National Institutes of Health). *Circulation* 2008;118:1080-111.
8. Udwadia ZF, Doshi AV, Lonkar SG, Singh CI. Prevalence of sleep-disordered breathing and sleep apnea in middle-aged urban Indian men. *Am J Respir Crit Care Med* 2004;169:168-73.
9. Reddy EV, Kadiravan T, Mishra HK, et al. Prevalence and risk factors of obstructive sleep apnea among middle-aged urban Indians: a community-based study. *Sleep Med* 2009;10:913-8.
10. Brady EM, Davies MJ, Hall AP, Talbot DC, Dick JL, Khunti K. An investigation into the relationship between sleep-disordered breathing, the metabolic syndrome, cardiovascular risk profiles, and inflammation between South Asians and Caucasians residing in the United Kingdom. *Metab Syndr Relat Disord* 2012;10:152-8.
11. Khoo SM, Tan WC, Ng TP, Ho CH. Risk factors associated with habitual snoring and sleep-disordered breathing in a multi-ethnic Asian population: a population-based study. *Respir Med* 2004;98:557-66.
12. Ng TP, Seow A, Tan WC. Prevalence of snoring and sleep breathing-related disorders in Chinese, Malay and Indian adults in Singapore. *Eur Respir J* 1998;12:198-203.
13. Leong WB, Arora T, Jenkinson D, et al. The prevalence and severity of obstructive sleep apnea in severe obesity: the impact of ethnicity. *J Clin Sleep Med* 2013;9:853-8.
14. Lear SA, Chockalingam A, Kohli S, Richardson CG, Humphries KH. Elevation in cardiovascular disease risk in South Asians is mediated by differences in visceral adipose tissue. *Obesity (Silver Spring)* 2012;20:1293-300.
15. Chandalia M, Lin P, Seenivasan T, et al. Insulin resistance and body fat distribution in South Asian men compared to Caucasian men. *PLoS One* 2007;2:e812.
16. Gray LJ, Yates T, Davies MJ, et al. Defining obesity cut-off points for migrant South Asians. *PLoS One* 2011;6:e26464.
17. Alberti KG, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;120:1640-5.
18. Wong ML, Sandham A, Ang PK, Wong DC, Tan WC, Huggare J. Craniofacial morphology, head posture, and nasal respiratory resistance in obstructive sleep apnoea: an inter-ethnic comparison. *Eur J Orthod* 2005;27:91-7.
19. O'Connor GT, Lind BK, Lee ET, et al. Variation in symptoms of sleep-disordered breathing with race and ethnicity: the Sleep Heart Health Study. *Sleep* 2003;26:74-9.
20. Hayes AL, Spilsbury JC, Patel SR. The Epworth score in African American populations. *J Clin Sleep Med* 2009;5:344-8.
21. Nikodemova M, Finn L, Mignot E, Salzieder N, Peppard PE. Association of sleep disordered breathing and cognitive deficit in APOE epsilon4 carriers. *Sleep* 2013;36:873-80.
22. Marcus CL, Moore RH, Rosen CL, et al. A randomized trial of adenotonsillectomy for childhood sleep apnea. *N Engl J Med* 2013;368:2366-76.
23. Billings ME, Auckley D, Benca R, et al. Race and residential socioeconomic status as predictors of CPAP adherence. *Sleep* 2011;34:1653-8.
24. Bakker JP, O'Keefe KM, Neill AM, Campbell AJ. Ethnic disparities in CPAP adherence in New Zealand: effects of socioeconomic status, health literacy and self-efficacy. *Sleep* 2011;34:1595-603.
25. Sharma SK, Agrawal S, Damodaran D, et al. CPAP for the metabolic syndrome in patients with obstructive sleep apnea. *N Engl J Med* 2011;365:2277-86.
26. Hoyos CM, Sullivan DR, Liu PY. Effect of CPAP on the metabolic syndrome: a randomised sham-controlled study. *Thorax* 2013;68:588-9.
27. Coughlin SR, Mawdsley L, Mugarza JA, Wilding JP, Calverley PM. Cardiovascular and metabolic effects of CPAP in obese males with OSA. *Eur Respir J* 2007;29:720-7.

SUBMISSION & CORRESPONDENCE INFORMATION

Submitted for publication July, 2013

Accepted for publication July, 2013

Address correspondence to: Sanjay R. Patel, M.D., Brigham and Women's Hospital, Division of Sleep Medicine, 221 Longwood Avenue, Room 225C, Boston, MA 02115; Tel: (857) 307-0347; Fax: (617) 278-6946; E-mail: spatel@partners.org

DISCLOSURE STATEMENT

The authors have indicated no financial conflicts of interest.