New Record of the Genus *Calyprella* from Korea

Chan Jung Lee¹, Wan Gyu Kim², Chang Sung Jhune¹, Jong Chun Cheong¹ and Soon Ja Seok^{3*}

¹Mushroom Research Division, National Institute of Horticultural & Herbal Science, RDA, Suwon 441-707, Korea

²Agricultural Microbiology Division, National Academy of Agricultural Science, RDA, Suwon 441-707, Korea

³National Agrobiodiversity Center, National Academy of Agricultural Science, RDA, Suwon 441-707, Korea

(Received February 25, 2009. Accepted March 24, 2009)

Calyprella capula is reported for the first time in Korea. It was found in spring in Chungcheongbuk-do, central Korea, growing in a dead stalk of *Sesamum indicum*. A description and molecular phylogenetic analyses are reported.

KEYWORDS : *Calyprella capula*, Cyphelloid, Distribution, New record

Homobasidiomycetes are the most conspicuous group of fungi and includes approximately 16,000 described species of mushroom-forming fungi and related forms (Kirk *et al.*, 2001). In Korea, 1424 species of Basidiomycotina from 330 genera, 82 families and 15 orders have been recorded (Lee, 2005). Members of the Homobasidiomycetes include gilled mushrooms, boletes, polypores, and puffballs. The Homobasidiomycetes also contain a relatively obscure group called the cyphelloid fungi. Cyphelloid fungi include approximately 120 species in about 40 genera, but there is an estimated actual diversity of ca. 500 species (Agerer, 1978; Cooke, 1962; Donk, 1959; Reid, 1964; Singer, 1986). *Calyprella* species have been included in the cyphelloid fungi. Generally, cyphelloid fungi have minute, cup- to tube-shaped, often pendant fruiting bodies that are less than 2 mm in size. Cyphelloid fruiting bodies usually have a smooth, even hymenophore that lines their concave inner surface (Agerer, 1978a, 1983a; Donk, 1951, 1959, 1966).

In spring of 2006, a *Calyprella* species was collected by an author from a gingili plant at Chungcheongbuk-do in Korea, and identified as *Calyprella capula* (Holmsk.) Quélet following the descriptions of Redhead and Traquair (1981) and Breitenbach and Kränzlin (1986). To the best of our knowledge, this genus has not been reported in Korea. The morphology and genotypic characteristics of the isolate were studied.

Materials and Methods

Fungal isolates and morphology. Macroscopic and microscopic descriptions are based on fresh specimens. Color nomenclature refers to Manual of Color Names (Japan Color Research Institute 1973) and color codes in parentheses are from The Munsell Book of Color (Mun-

sell Color, 1963). Microscopic characters were examined from material mounted in 5% KOH or 3% KOH + Congo red. A total of 50 basidiospores from mounted lamellae were examined for size. A minimum of 20 basidiospores, basidia and other structures were measured for each collection we examined. The strain is maintained in Herbarium Conservation Center Nias (HCCN). Cultures were grown in a biochemical oxygen-demand incubator at 25°C on potato dextrose agar.

Genomic DNA isolation, PCR and DNA sequencing.

DNA was isolated from pure cultures of fungus using a SDS-CTAB (sodium dodecyl sulfate-cetyltrimethyl-ammonium bromide) method (Kim *et al.*, 1990). ITS1 and ITS4 primer sets were used for amplification of internal transcribed spacer (ITS) regions (White *et al.*, 1990). PCR amplifications were carried out in a Techne thermocycler (Techne LTD, Duxford, Cambridge, U.K.). PCR consisted of 35 cycles of 1 min at 94°C, 1 min at 58°C, and 2 min at 72°C. Contaminating primers and dNTPs were removed from PCR products using the High Pure PCR Product Purification Kit (Bioneer Co., Chungbuk, Korea). Big Dye Terminator Kit and ABI Prism 310 Genetic Analyzer (Perkin Elmer, New Jersey, USA) was used for sequencing.

Molecular phylogenetic analysis. The partial ITS region sequence (649nt) of strain CH3 was compared to the ITS region sequence available from GenBank using the BLAST program to determine an approximate phylogenetic affiliation. The gene sequence was then multiply aligned with those of closely related species using the program MEGALIGN (DNASTAR). Phylogenetic trees were constructed using the neighbour-joining method (Saitou and Nei, 1987) with the MEGA 3.0 program (Kumar *et al.*, 2004); bootstrap percentages were based on 1000 replications (Felsenstein, 1985). The partial ITS

*Corresponding author <E-mail : mycena@rda.go.kr>

Fig. 1. Basidiomata of *Calyptella capula* (HCCN00010) in a habitat.

region sequence determined in this study has been deposited in the GenBank Database under accession numbers of EF374132 for strain CH3.

Results and Discussion

Genus *Calyptella* Quélet, *Enchir. F.*, p 216. 1886.

(**Korean name:** Eun-Bang-Ul-Beo-Seot-sok)

Type species: *Calyptella capula* (Holmsk.) Quélet

Calyptella capula (Holmsk.) Quélet, *Fl. mycol. France* (Paris): 25 (1886)

(**Korean name:** Eun-Bang-Ul-Beo-Seot)

Synonymy:

Calyptella laeta (Fr.) W.B. Cooke, *Beih. Sydowia* 4: 40 (1961)

Chaetocypha capula (Holmsk.) Kuntze, *Revis. gen. pl.* (Leipzig) 2: 847 (1891)

Chaetocypha laeta (Fr.) Kuntze, *Revis. gen. pl.* (Leipzig) 2: 847 (1891)

Chaetocypha pimii (W. Phillips) Kuntze, *Revis. gen. pl.* (Leipzig) 2: 847 (1891)

Cyphella capula (Holmsk.) Fr., *Epicer. syst. mycol.* (Upsaliae): 568 (1838)

Cyphella capula var. *flavescens* Pat., *Tabl. analyt. Fung. France* (Paris): 56 (1883)

Cyphella laeta Fr., *Epicer. syst. mycol.* (Upsaliae): 568 (1838)

Cyphella pimii W. Phillips, *Grevillea* 13(no. 66): 49 (1884)

Cyphella velenovskyi Pilát, *Annls mycol.* 22: 206 (1924)

Peziza capula Holmsk., 1: 286 (1799)

Basidiomes. Basically cupulate, dorsally stipitate, extremely fragile, solitary to gregarious, outer surface smooth, white to cream-colored, inner surface with the hymenium likewise smooth, white to cream-colored, discoid when young, becoming conical to campanulate, 1~4 mm wide, 1~3 mm deep, externally whitish to pale mouse grey, with age becoming blackish, glabrous or appearing frosted when magnified slightly, with incurved margins which

Fig. 2. Microscopic structures of *Calyptella capula* (HCCN00010). A, basidia; B, basidiospores; C, D, generative hyphae; E, hyphal ends of generative hyphae (Scale bar: 5 μ m).

flair with age and become slightly eroded.

Hymenium. 0.2~2.0 mm long, smooth to radially furrowed or folded, whitish to pale grey or yellowish grey. Pseudostipe; short to elongate, usually eccentric, solid, often frosted to pubescent, concolorous or darker than the receptacle.

Basidia. 20~25 $\times$ 5~8 μ m, cylindrical when young, becoming clavate, 4-sterigmata, basal clamped.

Basidiospores. 6~9 $\times$ 3~4.5 μ m, ellipsoid to obovoid, inequilateral, somewhat flattened on one side, thin-walled, hyaline, nonamyloid, smooth, with a prominent apiculus.

Fig. 3. Neighbor joining phylogenetic dendrogram of *Calyprella* species based on internal transcribed spacer (ITS) sequence. Numbers on the branches represent percentage of 1000 bootstrap frequencies.

Cystidia. Cystidia not seen.

Hyphal system. Monomitic, hyphae in the subhymenium 2–4 μm across, septa with clamps, hyphae in the stalk and in the trama up to 15 μm across, hyphal ends in the marginal part of the cup sinuous-gnarled with outgrowths, some exserted.

Habitat. Spring, dead stems on sesame (*Sesamum indicum*).

Materials examined. Eumseong-gun, Chungbuk, 22 May 2006. (Herbarium No. : HCCN00010).

Molecular characterization. The rDNA ITS region of the *Calyprella* isolate was amplified with conserved fungal primers ITS1 and ITS4. The size of the ITS fragment was approximately 649 bp, which includes ITS1, 5.8S and ITS4 regions. The ITS region was sequenced and the sequence data was submitted to GenBank nucleotide database (accession number EF374132). The sequence was compared with the published Cyphelloid fungi sequences available in the NCBI database. The strain is closely related to strains of the type species of *C. capula* (Fig. 3). This strain showed 96% ITS region sequence similarity to *C. capula*, according to blast search for homologous. Bodensteiner *et al.* (2001) reported Cyphelloid species *C. capula* (the type species of *Calyprella*) is the sister group of the resupinatus clade, but this placement received weak bootstrap support. *C. capula* produces cupulate fruiting

bodies covered by surface hyphae with multiply branched, coralloid excrescences that lack a crystal covering (Cooke, 1962; Donk, 1951; Reid, 1961; Singer, 1962, 1986).

References

- Agerer, R. 1978. *Lachnella-Crinipellis, Stigmatolemma-Fistulina*: zwei Verwandtschaftsreihen. *Z. Mykol.* 44:51-70.
- Agerer, R. 1978. Cyphelloide Pilze aus Teneriffa. *Nova Hedwigia* 30:295-341.
- Agerer, R. 1983. Beitrag zur Flora cyphelloider Pilze aus der Neotropis-V. Zwei neue Gattungen: Incrustocalyprella und Metulocyphella. *Z. Mykol.* 49:155-164.
- Bodensteiner, P., Agerer, R., Desjardin, D. E., and Horak, E. 2001. A new species of *Calathella* from Bali. *Mycologia* 93: 1010-1013.
- Breitenbach, J. and Kränzlin, F. 1986. Fungi of Switzerland (English version), vol. 2. Verlag Mykologia, Luzern, pp. 200-201.
- Cooke, W. B. 1962. The cyphellaceous fungi. A study in the Porotheliales. *Beih. Sydowia* 4:1-144.
- Donk, M. A. 1951. The generic names proposed for Hymenomycetes-I, "Cyphellaceae". *Reinwardtia* 1:199-220.
- Donk, M. A. 1959. Notes on "Cyphellaceae."-I. *Persoonia* 1: 25-110.
- Donk, M. A. 1966. A reassessment of the Cyphellaceae. *Acta Bot. Neerl.* 15:95-101.
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39:783-791.
- Kim, W. K., Mauthe, W., Hausner, G. and Klassen, G. R. 1990. Isolation of high molecular weight DNA and double-stranded RNAs from fungi. *Can. J. Bot.* 68:1898-1902.
- Kirk, P. M., Cannon, P. F. and David, J. C. 2001. Ainsworth and

- Bisby's Dictionary of Fungi, ninth ed. CAB International University Press, Wallingford, UK.
- Kumar, S., Tamura, K. and Nei, M. 2004. MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform.* 5:150-163.
- Lee, J. Y. 2005. Additional list of recorded mushrooms in Korea (from 2001 to 2004). *Kor. J. Mycol.* 33: 54-57.
- National Mycological Herbarium, Biosystematics Research Institute, Ottawa, Ontario. 2p.
- Redhead, S. A. and Traquair, J. A. 1981. *Calyprella capula*. Fungi Canadenses No. 202.
- Reid, D. A. 1961. Fungi Venezuelani V. The Cyphellaceae of Venezuela. *Kew Bull.* 15:261-275.
- Reid, D. A. 1964. Notes on some fungi in Michigan-I. "Cyphellaceae". *Persoonia* 3:97-154.
- Saitou, N. and Nei, M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4:406-425.
- Singer, R. 1962. The Agaricales in modern taxonomy. Second ed., Cramer, Lehre.
- Singer, R. 1986. The Agaricales in modern taxonomy, fourth ed. Koeltz Scientific Books, Königstein, Germany.
- White, T. J., Bruns, T., Lee S. and Taylor, J. W. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: PCR Protocols: A Guide to Methods and Applications, pp. 315-322. Eds. M. A. Innis, D. H. Gelfand, J. J. Sninsky and T. J. White. Academic Press, Inc., New York.