Published in final edited form as:

Biochem Biophys Res Commun. 2012 August 3; 424(3): 404–408. doi:10.1016/j.bbrc.2012.06.120.

TAA repeat variation in the *GRIK2* gene does not influence age at onset in Huntington's disease

Ji-Hyun Lee^{1,2}, Jong-Min Lee¹, Eliana Marisa Ramos^{1,3}, Tammy Gillis¹, Jayalakshmi S. Mysore¹, Shotaro Kishikawa¹, Tiffany Hadzi⁴, Audrey E. Hendricks⁵, Michael R. Hayden⁶, Patrick J. Morrison⁷, Martha Nance⁸, Christopher A. Ross⁹, Russell L. Margolis⁹, Ferdinando Squitieri¹⁰, Cinzia Gellera¹¹, Estrella Gomez-Tortosa¹², Carmen Ayuso¹³, Oksana Suchowersky¹⁴, Ronald J. Trent¹⁵, Elizabeth McCusker¹⁶, Andrea Novelletto¹⁷, Marina Frontali¹⁸, Randi Jones¹⁹, Tetsuo Ashizawa²⁰, Samuel Frank⁴, Marie-Helene Saint-Hilaire⁴, Steven M. Hersch²¹, Herminia D. Rosas²¹, Diane Lucente¹, Madaline B. Harrison²², Andrea Zanko²³, Ruth K. Abramson²⁴, Karen Marder²⁵, Jorge Sequeiros^{3,26}, G. Bernhard Landwehrmeyer²⁷, on behalf of the Registry Study of the European Huntington's Disease Network, Ira Shoulson Network²⁸, on behalf of the Huntington Study Group COHORT project, Richard H. Myers⁴, Marcy E. MacDonald¹, and James F. Gusella^{1,29}

¹Molecular Neurogenetics Unit, Center for Human Genetic Research, Massachusetts General Hospital, Boston, MA 02114, USA

²Department of Pharmacology, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 120-752, Korea

³UnIGENE, IBMC – Institute for Molecular and Cell Biology, Universidade do Porto, 4150-180, Porto, Portugal

⁴Department of Neurology, Boston University School of Medicine, Boston, MA, 02118, USA

⁵Department of Biostatistics, Boston University School of Public Health, Boston, MA, 02118, USA

⁶University of British Columbia, Center for Molecular Medicine and Therapeutics, Vancouver BC, Canada, V5Z 4H4

⁷Regional Medical Genetics Centre, Belfast HSC Trust, Belfast, BT9 7AB, and University of Ulster, Cromore Road, Coleraine, BT52 15A, United Kingdom

⁸Hennepin County Medical Center, 701 Park Avenue, Minneapolis, MN, 55415, USA

⁹Johns Hopkins University, Department of Psychiatry and Behavioral Sciences, Baltimore, MD, 21287, USA

© 2012 Elsevier Inc. All rights reserved.

Corresponding author: Ji Hyun Lee, PhD, Department of Pharmacology, Pharmacogenomic Research Center for Membrane, Transporters and Research Center for Human Natural Defense System, Yonsei University, College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 120-752, Korea, Tel: +82 2 2228 1743/Fax: +82 2 313 1894/ jihyni@yuhs.ac.

Competing interests: The authors declare that they have no competing interests.

Authors' contributions: JHL participated in the design of the study, analyzed the data statistically and drafted the manuscript. JML assisted in statistical analysis of the data and critical revision of the manuscript. EMR, TG, JSM and SK generated the molecular data while TH, AEH, MRH, PM, MN, CAR, RLM, FS, CG, EGT, CA, OS, RJT, EM, AN, MF, RJ, TA, SF, MHSH, SMH, HDR, DL, MBH, AZ, RKA, KM, JS, GBL and IS participated in design of the study and helped to generate the clinical data. RHM, MEM and JFG conceived the study, participated in its design and critically revised the manuscript. All authors have read and approved the final manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

- ¹⁰Centre for Neurogenetics and Rare Diseases, IRCCS Neuromed, 86077, Pozzilli (IS), Italy
- ¹¹Istituto Nazionale Neurologico C. Besta, 20133, Milan, Italy
- ¹²Department of Neurology, Fundación Jiménez Díaz, Madrid 28040, Spain
- ¹³Department of Genetics, Fundación Jiménez Díaz, Madrid 28040, Spain
- ¹⁴Division of Neurology, University of Alberta, Edmonton, Alberta T6G 2B7
- ¹⁵Sydney Medical School, University of Sydney, NSW 2006, Australia
- ¹⁶Department of Neurology, Westmead Hospital, Westmead Sydney, NSW 2145, Australia
- ¹⁷Department of Biology, University Tor Vergata, 00133 Rome, Italy
- ¹⁸Institute of Neurobiology and Molecular Medicine, 00133 Rome, Italy
- ¹⁹Emory University School of Medicine, Center for Neurodegenerative Disease, Atlanta GA, 30322, USA
- ²⁰Department of Neurology, University of Florida, Gainesville, FL, 32610, USA
- ²¹MassGeneral Institute for Neurodegenerative Disorders, Massachusetts General Hospital, 114 16th Street, Charlestown, MA, 02129, USA
- ²²Department of Neurology, University of Virginia, Charlottesville, VA, 22908, USA
- ²³Memory and Aging Center, University of California, San Francisco, CA, 94143, USA
- ²⁴Department of Neuropsychiatry and Behavioral Science, University of South Carolina School of Medicine, Columbia, SC, 29209, USA
- ²⁵Department of Neurology, Columbia University Medical Center, New York, NY, 10032, USA
- ²⁶ICBAS, Universidade do Porto, 4099-003, Porto, Portugal
- ²⁷Department of Neurology, Ulm University, 89081 Ulm, Germany
- ²⁸Departments of Neurology, Pharmacology and Human Science, Georgetown University Medical Center, Washington, DC, 20007, USA
- ²⁹Department of Genetics, Harvard Medical School, Boston MA 02115

Abstract

Huntington's disease is a neurodegenerative disorder caused by an expanded CAG trinucleotide repeat whose length is the major determinant of age at onset but remaining variation appears to be due in part to the effect of genetic modifiers. *GRIK2*, which encodes GluR6, a mediator of excitatory neurotransmission in the brain, has been suggested in several studies to be a modifier gene based upon a 3' untranslated region TAA trinucleotide repeat polymorphism. Prior to investing in detailed studies of the functional impact of this polymorphism, we sought to confirm its effect on age at onset in a much larger dataset than in previous investigations. We genotyped the *HD* CAG repeat and the *GRIK2* TAA repeat in DNA samples from 2,911 Huntington's disease subjects with known age at onset, and tested for a potential modifier effect of *GRIK2* using a variety of statistical approaches. Unlike previous reports, we detected no evidence of an influence of the *GRIK2* TAA repeat polymorphism on age at motor onset. Similarly, the *GRIK2* polymorphism did not show significant modifier effect on psychiatric and cognitive age at onset in HD. Comprehensive analytical methods applied to a much larger sample than in previous studies do not support a role for *GRIK2* as a genetic modifier of age at onset of clinical symptoms in Huntington's disease.

Keywords

Huntington's disease (HD); Age at onset; *GRIK2*; Genetic modifier

Introduction

Huntington's disease (HD) is an autosomal dominant, progressive neurodegenerative disorder that presents with motor dysfunction, cognitive decline and psychiatric disturbances due to expansion of a trinucleotide CAG repeat encoding a polyglutamine tract in the huntingtin protein [1,2]. There is a strong inverse correlation between age at diagnosis by onset of motor signs and the CAG repeat length, which accounts for up to 67% of the overall variance [3-5]. The remaining variation is strongly heritable, indicating a contribution of modifier genes to age at onset [6-8]

GRIK2 (Glutamate receptor, ionotropic kainate 2), encoding the GluR6 subunit of the predominant excitatory neurotransmitter receptor family in the human brain, is an attractive candidate as an HD modifier because of its potential role in excitotoxic cell death [9]. A TAA trinucleotide repeat polymorphism in the 3' untranslated region (3'UTR) of *GRIK2* mRNA was reported to be associated with age at onset of diagnostic motor signs in small studies from the United Kingdom, New England, France, India, and Italy [10-14], the largest of which comprised less than 300 subjects. The two initial studies suggested that the modifier effect was due to genetic variation on chromosomes carrying a 16 TAA repeat allele. The genetic variation responsible for the earlier than expected age at onset was implicated as the TAA repeat itself by subsequent haplotype analysis [15]. However, the mechanism by which this polymorphism could act is not certain and might conceivably include effects on alternative splicing, editing, stability or translational regulation of the *GRIK2* mRNA. Consequently, before embarking on extensive molecular analyses to define the mechanism of action, we carried out a comprehensive analysis of a much larger set of HD subjects than has previously been examined to confirm *GRIK2* as a genetic modifier of HD pathogenesis.

Methods and materials

HD samples

HD patient DNA samples from individuals with known age at motor, psychiatric and/or cognitive onset were obtained from ongoing genetic studies at the MGH HD Center Without Walls, members of the HD-MAPS collaboration, post-mortem brain specimens (Harvard Brain Tissue Resource Center and the UCLA Brain Bank), and two large observational studies: the Huntington Study Group's COHORT project and the European Huntington Disease Network's REGISTRY study. In total, 2,911 HD heterozygote subjects with one expanded HD allele and known age of motor onset (2,362 individuals), psychiatric onset (547 individuals) and/or cognitive onset (210 individuals) were genotyped for the *GRIK2* TAA repeat polymorphism and for the *HD* CAG repeat as described previously [9,16]. The mean age at onset was 42.6 (range, 4-92) and the mean expanded allele *HD* CAG repeat length was 45.1 (range, 36-98). This study used de-identified DNA samples and was approved by the Institutional Review Board of the Partners HealthCare System.

Data analysis

This study utilized samples from 2,911 HD heterozygote subjects with one expanded HD allele and known age of motor onset (2,365 individuals), psychiatric onset (547 individuals) and/or cognitive onset (210 individuals). Potential modifier effects were explored by adding the factor to a linear regression model relating the natural log-transformed age at onset to

HD CAG repeat length and determining the degree of improvement in goodness-of-fit. Since the regression plots for onset of symptoms in the three different domains show quite different relationships between of natural log-transformed age at onset and *HD* CAG repeat length, each category of age at onset was analyzed separately. SPSS 11.5 (SPSS Inc., Chicago, IL, USA) was used for all statistical analyses.

Results

The *GRIK2* TAA repeat displays 7 alleles, ranging from 12 to 17 TAAs. The 16 TAA allele was reported previously to be associated with earlier age of onset in HD [10,11]. We initially tested potential modifier effects by determining whether adding the *GRIK2* genotype (dominant model) had a significant impact on a linear regression model relating the natural log-transformed age at onset to HD CAG repeat length. Separate tests were performed for 1) presence of at least one 16 TAA allele, 2) presence of at least one of the two longest alleles (16 or 17 TAAs), and 3) the *GRIK2* TAA repeat as a continuous trait, based upon the larger of the two alleles present in each individual. Figure 1A shows the relationship in 2,362 HD subjects of age at onset of diagnostic motor signs with CAG repeat length and Figure 1B reveals a lack of any significant impact of adding *GRIK2* to the model on improving the R^2 value.

To avoid the potential disproportionate impact of subjects with extreme CAG lengths reported by Lee et al. 2012 [17], we calculated standardized residuals for the 2,205 subjects with CAG repeats in the 40-53 range, relative to the regression line generated in that report. This analysis (Figure 1C-D) again shows no evidence of an effect of *GRIK2* genotype on motor onset. We also performed a number of analyses in which we subdivided these samples into HD subjects with extremes of age at onset associated with each CAG repeat length using different cut-offs for inclusion or exclusion and in no case did comparison of *GRIK2* genotypes between the two extreme groups reveal a significant difference. Thus, our findings in this large dataset indicate that the *GRIK2* TAA repeat polymorphism, previously thought to be an HD modifier, is not significantly associated with deviation in HD age of motor onset from that expected based upon CAG repeat length.

Although HD is typically diagnosed based upon onset of motor signs, the age at onset for cognitive and/or psychiatric symptoms is available in a subset of this sample. Directly comparable to Figure 1, Figure 2 shows no significant effect of the *GRIK2* TAA repeat polymorphism on age at onset of either psychiatric (Figure 2A-B; 547 subjects) or cognitive (Figure 2C-D; 210 subjects) symptoms.

Discussion

It is well established that the expanded CAG repeat in the *HD* gene is the primary determinant of age of onset of clinical symptoms, but that the mutation alone does not explain all of the variation observed. The remaining variance displays a high degree of heritability, supporting the existence of genetic modifiers, factors whose polymorphic variation influences the course of HD. Consequently, identification of genetic modifiers from human patients represents a potential route to discovering validated targets whose modulation would be expected to alter HD pathogenesis. To date, candidate gene strategies have produced a number of reports of potential modifiers from pathways postulated to be involved in HD pathogenesis [7,12,18-21], including several in which the *GRIK2* TAA repeat polymorphism was associated with altered onset age [10,11,15]. *GRIK2* encodes a protein integrally involved in mediating excitatory neurotransmission in the brain and its status as a genetic modifier would bolster the excitotoxicity model of HD. However, detailed

investigation of its mode of action would entail a major investment of labor and research funds.

As genetic analysis of functional variants has argued strongly against other attractive modifier candidates, such as *BDNF* which showed no effect on age at onset in two large studies [22,23], we felt it important to confirm the *GRIK2* modifier effect in a much larger study sample than was previously tested. The increased power afforded by the much larger sample size and more comprehensive analytical methods in our current study argue strongly that *GRIK2* has no significant effect on HD pathogenesis leading to age of onset of motor symptoms. Albeit with smaller sample sizes, our study also does not support modifier effects on age of onset psychiatric or cognitive symptoms. These findings argue against the pursuit of detailed molecular and biological analysis of any putative functional effect of this *GRIK2* polymorphism as a clue to how to alter HD pathogenesis. In this regard, they are consistent with negative findings by Metzger et al. [24] in a German HD population and by Andresen et al. [25] in the Venezuela HD population, although the latter had relatively few 16 TAA alleles.

Our findings present a cautionary note for HD modifier studies in general since an apparent *GRIK2* effect was noted previously for motor onset in multiple studies [10-15]. These apparent positive associations may have been due to small, possibly unrepresentative samples and may have been disproportionately influenced by genotypic or phenotypic outliers.

In conclusion, our study results showed that the previously studied TAA repeat variation in the *GRIK2* gene does not influence age at onset in Huntington's disease using the largest dataset assembled to date and suggest that similarly large-scale investigations may be required to adequately assess the potential modifier effects of other reported candidates.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank contributors to the HD-MAPS, COHORT and Registry studies, listed in Appendix 1. This work was supported by grants from the National Institutes of Health NINDS Huntington's Disease Center Without Walls NS16367, the CHDI Foundation Inc., and the Huntington's Disease Society of America's Coalition for the Cure. JHL received support from a National Research Foundation of Korea Grant funded by the Korean Government [NRF- 2009-352-E00010]. EMR is the recipient of a scholarship from FCT (SFRH/BD/44335/2008). The Huntington Study Group COHORT project was supported by the CHDI Foundation, Inc. and the REGISTRY study of the European Huntington's Disease Network was supported by the High Q Foundation.

References

1. Walker FO. Huntington's disease. *The Lancet*. 2007; 369:218–228.
2. Huntington's Disease Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. The Huntington's Disease Collaborative Research Group. *Cell*. 1993; 72:971–983. [PubMed: 8458085]
3. Duyao M, Ambrose C, Myers R, Novelletto A, Persichetti F, Frontali M, Folstein S, Ross C, Franz M, Abbott M. Trinucleotide repeat length instability and age of onset in Huntington's disease. *Nature Genetics*. 1993; 4:387–392. [PubMed: 8401587]
4. Andrew SE, Goldberg YP, Kremer B, Telenius H, Theilmann J, Adam S, Starr E, Squitieri F, Lin B, Kalchman MA. The relationship between trinucleotide (CAG) repeat length and clinical features of Huntington's disease. *Nature Genetics*. 1993; 4:398–403. [PubMed: 8401589]

5. Snell RG, MacMillan JC, Cheadle JP, Fenton I, Lazarou LP, Davies P, MacDonald ME, Gusella JF, Harper PS, Shaw DJ. Relationship between trinucleotide repeat expansion and phenotypic variation in Huntington's disease. *Nature Genetics*. 1993; 4:393–397. [PubMed: 8401588]
6. Djousse L, Knowlton B, Hayden MR, Almqvist EW, Brinkman RR, Ross CA, Margolis RL, Rosenblatt A, Durr A, Dode C, Morrison PJ, Novelletto A, Frontali M, Trent RJ, McCusker E, Gomez-Tortosa E, Mayo Cabrero D, Jones R, Zanko A, Nance M, Abramson RK, Suchowersky O, Paulsen JS, Harrison MB, Yang Q, Cupples LA, Mysore J, Gusella JF, MacDonald ME, Myers RH. Evidence for a modifier of onset age in Huntington disease linked to the HD gene in 4p16. *Neurogenetics*. 2004; 5:109–114. [PubMed: 15029481]
7. Gusella JF, MacDonald ME. Huntington's disease: the case for genetic modifiers. *Genome Med*. 2009; 1:80. [PubMed: 19725930]
8. Wexler NS, Lorimer J, Porter J, Gomez F, Moskowitz C, Shackell E, Marder K, Penchaszadeh G, Roberts SA, Gayn J, Brocklebank D, Cherny S, Cardon LR, Gray J, Dlouhy SR, Wiktorski S, Hodes ME, Conneally PM, Penney JB, Gusella J, Cha J, Irizarry M, Rosas D, Hersch S, Hollingsworth Z, MacDonald M, Young AB, Andresen JM, Housman DE, De Young MM, Bonilla E, Stillings T, Negrette A, Snodgrass SR, Martinez-Jaurieta MD, Ramos-Arroyo MA, Bickham J, Ramos JS, Marshall F, Shoulson I, Rey GJ, Feigin A, Arnheim N, Acevedo-Cruz A, Acosta L, Alvir J, Fischbeck K, Thompson LM, Young A, Dure L, O'Brien CJ, Paulsen J, Brickman A, Krch D, Peery S, Hogarth P, Higgins DS, Landwehrmeyer B. Venezuelan kindreds reveal that genetic and environmental factors modulate Huntington's disease age of onset. *Proceedings of the National Academy of Sciences of the United States of America*. 2004; 101:3498–3503. [PubMed: 14993615]
9. Paschen W, Blackstone CD, Haganir RL, Ross CA. Human GluR6 kainate receptor (GRIK2): molecular cloning, expression, polymorphism, and chromosomal assignment. *Genomics*. 1994; 20:435–440. [PubMed: 8034316]
10. Rubinsztein DC, Leggo J, Chiano M, Dodge A, Norbury G, Rosser E, Craufurd D. Genotypes at the GluR6 kainate receptor locus are associated with variation in the age of onset of Huntington disease. *Proceedings of the National Academy of Sciences of the United States of America*. 1997; 94:3872–3876. [PubMed: 9108071]
11. MacDonald ME, Vonsattel JP, Shrinidhi J, Couropmitree NN, Cupples LA, Bird ED, Gusella JF, Myers RH. Evidence for the GluR6 gene associated with younger onset age of Huntington's disease. *Neurology*. 1999; 53:1330–1332. [PubMed: 10522893]
12. Naz P, Vuillaume I, Deste A, Pasquier F, Sablonniere B. Mutation analysis and association studies of the ubiquitin carboxy-terminal hydrolase L1 gene in Huntington's disease. *Neuroscience Letters*. 2002; 328:1–4. [PubMed: 12123845]
13. Chattopadhyay B, Ghosh S, Gangopadhyay PK, Das SK, Roy T, Sinha K, Jha DK, Mukherjee SC, Chakraborty A, Singhal BS, Bhattacharya AK, Bhattacharyya NP. Modulation of age at onset in Huntington's disease and spinocerebellar ataxia type 2 patients originated from eastern India. *Neuroscience Letters*. 2003; 345:93–96. [PubMed: 12821179]
14. Cannella M, Gellera C, Maglione V, Giallonardo P, Cislighi G, Muglia M, Quattrone A, Pierelli F, Di Donato S, Squitieri F. The gender effect in juvenile Huntington disease patients of Italian origin, *American journal of medical genetics. Part B. Neuropsychiatric genetics*. 2004; 125B:92–98. [PubMed: 14755452]
15. Zeng W, Gillis T, Hakky M, Djousse L, Myers RH, MacDonald ME, Gusella JF. Genetic analysis of the GRIK2 modifier effect in Huntington's disease. *BMC Neurosci*. 2006; 7:62. [PubMed: 16959037]
16. Warner JP, Barron LH, Brock DJ. A new polymerase chain reaction (PCR) assay for the trinucleotide repeat that is unstable and expanded on Huntington's disease chromosomes. *Mol Cell Probes*. 1993; 7:235–239. [PubMed: 8366869]
17. Lee JM, Ramos EM, Lee JH, Gillis T, Mysore JS, Hayden MR, Warby SC, Morrison P, Nance M, Ross CA, Margolis RL, Squitieri F, Orobello S, Di Donato S, Gomez-Tortosa E, Ayuso C, Suchowersky O, Trent RJ, McCusker E, Novelletto A, Frontali M, Jones R, Ashizawa T, Frank S, Saint-Hilaire MH, Hersch SM, Rosas HD, Lucente D, Harrison MB, Zanko A, Abramson RK, Marder K, Sequeiros J, Paulsen JS, Landwehrmeyer GB, Myers RH, Macdonald ME, Gusella JF. CAG repeat expansion in Huntington disease determines age at onset in a fully dominant fashion. *Neurology*. 2012

18. Arning L, Mont D, Hansen W, Wieczorek S, Jagiello P, Akkad DA, Andrich J, Kraus PH, Saft C, Epplen JT. ASK1 and MAP2K6 as modifiers of age at onset in Huntington's disease. *Journal of Molecular Medicine*. 2008; 86:485–490. [PubMed: 18327563]
19. Dhaenens C, Burnouf S, Simonin C, Van Brussel E, Duhamel A, Defebvre L, Duru C, Vuillaume I, Cazeneuve C, Charles P, Maison P, Debruxelles S, Verny C, Gervais H, Azulay J, Tranchant C, Bachoud-Levi A, Drr A, Bue L, Krystkowiak P, Sablonniere B, Blum D. A genetic variation in the ADORA2A gene modifies age at onset in Huntington's disease. *Neurobiology of disease*. 2009; 35:474–476. [PubMed: 19591938]
20. Metzger S, Bauer P, Tomiuk J, Laccone F, Didonato S, Gellera C, Soliveri P, Lange HW, Weirich-Schwaiger H, Wenning GK, Melegh B, Havasi V, Balik L, Wieczorek S, Arning L, Zaremba J, Sulek A, Hoffman-Zacharska D, Basak AN, Ersoy N, Zidovska J, Kebrdlova V, Pandolfo M, Riba P, Kadasi L, Kvasnicova M, Weber BH, Kreuz F, Dose M, Stuhmann M, Riess O. The S18Y polymorphism in the UCHL1 gene is a genetic modifier in Huntington's disease. *Neurogenetics*. 2006; 7:27–30. [PubMed: 16369839]
21. S C, Taherzadeh-Fard E, Andrich J, Wieczorek S, Arning L. PGC-1alpha as modifier of onset age in Huntington disease. *Molecular neurodegeneration*. 2009 Feb 6.4:10–10. [PubMed: 19200361]
22. Kishikawa S, Li J, Gillis T, Hakky M, Warby S, Hayden M, MacDonald ME, Myers RH, Gusella JF. Brain-derived neurotrophic factor does not influence age at neurologic onset of Huntington's disease. *Neurobiology of disease*. 2006; 24:280–285. [PubMed: 16962786]
23. Di Maria E, Marasco A, Tartari M, Ciotti P, Abbruzzese G, Novelli G, Bellone E, Cattaneo E, Mandich P. No evidence of association between BDNF gene variants and age-at-onset of Huntington's disease. *Neurobiology of disease*. 2006; 24:274–279. [PubMed: 16905325]
24. Metzger S, Bauer P, Tomiuk J, Laccone F, Didonato S, Gellera C, Mariotti C, Lange HW, Weirich-Schwaiger H, Wenning GK, Seppi K, Melegh B, Havasi V, Balik L, Wieczorek S, Zaremba J, Hoffman-Zacharska D, Sulek A, Basak AN, Soydan E, Zidovska J, Kebrdlova V, Pandolfo M, Riba P, Kadasi L, Kvasnicova M, Weber BH, Kreuz F, Dose M, Stuhmann M, Riess O. Genetic analysis of candidate genes modifying the age-at-onset in Huntington's disease. *Human Genetics*. 2006; 120:285–292. [PubMed: 16847693]
25. Andresen JM, Gayn J, Cherny SS, Brocklebank D, Alkorta-Aranburu G, Addis EA, Cardon LR, Housman DE, Wexler NS. Replication of twelve association studies for Huntington's disease residual age of onset in large Venezuelan kindreds. *Journal of Medical Genetics*. 2007; 44:44–50. [PubMed: 17018562]

Highlights

- The exact identification of genetic modifiers of HD is very important.
- We investigated *GRIK2* gene for a role in determining age at onset in HD.
- We used a much larger sample from multiple cohorts than in previous studies.
- we also applied the comprehensive analytical methods
- We detected no evidence of an influence of the *GRIK2* gene polymorphism on AO in HD.

Figure 1. *GRIK2* modifier analysis in 2,362 HD subjects with known age at motor onset
For the *GRIK2*TAA repeat polymorphism, subjects with at least one 16 or 17 TAA allele are displayed as closed orange and red circles, respectively, with all other subjects being represented by open black circles.
A. The relationship between *HD* expanded allele and log transformed age at onset of diagnostic motor signs.
B. Summary statistics for linear regression analysis of data shown in A, in each dominantly encoded *GRIK2* allele mode. The P value for each *GRIK2* allele mode tests for significant improvement in the model over use of the expanded CAG repeat length alone.
C. The distribution of the standardized residual (Z-Score) for subjects with 40-53 CAG repeats, calculated for each subject based upon the standard curve generated from 3,674 HD subjects in [17], is plotted against expanded CAG repeat length. Subjects with a 16 or 17 TAA allele are displayed as closed orange and red circles, respectively.
D. Summary statistics for global linear regression analysis of data shown in C, for each *GRIK2* genotype mode. The P value for each *GRIK2* allele mode tests for association with the residual variance in age at onset after accounting for the contribution of the CAG repeat length.

Figure 2. *GRIK2* modifier analysis in 547 HD subjects with known age at psychiatric onset and in 210 HD subjects with known age at cognitive onset

For the *GRIK2* TAA repeat polymorphism, subjects with at least one 16 or 17 TAA allele are displayed as closed orange and red circles, respectively, with all other subjects being represented by open black circles.

A. The relationship between *HD* expanded allele and log transformation of age at onset of psychiatric signs.

B. Summary statistics for linear regression analysis for data shown in A, in each *GRIK2* allele mode. The near-significant result for analysis of the TAA repeat as a continuous trait is skewed by the small number of extreme CAG repeat samples, as identical analysis of the bulk of subjects with CAG repeats of 40-53 yielded $P=0.503$

C. The relationship between *HD* expanded allele and log transformation of age at onset of cognitive signs.

D. Summary statistics for linear regression analysis for data shown in C, in each *GRIK2* allele mode.