Published in final edited form as:

Biol Psychiatry. 2013 May 1; 73(9): 836–842. doi:10.1016/j.biopsych.2013.01.002.

Does a shared neurobiology for foods and drugs of abuse contribute to extremes of food ingestion in anorexia and bulimia nervosa?

Walter H Kaye, MD¹, Christina E Wierenga, PhD¹, Ursula F Bailer, MD^{1,4}, Alan N Simmons, PhD^{3,2,1}, Angela Wagner, MD, PhD¹, and Amanda Bischoff-Grethe, PhD¹

¹University of California San Diego, Department of Psychiatry, La Jolla CA

²Veterans Affairs San Diego Healthcare System, Psychiatry Service, San Diego CA

³Veterans Affairs San Diego Healthcare System, Research Service, San Diego CA

⁴Medical University of Vienna, Department of Psychiatry and Psychotherapy, Division of Biological Psychiatry, Vienna, Austria

Abstract

Is starvation in anorexia nervosa (AN) or overeating in bulimia nervosa (BN) a form of addiction? Alternatively, why are individuals with BN more vulnerable and AN protected from substance abuse? Such questions have been generated by recent studies that suggest that there are overlapping neural circuits for foods and drugs of abuse.

In order to determine whether a shared neurobiology contributes to eating disorders (EDs) and substance abuse, this review focused on imaging studies that investigated response to tastes of food and tasks designed to characterize reward and behavioral inhibition in AN and BN.

BN and those with substance abuse disorders may share dopamine D2 receptor related vulnerabilities, and opposite findings may contribute to “protection” from substance abuse in AN. Moreover, imaging studies provide insights into executive cortico-striatal processes related to extraordinary inhibition and self-control in AN and diminished inhibitory self-control in BN that may influence the rewarding aspect of palatable foods and likely other consummatory behaviors.

AN and BN tend to have premorbid traits, such as perfectionism and anxiety that make them vulnerable to employing extremes of food ingestion which serve to reduce negative mood states. Dysregulation within and/or between limbic and executive cortico-striatal circuits contributes to such symptoms. Limited data support the hypothesis that reward and inhibitory processes may contribute to symptoms in ED and addictive disorders, but little is known about the molecular biology of such mechanisms in terms of shared or independent processes.

© 2013 Society of Biological Psychiatry. Published by Elsevier Inc. All rights reserved.

Corresponding Author: Walter Kaye, MD, Professor of Psychiatry, Director, UCSD Eating Disorders Treatment and Research Program, UCSD Department of Psychiatry, 8950 Villa La Jolla Dr. Suite C207, La Jolla, CA 92037, <http://eatingdisorders.ucsd.edu>, wkaye@ucsd.edu, Phone: 858 205 7293, Fax: 858 534 6727.

Financial Disclosures

The authors report no biomedical financial interests or potential conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

imaging; eating disorders; anorexia nervosa; bulimia nervosa; PET; fMRI; addictive Disorders

Clinical presentation of anorexia and bulimia nervosa

Anorexia nervosa (AN) and bulimia nervosa (BN) are disorders that are characterized by pathological eating behaviors and distorted body image. They have a narrow range of age of onset (early adolescence), stereotypic presentation of symptoms and course, and tend to be female gender specific. Although often considered to be caused by psychosocial factors, recent studies have shown that genetic heritability accounts for approximately 50–80% of the risk of developing AN and BN(1, 2) and contributes to neurobiological factors underlying these eating disorders (EDs).(3)

Two types of consummatory behavior are seen in AN and BN. Restricting-type anorexics (AN) lose weight purely by restricted dieting. Individuals with BN, who tend to maintain a normal body weight, alternate restricting with episodic binge eating and/or purging. In addition, some individuals have both AN and BN (AN-BN). For AN, there is an anxiety-reducing character to dietary restraint and reduced daily caloric intake(4-6), whereas eating stimulates dysphoric mood.(7) For BN, negative affect, mood lability, and stress may trigger binge eating episodes, and the binge and purge cycle may then serve to reduce dysphoria and/or anxiety.(8-13)

It has been argued that AN and BN share some common risk and liability factors because these disorders are often cross-transmitted in families, and transitions between AN and BN occur.(14-17) Importantly, AN and BN tend to share certain temperament and personality traits, which often first occur in childhood before the onset of an ED, and may create a vulnerability to develop an ED. These traits include anxiety(18), harm avoidance(19), perfectionism(20), obsessiveness(21, 22), and interoceptive deficits.(23) Moreover, these traits persist after recovery.(24)

In contrast, AN and BN tend to have opposite extremes of inhibitory self-control. Pure restrictor-type AN tend to be over-controlled and over-concerned about consequences. They have long been noted to be anhedonic and ascetic, able to sustain self-denial of food as well as most comforts and pleasures in life.(25) For example, AN have an enhanced ability to delay reward (i.e., show less reduction in the value of a monetary reward over time) compared to healthy volunteers.(26) This enhanced cognitive control and ability to delay reward may help to maintain persistent food restriction. In contrast, individuals with BN and AN-BN tend to have poor impulse control, engage in greater novelty, pleasure, and stimuli seeking behavior, and are less paralyzed by concerns with future consequences.(23, 24, 27-29)

From another perspective, there have been efforts to characterize behavioral domains that represent these symptoms. A considerable literature shows that harm avoidance (HA), a temperament trait (30) that contains elements of anxiety, inhibition, and inflexibility, is elevated in individuals who are ill and recovered from AN and BN.(27) In addition, AN and BN have increased sensitivity to punishment,(31) but AN have reduced sensitivity and BN have increased sensitivity to reward. ED individuals with bingeing and/or purging behavior also endorse greater sensation seeking than AN and controls, whereas AN tend to score lower on novelty seeking than BN, providing further evidence that differences in reward sensitivity and/or behavioral inhibition may differentiate AN from BN, though what distinguishes shared and independent traits remains unclear.(27)

Relationship of ED traits to substance use

This issue explores the relationship between food and addiction and overlapping neural circuits of reward and self-control. AN and BN show patterns of extremes of self-control and food consumption that appear to extend to their use of alcohol and illicit drugs. That is, meta-analyses(32, 33) show increased rates of drug and alcohol abuse in BN and decreased rates in AN. Because of space limitations, this review will focus on two areas of research in ED. First, we discuss dopamine (DA) positron emission tomography (PET) studies, because DA is considered a key to the rewarding effects of natural and drug rewards.(34) Second, in order to characterize the role of reward, interoception, and executive control in AN and BN, this review will focus on functional magnetic resonance imaging (fMRI) studies that have used taste of foods, reward, or inhibition tasks.

Difficulties untangling cause from consequence in EDs

Studies of ED must consider the impact of malnutrition on brain function. For example, malnutrition in AN(3) is associated with changes in brain structure (e.g., reduction in gray matter, greater cerebrospinal fluid (CSF) volume, altered white matter integrity) and profound metabolic, electrolyte, and endocrine disturbances. Furthermore, studies in animals suggest that diet and weight can influence DA metabolism.(35, 36) One method of avoiding the confounding effects of abnormal nutritional status is to study those who have recovered (REC) from AN and BN, although it remains conjectural whether abnormal findings reflect traits or scars. Thus, cause and consequence of pathological eating and the impact of malnutrition on neural processes in AN and BN remain a major methodological question.

DA function in AN and BN

Several lines of evidence suggest that individuals with ED have altered striatal DA function. PET studies with the radioligand [¹¹C]raclopride find that REC AN have increased binding of DA D2/D3 receptors at baseline in the anterior ventral striatum (AVS) relative to controls.(25, 37) Because PET measures of [¹¹C]raclopride binding are sensitive to endogenous DA concentrations,(38) this difference could be due to either a reduction in the intrasynaptic DA concentration or an elevation of the density and/or affinity of the D2/D3 receptors in this region. The former is supported by data showing that REC AN have a reduction of the DA metabolite homovanillic acid (HVA) in CSF. To our knowledge, no PET DA studies have been reported in ill AN.

Dorsal caudate and dorsal putamen DA D2/D3 binding potential (BP_{ND}) was positively and significantly associated with HA in REC AN/AN-BN(25) and in REC AN/AN-BN/BN.(37) Importantly, rats characterized as risk averse(39) showed greater D2 mRNA expression in the dorsal caudate, supporting human studies suggesting that DA signaling in executive cortical striatal circuitry is related to risk of adverse consequences or inhibitory control(40, 41) or anxiety.(42) A study(43) using PET [¹¹C]raclopride binding with amphetamine to assess endogenous DA production found that DA release in the precommissural dorsal caudate (preDCA) was associated with increased anxiety in REC AN. In contrast, controls showed euphoria associated with AVS DA release, consistent with many, but not all control responses to amphetamine.(44-46) Ingestion of palatable food is associated with striatal endogenous DA release.(47) If AN experience endogenous DA release as anxiogenic, rather than hedonic, it may explain their pursuit of starvation, because food refusal may be an effective means of diminishing such anxious feelings in AN individuals.

In terms of DA studies in ill BN individuals, while CSF HVA levels were normal, two studies found reduced CSF HVA(48, 49) was associated with increased frequency of binge behaviors in ill BN. A stimulant PET [¹¹C]raclopride paradigm(50) in ill BN found a trend

toward decreased mean D2/D3 receptor BP_{ND} in the posterior putamen and caudate, and a significantly reduced/blunted DA release to the psychostimulant challenge in the anterior and posterior putamen. Similar to the CSF HVA studies, there was a statistically significant negative association between the frequencies of binge eating and vomiting and the striatal DA response. Studies overfeeding animals with a palatable high-energy diet have shown progressively worsening reward deficit with lower basal and evoked DA levels in the nucleus accumbens and reduced D2 striatal levels.(for review see Kenny(51)) Thus studies showing that binge behavior in ill BN is associated with reduced CSF HVA and DA release are consistent with these rodent studies. Importantly, REC BN and AN-BN have normal DA D2/D3 striatal binding.(37) and CSF HVA levels supporting the notion of being “addicted” to food.(52) However, a study using a catecholamine depletion design, points to reduced reward learning in REC BN.(53) Whether this is a trait vulnerability, or a scar creating a persistent dysregulation of the central catecholaminergic neurotransmitter systems is unknown.

There have been relatively consistent findings (see(54, 55)) of reduced PET [^{11}C]raclopride binding potential in the striatum, including the ventral striatum in those with substance abuse disorders. It is of potential relevance that decreased D2/D3 receptor binding has been found in obese subjects (see(51, 56)). There is some indication that ill BN individuals may have reduced striatal [^{11}C]raclopride binding, which raises the intriguing possibility that BN and substance abuse or obesity share DA D2 receptor related vulnerabilities. As noted above, three studies have found correlations between reduced DA activity and increased binge and vomit frequency, although the cause and consequence between consummatory patterns and DA function remain conjectural. While data are also very limited, REC restricting type AN have increased AVS [^{11}C]raclopride binding. This raises the provocative question of whether such opposite findings contribute to “protection” from substance abuse in AN. The reasons for possible striatal regional differences between AN and BN are not known. DA D2/D3 receptors are just one small part of a complex DA system that involves the interaction of a number of DA receptors and other molecules. There are differences within striatal regions in terms of the relative balance of D2 and D3 receptors (57), expression of the DA transporter (58), and DA D1 receptor density.(59) Moreover, there are not homogenous regions in terms of function. For example, DA D2/D3 receptors show opposite roles in subregions of the nucleus accumbens,(60) Finally, there is uncertainty regarding state-effects of illness and whether these are secondary to extremes of dietary intake, premorbid and persistent traits, or scars.

Higher order neural processes modulating reward, emotionality, and inhibition

Data support the hypothesis(3) that AN and BN individuals have an imbalance within and/or between ventral limbic and dorsal cognitive circuits. Specifically, a ventral limbic neural circuit, which includes the amygdala, anterior insula (AI), AVS, ventral regions of the anterior cingulate cortex (ACC) and the orbitofrontal cortex (OFC), is involved in identifying rewarding and emotionally significant stimuli and for generating affective responses to these stimuli.(61, 62) A dorsal executive function neural circuit, which includes dorsal regions of the caudate, dorsolateral prefrontal cortex (DLPFC), parietal cortex, and other regions, is thought to modulate selective attention, planning, and effortful regulation of affective states.(61, 62) The ventral limbic and dorsal cognitive neural circuits are critically involved in inhibitory decision-making processes, especially involving reward-related behaviors. Together they process the reward value and/or affective valence of environmental stimuli, assess the future consequences of one’s own actions (response selection) and inhibit inappropriate behaviors (response inhibition). Dysfunction within these regions has been

proposed to be a key neural mechanism underlying altered behavioral regulation, reward regulation, or cognition found in addiction.(63, 64)

Reward and interoceptive function in regards to tasting food in AN and BN

fMRI imaging studies of appetitive behaviors in ED have used designs that either administer a taste of some food or present images of food. This review will focus on studies of taste because much is known about how tastes of food activate reward and interoceptive circuitry in healthy subjects.(34, 65) In brief, the AI is the primary gustatory taste cortex, and thus responds to tastes of food (see(3)). The AI, as well as the ACC and OFC, code the sensory-hedonic response to taste and innervate a broad region of the rostral ventral-central striatum, (66-68) where behavioral repertoires are then computed. Thus this network may play a crucial role in linking sensory-hedonic experiences to the motivational components of reward(66) as well as emotionality, providing conscious awareness of these urges.(34, 62)

While there are relatively few neuroimaging studies of taste and interoceptive function in ED, the literature to date is relatively consistent between the ill and REC states, supporting the possibility that reward and interoceptive processing are trait characteristics. A study(69) which administered tastes of sucrose or water found that REC AN showed lower neural activation of the insula, including the primary cortical taste region, and ventral and dorsal striatum to both sucrose and water compared to controls. Insular neural activity correlated with pleasantness ratings for sucrose in CW, but not in AN subjects. A study(70) that used response to chocolate milk when hungry and satiated, found that ill AN had minimal insula response to the tastes of food when hungry. Administration of boluses of cream, water, and a non-caloric viscous taste(71) found that REC BN showed an exaggerated AVS response for the cream/water contrast in comparison to REC AN or CW. Considered together, these findings raise the intriguing possibility that individuals who undereat or overeat have an altered set-point, and/or altered sensitivity for sensory-interoceptive-reward processes, when consuming palatable foods.(65) Thus the set-point for undereaters may mimic a continuous state of satiety that limits interoceptive and reward processing, whereas overeaters may experience a more tonic state of subjective hunger and thus hyper-activate related brain circuits.

Recent studies in the field of obesity have highlighted the importance of anticipation in the neural response to food stimuli. For example, animal studies(72) show that following stimulus conditioning, DA neurons shift firing from the consumption of food to the anticipated consumption of food or to cues associated with food consumption. There are a number of studies using tastants in ED in which the design might elicit anticipatory responses. An associative learning task between conditioned visual stimuli and unconditioned sucrose taste stimuli(73) found that reward learning signals were greater in the ill AN and less in the obese participants compared to the controls in the AVS, insula, and OFC, suggesting that brain reward circuits were more responsive in AN and less responsive in obese participants. In a complex design viewing and tasting alone/together pleasant/unpleasant stimuli that appeared to mix anticipatory and consummatory response,(74) REC AN had increased ventral striatal activity in response to sights and flavors of a pleasant stimuli (chocolate) and increased insula and posterior dorsal caudate response to flavors and sights of aversive foods when compared to controls. Taken together these studies suggest AN may be highly responsive to stimulus cues, perhaps as a mechanism to predict and control the anxiety produced by stimuli that may be associated with subjective unpleasantness, similar to anticipatory sensitivity connected with stimulus avoidance found in highly anxious individuals.(75)

Interestingly, a reciprocal processing pattern has been observed in BN individuals. Using a response to anticipated or tastes of chocolate milkshake (vs. tasteless solution)(76) ill BN women, in contrast to healthy controls, showed trends for hypo-activation in the right anterior insula in response to anticipated and in the left middle frontal gyrus, right posterior insula, right precentral gyrus, and right mid dorsal insula in response to consumption. A conditioned visual stimuli and unconditioned sucrose taste stimuli paradigm(77) found that ill BN individuals showed reduced brain response compared with controls for unexpected receipt and omission of taste stimuli, as well as reduced brain regression response to the temporal difference learning computer model generated reward values in insula, ventral putamen, amygdala, and OFC. These studies suggest that in contrast to AN, BN may be less sensitive to food cues, which may contribute to their tendency to overeat during binge episodes.

The imbalance of stimulus consumption and anticipation in AN and BN may suggest dysfunctional stimulus integration that could relate to the clinically observed disconnect between reported and actual interoceptive states. Key in this process is the insula which is thought to code interoceptive prediction error, signaling mismatch between actual and anticipated bodily arousal, which in turn elicits subjective anxiety and avoidance behavior. (78) A study of response to pain confirms a mismatch between anticipation and objective responses in REC AN.(79) That is, REC AN compared to controls, showed greater activation within right AI, DLPFC and cingulate cortex during pain anticipation, and greater activation within DLPFC and decreased activation within posterior insula during painful stimulation. Greater anticipatory AI activation correlated positively with alexithymia (difficulty identifying emotions) in REC AN subjects, suggesting an altered ability to accurately perceive bodily signals persists after recovery. Such disturbances could be tied to the preponderance of alexithymia in AN,(80) which also is related to increased insula activation and is thought to be tied to deficits in emotional processing.(81) Given that food is an inherently emotional stimulus in AN and BN, a mismatch between limbic hyperarousal and cognitive control may be tied to both interoceptive and emotional processing deficits in this population.

Reward and inhibition studies

A recent series of fMRI studies based on reward and inhibition tasks has explored the response of ventral and dorsal corticostriatal systems in AN and BN in comparison to controls to better understand the modulation of reward, emotionality, and behavioral inhibition. Using a monetary choice task to investigate response to positive and negative feedback, REC AN(82) and REC BN(83) had a similar abnormal AVS response in that they failed to differentiate wins and losses in comparison to controls who distinguished positive and negative feedback. Animal studies show that the AVS processes motivational aspects toward stimuli by modulating the influence of limbic inputs on striatal activity(84, 85) so that even abstract rewards (money) activate the AVS in proportion to the reward amount or the deviation from an expected payoff.(86) This may indicate a failure in REC AN and REC BN to appropriately bind, scale, or discriminate responses to salient stimuli, suggesting an impaired ability to identify the emotional significance of stimuli.(83)

In addition REC AN also exhibited(82) an exaggerated response to positive and negative feedback in the dorsal caudate relative to CW as well as elevated activity in executive cortical regions (dorsal lateral prefrontal and parietal) which project to the dorsal caudate(67) and which was positively associated with baseline trait anxiety for both wins and losses. Using a set shifting fMRI paradigm(87) ill AN showed predominant activation of frontoparietal networks indicative of excessive effortful and supervisory cognitive control during task performance, but hypoactivation in the ventral anterior cingulate-striato-thalamic

loop involved in motivation related behavior. Similarly, an increase in brain response in posterior visual and inferior parietal cortex regions was found in ill restricting type AN that correlated with correctly inhibited no-go trials.(88) Using a Stop signal task(89) REC AN subjects had altered task-related activation in the medial prefrontal cortex (mPFC), a critical node of the inhibitory control network.

Together, these results support the possibility that AN have an imbalance in information processing, with impaired ability to identify the emotional significance of a stimulus but increased traffic in neurocircuits concerned with planning and consequences and which is associated with anxiety. This over-reliance on cognitive brain circuits involved in linking action to outcome may constitute an attempt at “strategic” (as opposed to hedonic) means of responding to reward stimuli. Does this protect AN against substance abuse disorders? It is noteworthy that cognitive strategies that have been shown to effectively regulate craving are associated with increased brain response in dorsal-frontal-striatal regions associated with cognitive control and decreased brain response in ventral limbic striatal regions.(90)

In contrast to the possibility that excessive inhibitory control may characterize AN, inhibitory control processes may be impaired in women with BN, likely because of their failure to engage frontostriatal circuits appropriately. For example, during correct responding on incongruent trials of the Simon Spatial Incompatibility task,(29, 91) ill BN adolescents and adults failed to activate the left inferiolateral prefrontal cortex, bilateral inferior frontal gyrus, lenticular and caudate nuclei, and ACC. They also demonstrated aberrant function of the dorsal ACC, activating it more when making errors than when responding correctly. In support of such a possibility, REC BN(83) had abnormal responses to positive and negative feedback in the dorsal caudate. In comparison, ill AN-BN(88) had increased activation of the right DLPFC using a Go-NoGo task, so that it remains to be determined whether AN-BN and BN individuals have discrepant findings in terms of dorsal circuit activity. In summary, these findings provide evidence of functional abnormalities in BN within the neural system that subserves self-regulatory control, which may contribute to binge eating and other impulsive behaviors.

Summary

It remains conjectural whether years of pathological eating is the sole cause of aberrant neurobiology in AN. Whereas most people find starvation to be unpleasant, and recidivism with dieting is high, AN individuals find that food is anxiety provoking and starvation comforting. If AN were merely a consequence of an out of control diet, the rate of AN in this society would probably be much higher. Alternatively, underlying traits may be contributory. For example, many of the characteristic temperament and personality behaviors typical of AN and BN are expressed in childhood, years before the onset of an ED, suggesting the importance of certain vulnerabilities. It is also possible that altered brain function results from both underlying neurobiological traits as well as years of pathological behavior. There have been relatively few imaging studies published in ED so there is a limited body of data regarding neural processes in ED. Very limited data suggest that some abnormal brain processes are relatively similar in the ill and REC state, suggesting little influence of illness state. However, the prospective and longitudinal studies necessary to definitively answer these questions have not been done.

While review studies raise the possibility that individuals with AN might be protected from, and BN vulnerable to substance abuse, it is not known whether this is related to a shared biology with appetite regulation. Although few studies have been done, restricting type AN have increased AVS [¹¹C]raclopride binding whereas there are suggestions that BN have reduced striatal [¹¹C]raclopride binding. This raises the intriguing possibility that BN and

substance abuse share DA D2 receptor related vulnerabilities to the rewarding aspect of palatable foods or substance use, whereas opposite findings may contribute to “protection” from substance abuse in AN. In addition, AN have extraordinary inhibition and self-control, whereas BN have diminished inhibitory self-control. fMRI studies consistently show that AN and BN have altered activity of dorsal executive/cognitive circuitry, that tends to be consistent with the possibility that AN have enhanced higher-order inhibitory function and BN have reduced inhibition. We hypothesize that AN are able to inhibit consummatory drives and have extraordinary self-control because they have exaggerated dorsal cognitive circuit function, whereas BN individuals are vulnerable to overeating when hungry and use substances because they have less ability to self-regulate and control their impulses.

Despite these advances in the neurocircuitry of appetite, reward, and cognitive control, several questions remain regarding implications for EDs in the ill and recovered state. For example, what is the cause and effect between reward and inhibition? Studies of food tastes and negative and positive feedback consistently show altered ventral limbic circuitry in AN and BN. It is possible that both AN and BN have an inability to precisely identify and/or modulate emotionality and reward in response to salient stimuli. Poor context separation may make them vulnerable to inappropriately responding to the rewarding aspects of consummatory stimuli. Do AN have “normal” cognitive association networks in the dorsal neurocircuit (e.g. DLPFC-to dorsal striatum) that directs motivated actions in the face of impaired ability of the ventral striatal pathways to direct more ‘automatic’ or intuitive motivated responses? Alternatively, is otherwise adequate ventral limbic-striatal circuitry too strongly inhibited by “hyperactive” inputs from cognitive domains such as the DLPFC and the parietal cortex? For BN, is there hypoactivity of frontal cognitive pathways that is not sufficient to inhibit “normal” hedonic/incentive motivation impulses in “reward” neural circuits, or are there exaggerated hedonic/incentive motivation impulses in “reward” neural circuits that overwhelm normal cognitive inhibition? Discovering answers to some of these questions will bring us closer to developing effective pharmacological and behavioral treatments for these deadly disorders.

Finally, the future of ED may well be in taking a dimensional approach using the Research Domains.⁽⁹²⁾ ED research has shown functional brain deficits in all five of the primary Research Domains: Negative Valence Systems (i.e., increased anxiety), Positive Valence Systems (i.e., increased habit formation and decreased reward processing), Cognitive Systems (i.e., increased behavioral inhibition), Systems for Social Processes (i.e., increased self and social evaluations), and Arousal/Regulatory Systems (i.e., interoceptive deficits). Future research should take into account the changing landscape of how psychiatric disorders are conceptualized, for both the DSM-V and beyond, and should aim to measure the behavioral aspects of ED in a dimensional way such that these conceptual models can be better quantified and understood.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

W Kaye: UCSD, National Institute of Mental Health (NIMH) MH042984 and MH092793, and the Price Foundation.

C Wierenga: UCSD, National Institute of Mental Health (NIMH) MH042984 and MH092793

U Bailer: UCSD, National Institute of Mental Health (NIMH) MH092793, Medical University of Vienna, Austria, UCSD, the Hilda & Preston Davis Foundation

A Simmons: CESAMH, National Institute of Mental Health (NIMH) MH042984

A Bischoff-Grethe: MH086017 and MH042984, the Price Foundation

References

1. Bulik C, Sullivan PF, Tozzi F, Furberg H, Lichtenstein P, Pedersen NL. Prevalence, heritability and prospective risk factors for anorexia nervosa. *Archives of General Psychiatry*. 2006; 63:305–312. [PubMed: 16520436]
2. Kaye WH, Bulik C, Plotnicov K, Thornton L, Devlin B, Fichter M, et al. The genetics of anorexia nervosa collaborative study: Methods and sample description. *Int J Eat Disord*. 2008; 41:289–300. [PubMed: 18236451]
3. Kaye W, Fudge J, Paulus M. New insight into symptoms and neurocircuit function of anorexia nervosa. *Nat Rev Neurosci*. 2009; 10:573–584. [PubMed: 19603056]
4. Kaye, W.; Strober, M.; Klump, KL. Neurobiology of eating disorders. In: Martin, A., et al., editors. *Pediatric Psychopharmacology, Principles & Practice*. New York: Oxford University Press; 2003. p. 224–237.
5. Vitousek K, Manke F. Personality variables and disorders in anorexia nervosa and bulimia nervosa. *J Abnorm Psychol*. 1994; 103:137–147. [PubMed: 8040475]
6. Steinglass J, Sysko R, Mayer L, Berner L, Schebendach J, Wang Y, et al. Pre-meal anxiety and food intake in anorexia nervosa. *Appetite*. 2010; 55:214–218. [PubMed: 20570701]
7. Frank G, Kaye W. Current status of functional imaging in eating disorders. *Int J Eat Disord*. 2012; 45:723–736. [PubMed: 22532388]
8. Abraham S, Beaumont P. How patients describe bulimia or binge eating. *Psychol Med*. 1982; 12:625–635. [PubMed: 6957905]
9. Johnson C, Larson R. Bulimia: an analysis of mood and behavior. *Psychosom Med*. 1982; 44:341–351. [PubMed: 6959174]
10. Kaye WH, Gwirtsman HE, George DT, Weiss SR, Jimerson DC. Relationship of mood alterations to bingeing behaviour in bulimia. *British Journal of Psychiatry*. 1986; 149:479–485. [PubMed: 3814933]
11. Smyth J, Wonderlich S, Heron K, Sliwinski M, Crosby R, Mitchell J, et al. Daily and momentary mood and stress are associated with binge eating and vomiting in bulimia nervosa patients in the natural environment. *J Cons Clin Psychol*. 2007; 75:629–638.
12. Crosby R, Wonderlich S, Engel S, Simonich H, Smyth J, Mitchell J. Daily mood patterns and bulimic behaviors in the natural environment. *Behav Res Ther*. 2009; 47:181–188. [PubMed: 19152874]
13. Haedt-Matt A, Keel P. Revisiting the affect regulation model of binge eating: a meta-analysis of studies using ecological momentary assessment. *Psychol Bull*. 2011; 137:660–681. [PubMed: 21574678]
14. Kendler KS, MacLean C, Neale M, Kessler R, Heath A, Eaves L. The genetic epidemiology of bulimia nervosa. *Am J Psychiatry*. 1991; 148:1627–1637. [PubMed: 1842216]
15. Lilenfeld LR, Kaye WH, Greeno CG, Merikangas KR, Plotnicov K, Pollice C, et al. A controlled family study of anorexia nervosa and bulimia nervosa: psychiatric disorders in first-degree relatives and effects of proband comorbidity. *Arch Gen Psychiatry*. 1998; 55:603–610. [PubMed: 9672050]
16. Strober M, Freeman R, Lampert C, Diamond J, Kaye W. Controlled family study of anorexia nervosa and bulimia nervosa: evidence of shared liability and transmission of partial syndromes. *Am J Psychiatry*. 2000; 157:393–401. [PubMed: 10698815]
17. Walters EE, Kendler KS. Anorexia nervosa and anorexic-like syndromes in a population-based female twin sample. *American Journal of Psychiatry*. 1995; 152:64–71. [PubMed: 7802123]
18. Kaye W, Bulik C, Thornton L, Barbarich N, Masters K, Fichter M, et al. Comorbidity of anxiety disorders with anorexia and bulimia nervosa. *Am J Psychiatry*. 2004; 161:2215–2221. [PubMed: 15569892]

19. Fassino S, Abbate-Daga G, Amianto F, Leombruni P, Boggio S, Rovera GG. Temperament and character profile of eating disorders: a controlled study with the temperament and character inventory. *Int J Eat Disord*. 2002; 32:412–425. [PubMed: 12386906]
20. Friederich H, Herzog D. Cognitive-behavioral flexibility in anorexia nervosa. *Curr Top Behav Neurosci*. 2011; 6:111–123. [PubMed: 21243473]
21. Anderluh MB, Tchanturia K, Rabe-Hesketh S, Treasure J. Childhood obsessive-compulsive personality traits in adult women with eating disorders: defining a broader eating disorder phenotype. *American Journal of Psychiatry*. 2003; 160:242–247. [PubMed: 12562569]
22. van den Heuvel O, Veltman D, Groenewegen H, Witter M, Merkelbach J, Cath D, et al. Disorder-specific neuroanatomical correlates of attentional bias in obsessive-compulsive disorder, panic disorder, and hypochondriasis. *Arch Gen Psychiatry*. 2005; 62:922–933. [PubMed: 16061770]
23. Lilenfeld L, Wonderlich S, Riso LP, Crosby R, Mitchell J. Eating disorders and personality: a methodological and empirical review. *Clinical Psychology Review*. 2006; 26:299–320. [PubMed: 16330138]
24. Wagner A, Barbarich N, Frank G, Bailer U, Weissfeld L, Henry S, et al. Personality traits after recovery from eating disorders: Do subtypes differ? *International Journal of Eating Disorders*. 2006; 39:276–284. [PubMed: 16528697]
25. Frank G, Bailer UF, Henry S, Drevets W, Meltzer CC, Price JC, et al. Increased dopamine D2/D3 receptor binding after recovery from anorexia nervosa measured by positron emission tomography and [¹¹C]raclopride. *Biological Psychiatry*. 2005; 58:908–912. [PubMed: 15992780]
26. Steinglass J, Albano A, Simpson H, Carpenter K, Schebendach J, Attia E. Fear of food as a treatment target: Exposure and response prevention for anorexia nervosa in an open series. *Int J Eat Disord*. 2012; 45:615–621. [PubMed: 21541979]
27. Cassin S, von Ranson K. Personality and eating disorders: a decade in review. *Clinical Psychology Review*. 2005; 25:895–916. [PubMed: 16099563]
28. Claes L, Nederkoorn C, Vandereycken W, Guerrieri R, Vertommen H. Impulsiveness and lack of inhibitory control in eating disorders. *Eat Behav*. 2006; 7:196–203. [PubMed: 16843221]
29. Marsh R, Steinglass J, Gerber A, Graziano O'Leary K, Wang Z, Murphy D, et al. Deficient activity in the neural systems that mediate self-regulatory control in bulimia nervosa. *Arch Gen Psychiatry*. 2009; 66:51–63. [PubMed: 19124688]
30. Cloninger, C.; Przybeck, T.; Svrakic, D.; Wetzel, R. *The Temperament and Character Inventory (TCI): A guide to its development and use*. St Louis: 1994. p. 19-28.
31. Harrison A, O'Brien N, Lopez C, Treasure J. Sensitivity to reward and punishment in eating disorders. *Psy Res*. 2010; 177:1–11.
32. Calero-Elvira A, Krug I, Davis K, Lopez C, Fernandez-Aranda F, Treasure J. Meta-analysis on drugs in people with eating disorders. *Eur Eat Disord Rev*. 2009; 17:243–259. [PubMed: 19475697]
33. Gadalla T, Piran N. Co-occurrence of eating disorders and alcohol use disorders in women: a meta analysis. *Arch Womens Ment Health*. 2007; 10:133–140. [PubMed: 17533558]
34. Volkow N, Wang G, Fowler J, Tomasi A, Baler R. Food and drug reward: overlapping circuits in human obesity and addiction. *Curr Top Behav Neurosci*. 2012; 11:1–24. [PubMed: 22016109]
35. Avena N, Borcarys M, Hoebel B. Animal models of sugar and fat bingeing: relationship to food addiction and increased body weight. *Methods Mol Biol*. 2012; 829:351–365. [PubMed: 22231826]
36. Johnson P, Kenny P. Dopamine D2 receptors in addiction-like reward dysfunction and compulsive eating in obese rats. *Nat Neurosci*. 2010; 13:635–641. [PubMed: 20348917]
37. Bailer U, Frank G, Price J, Meltzer C, Becker C, Mathis C, et al. Interaction between serotonin transporter and dopamine D2/D3 receptor radioligand measures is associated with harm avoidant symptoms in anorexia and bulimia nervosa. *Psych Res Neuroimaging*. 2012 In Press.
38. Drevets W, Gautier C, Price J, Kupfer D, Kinahan P, Grace A, et al. Amphetamine-induced dopamine release in human ventral striatum correlates with euphoria. *Biological Psychiatry*. 2001; 49:81–96. [PubMed: 11164755]
39. Simon N, Montgomery K, Beas B, Mitchell M, LaSarge C, Mendez I, et al. Dopaminergic modulation of risky decision-making. *J Neuroscience*. 2011; 31:17460–17470.

40. Ghahremani D, Lee B, Robertson C, Tabibnia G, Morgan A, De Shelter N, et al. Striatal Dopamine D2/D3 Receptors Mediate Response Inhibition and Related Activity in Frontostriatal Neural Circuitry in Humans. *J Neuroscience*. 2012; 32:7316–7324.
41. Kaasinen V, Nurmi E, Bergman J, Eskola J, Solin O, Sonninen P, et al. Personality traits and brain dopaminergic function in Parkinson's disease. *Proceedings of the National Academy of Science USA*. 2001; 98:13272–13277.
42. Laakso A, Wallius E, Kajander J, Bergman J, Eskola O, Solin O, et al. Personality traits and striatal dopamine synthesis capacity in healthy subjects. *American Journal of Psychiatry*. 2003; 160:904–910. [PubMed: 12727694]
43. Bailer U, Narendran R, Frankle W, Himes M, Duvvuri V, Mathis C, et al. Amphetamine induced dopamine release increases anxiety in individuals recovered from anorexia nervosa. *Int J Eat Disord*. 2012; 45:263–271. [PubMed: 21541980]
44. Brauer L, de Wit H. Subjective responses to D-amphetamine alone and after pimozide pretreatment in normal, healthy volunteers. *Biol Psych*. 1996; 39:26–31.
45. de Wit H, Uhlenhuth E, Johanson C. Individual differences in the reinforcing and subjective effects of amphetamine and diazepam. *Drug Alcohol Depend*. 1986; 16:341–360. [PubMed: 3698813]
46. Gabbay F. Variations in affect following amphetamine and placebo: markers of stimulant drug preference. *Exp Clin Psychopharmacol*. 2003; 11:91–101. [PubMed: 12622347]
47. Bassareo V, Di Chiara G. Differential responsiveness of dopamine transmission to food-stimuli in nucleus accumbens shell/core compartments. *Neuroscience*. 1999; 89:637–641. [PubMed: 10199600]
48. Jimerson DC, Lesem MD, Kaye WH, Brewerton TD. Low serotonin and dopamine metabolite concentrations in cerebrospinal fluid from bulimic patients with frequent binge episodes. *Archives of General Psychiatry*. 1992; 49:132–138. [PubMed: 1372494]
49. Kaye WH, Ballenger JC, Lydiard RB, Stuart GW, Laraia MT, O'Neil P, et al. CSF monoamine levels in normal-weight bulimia: evidence for abnormal noradrenergic activity. *American Journal of Psychiatry*. 1990; 147:225–229. [PubMed: 1689122]
50. Broft A, Shingleton R, Kaufman J, Liu F, Kumar D, Slifstein M, et al. Striatal dopamine in bulimia nervosa: A PET imaging study. *Int J Eat Disord*. 2012; 45:648–656. [PubMed: 22331810]
51. Kenny P. Reward mechanisms in obesity: new insights and future directions. *Neuron*. 2011; 69:664–679. [PubMed: 21338878]
52. Kaye WH, Frank GK, McConaha C. Altered dopamine activity after recovery from restricting-type anorexia nervosa. *Neuropsychopharmacology*. 1999; 21:503–506. [PubMed: 10481833]
53. Grob S, Pizzagalli D, Dutra S, Stern J, Morgeli H, Milos G, et al. Dopamine-related deficit in reward learning after catecholamine depletion in unmedicated, remitted subjects with bulimia nervosa. *Neuropsychopharm*. 2012; 37:1945–1952.
54. Volkow N, Fowler J, Wang G, Baler R, Telang F. Imaging dopamine's role in drug abuse and addiction. *Neuropharm*. 2009; 56(Suppl 1):3–8.
55. Volkow N, Wang G, Fowler J, Tomasi A. Addiction circuitry in the human brain. *Ann Rev Pharmacol Toxicol*. 2012; 52:321–336. [PubMed: 21961707]
56. Volkow N, Wang G, Baler R. Reward, dopamine and the control of food intake: implications for obesity. *Trends Cogn Sci*. 2011; 15:37–46. [PubMed: 21109477]
57. Gurevich E, Joyce J. Distribution of dopamine D3 receptor expressing neurons in the human forebrain: Comparison with D2 receptor expressing neurons. *Neuropsychopharmacology*. 1999; 20:60–80. [PubMed: 9885786]
58. Martinez D, Slifstein M, Broft A, Mawlawi O, Hwang D, Huang Y, et al. Imaging human mesolimbic dopamine transmission with positron emission tomography. Part II: amphetamine-induced dopamine release in the functional subdivisions of the striatum. *Journal of Cerebral Blood Flow and Metabolism*. 2003; 23:285–300. [PubMed: 12621304]
59. Muly E, Maddox M, Khan Z. Distribution of D(1) and D(5) dopamine receptors in the primate nucleus accumbens. *Neuroscience*. 2010; 169:1557–1566. [PubMed: 20600664]
60. Besson M, Belin D, McNamara R, Theobald D, C A, Beckett V, C BM, Newman A, et al. Dissociable control of impulsivity in rats by dopamine D2/3 receptors in the core and shell

- subregions of the nucleus accumbens. *Neuropsychopharmacology*. 2010; 35:560–569. [PubMed: 19847161]
61. Phillips M, Drevets W, Rauch S. Neurobiology of emotion perception II: implications for major psychiatric disorders. *Biol Psych*. 2003; 54:515–528.
 62. Phillips M, Drevets W, R SL, Lane R. Neurobiology of emotion perception I: The neural basis of normal emotion perception. *Biol Psych*. 2003; 54:504–514.
 63. Goldstein R, Volkow ND. Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *Am J Psychiatry*. 2002; 159:1642–1652. [PubMed: 12359667]
 64. Feil J, Sheppard D, Fitzgerald P, Yucel M, Lubman D, Bradshaw J. Addiction, compulsive drug seeking, and the role of frontostriatal mechanisms in regulating inhibitory control. *Neurosci Biobehav Rev*. 2012; 35:248–275. [PubMed: 20223263]
 65. Small D. Individual differences in the neurophysiology of reward and the obesity epidemic. *Int J Obes*. 2009; 33(Suppl 2):S44–48.
 66. Devinsky O, Morrell MJ, Vogt BA. Contributions of anterior cingulate cortex to behaviour. *Brain*. 1995; 118:279–306. [PubMed: 7895011]
 67. Haber SN, Kim K, Mailly P, Calzavara R. Reward-related cortical inputs define a large striatal region in primates that interface with associative cortical connections, providing a substrate for incentive-based learning. *J Neurosci*. 2006; 26:8368–8376. [PubMed: 16899732]
 68. Rolls ET. Taste, olfactory, and food texture processing in the brain, and the control of food intake. *Physiol Behav*. 2005; 85:45–56. [PubMed: 15924905]
 69. Wagner A, Aizenstein H, Frank GK, Figurski J, May JC, Putnam K, et al. Altered insula response to a taste stimulus in individuals recovered from restricting-type anorexia nervosa. *Neuropsychopharmacology*. 2008; 33:513–523. [PubMed: 17487228]
 70. Vocks S, Herpertz S, Rosenberger C, Senf W, Gizewski E. Effects of gustatory stimulation on brain activity during hunger and satiety in females with restricting-type anorexia nervosa: An fMRI study. *J Psychiatr Res*. 2011; 45:395–403. [PubMed: 20709330]
 71. Radeloff D, Willmann K, Otto L, Lindner M, Putnam K, van Leeuwen S, et al. High-fat taste challenge reveals altered striatal response in women recovered from bulimia nervosa - a pilot study. *World J Biol Psychiatry*. 2012 Epub ahead of print.
 72. Stice E, Spoor S, Bohon C, Small D. Relation between obesity and blunted striatal response to foods is moderated by TaqIA A1 allele. *Science*. 2008; 322:449–452. [PubMed: 18927395]
 73. Frank G, Reynolds J, Shott M, Jappe L, Yang T, Tregellas J, et al. Anorexia nervosa and obesity are associated with opposite brain reward response. *Neuropsychopharm*. 2012; 37:2031–2046.
 74. Cowdrey F, Park R, Harmer C, McCabe C. Increased neural processing of rewarding and aversive food stimuli in recovered anorexia nervosa. *Biol Psych*. 2011; 70:736–743.
 75. Simmons A, Strigo I, Matthews S, Paulus M, Stein M. Anticipation of aversive visual stimuli is associated with increased insula activation in anxiety-prone subjects. *Biol Psychiatry*. 2006; 60:402–409. [PubMed: 16919527]
 76. Bohon C, Stice E. Reward abnormalities among women with full and subthreshold bulimia nervosa: a functional magnetic resonance imaging study. *Int J Eat Disord*. 2011; 44:585–595. [PubMed: 21997421]
 77. Frank G, Reynolds J, Shott M, O'Reilly R. Altered temporal difference learning in bulimia nervosa. *Biol Psych*. 2011; 70:728–735.
 78. Paulus M, Stein MB. An insular view of anxiety. *Biol Psychiatry*. 2006; 60:383–387. [PubMed: 16780813]
 79. Strigo I, Matthews S, Simmons A, Oberndorfer T, Reinhardt L, Klabunde M, et al. Altered insula activation during pain anticipation in individuals recovered from anorexia nervosa: evidence of interoceptive dysregulation. *Int J Eat Disord*. 2012 In Press. 10.1002/eat.22045
 80. Sexton M, Sunday S, Hurt S, Halmi K. The relationship between alexithymia, depression, and axis II psychopathology in eating disorder inpatients. *Int J Eat Disord*. 1998; 23:277–286. [PubMed: 9547662]

81. Heinzel A, Schafer R, M H, Schieffer A, Ingenhag A. Differential modulation of valence and arousal in high-alexithymic and low-alexithymic individuals. *Neuroreport*. 2010; 21:998–1002. [PubMed: 20811301]
82. Wagner A, Aizenstein H, Venkatraman M, Fudge J, May J, Mazurkewicz L, et al. Altered reward processing in women recovered from anorexia nervosa. *Am J Psychiatry*. 2007; 164:1842–1849. [PubMed: 18056239]
83. Wagner A, Aizenstein H, Venkatraman V, Bischoff-Grethe A, Fudge J, May J, et al. Altered striatal response to reward in bulimia nervosa after recovery. *Int J Eat Disord*. 2010; 43:289–294. [PubMed: 19434606]
84. Schultz W. Neural coding of basic reward terms of animal learning theory, game theory, microeconomics and behavioural ecology. *Science*. 2004; 14:139–147.
85. Yin H, Knowlton B. The role of the basal ganglia in habit formation. *Nature Neuroscience Rev*. 2006; 7:464–476.
86. Montague R, Hyman S, Cohen J. Computational roles for dopamine in behavioural control. *Nature*. 2004; 431:760–767. [PubMed: 15483596]
87. Zastrow A, Kaiser S, S C, Walthe S, Herzog W, Tchanturia K, Belger A, et al. Neural correlates of impaired cognitive-behavioral flexibility in anorexia nervosa. *Am J Psychiatry*. 2009; 166:608–616. [PubMed: 19223435]
88. Lock J, Garrett A, Beenhakker J, Reiss A. Aberrant brain activation during a response inhibition task in adolescent eating disorder subtypes. *Am J Psychiatry*. 2011; 168:55–64. [PubMed: 21123315]
89. Oberndorfer T, Kaye W, Simmons A, Strigo I, Matthews S. Demand-specific alteration of medial prefrontal cortex response during an inhibition task in recovered anorexic women. *Int J Eat Disord*. 2011; 44:1–8. [PubMed: 20127942]
90. Kober H, Mende-Siedlecki P, Kross E, Weber J, Mischel W, Hart C, et al. Prefrontal-striatal pathway underlies cognitive regulation of craving. *Proc Natl Acad Sci USA*. 2010; 107:14811–14816. [PubMed: 20679212]
91. Marsh R, Horga G, Wang Z, Wang P, Klahr K, Berner L, et al. An FMRI study of self-regulatory control and conflict resolution in adolescents with bulimia nervosa. *Am J Psychiatry*. 2011; 168:1210–1220. [PubMed: 21676991]
92. Insel T, Cuthbert B, Garvey M, Heinssen R, Pine D, Quinn K, et al. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *Am J Psychiatry*. 2010; 167:748–751. [PubMed: 20595427]