Published in final edited form as:

J Am Chem Soc. 2013 April 3; 135(13): 4970–4973. doi:10.1021/ja401606e.

Pd-Catalyzed Asymmetric β -Hydride Elimination En Route to Chiral Allenes

Ian T. Crouch, Robynne K. Neff, and Doug E. Frantz*

Department of Chemistry, The University of Texas at San Antonio, San Antonio, TX 78249

Abstract

We wish to report our preliminary results on the discovery and development of a catalytic, asymmetric β -hydride elimination from vinyl Pd(II)-complexes derived from enol triflates to access chiral allenes. To achieve this, we developed a class of chiral phosphite ligands that demonstrate high enantioselectivity, allow access of either allene enantiomer, and are readily synthesized. The methodology is demonstrated on over 20 substrates and application to the formal asymmetric total synthesis of the natural product, (+)-epibatidine, is also provided.

Organic chemists are on a perpetual path of discovery to identify new methods for asymmetric synthesis. Historically, much of this effort has focused on the generation and control of point chirality (i.e. stereogenic centers) and as a result, approaches to engender axial chirality have not been as forthcoming. A perfect example of this unbalanced dichotomy is in the synthesis of chiral allenes. While allenes continue to be exploited as substrates in a multitude of synthetic methodologies,¹ as chiral ligands in asymmetric catalysis² and exist as structural motifs in ~150 natural products (usually in enantioenriched form),³ robust methods to synthesize chiral allenes directly from prochiral substrates with high stereochemical fidelity are few and far between.^{4–9}

Our own interest in this area was sparked by our ability to exploit stereodefined enol triflates as pluripotent substrates in organic synthesis.^{10,11} In previous work, we reported the catalytic elimination/isomerization of enol triflates to 1,3-dienes using Pd(P(*t*Bu)₃)₂ as the ideal catalyst.¹² A viable mechanism for this reaction involves an initial β -hydride elimination from a cationic vinyl Pd(II)-complex generating the corresponding allene intermediate that subsequently isomerizes to the 1,3-diene product (Scheme 1). We believed that it might be possible to interrupt this catalytic cycle under attenuated reaction conditions to allow the allene to be isolated as the terminal product in these reactions.

More importantly, we envisioned that this pathway might lead to a viable approach to access enantioenriched allenes using chiral ligands associated with the Pd-catalyst via an asymmetric β -hydride elimination; a reaction that has no precedence in the literature as far as we are aware.¹³

Our initial experiments began with (*E*)-enol triflate **1** as our model substrate and Pd₂dba₃ as our preferred source of palladium (eq. 1). A preliminary survey of ~60 commercially available chiral phosphorous-based ligands to promote the asymmetric β -hydride elimination

Corresponding Author, doug.frantz@utsa.edu.

ASSOCIATED CONTENT

Supporting Information. Detailed experimental procedures and ligand screen, ¹H and ¹³C NMR spectra and chiral HPLC traces. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors claim no competing financial interest.

en route to chiral allene **2** provided only partial success with respect to enantioselectivity. Fortuitously, however, we were able to determine that several of these ligands were in fact able to induce moderate enantioselectivity initially but competing racemization over the course of the reaction was responsible for the observed low enantioselectivity at higher conversions.¹⁴ Upon closer inspection, a general trend was noticed that the rate of racemization was slower with phosphite-based ligands than phosphine-based ligands. Thus, we turned our attention to the design of our own chiral phosphite ligands. After an extensive screen of ~50 novel phosphite ligands, we found ligands based on the BINOL scaffold in combination with menthol-derived alcohols proved most effective. A highly condensed summary of our optimization studies with a selected set of these ligands is provided in Table 1.¹⁵

(1)

With two chiral phosphite ligands capable of providing either allene enantiomer in high excess (Table 1, entries 5 and 6), we turned our attention to the robust synthesis of these ligands that would facilitate their utility in asymmetric β -hydride eliminations. Their respective syntheses are outlined in Scheme 2. Both routes developed are easily scalable providing multi-gram quantities of **4** and **5** in 80% and 23% overall yield (respectively) from commercially available starting materials.

The applicability of this methodology to a broad range of disubstituted (*E*)-enol triflates is shown in Table 2. The preferred reaction conditions include using isopropyl acetate as the solvent (0.2 M), 5 mol% Pd_2dba_3 , 10 mol% chiral ligand, Hünig's base (4 eq), and performing the reaction at room temperature. Under these conditions, the catalytic system tolerates multiple functionalities providing good yields of the corresponding chiral allenes in moderate to excellent enantioselectivities. Absolute configuration assignments in Table 2 are based on the observed optical rotation and application of the rules developed by both Brewster and Lowe for chiral allenes.¹⁶

During the course of these studies we did observe differences with respect to substrate specificity between ligands **4** and **5** that we wish to highlight. In general, higher enantioselectivities are obtained using ligand **4** with enol triflates that contain bulky carbonyl groups (compare **6** and **7** for both ligands). In contrast, higher enantioselectivities are observed with ligand **5** with less hindered carbonyl groups (Table 1, entries 5 vs. 6). We cannot yet formulate a stereochemical model for these reactions that would rationalize the apparent substrate preferences between the two ligands. Nonetheless, we believe the identification of two highly selective, complementary ligands is an advantage that broadens the scope of viable substrates at this early stage of development.

There are several limitations of this methodology that we would like to disclose (Scheme 3). First, the corresponding (*Z*)-enol triflates do not participate in this reaction. Based on our previous work, we believe this lack of reactivity can be attributed to the internal chelation of the carbonyl group after stereospecific oxidative addition that attenuates the agostic interaction with the β -hydrogens in these cationic Pd(II) -complexes.¹² Even the addition of Lewis acids (i.e., TMSOTf) did not recover reactivity of (*Z*)-enol triflates as had been

observed previously.¹² Second, fully substituted (*E*)-enol triflates such as **28** that would provide trisubstituted chiral allenes are poor substrates for this methodology. In this particular example, we were only able to observe ~30% conversion to the chiral allene **29** in 20% ee. While we view these as significant limitations that warrant disclosure, we are confident that further optimization will ultimately provide a solution for these substrates.

The application of this methodology to the formal asymmetric total synthesis of (+)-epibatidine is shown in Scheme 4.¹⁷ Originally discovered in 1976 from the skin extract of the Ecaudorian poison frog, *Epipedobates tricolor*, interest in epibatidine did not peak until its structure was reported in 1992 along with its exquisite biological activity as a powerful analgesic (~200× over morphine).¹⁸ It is now known that epibatidine is a highly potent agonist at multiple neuronal nicotinic acetylcholine receptor (nAChR) subtypes.¹⁹ However, the lack of receptor specificity and dose-limiting toxicity have led to significant efforts to identify analogs that overcome these limitations that still continue today.²⁰

Our approach to (+)-epibatidine began with enol triflate (*E*)-**30** readily available from the corresponding acetoacetate derivative.¹⁰ Catalytic, asymmetric β-hydride elimination using ligand **5** gave (+)-(*S*)-**19** in 75% yield and 96% ee on a 3 mmol scale. Lewis acid catalyzed Diels-Alder reaction with excess *N*-Boc pyr-ole gave the corresponding *endo*-product **31** in 57% yield.²¹ Low-pressure chemoselective hydrogenation followed by ozonolysis yielded β-keto ester **32** in 65% overall yield. Subsequent decarboxylation followed by reprotection gave ketone **33** in 70% yield and 89% ee. Previously, Merck has shown the conversion of racemic **33** to both enantiomers of epibatidine that completes the formal asymmetric total synthesis.²²

In summary, we have discovered a Pd-catalyzed, asymmetric β-hydride elimination from enol triflates to access chiral allenes using two optimized phosphite ligands that provide complementary substrate scope. Extension of this methodology to the formal total synthesis of (+)-epibatidine is also described. system to additional substrates is currently underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We like to thank the Welch Foundation (AX-1735) for supporting this work. We also thank Dr. Stephen Bach and Dr. David Black for high-resolution mass spectral data (supported by a grant from the National Institute on Minority Health and Health Disparities (G12MD007591) from the National Institutes of Health).

REFERENCES

1. For a recent review, see: Yu S, Ma S. *Angew. Chem. Int. Ed.* 2012; 51:3074.
2. (a) Cai F, Pu X, Qi X, Lynch V, Radha A, Ready JM. *J. Am. Chem. Soc.* 2011; 133:18066. [PubMed: 21972824] (b) Pu X, Qi X, Ready JM. *J. Am. Chem. Soc.* 2009; 131:10364. [PubMed: 19722613] (c) Löhr S, Averbek J, Schürmann M, Krause N. *Eur. J. Inorg. Chem.* 2008:552. (d) Sato I, Matsueda Y, Kadowaki K, Yonekubo S, Shibata T, Soai K. *Helv. Chim. Acta.* 2002; 85:3383.
3. Hoffmann-Röder A, Krause N. *Angew. Chem. Int. Ed.* 2004; 43:1196.
4. For reviews on the synthesis of allenes, see: Yu S, Ma S. *Chem. Commun.* 2011; 47:5384. Brummond KM, DeForest JE. *Synthesis.* 2007:795.
5. (a) Ye J, Fan W, Shenming M. *Chem. Eur. J.* 2013; 19:716. [PubMed: 23143854] (b) Periasamy M, Sanjeevakumar N, Dalai M, Gurubrahmam R, Reddy PO. *Org. Lett.* 2012; 14:2932. [PubMed: 22587711] (c) Ye J, Li S, Fan W, Kuang J, Liu J, Liu Y, Miao B, Wan B, Wang Y, Xie X, Yu Q,

- Yuan W, Ma S. *Org. Lett.* 2012; 14:1346. [PubMed: 22356094] (d) Li C-Y, Wang X-B, Sun X-L, Tang Y, Zheng J-C, Xu Z-H, Zhou Y-G, Dai L-X. *J. Am. Chem. Soc.* 2007; 129:1494. [PubMed: 17283983]
6. (a) Ogasawara M, Ikeda H, Nagano T, Hayashi T. *J. Am. Chem. Soc.* 2001; 123:2089. [PubMed: 11456845] (b) Han JW, Tokunaga N, Hayashi T. *J. Am. Chem. Soc.* 2001; 123:12915. [PubMed: 11749556]
7. Liu H, Leow D, Huang K-W, Tan C-H. *J. Am. Chem. Soc.* 2009; 131:7212. [PubMed: 19422238]
8. Li Z, Boyarskikh V, Hansen JH, Autschbach J, Musaev DG, Davies HML. *J. Am. Chem. Soc.* 2012; 134:15497. [PubMed: 22924394]
9. (a) Wan B, Ma S. *Angew. Chem. Int. Ed.* 2013; 52:441. (b) Trost BM, Fandrick DR, Dinh DC. *J. Am. Chem. Soc.* 2005; 127:14186. [PubMed: 16218604]
10. Babinski D, Soltani O, Frantz DE. *Org. Lett.* 2008; 10:2901. [PubMed: 18540630]
11. (a) Babinski DJ, Bao X, El Arba M, Chen B, Hrovat DA, Borden WT, Frantz DE. *J. Am. Chem. Soc.* 2012:16139. [PubMed: 22963196] (b) Babinski DJ, Aguilar HA, Still R, Frantz DE. *J. Org. Chem.* 2011; 76:5915. [PubMed: 21682322]
12. Crouch IT, Dreier T, Frantz DE. *Angew. Chem. Int. Ed.* 2011; 50:6128–6132.
13. The recent work by Fu and co-workers on the dehydrohalogenation of alkyl halides via Pd-catalyzed β -hydride elimination is worthy to note here. Bissember AC, Levina A, Fu GC. *J. Am. Chem. Soc.* 2012; 134:14232. [PubMed: 22905894]
14. Racemization of allenes in the presence of Pd(II)-salts is known and can also be useful for dynamic kinetic resolution. Deska J, del Pozo Ochoa C, Bäckvall JE. *Chem. Eur. J.* 2010; 16:4447. [PubMed: 20309981] Horváth A, Bäckvall JE. *Chem. Commun.* 2004; 40:964.
15. A detailed summary of our screening efforts can be found in the Supporting Information.
16. (a) Lowe G. *Chem. Commun.* 1965:411. (b) Brewster JH. *Top. Stereochem.* 1967; 2:1.
17. Recent asymmetric syntheses of epibatidine: Jensen KJ, Weise CF, Dickmeiss G, Morana F, David RL, Jørgensen KA. *Chem. Eur. J.* 2012; 18:11913. [PubMed: 22907652] (b) Aggarwal VK, Olofsson B. *Angew Chem. Int. Ed.* 2005; 44:5516. (c) Lee C-LK, Loh T-P. *Org. Lett.* 2005; 7:2965. [PubMed: 15987181] (d) Evans DA, Scheidt KA, Downey W. *Org. Lett.* 2001; 3:3009. [PubMed: 11554830]
18. Spande TF, Garraffo HM, Edwards MW, Yeh HJC, Pannell L, Daly JW. *J. Am. Chem. Soc.* 1992; 114:3475.
19. Holladay MW, Wasicak JT, Lin N-H, He Y, Ryther KB, Bannon AW, Buckley MJ, Kim DJB, Decker MW, Anderson DJ, Campbell JE, Kuntzweiler TA, Donnelly-Roberts DL, Piattoni-Kaplan M, Briggs CA, Williams M, Arneric SP. *J. Med. Chem.* 1998; 41:407. [PubMed: 9484491]
20. Liu J, Eaton B, Caldarone B, Lukas RJ, Kozikowski AP. *J. Med. Chem.* 2010; 53:6973. [PubMed: 20822184]
21. Kimura H, Fujiwara T, Katoh T, Nishide K, Kajimoto T, Node M. *Chem. Pharm. Bull.* 2006; 54:399. [PubMed: 16508202]
22. Fletcher SR, Baker R, Chambers MS, Herbert RH, Hobbs SC, Thomas SR, Verrier HM, Watt AP, Ball RG. *J. Org. Chem.* 1994; 59:1771.

Scheme 1.
Pd-Catalyzed Pathways to 1,3-Dienes or Allenes from Enol Triflates.

Scheme 2.
Synthesis of Ligands 4 and 5.

Scheme 3.
Known Limitations of the Methodology

Scheme 4.
Formal Total Synthesis of (+)-Epibatidine

Table 1

Selected Optimization Data for the Pd-Catalyzed Asymmetric β -Hydride Elimination of Enol Triflates to Chiral Allenes.

entry	R ¹	R ²	mol % Pd ₂ dba ₃ ligand	ee of 2 ^a	yield of 2 ^a
1	H		6% 12%	25%	69% ^{b,d,f}
2	Ph	"	3% 6%	54%	69% ^{b,d,f}
3		"	3% 6%	63%	74% ^{b,d,f}
4		"	3% 6%	65% 76%	31% ^{b,d,f} 59% ^{c,d,f}
5	"		5% 10%	85% 80%	^{c,e,g}
6	"		5% 10%	93% h	81% ^{c,e,g}

^a Enantiomeric excesses (ee) and yields reported were determined by quantitative chiral HPLC analysis.

^b Reaction performed in THF.

^c Reaction performed in *i*-PrOAc.

^d Reaction performed at 35 °C

^e Reaction performed at r.t.

^f 2 equiv of base.

^g 4 equiv of base.

^h Reaction performed using ligand based on (*S*)-BINOL instead of (*R*)-BINOL yielding the opposite enantiomer of **2**.

Table 2

Preliminary Scope of the Pd-Catalyzed, Asymmetric β -Hydride Elimination En Route to Chiral Allenes.^{a,b} ^aReported are isolated yields with enantiomeric excess (ee) or diastereomeric excess (de) determined by chiral HPLC analysis.^bAbsolute configuration assignments are based on the observed optical rotation and application of the rules developed by both Brewster and Lowe for chiral allenenes (see ref. 16).^cReaction performed with the enantiomer of ligand **4** provided (+)-**6** in equal yield and ee (87% yield, 95% ee).