Published in final edited form as:

Biochemistry. 2011 October 25; 50(42): 9114–9124. doi:10.1021/bi201260h.

Structural Insights into Complete Metal Ion Coordination from Ternary Complexes of B Family RB69 DNA Polymerase

Shuangluo Xia^{†,1}, Mina Wang^{†,1}, Gregor Blaha¹, William H. Konigsberg¹, and Jimin Wang^{1,*}

¹Department of Molecular Biophysics and Biochemistry, Yale University, 266 Whitney Avenue, New Haven, CT 06520-8114, USA

Abstract

We have captured a pre-insertion ternary complex of RB69 DNA polymerase (RB69pol) containing the 3' hydroxyl group at the terminus of an extendable primer ($p\text{O}3'$) and a non-hydrolyzable 2'-deoxyuridine 5'- α,β -substituted triphosphate, dUpXpp, where X is either NH or CH₂, opposite a complementary templating dA nucleotide residue. Here we report four structures of these complexes formed by three different RB69pol variants with catalytically inert Ca²⁺ and other four structures with catalytically competent Mn²⁺ or Mg²⁺. These structures provide new insights into why the complete divalent metal-ion coordination complexes at the A and B sites are required for nucleotidyl transfer. They show that the metal ion in the A site brings $p\text{O}3'$ close to the α -phosphorus atom (P α) of the incoming dNTP to enable phosphodiester bond formation through simultaneous coordination of both $p\text{O}3'$ and the non-bridging Sp oxygen of the dNTP's α -phosphate. The coordination bond length of metal-ion A as well as its ionic radius determines how close $p\text{O}3'$ can approach P α . These variables are expected to affect the rate of bond formation. The metal ion in the B site brings the pyrophosphate product close enough to P α enabling pyrophosphorolysis as well as assisting in the departure of the pyrophosphate. In these dUpXpp-containing complexes, $p\text{O}3'$ occupies the vertex of a distorted metalion A coordination octahedron. When $p\text{O}3'$ is placed into the vertex of a non-distorted, idealized metalion A octahedron, it is within bond formation distance to P α . This geometric relationship appears to be conserved among DNA polymerases of known structure.

DNA polymerases (pols) catalyze DNA synthesis using a two metal ion-mechanism (1–3). Metalion A has been proposed to activate the attacking 3'-OH of the primer-terminus ($p\text{O}3'$) by bringing it closer to the α -phosphorus atom (P α) of the incoming dNTP to enable phosphodiester bond formation. Metal-ion B accompanies the triphosphate moiety of the incoming dNTP, stabilizes the pentavalent transition state (TS), and assists in the departure of the pyrophosphate group. With pols, proper positioning of the two catalytic divalent metal ions leads to a putative triangular bipyramidal TS, attainable with substantially-lowered activation energy, which is inaccessible in the absence of a pol. Even though all pols provide conserved ligands, namely two carboxylates and a backbone carbonyl oxygen, for the two metal ions at their catalytic centers, the extent of TS stabilization varies from one class of pols to another due to additional interactions with other residues. For example, the stabilization of both the TS and the closed pre-insertion ternary complex of replicative RB69 DNA polymerase (RB69pol) is provided not only by metal ions and complementary base-pairing, but also by the conserved residues R482, K560, and N564 in the Fingers domain (4,

*Corresponding Author: Jimin Wang, Phone: (203)-432-5737; Fax: (203)-432-3282. jimin.wang@yale.edu.

[†]S. X. and M. W. made equal contributions to this work.

SUPPORTING INFORMATION AVAILABLE

Supporting information includes additional 9 figures available on the ACS website. This material is available free of charge via the Internet at <http://pubs.acs.org>.

5). This results in high incorporation efficiency for complementary dNMPs. Absent these interactions, i.e. when non-complementary incoming dNTPs are present, the closed ternary complex becomes unstable so that the Fingers reopen rapidly releasing non-complementary dNTPs (6).

Since transfer of two protons is associated with incorporation of complementary dNMPs by all pols that have been studied, this has ruled out that the large conformational change of the Fingers closing is rate-limiting (7, 8). Instead, the chemical step, or a step immediately before or after it, might be rate-limiting. Thus, it follows that TS stabilization determines dNMP incorporation rates. To obtain a more complete understanding of the pol-catalyzed nucleotidyl transfer reaction, it is essential to have structures of pre-insertion ternary complexes that approximate TS as closely as possible. This requirement stems from the lack of information about the optimal geometry of the TS as there has not been any pol TS-inhibitor identified to date. Using the α,β -substituted, non-hydrolyzable complementary dUpNpp, Wilson and colleagues have captured a pre-insertion ternary complex of pol β and at the first time visualized $p\text{O}3'$, a critical element necessary for the two metal-ion mechanism to function in nucleotidyl transfer (9). From their studies, followed by subsequent theoretical calculations, a model for putative TS involving three coordination polygons was proposed in attempt to explain the observed catalytic efficiency of dNMP incorporation as well as the base selectivity exhibited by pol β (10). However, whether this TS model from pol X family is applicable to other families of pols remains to be established. Since pol β is a repair enzyme and differs from high-fidelity pols including RB69pol, we require similar types of data for high-fidelity pols to establish the general principles that govern nucleotidyl transfer perhaps by all pols for correct incorporation of dNMPs. These general principles are critical for base selectivity because there is a strong correlation between the degree of base selectivity and the incorporation efficiency of complementary dNMPs among all naturally-occurring pols (11), and because there must be a shared, underlying structural basis that can account for these two properties.

We have previously determined the structure of RB69pol ternary complexes with a 2',3'-dideoxy-terminated primer/template (ddP/T) (4, 5, 12–16). In fact, nearly all structures of the pol ternary complexes were determined with a ddP/T with an exception of pol β (9). Although a $p\text{O}3'$ can be modeled into a P/T computationally, its precise functional role in metal-ion coordination and in TS stabilization cannot be determined this way. We also used the catalytically inert Ca^{2+} in place of catalytically competent Mg^{2+} or Mn^{2+} for crystallization so that the precise geometry of catalytically relevant coordination polygons could not be ascertained. With an extendable P/T and a complementary dNTP, truce contamination of Mn^{2+} or Mg^{2+} has always been a serious issue for capturing the pol preinsertion ternary complex, because the presence of these metal ions leads to unintended primer-extension during the time required for crystallization. This introduces heterogeneity and prevents crystallization. By relying on ddP/Ts, we were unable to establish the precise functional roles of $p\text{O}3'$ in the metal-ion coordination geometry within the ternary complex. Without including Mn^{2+} or Mg^{2+} in the ternary complex instead of Ca^{2+} for comparison, we could not explain why Ca^{2+} is catalytically inert.

Here we report structural studies on a high-fidelity B family replicative pol (RB69pol) and several of its variants using α,β -substituted non-hydrolyzable, complementary dUpXpp analogues of dUTP to capture the pre-insertion closed ternary complex with an extendable dP/T that has a $p\text{O}3'$. By comparing our results with those obtained with pol β , we have identified specific structural features of the catalytic metal-ion coordination geometry, namely three coordination polygons, that are common to all pols of known structure and suggest that the precise geometry of these polygons governs the efficiency of the nucleotidyl transfer reaction. We describe here a total of nine high-resolution structures of ternary

complexes formed by three RB69pol variants, which include the wild type (wt), a triple mutant (tm), L561A/S565G/Y567A, and a quadruple mutant (qm), L415A/L561A/S565G/Y567A. We have designed these mutants as well as other substitutions in the nascent-basepairing binding pockets for a series of crystallographic studies on RB69pol because pre-insertion ternary complexes formed with these variants were readily crystallized (4, 5, 12–15). The triple mutant as well as other nascent-basepairing-binding pocket mutants exhibits the same kinetic parameters for incorporation of complementary dNMPs as the wt (16, 17). However, substitutions of amino acid residues that directly interact with the triphosphate moiety of incoming dNTP in the Fingers domain have reduced incorporation efficient for complementary dNMPs (18). Moreover, the structures of the wt, tm, and qm were isomorphous to one another, and their overall structure was completely superimposable except for the restricted region at the sites of amino acid substitutions. Since this study focuses on the coordination geometry for incorporation of complementary dNMPs, we have properly selected three variants of RB69pol, wt, tm, and qm. Four of the structures presented here were determined in the presence of Ca^{2+} (Table 1) and five in the presence of Mn^{2+} or Mg^{2+} . Among these structures, we included one structure of ternary complex containing Mg^{2+} /dCTP opposite dG with a ddP/T as a control for the tm variant (Table 2). In the RB69pol tm, all three amino acid replacements are far from the pol active site (4, 5). In the qm, the fourth substitution, L415A, is near the pol active site, but we show that it has only a minimal effect on the metal-ion coordination octahedrons. Thus, the geometric relationship derived from these structures, namely the relationship between the putative triangular, bipyramidal TS polygon connected by two metal-ion coordination octahedrons, is found to be present in all the RB69pol variants. The relative position of these three polygons is defined by two shared carboxylates (D411 and D623 in RB69pol) and a shared non-bridging Sp oxygen (O2 α) of the incoming dNTP (footnote: the prochirality definition of dNTP is inverted in dUpXpp when X is NH or CH₂. To avoid confusion, we prefer the common name O2 α for this oxygen of dNpXpp analogues).

EXPERIMENTAL PROCEDURES

Chemicals

All chemicals were of the highest quality available; dUpNpp and dUpCpp were purchased from Jena Bioscience (Jena, Germany).

Enzymes

Wt, the tm, and the qm RB69pols in an exonuclease-deficient (exo^-) background (D222A and D327A) were over-expressed in *Escherichia coli*, purified, and stored as previously described (5, 16, 17).

P/T DNA Substrates

All oligonucleotides were synthesized by W. M. Keck Foundation Biotechnology Resource Laboratory (Yale University). The sequence of the first template strand used for cocrystallization of the eight dUpXpp-containing complexes was 5'-TCAAGTAAGCAGTCCGCG-3' and the sequence of the second template for the ninth dCTP-containing ternary complex differed in the fourth nucleotide where A was replaced with G to provide a complementary templating base for the incoming dCTP. The sequences of the primer strands for the dP/T and ddP/T were 5'-GCGGACTGCTTAC-3' and 5'-GCGGACTGCTTAddC-3', respectively, where ddC is a 2',3'-dideoxyterminal C. The complementary oligonucleotides were annealed by heating to 95°C and gradually cooling to 25°C to form the P/T or ddP/dT duplexes.

Crystallization of the RB69pol Ternary Complexes with Catalytically Inactive Ca²⁺

The ternary complexes were prepared by mixing an equimolar ratio of wt, tm or qm RB69pol (exo⁻) with a freshly annealed dP/T or ddP/T and a complementary incoming dUpNpp, dUpCpp or dCTP. After the solution had been mixed, the [ternary complex] was about 12 mg/ml in the excess of dUpXpp or dCTP with [dUpXpp] or [dCTP] at 0.5 mM. Crystals of the tm ternary complex were grown under oil in a microbatch procedure by mixing equal volumes of the ternary complex with a solution containing 100 mM sodium cacodylate buffer (pH 6.5), 150 mM CaCl₂, and 1–3% (w/v) polyethylene glycol 350 monomethyl ether (PEG 350 MME). Crystals of the wt ternary complex were grown under the same conditions except using 5% (w/v) PEG 350 MME. Rectangle-shaped and some cubic-like shaped crystals typically grew in 2 days at 20°C to a size of 0.1 mm X 0.1 mm X 0.1 mm. Crystals were stabilized and cryoprotected, by transferring them first to a stabilization solution containing 100 mM sodium cacodylate buffer (pH 6.5), 20% (w/v) and then to a solution of 30% PEG 350 MME (for cryoprotection) with 100 mM CaCl₂ before freezing them in liquid nitrogen.

Preparation of Crystals of the RB69pol Ternary Complexes with Catalytically Active Mn²⁺ or Mg²⁺

The crystals of the Mn²⁺ or Mg²⁺-containing complexes were prepared using soaking-replacement methods. The crystals of the Ca²⁺-containing ternary complexes were stabilized in the same initial solution of 20% (w/v) PEG 350 MME containing 100 mM MnCl₂ or MgCl₂, to replace CaCl₂, for 30 min changing the buffer twice. They were then transferred to a solution made up of 30% (w/v) PEG 350 MME with 100 mM MnCl₂ or MgCl₂ for freezing. Complete replacement was verified using anomalous difference Fourier maps for Mn²⁺ or from the coordination geometry for Mg²⁺.

X-ray Diffraction Data Collection, Structure Determination, and Refinement

X-ray diffraction data were collected at various wavelengths at 110 K, at beamline 24ID-E, the Northeast Collaborative Access Team (NECAT), Advanced Photon Source (APS), Argonne National Laboratory (ANL), Argonne, IL as well as at beamline X29A, National Synchrotron Light Source (NSLS), Brookhaven National Laboratories (BNL), NY. The data were processed as summarized in Tables 1 & 2 using the 8 HKL2000 program suite (19). The structure was determined by the automated molecular replacement method Phaser as implemented in CCP4 (20, 21), starting with the wt RB69pol structure of the ternary complex [PDB entry 3NCI (5)]. The P/T DNA duplex and the incoming dUpXpp were built into residual electron density maps using COOT (22). The structure was refined as summarized in Tables 1 & 2 using Refmac5 (23). All figures were made using Ribbons (24). All structural comparison was made using superposition of Cα coordinates of the entire pol structure using COOT (22) unless otherwise noted.

Geometric Relationship between the pO3'-Pa Distance and the Ionic Radius of Idealized Coordination Polygons

To establish the numerical relationship between the pO3'-Pa distance and the ionic radius of divalent metal ions, we have assumed that pO3' occupies at a vertex of an idealized octahedron whose coordination bond length to the metal ion is defined by the ionic radius, and that at the hypothetical TS, O5', O1α and O2α are coplanar with Pa with fixed inter-atomic distance of 1.52 Å, which is slightly longer than their typical bond distances (10). Because pO3', Pa, and O2α form a rightangle triangle, the pO3'-Pa distance is a function of the pO3'-O2α edge length, which is also the edge length of the idealized metal-ion coordination octahedron. We numerically calculated the pO3'-Pa distance as a function of

the metal ion coordination length from 2.0 to 3.0 Å, plotted it, and estimated the slope of this relationship to be 1.58.

PDB Entry

Coordinates and structure factors have been deposited in the Protein Data Bank as entries 3SQ1, 3S9H, 3SCX, and 3SPZ for Ca²⁺/dUpXpp-containing complexes (Table 1) and 3SJJ, 3SQ0, 3SI6, 3SPY, and 3SNN for Mn²⁺ or Mg²⁺/dUpXpp or dCTP-containing complexes (Table 2). The entry 3SQ1 was with wt RB69pol, the entries 3SPZ and 3SPY were with the qm, and other all entries were with the tm. The entry 3SNN was with dCTP, the entries 3SPZ and 3SPY were with dUpCpp, and all other entries with dUpNpp. The entries 3SJJ and 3SQ0 were with Mn²⁺, and the entries 3SI6, 3SPY and 3SNN were with Mg²⁺.

Measurement of K_d for dUpNpp in Competition with dTTP and Analysis of Kinetic Data

Competition experiments were performed under single-turnover conditions at 23°C using a KenTek model RQF-3 instrument with a buffer solution of 66 mM Tris-HCL (pH 7.4). Briefly, the enzyme and the P/T DNA duplex from one syringe were rapidly mixed with Mg²⁺ (or Mn²⁺), dTTP and various [dUpNpp] from the other syringe for times ranging from 0 s to 0.06 s. The final concentrations after mixing were as follows: 1.2 μM enzyme, 83 nM P/T, 25 μM dTTP, and 10 mM Mg²⁺ (or Mn²⁺). Reactions were quenched with 0.5 M EDTA (pH 8.0). Substrates and products were separated by 19:1% (w/v) PAGE containing 8 M urea. Imaging was performed using a phosphorimager, Molecular Dynamics Storm 860, and quantified using ImageQuaNT software.

The amount of product formed versus time was plotted for each [dUpNpp] and fitted by nonlinear regression to the general form of Eq. 1 to obtain observed rates for product formation.

$$Y = \sum_i^n A_i e^{-k_i t} + C \quad \text{Eq.1}$$

where Y is a value corresponding to the concentration of the DNA product, C is the offset, A_i is the observed amplitude of product formation, and k_{obs} or k_i (with $i=1$ for single-phase turnover) is the observed rate constant. The $K_{d,app}^{dUpNpp}$ was obtained by plotting k_{obs} versus [dUpNpp] using Eq. 2:

$$k_{obs} = A \frac{[dUpNpp]}{K_{d,app}^{dUpNpp} + [dUpNpp]} + C \quad \text{Eq.2}$$

where A is the overall change in the observed rate over the concentration series, and C is the Y -axis intercept. The ground-state association constant for dUpNpp was obtained according to the Eq. 3:

$$K_d^{dUpNpp} = K_{d,app}^{dUpNpp} / (1 + \frac{[dTTP]}{K_d^{dTTP}}) \quad \text{Eq.3}$$

where [dTTP] is the concentration of dTTP used in the assay, and K_d^{dTTP} is the ground-state binding constant for dTTP.

RESULTS AND DISCUSSION

Overview of the Structures and Metal Ion-Coordination of RB69pol Ternary Complexes

We have determined nine high-resolution isomorphous crystal structures of pre-insertion ternary complexes of the wt RB69pol and two RB69pol mutant variants (tm, and qm). Four of the structures contained Ca^{2+} (Table 1) and five contained Mn^{2+} or Mg^{2+} (Table 2). Among these complexes, six included dUpNpp with each of the three RB69pol variants, two had dUpCpp with qm, and one had dCTP with the tm. All incoming dNTPs (dCTP or dUpXpp) were complementary to the respective templating bases. Among these complexes, six were formed with tm, two with qm, and one with wt. The overall structures of these complexes were nearly identical albeit with subtle variations in the hydration networks near the amino acid substitution sites. Of the nine structures, the wt RB69pol ternary complex was determined at 1.83-Å resolution (the highest resolution) with free R-factor of 21.2% (Table 1). The resolution of all of these structures described here varied from 2.43 to 1.83 Å with free R-factors ranging from 21.2% to 26.5%. With the exception of the two lowest-resolution structures at about 2.4 Å (3SPZ and 3SJJ), all others had sufficiently high-resolution to accurately visualize the network of ordered water molecules, particularly those near the pol active site and serving as metal ion coordination ligands.

The triphosphate tail of the incoming dUpXpp or dCTP in eight of the nine complexes was coordinated to the B metal ion in the classic α,β,γ -tridentate state with the exception of the ninth Mg^{2+} /dUpNpp complex (3SI6 in Table 2). In all of these eight complexes, both A and B metal ions were clearly visible. Metal ion A is coordinated by the carboxylate side chains of D411, D623, the 3'-hydroxyl group of the priming nucleotide residue ($p\text{O}3'$), and O2a (i.e., a non-bridging Sp oxygen of the incoming dNTP's $\text{P}\alpha$) in addition to ordered water molecules. Metal ion B is also coordinated by D411, D623, and by the backbone carbonyl oxygen of L412 as well as by three oxygen ligands from the triphosphate tail of the incoming dNTP. The Mg^{2+} /dUpNpp-containing tm ternary complex however was in a β,γ -bidentate coordination state where Mg^{2+} was present only in the B site (Fig. S1). This coordination state was also observed when another structure of the same complex was determined at a lower resolution (2.2-Å) providing further confirmation for the β,γ -bidentate complex. The occurrence of this β,γ coordination state prompted us to extend our studies using the α,β CH₂-substituted nonhydrolyzable dUpCpp in an attempt to obtain an α,β,γ -tridentate Mg^{2+} coordination state which we believed would be on the reaction pathway and closer to the TS (Fig. S2).

Within this set of crystal structures (5, 12–15), we found a number of commonly occurring variations, including different conformations of an exonuclease loop and of the C-terminal tail of RB69pol as well as different structures of the 5' template strand overhang associated with slightly different packing of the complexes (Fig. S3). None of these variations have any effects on the coordination geometry since all three amino acid substitutions in tm are far from the metal ion coordination site although they have altered the structure of ordered molecules near the substitution sites (Fig. S4, Fig. S5). The L415A substitution in qm had a minimal effect on metal ion coordination geometry because it allowed an insertion of an ordered water molecule into a hydrophobic core near the catalytic carboxylate (W-101 in Fig. S6). This water molecule forms a hydrogen bond with the backbone carboxyl of T622 and causes a subtle repositioning of the D623 backbone in the Ca^{2+} -containing complex, but not in the Mg^{2+} -containing complex.

Competitive Inhibition of dUpNpp against dTMP Incorporation

Competitive inhibition assays for the incorporation of complementary dTMP with both wt and tm RB69pols by dUpNpp have shown that the ground-state binding affinities for

dUpNpp (Table 3) are with the same range as complementary dNTPs when Mg^{2+} is the divalent metal ion (17). RB69pol and many other pols can use both Mg^{2+} and Mn^{2+} for incorporation of complementary dNMPs with similar catalytic efficiency but often with increased incorporation efficiency of non-complementary dNMPs by Mn^{2+} relative to Mg^{2+} (25–29). The observation of the same α,β,γ -tridentate coordination of dUpNpp (see below but with one exception, 3SI6) as that of dCTP of tm RB69pol (3SNN) and wt RB69pol (3NCI, 5) is consistent with the nature of competitive inhibition of dUpNpp for dTTP. RB69pol and many other pols cannot use Ca^{2+} for incorporation of any dNMPs (29).

Ca^{2+} Coordination in dUpNpp Ternary Complexes

We have determined four structures of Ca^{2+} /dUpXpp ternary complexes formed by wt, tm and qm variants of RB69pol with a dP/T (Table 1). A common feature of these complexes is that Ca^{2+} prefers to have more than 6 ligands due to its large ionic radius (Fig. 1). This preference is not influenced by amino acid substitutions in tm or qm. The coordination polygons and pol active sites are completely superimposable between wt and tm Ca^{2+} /dUpNpp ternary complexes (Fig. 1B). The coordination preference for Ca^{2+} is also unaffected by the type of complementary incoming dNTP (dUpXpp or dCTP) because the Ca^{2+} /dCTP and the Ca^{2+} /dUpNpp wt ternary complexes are also completely superimposable (Fig. 1F).

There are typically 8 ligands for the Ca^{2+} bound in the B site (CaB) in the complexes containing a ddP/T. These ligands include O δ 1 of D411, O δ 1 of D623, the backbone carbonyl oxygen atom of L412, and three oxygen ligands from the triphosphate tail of the incoming dNTPs as found for MnB and MgB. Two additional ligands are the O δ 2 of D411 and W-4, both of which are shared by CaB and the Ca^{2+} bound in the A site (CaA) and are responsible for reducing the CaA-CaB distance to 3.9 Å in the ddP/T complexes (Fig. 1) relative to the corresponding CaA-CaB, MnA-MnB, or MgA-MgB distances in the dP/T complexes (see below). The average coordination bond length for CaB is 2.48 ± 0.30 Å and it is 2.62 ± 0.20 Å for CaA (Fig. 1). This coordination bond length is consistent with the ionic radius of Ca^{2+} and its coordination preferences are the same as found in structures of Ca^{2+} -containing small molecules as well as other protein structures (30, 31). CaA in the ddP/T complexes typically has two empty coordination sites that are occupied by *p*O3' and W-7 in the dP/T complexes (Fig. 1).

When the Ca^{2+} /dUpNpp tm ternary complex is formed with an extendable primer (dP/T), the structure of ordered water molecules within the Ca^{2+} coordination sphere was altered, but not the binding mode of either the P/T or dUpNpp (Fig. 1D). Due to the interaction between *p*O3' and CaA (with an inter-atomic distance of 3.3 Å), CaA is shifted further from CaB (Fig. 1C). In this coordination complex, W-4 is lost, and D411 is now reoriented so that it makes one-to-one interactions with CaA and CaB using its O δ 1 and O δ 2, respectively. The displacement of CaA towards *p*O3' increases the CaA-CaB distance to 4.1 Å but the *p*O3'-Pa distance remains at 4.0 Å. The relatively large displacement of CaA caused by the presence of *p*O3' compared to MgA suggests that Ca^{2+} binding in the A site is much weaker than Mg^{2+} . The weaker binding of Ca^{2+} also reduces its ability to pull *p*O3' towards Pa. Due to its relatively large ionic radius, Ca^{2+} can accommodate more ligands than Mn^{2+} and Mg^{2+} . However, the interactions between Ca^{2+} and an individual ligand are generally weaker than those of either Mn^{2+} or Mg^{2+} so that same ligands of the carboxylate side chains bind Ca^{2+} less tightly than Mn^{2+} or Mg^{2+} . The conclusion about the CaA-CaB and *p*O3'-Pa distances within the Ca^{2+} /dUpXpp complexes observed here can be extrapolated into the complementary incoming Ca^{2+} /dNTP complexes because the Ca^{2+} /dNTP and Ca^{2+} /dUpNpp wt ternary complexes are completely superimposable (Fig. 1D). Reexamination of our previous Ca^{2+} /dNTP wt ternary complexes (5) further supports the notion that Ca^{2+} binding to the A site is never very strong, as evident from its large temperature B-factor. B-

factor for Ca^{2+} in the A site sometimes was so large that its presence could not always be confirmed.

Mn^{2+} Coordination in dUpNpp Ternary Complexes

When Mn^{2+} was soaked into crystals of the Ca^{2+} /dUpNpp tm ternary complex, complete replacement of Ca^{2+} by Mn^{2+} was evident from the strong anomalous difference Fourier signals from Mn^{2+} as well as from the details of its coordination geometry (Fig. 2A). In this complex, Mn^{2+} is present in both A and B metal ion sites, and each Mn^{2+} has 6 ligands. Mn^{2+} in the B site is the center of a nearly perfect octahedron that has an average coordination bond length of 2.16 ± 0.06 Å (averaged from two independent Mn^{2+} -containing structures). This Mn^{2+} has an average coordination bond angle of $88 \pm 9^\circ$ with respect to any two adjacent ligands (Fig. 2). However, MnA is in the center of a distorted octahedron caused mainly by the displacement of $p\text{O}3'$ so that the $p\text{O}3'$ -MnA distance is 2.44 Å whereas the average coordination bond length for the other five ligands is 2.17 Å (Fig. 2B, 2D). This displacement is also reflected in the $p\text{O}3'$ -MnA-O2 α coordination bond angle of 104° rather than the optimal octahedral angle of 90° (Fig. 2D).

When the Mn^{2+} /dUpNpp tm ternary complex was prepared with a ddP/T (3SQ0 entry in Table 2), we were able to extend the resolution to 2.0 Å. Comparison of two Mn^{2+} /dUpNpp complexes (Fig. 2), one with a dP/T and the other with a ddP/T, shows that the absence of $p\text{O}3'$ caused only a small displacement of Mn^{2+} in the A site. Because the displacement of Mn^{2+} in the A site was accompanied by the same shifts W-5 and W-6 ordered water molecules, its coordination octahedron remained unchanged (Fig. 2F). The fact that the absence of $p\text{O}3'$ did not substantially alter the network of ordered water molecules near the pol active site, nor the number of ligands for Mn^{2+} in both A and B sites in the Mn^{2+} /dUpNpp complexes stands in sharp contrast to the situation with the Ca^{2+} /dUpNpp complexes (Fig. 1).

Comparison of the Mn^{2+} /dUpNpp and the Ca^{2+} /dUpNpp complexes shows that there is an expansion of both B and A metal-ion coordination polygons in the latter complex due to the larger ionic radius of Ca^{2+} than Mn^{2+} (Fig. 2C). In addition to the different number of coordination ligands between the two complexes as described above (Fig. 1 & Fig. 2), the expansion of the B site polygon is accomplished through small displacements of the carboxylate side chains as well as the triphosphate ligands. The₁₅ expansion of the A site polygon is accomplished through displacements of both $p\text{O}3'$ and Pa in opposite directions so that the distance between them increases. In this case, the $p\text{O}3'$ -Pa distance in the Ca^{2+} /dUpNpp complex is 4.0 Å (Fig. 1C), which is greater than the 3.6-Å distance observed in the Mn^{2+} /dUpNpp complex (Fig. 2D). It is worth noting that even in the Mn^{2+} /dUpNpp complex, the $p\text{O}3'$ -Pa distance is still too great for phosphodiester bond formation and cannot account for the differences in divalent metal-ion reactivity involving Mn^{2+} (25–28). Although these complexes do not accurately represent the catalytically active state for nucleotidyl transfer, they are reasonable starting points for predicting the nature of the TS involving the corresponding divalent metal ions, as has previously been done for pol β (9, 10).

In the Mn^{2+} /dUpNpp complex, the distance between Mn^{2+} in the A site and O2 α , or between the Mn^{2+} and O62 of both D411 and D623, are all about 2.2 Å (Fig. 2B), whereas the corresponding distances are about 2.6 Å in the Ca^{2+} /dUpNpp complex (Fig. 1C). The MnA-O2 α and MnA-O62 distances are shorter than the corresponding CaA-O2 α and CaA-O62 distances, suggesting that Mn^{2+} binding to the A site is much tighter than Ca^{2+} . Thus, the location of MnA relative to the triphosphate tail and carboxylates is more precisely defined than that of the corresponding CaA. In the Mn^{2+} /dUpNpp complex, the inclusion of $p\text{O}3'$ has resulted in a larger displacement of the priming nucleotide residue (along with the

*p*O3') than the displacement of MnA itself (Fig. 2F). By comparison, the inclusion of *p*O3' in the Ca²⁺/dUpNpp complex resulted only in displacements of CaA, but not the priming nucleotide residue (Fig. 1D). These results further reinforce the notion that Ca²⁺ binds weakly to the A site and has little influence on the location of the *p*O3' or on the priming nucleotide residue. In contrast, Mn²⁺ binds strongly to the A site and determines the location of the *p*O3' and the priming nucleotide residue should be located. As a consequence, the MnA-*p*O3' distance is 2.4 Å, but the CaA-*p*O3' distance is 3.3 Å. These metal-ion coordination bond-length differences also affect the *p*O3'-Pa distance, which is reduced by both Mn²⁺ and Mg²⁺ as discussed further above, but not by Ca²⁺. The *p*O3'-Pa distance in the Mn²⁺/dUpNpp complex is 3.6 Å, and it is 4.0 Å in the Ca²⁺/dUpNpp complex (Figs. 1 & 2).

Mg²⁺ Coordination in dUpXpp or dCTP Ternary Complexes

In the structure of the Mg²⁺/dUpNpp tm ternary complex, determined at 1.85-Å resolution (Table 2), we were surprised to find that Mg²⁺ was present only in the B site and that it formed α,β,γ -bidendate coordination complex with the triphosphate tail of dUpNpp (Fig. S1). Whether this coordination state is related to the kinetics of correct dNMP incorporation in the presence of Mg²⁺ is beyond the scope of this study. However, when dCTP or dUpCpp was part of the ternary complex with Mg²⁺, we observed the α,β,γ -tridentate complex. The structure of the Mg²⁺/dCTP tm ternary complex was determined at 2.0-Å resolution, and the structure of the Mg²⁺/dUpCpp qm ternary complex was determined at 2.15-Å resolution (Table 2).

Similar to the Mn²⁺/dUpNpp tm ternary complex, Mg²⁺ in the B site of the Mg²⁺/dUpCpp qm ternary complex is at the center of a nearly perfect octahedron and Mg²⁺ in the A site is at the center of a distorted octahedron (Fig. 3, Fig. S2). The distortion also occurs at the vertex occupied by *p*O3' as seen in the Mn²⁺ complexes. The average coordination bond length for MgB and MgA is about 2.17 and 2.20 Å (Fig. 3), respectively, in the Mg²⁺/dUpCpp-containing qm ternary complex (entry 3SPY in Table 2). These distances are similar to those in the Mn²⁺/dUpNpp complex (entry 3SJJ in Table 2). However, we observed some small variations in the average bond lengths for MgB, depending on the type of incoming dNTP: 2.22±0.19 Å for the Mg²⁺/dCTP complex (entry 3SNN), 2.06±0.15 Å for the Mg²⁺/dUpNpp complex (3SI6), and 2.17±0.23 Å for another Mg²⁺/dUpNpp complex determined at 2.2-Å resolution (which is lower than the entry 3SI6 at 1.85-Å resolution described here in Table 2). All of these distances are within the range of typical Mg²⁺ coordination bond lengths (30, 31).

There were subtle conformational differences between dUpNpp and dUpCpp in these ternary complexes. Judging from inter-atomic hydrogen-bonding distances, dUpCpp appears to have a stronger intra-molecular interaction between O3' and the β -phosphate (2.5 Å), but a weaker inter-molecular interaction between *p*O3' and O2 α of Pa (3.5 Å), than the corresponding distances of 2.8 and 3.0 Å in the dUpNpp complex, respectively (Fig. 3B). It is also likely that the triphosphate tail of dUpCpp is more rigid than dUpNpp due to the stronger intra-molecular hydrogen bond and an extra hydrogen atom in the CH₂ substituent relative the NH substituent. This may have prevented the triphosphate tail of dUpCpp from adopting the optimal conformation for coordination to metal ion B than dUpNpp, resulting in slightly larger Mg²⁺-centered coordination octahedrons than Mn²⁺-centered octahedrons in contrast to their ionic radii. In the Mg²⁺/dUpCpp complex, the *p*O3'-Pa distance and the MgA-MgB distance are 3.8 Å and 3.7 Å, respectively (Fig. 3), which are also longer than the corresponding distances of 3.6 and 3.4 Å in the Mn²⁺/dUpNpp complex (Fig. 2). In addition to the differences in rigidity between dUpCpp and dUpNpp in corresponding complexes, the fourth L415A substitution in the qm relative to the tm could have some effect on these distances. Nevertheless, the metal-ion coordination geometry that determines

the efficiency of the nucleotidyl transfer reaction, as discussed below, is nearly identical in both the $\text{Mg}^{2+}/\text{dUpCpp}$ and $\text{Mn}^{2+}/\text{dUpNpp}$ complexes (Figs. 2 & 3).

We noted that the reorientation of the D411 side chain occurred in the $\text{Ca}^{2+}/\text{dUpCpp}$ qm ternary complex (Fig. 4A), but not in the $\text{Mg}^{2+}/\text{dUpCpp}$ complex (Fig. 3). In this new orientation, O82 of D411 interacts with both CaB and CaA. As discussed above, the CaA binding is relatively weak and its location is distributed wide a wide range. The observed CaA in the $\text{Ca}^{2+}/\text{dUpCpp}$ qm ternary complex is at the extreme end of this distribution, which is evident when comparing with the Mg^{2+} ternary complex, but less obvious in pairwise comparison with other Ca^{2+} complexes (Fig. 4). Between the $\text{Ca}^{2+}/\text{dUpCpp}$ and $\text{Mg}^{2+}/\text{dUpCpp}$ qm ternary complexes, the displacement of metal ion A is as large as 1.6 Å. This may result from the added flexibility of catalytic carboxylates due to the L415A substitution in qm. However, this does not affect the Mg^{2+} coordination state in the $\text{Mg}^{2+}/\text{dUpCpp}$ complex, nor the mechanistic implications for incorporation of complementary dNMPs.

Mechanistic Implications of α,β,γ -Tridendate Mg^{2+} and Mn^{2+} Coordination Geometry for Catalysis and Its Conservation Among Various Pols

The α,β,γ -tridendate metal-ion coordination geometry observed here is nearly identical for both Mn^{2+} and Mg^{2+} (Fig. 2D, Fig. 3C). In both cases, the distorted vertex in the metal-ion A coordination octahedron is occupied by $p\text{O}3'$, whereas the metal-ion B coordination octahedron is nearly ideal (Fig. 5A). The relative position of these two octahedrons is defined by the shared O2 α (Sp oxygen ligand of dNTPs) and the two conserved catalytic carboxylates D411 and D623. For a given ionic radius of the divalent metal ion and a fixed size of the coordination octahedron, the distance between the two coordination centers is invariant (Fig. 5BC). One possible trajectory that can reduce the $p\text{O}3'$ -Pa distance is a positional readjustment of the $p\text{O}3'$ along with the priming nucleotide residue. If $p\text{O}3'$ is computationally modeled in the non-distorted vertex of an idealized octahedron as was done in pol β (10), it is within the expected distance for bond formation (Fig. 5A). This modeling suggests that metal ion A is likely to be responsible for pulling $p\text{O}3'$ towards Pa through shared coordination.

We can now propose a model for the TS for nucleotidyl transfer catalyzed by RB69pol, similar to the one for pol β (10), based on the newly defined functional roles of $p\text{O}3'$ and on the relationship between the two metal ion-centered and one Pa-centered coordination polygons. At the mid-point of the hypothetical Pa inversion during nucleotidyl transfer in this TS model, three equatorial oxygen atoms on Pa (namely, O1 α , O2 α , and O5') become coplanar, and the two axial oxygen atoms maintain the corresponding distance to the Pa phosphorus center for spontaneous bond formation between $p\text{O}3'$ and Pa, and bond breakage between O3 α and Pa (Fig. 5A). To approach the TS, metal ion A should reduce the $p\text{O}3'$ -Pa distance. After computer modeling to place the $p\text{O}3'$ at the vertex of an ideal metal-ion A-centered coordination octahedron with Mg^{2+} or Mn^{2+} , the $p\text{O}3'$ -Pa distance will be reduced to a value consistent with spontaneous bond formation (10). With the two ideal metal ion-centered coordination octahedrons and the pentavalent Pa-centered TS bipyrimidal polygon, we can establish a linear relationship between the $p\text{O}3'$ -Pa distance and the ionic radius of any divalent metal ion where the slope is about 1.58. With this slope and the average coordination bond lengths observed here, the calculated $p\text{O}3'$ -Pa distance for Mg^{2+} and Mn^{2+} is about 2.1–2.2 Å. We have calculated the $p\text{O}3'$ -Pa distance for Ca^{2+} to be about 2.9 Å based on the proposed TS model, which is far beyond the expected distance for spontaneous bond formation (10). Because we observed that Ca^{2+} has little influence on the position of $p\text{O}3'$, the actual $p\text{O}3'$ -Pa distance cannot be reduced at all by any coordination effects of Ca^{2+} . We note that divalent metal ion geometry is not completely symmetrical for nucleotidyl transfer and pyrophosphorolysis in this hypothetical

TS model because metal ion B is not directly involved in interaction with the α,β -bridging oxygen of the incoming dNTPs.

The proposed TS model provides insights into how the TS might be stabilized within the closed ternary complex. In the TS, $p\text{O}3'$ has one negative charge after deprotonation and is attracted to the partial positive charges of Pa . All four ligands of Pa ($\text{O}5'$, $\text{O}1\alpha$, $\text{O}2\alpha$, and $\text{O}3\alpha$) have extra partial negative charges after drawing electrons from Pa . Based on the observed geometry, we propose that W-5, metal ion A, and C6-H/C8-H of the incoming base neutralize partial negative charges on $\text{O}1\alpha$, $\text{O}2\alpha$, and $\text{O}5'$, respectively (5), and that ϵ -amino group of K560 neutralizes partial negative charges on $\text{O}3\alpha$ by providing a proton in nucleotidyl transfer, a step that has already been established from biochemical experiments (8, 18).

To address the relevance of the proposed TS for nucleotidyl transfer with respect to other pols, we compared the metal ion-centered octahedrons observed here in RB69pol with octahedrons in other pols. However, most of the ternary complexes for other pols, with the exception of pol β (9), have been obtained with incomplete metal-ion coordination geometry due to missing $p\text{O}3'$ or to the use of Ca^{2+} . With RB69pol, two divalent catalytic metal ions have two shared carboxylates DX and DY and the Sp oxygen ligand of dNTP's Pa ($\text{O}2\alpha$), where DX and DY are D411 and D623, respectively (Fig. 5). In addition to DX and DY, three other oxygen ligands for metal ion B come from the triphosphate tail of the incoming dNTP, one from each phosphate, and the sixth ligand from the backbone carbonyl of a hydrophobic residue Z, where Z is L412. These ligands are conserved in T7 pol, HIV reverse transcriptase, and Dpo4 (31–33), with corresponding DX/DY/Z being D475/D654/A476, D110/D185/V111, and D7/D105/F8, respectively (Fig. 5E, Fig. S7). When metal ions and corresponding ligands were used for superposition between RB69pol and each of these pols, we observed the same relationship of the three polygons, suggesting the essential functional role of metal ion A for reducing the $p\text{O}3'$ - Pa distance in a conserved nucleotidyl transfer mechanism (Fig. 5). In addition, the structural basis as to why Ca^{2+} is catalytically inert, and why Mg^{2+} and Mn^{2+} are catalytically competent in the case of RB69pol should be universally applicable to all DNA pols.

The geometric conservation of three polygons observed in RB69pol here also extends to pol β , whose Palm topology is completely unrelated to RB69pol (Fig. 5E). In pol β , DX and DY are D190 and D192, respectively. Interestingly, pol β has another carboxylate, D256, serving as a ligand to metal ion A replacing an ordered water molecule that is present in RB69pol. However, pol β also has an ordered water molecule serving as a sixth ligand to metal ion B replacing Z (9). Therefore, ligand variations exist for two metal ions in the pol active site of these two pols, which are the only two pols up to date that have complete metal-ion coordination in pre-insertion ternary complex. Nevertheless, the relationship among three polygons for nucleotidyl transfer is exactly the same between these two pols. The conservation of this relationship in RB69pol, pol β , and other pols suggests a conserved catalytic mechanism for all pols (9, 31–34), and supports of an earlier proposal using the conserved “catalytic entity” for structural alignment of non-homologous pols, where the “catalytic entity” referred to the metal ion coordination octahedrons that were only partially available at the time (35).

Although three polygons centered at divalent metal ions in the hypothetical TS for nucleotidyl transfer TS appear to be conserved among various pols, the specific motions involved in the ternary complexes approaching the TS may not be similar because the torsion angles of incoming dNTPs differ from one pol family to the next (Fig. S8). Similarly, the torsion angles and sugar puckering of priming nucleotide residues also vary, even though the bases of the priming nucleotide residues and of the incoming dNTPs are

always stacked perfectly. Torsion angle variations of incoming dNTPs can be eliminated when comparing pre-insertion ternary complexes of two pols from the same pol B family, for example, between RB69pol and ϕ 29 DNA polymerase (5, 34). In this case, the starting point of the ternary complex and the ending point of the TS are identical, and comparison between the two pols should further reveal new insights into specific motions in the reaction trajectory. When the metal ion-centered coordination octahedrons are used for superposition, we are surprised to learn that the orientation of the P/T duplexes in ϕ 29 pol and RB69pol differ from the situation where when Palm domains are used for superposition (Fig. S9). This finding suggests that the reorientation of the P/T duplexes may occur when the closed ternary complex approaches the TS, because $p\text{O}3'$ is a part of the P/T and cannot be displaced independent of it.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was in part funded by NIH R01 GM063276-9 and 3R01GM063276-08S1 to W. K., and Steitz Center for Structural Biology, Gwangju Institute of Science and Technology (SCSB-GIST), Republic of Korea to J.W.

ABBREVIATIONS

pol	DNA polymerase
RB69pol	bacteriophage RB69 DNA polymerase
wt	wild type
tm	triple mutant with three substitutions L561A, S565G, and Y567A
qm	quadruple mutant with four substitutions L415A, L561A, S565G, and Y567A
dCTP or dNTP	2'-deoxycytosine or 2'-deoxynucleoside 5'-triphosphate
dUpNpp	non-hydrolyzable 2'-deoxyuridine 5'- α,β -imino triphosphate analogue
dUpCpp	non-hydrolyzable 2'-deoxyuridine 5'- α,β -methylene triphosphate analogue
dNMP	deoxynucleoside monophosphate
TS	transition state
$p\text{O}3'$	primer-terminal 3'-hydroxyl
Pa	α -phosphorus atom of the triphosphate moiety of incoming dNTPs or dUpXpp where X is NH or CH ₂
ddC	2',3'-dideoxy-terminated non-extendable primer without $p\text{O}3'$ in comparison with dC, the standard extendable primer with $p\text{O}3'$
ddP/T or dP/T (P/T) duplex	2',3'- dideoxy-terminated or 2'-deoxy-terminated primer/template duplex, respectively
PEG 350 MME	polyethylene glycol 350 monomethyl ether

REFERENCES

1. Beese LS, Steitz TA. Structural basis for the 3'-5' exonuclease activity of Escherichia coli DNA polymerase I: a two metal ion mechanism. *EMBO J.* 1991; 10:25–33. [PubMed: 1989886]
2. Steitz TA. DNA- and RNA-dependent DNA polymerases. *Curr. Opin. Struct. Biol.* 1993; 3:31–38.
3. Joyce CM, Steitz TA. Function and Structure relationship in DNA polymerases. *Ann. Rev. Biochem.* 1994; 63:777–822. [PubMed: 7526780]
4. Franklin MC, Wang J, Steitz TA. Structure of the replicating complex of a pol alpha family DNA polymerase. *Cell.* 2001; 105:657–667. [PubMed: 11389835]
5. Wang M, Xia S, Blaha G, Steitz TA, Konigsberg WH, Wang J. Insights into base selectivity from the 1.8 Å resolution structure of an RB69 DNA polymerase ternary complex. *Biochemistry.* 2011; 50:581–590. [PubMed: 21158418]
6. Tsai YC, Johnson KA. A new paradigm for DNA polymerase specificity. *Biochemistry.* 2006; 45:9675–9685. [PubMed: 16893169]
7. Castro C, Smidansky ED, Maksimchuk KR, Arnold JJ, Korneeva VS, Gotte M, Konigsberg W, Cameron CE. Two proton transfers in the transition state for nucleotidyl transfer catalyzed by RNA- and DNA-dependent RNA and DNA polymerases. *Proc. Natl. Acad. Sci. USA.* 2007; 104:4267–4272. [PubMed: 17360513]
8. Castro C, Smidansky ED, Arnold JJ, Maksimchuk KR, Moustafa I, Uchida A, Gotte M, Konigsberg W, Cameron CE. Nucleic acid polymerase use a general acid for nucleotidyl transfer. *Nat. Struct. Mol. Biol.* 2009; 16:212–218. [PubMed: 19151724]
9. Batra VK, Beard WA, Shock DD, Krahn JM, Pedersen LC, Wilson SH. Magnesium-induced assembly of a complete DNA polymerase catalytic complex. *Structure.* 2006; 14:757–758. [PubMed: 16615916]
10. Lin P, Pedersen LC, Batra VK, Beard WA, Wilson SH, Pedersen LG. Energy analysis of chemistry for correct insertion by DNA polymerase beta. *Proc. Natl. Acad. Sci. USA.* 2006; 103:13294–13299. [PubMed: 16938895]
11. Beard WA, Shock DD, Vande Berg JJ, Wilson SH. Efficiency of correct nucleotide insertion governs DNA polymerase fidelity. *J. Biol. Chem.* 2002; 277:47393–47398. [PubMed: 12370169]
12. Beckman J, Wang M, Blaha G, Wang J, Konigsberg WH. Substitution of Ala for Tyr567 in RB69 DNA polymerase allows dAMP to be inserted opposite 7,8-dihydro-8-oxoguanine. *Biochemistry.* 2010; 49:4116–4125. [PubMed: 20411947]
13. Beckman J, Wang M, Blaha G, Wang J, Konigsberg WH. Substitution of Ala for Tyr567 in RB69 DNA polymerase allows dAMP and dGMP to be inserted opposite 7-guanidinohydroxydantoin. *Biochemistry.* 2010; 49:8554–8563. [PubMed: 20795733]
14. Xia S, Wang M, Lee HR, Sinha A, Blaha G, Christian T, Wang J, Konigsberg W. Variation in mutation rates caused by RB69pol fidelity mutants can be rationalized on the basis of their kinetic behavior and crystal structures. *J. Mol. Biol.* 2011; 406:558–570. [PubMed: 21216248]
15. Xia S, Konigsberg WH, Wang J. Hydrogen-bonding capability of a templating difluorotoluene nucleotide residue in an RNA DNA polymerase ternary complex. *J. Am. Chem. Soc.* 2011; 133:10003–10005. [PubMed: 21667997]
16. Zhang H, Rhee C, Benenek A, Drake JW, Wang J, Konigsberg W. The L561A substitution in the nascent base-pair binding pocket of RB69 DNA polymerase reduce base selectivity. *Biochemistry.* 2006; 45:2211–2220. [PubMed: 16475809]
17. Zhang H, Beckman J, Wang J, Konigsberg W. RB69 DNA polymerase mutants with expanded nascent base-pair binding pockets are high efficient but have reduced base selectivity. *Biochemistry.* 2009; 48:6940–6950. [PubMed: 19522539]
18. Yang G, Franklin M, Lin TC, Konigsberg W. Correlation of the kinetics of finger domain mutants in RB69 DNA polymerase with its structure. *Biochemistry.* 2002; 41:2526–2534. [PubMed: 11851399]
19. Otwinowski, Z.; Minor, W. Processing of X-ray diffraction data collected in oscillation mode. In: Carter, CW., Jr; Sweet, RM., editors. *Methods in Enzymology*. Vol. 276. New York: Academic Press; 1997. p. 307-326. *Macromolecular Crystallography, part A*

20. McCoy AJ, Grosse-Kunstev RW, Adams PD, Win LC, Storoni LC, Read RJ. Phaser crystallographic software. *J. Appl. Cryst.* 2007; 40:458–674.
21. Pottertton E, McNicholas S, Krissinel E, Cowtan K, Noble M. The CCP4 molecular-graphics project. *Acta Cryst.* 2002; D58:1955–1957.
22. Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. *Acta Cryst.* 2004; D60:2126–2132.
23. Murshudov GN, Vagin AA, Dodson EJ. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Cryst.* 1997; D53:240–255.
24. Carson M. Ribbons 2.0. *J. Appl. Cryst.* 1991; 24:958–961.
25. Goodman MF, Keener S, Guidotti S, Branscomb EW. On the enzymatic basis for mutagenesis by manganese. *J. Biol. Chem.* 1983; 258:3469–3475. [PubMed: 6833210]
26. Hays H, Berdis AJ. Manganese substantially alters the dynamics of translesion DNA synthesis. *Biochemistry.* 2002; 41:4771–4778. [PubMed: 11939771]
27. Reineks EZ, Berdis AJ. Evaluating the effects of enhanced processivity and metal ions on translesion DNA replication catalyzed by the bacteriophage R4 DNA polymerase. *J. Mol. Biol.* 2003; 328:1027–1045. [PubMed: 12729739]
28. Zakharova E, Wang J, Konigsberg W. The activity of selected RB69 DNA polymerase mutants can be restored by manganese ions: the existence of alternative metal ion ligands used during the polymerization cycle. *Biochemistry.* 2004; 43:6587–6595. [PubMed: 15157091]
29. Pelletier H, Sawaya MR, Wolfle W, Wilson SH, Kraut J. A structural basis for metal ion mutagenicity and nucleotide selectivity in human polymerase beta. *Biochemistry.* 1996; 35:12762–12777. [PubMed: 8841119]
30. Shannon RD. Revised effective ionic radii and systematic studies of interatomic distances in halides and chalcogenides. *Acta Cryst.* 1976; A32:751–767.
31. Harding MM. Geometry of metal-ligand interactions in proteins. *Acta Cryst.* 2001; D57:401–411.
32. Doublet S, Tabor S, Long AM, Richardson CC, Ellenberger T. Crystal structure of a bacteriophage T7 DNA replication complex at 2.2 Å resolution. *Nature.* 1998; 391:251–258. [PubMed: 9440688]
33. Huang H, Chopra R, Verdine GL, Harrison SC. Structure of a covalently trapped catalytic complex of HIV-1 reverse transcriptase: implications for drug resistance. *Science.* 1998; 282:1669–1675. [PubMed: 9831551]
34. Ling H, Boudsocq F, Woodgate R, Yang W. Crystal structure of a Y-family DNA polymerase in action: a mechanism for error-prone and lesion-bypass replication. *Cell.* 2001; 107:91–102. [PubMed: 11595188]
35. Berman AJ, Kamtekar S, Goodman JL, Lazaro JM, deVega M, Blanco L, Salas M, Steitz TA. Structures of phi29 DNA polymerase complexed with substrate: the mechanism of translocation in B-family polymerases. *EMBO J.* 2007; 26:3496–3505.
36. Steitz TA, Smerdon SJ, Jager J, Joyce CM. A unified polymerase mechanism for nonhomologous DNA and RNA polymerases. *Science.* 1994; 266:2022–2025. [PubMed: 7528445]

Figure 1.

Ca^{2+} coordination in the dUpNpp-containing ternary complexes. (A) The wt complex in the presence of a 2',3'-dideoxy-terminated primer (ddC). (B) The Ca^{2+} coordination in the tm complex (golden) is exactly the same as in the wt complex (multi-colors as in A). (C) The tm complex in the presence of an extendable primer (dC). (D) Coordination comparison of the ternary complexes in the presence of dC (yellow) and ddC (golden) shows that the removal of *ptO3'* in ddC is responsible for introduction of more than 6 ligands for Ca^{2+} coordination, for example, for the addition of W-4 in (A). (E) The dCTP-containing ternary complex of wt RB69pol published previously (pdb entry, 3NCI) shows a new water molecule W-7 coordinated to CaA, but missing the bridging water molecules W-4. (F)

Coordination comparison of the dCTP (silver) and dUpNpp-containing (multi-colors) wt ternary complexes shows altered water molecule ligands, where W-4 is part of hydration network connecting to the α,β -bridging NH of dUpNpp.

Figure 2.

Mn^{2+} coordination in the dUpNpp-containing ternary complex. (A) Anomalous difference Fourier maps contoured at 6.5σ (red) and the omitted $F_{obs}-F_{calc}$ maps contoured at 1σ (cyan) in stereodiagram show well defined Mn^{2+} coordination. (B) Mn^{2+} coordination details: B site, nearly a perfect octahedron; A site, a distorted octahedron by $ptO3'$. (C) Coordination comparison of the ternary complexes in the presence of Mn^{2+} (multi-colors) and Ca^{2+} (silver) shows that the smaller ionic radius of Mn^{2+} than Ca^{2+} causes a contraction in both A and B sites, adding a stronger interaction between $ptO3'$ and MnA than between $ptO3'$ and CaA. (D) Details of Mn^{2+} coordination in stereodiagram with outlined octahedrons. (E) The Mn^{2+} /dUpNpp-containing tm ternary complex in the absence of $ptO3'$.

(F) Comparison of the Mn^{2+} /dUpNpp-containing tm ternary complexes in the presence (silver) and absence of $p\text{O}3'$ (mutli-color as in E).

Figure 3.

Mg²⁺ coordination in the dUpCcpp-containing qm ternary complex. (A) Complete coordination of the Mg²⁺/dUpCcpp-containing qm ternary complex. (B) Comparison of the Mg²⁺/dUpCcpp-containing qm ternary complex (multicolor) with the Mn²⁺/dUpNpp-containing tm ternary complex (silver). (C) Stereodiagram of the Mg²⁺ coordination.

Figure 4.

The Ca^{2+} /dUpCpp qm ternary complexes. (A) The Ca^{2+} /dUpCpp-containing qm ternary complex. (B-C) Two orthogonal views of comparison of the Ca^{2+} /dUpCpp-containing qm ternary complex (multicolor) with the Ca^{2+} /dUpNpp-containing tm ternary complex (silver). (D) Comparison of the Ca^{2+} /dUpCpp-containing qm ternary complex (multicolor) with the Ca^{2+} /dUpNpp-containing wt ternary complex (silver). (E-F) Two orthogonal views of comparison of Ca^{2+} /dUpCpp-containing qm ternary complex (multicolor) with the Mg^{2+} /dUpCpp-containing qm ternary complex (silver).

Figure 5.

Mechanistic implications from the α, β, γ -tridentate Mg^{2+}/Mn^{2+} coordination complex. (A) A displacement of $ptO3'$ into the vertex of a non-distorted, idealized metal-ion A octahedron (as indicated in black arrows and octahedron outlines) would place it in a distance for bond formation with $P\alpha$ and in a hypothetical TS model. $P\alpha$ is also displaced towards $ptO3'$ for the hypothetical $P\alpha$ -center inversion (see text). (B) Two idealized metal-ion coordination octahedrons linked by shared ligands. (C) Geometric relationship between the two idealized metal-ion coordination octahedrons (cyan and yellow) and a hypothetical pentavalent TS (magenta). (D) Coordination comparison of the dUpNpp-containing ternary complexes of RB69pol (golden) and pol β (yellow) upon superposition using metal ion ligands. Second

residue labels are for corresponding catalytic carboxylates of pol β . (E) Coordination comparison of the dUpNpp-containing ternary complex of RB69pol (yellow) with dNTP-containing ternary complexes of other pols: T7 DNA polymerase (green), HIV reverse transcriptase (cyan), and Dpo4 (magenta). Three B site ligands are backbone carbonyl (Z, L412, A476, V111, and F8, respectively, for the above-mentioned four pols) and two carboxylates (DX/DY, D411/D623, D475/D654, D110/D185, and D7/D105).

Table 1Summary of crystallographic data and refinement statistics for Ca²⁺-containing ternary complexes

RB69pol Nascent base pair	wt (ddC, Ca²⁺) dUpNpp:dA	tm (ddC, Ca²⁺) dUpNpp:dA	tm (dC, Ca²⁺) dUpUpp:dA	qm (dC, Ca²⁺) dUpCpp:dA
Space group	P2 ₁ 2 ₁ 2 ₁	P2 ₁ 2 ₁ 2 ₁	P2 ₁ 2 ₁ 2 ₁	P2 ₁ 2 ₁ 2 ₁
Unit cell dimensions [a, b, c (Å)]	75.10, 120.47, 130.56	75.05, 119.82, 130.79	75.02, 119.33, 130.66	78.17, 119.06, 130.21
Resolution range ^a (Å)	50–1.83 (1.86–1.83)	90–195 (2.02–1.95)	88–2.35 (2.43–2.35)	50–2.43 (2.52–2.43)
Number of reflections				
Unique	105,414	84,003	49,563	43,626
Redundancy	4.0 (3.3)	3.3 (1.5)	6.7 (6.6)	3.6 (3.6)
Completeness (%)	99.8 (97.3)	94.3 (57.9)	100 (100)	99.7 (99.9)
R _{merge} (%) ^b	8.6 (85.0)	9.3 (71.9)	16.8 (>100)	12.0 (85.8)
I/σ	13.7 (1.0)	12.7 (0.93)	9.2 (1.1)	9.0 (1.1)
Final model				
Amino acid residues	903	903	903	903
Water molecules	857	1,153	345	229
Metal ions	5	7	4	4
Template nucleotides	18	18	18	18
Primer nucleotides	13	13	13	13
dNTP molecules	1	1	1	1
Refinement Statistics				
Reflections	99,798	79,392	46,983	46,149
R _{work} (%) ^c	18.2 (31.1)	17.1 (30.6)	21.4 (29.5)	20.0 (26.8)
R _{free} (%) ^d	21.2 (33.5)	21.3 (35.0)	25.9 (40.0)	25.6 (34.8)
R _{o2p} ^e	2.78	2.15	1.40	1.33
r.m.s.d^f				
Bond length (Å)	0.007	0.0042	0.0066	0.008
Bond angle (°)	1.082	0.87	1.10	1.169
PDB access code	3SQ1	3S9H	3SCX	3SPZ

^aThe highest resolution shell statistics are in parenthesis.^bR_{merge} = $\langle \sum_{hk\ell} |I(hk\ell) - \langle I(hk\ell) \rangle| / \langle I(hk\ell) \rangle \rangle$, merging statistics for all symmetry-mates.^cR_{work} = $\sum_{hk\ell} |F_{obs}(hk\ell) - F_{calc}(hk\ell)| / \sum_{hk\ell} F_{obs}(hk\ell)$, crystallographic R-factor.^dR_{free} Cross-validation R-factor for ~5% of the total unique reflections that have been randomly selected.^eThe observation-to-parameter ratio is defined by the ratio between the unique number of observations for refinement and the number of variables, which is 4 four times of the number of atoms.^fr.m.s.d.: root-mean-square deviation from ideal values.

Table 2

Summary of crystallographic data and refinement statistics for Mn²⁺ or Mg²⁺-containing ternary complexes

RB09poll Nascent base pair	tm (dC, Mn ²⁺) dUpNpp:dA	tm (ddC, Mn ²⁺) dUpNpp:dA	tm (dC, Mg ²⁺) dUpCpp:dA	tm (ddC, Mg ²⁺) dCpTP:dG
Space group	P2 ₁ 2 ₁ 2 ₁	P2 ₁ 2 ₁ 2 ₁	P2 ₁ 2 ₁ 2 ₁	P2 ₁ 2 ₁ 2 ₁
Unit cell dimensions [a, b, c (Å)]	78.55, 118.22, 130.11	78.15, 119.67, 130.64	75.47, 120.20, 131.21	75.33, 119.53, 130.74
Resolution range ^a (Å)	47-2.38 (2.38-2.43)	50-2.00 (2.07-2.00)	89-1.85 (1.90-1.85)	20-2.00 (2.07-2.00)
Number of reflections				
Unique	45,847	82,500	65,531	77,923
Redundancy	4.1 (2.3)	3.2 (3.2)	3.8 (3.6)	8.6 (5.3)
Completeness (%)	89.4 (77.9)	99.2 (99.9)	99.5 (99.5)	98.2 (99.6)
R _{merge} (%) ^b	12.3 (77.8)	7.6 (89.8)	12.4 (82.7)	13.9 (>100)
I/σ	10.0 (1.0)	12.7 (1.2)	10.1 (1.3)	12.3 (1.3)
Final model				
Amino acid residues	901	903	903	903
Water molecules	138	302	1,116	585
Metal ions	7	5	1	5
Template nucleotides	18	18	18	18
Primer nucleotides	13	13	13	13
dNTP molecules	1	1	1	1
Refinement Statistics				
Reflections	42,465	78,132	95,428	74,966
R _{work} (%) ^c	20.2 (28.2)	20.8 (29.4)	18.3 (26.5)	19.3 (29.7)
R _{free} (%) ^d	26.5 (37.3)	24.5 (32.0)	21.9 (28.9)	22.2 (34.6)
R _{exp} ^e	1.30	2.34	2.60	2.17
r.m.s.d^f				
Bond length (Å)	0.0074	0.008	0.008	0.0068
Bond angle (°)	1.12	1.10	0.85	1.147
PDB access code	3SIJ	3SQ0	3SI6	3SPY
				3SNJ

^aThe highest resolution shell statistics are in parenthesis.

^b $R_{\text{merge}} = \langle \sum h k l \bar{I} / \sum h k l \rangle - \langle I(h k l) \rangle / \langle I(h k l) \rangle$, merging statistics for all symmetry-mates.

^c $R_{\text{work}} = \sum h k l / F_{\text{obs}}(h k l) - F_{\text{calc}}(h k l) / \sum h k l / F_{\text{obs}}(h k l)$, crystallographic R-factor.

^d R_{free} Cross-validation R-factor for ~5% of the total unique reflections that have been randomly selected.

^eThe observation-to-parameter ratio is defined by the ratio between the unique number of observations for refinement and the number of variables, which is 4 four times of the number of atoms.

^fr.m.s.d.: root-mean-square deviation from ideal values.

Table 3

Competitive inhibition constants of dUpNpp against dTMP incorporation

$K_d^{\text{dUpNpp}}(\mu\text{M})$	10mM Mg^{2+}	10mM Mn^{2+}
tm RB69pol	6	10
wt RB69pol	24	37