Morphological Variants of Coliphage P1¹

DONALD H. WALKER, JR.,² AND THOMAS F. ANDERSON

Institute for Cancer Research, Philadelphia, Pennsylvania 19111

Received for publication 10 March 1970

Lysates of P1 from all hosts tested contained at least three morphological variants with respect to head size. These were termed "big" (P1B), "small" (P1S), and "minute" (P1M). Since successive clonings of plaques isolated on many different hosts failed to change the proportions of the variants, we concluded that the production of variants was a function of the P1 genome rather than that of the host. In the electron microscope, the heads appeared to be icosadeltahedra, having face-to-face head diameters of 86 ± 2 nm, 65 ± 2 nm, and 47 ± 2 nm. Assuming the head capsids to be composed of the same protein subunits, these diameters were compatible with $T = 16$, 9, and 4 with a lattice constant (intercapsomere distance) of 12 to 13 nm. The tails of all variants were morphologically indistinguishable. Each consisted of a hollow tail tube surrounded by a contractile sheath attached to the head by means of a "head-neck connector" which could be a specialized vertex capsomere. In CsCl gradients, a number of bands were observed. One band contained the majority of P1B particles and 99% of the plaque-forming units. Two other bands contained P1S particles whose densities suggested a content of about 40 and 60% of the complete P1B genome. The less dense of these two bands also contained defective P1B particles with a calculated content of only 60% of the complete genome. The P1S particles tested injected their deoxyribonucleic acid (DNA) into host cells and killed them. Genetic markers contained in this band could be rescued by infectious P1B particles, confirming the evidence of Ikeda and Tomizawa that this fraction contains P1 DNA.

In 1960, Eduard Kellenberger brought a number of strains of P1 to this laboratory for examination in the electron microscope (EM). The heads of these particles appeared as regular hexagons, but they occurred in two sizes (3). Every single plaque that was examined contained particles of both sizes. This suggested that we were dealing not with two different strains of P1 but with some unexpected type of aberration in the assembly of the particle, because, in general, progeny virions resulting from single infection had been reported to be essentially identical to the parent virion in size and structure. It was, therefore, of considerable interest to understand how a virion can produce morphological variants which differ in size or shape.

Phage P1 had originally been isolated by Bertani (7) from the Lisbonne-Carrère strain of *Escherichia coli* and had been observed in the EM by Anderson (2) and by Labaw and Mosley (30). Particles with two different head sizes were subsequently shown to be separable in CsCl den-

sity gradients by Anderson and Walker (6). Ikeda and Tomizawa (19, 20) have since shown that both types of particle contain deoxyribonucleic acid (DNA) having a density slightly less than that of the host DNA. Other properties of these particles have been investigated by Karamata (Certificate Report, Inst. of Mol. Biol., Univ. of Geneva, Switzerland, 1964) and Walker (Ph.D. Thesis, Univ. of Pennsylvania, Philadelphia, 1966).

The big-headed form (P1B) is the infectious particle and is always in the majority. The small-headed particle (P1S) will not form plaques on any known hosts of P1B. Extensive EM observation of P1 lysates have revealed a third morphological variant, a particle with a "minute" head which has been termed P1M (Walker, 1966). Figure 1 shows examples of P1B, P1S, and P1M. Electron microscopy shows that all three head types are isometric polyhedra and that all possess tails which are visually indistinguishable.

Morphological variants have thus far been observed for coliphage T4. The normal T4 head is an elongated polyhedron, but lysates of T4 also contain a short-headed form which has the same width but which is isometric and, therefore, shows a regular hexagonal outline in the EM. These

¹ Presented in part by Donald H. Walker, Jr., in partial fulfillment of the requirements for the Ph.D. degree in Molecular Biology from the University of Pennsylvania.

² Present address: Department of Microbiology, School of Medicine, University of Iowa, Iowa City, Iowa 52240.

FIG. 1. Electron micrograph of a lysate of P1-Sh showing particles of three head sizes. Arrows point to the heads of P1B, P1S, and P1M, all of which have tails with contracted sheaths. Stained with silicotungstate. The bar in this and subsequent micrographs represents 100 nm.

particles carry two-thirds fragments of the normal T4 genome that are randomly permuted (35; J. Carnighan and G. Mosig, *personal communication*). Other cases of T4 morphological variants have been observed but are quite rare: normal (prolate) or abnormal (biprolate) heads to which two or even three tails may be attached (8, 32, 38). None of these T4 variants should be confused with the "conformational variations" of T-even phages reported by Cummings and Kozloff (14); these are reversible changes in head shape that result from variations in pH and ionic strength.

We have determined the densities in CsCl gradients of a number of forms of P1 and were thus able to estimate their contents of DNA. The biological properties of the P1S particles have also been determined after their isolation from CsCl gradients. Although P1S do not form plaques, they can kill their host cells. They are capable of adsorbing to and injecting their DNA into host cells. At least one genetic marker can be rescued from defective P1 particles by coinfecting normal P1B particles.

MATERIALS AND METHODS

Phage strains. P1 [P1*kc* of Lennox (31)] was a gift from S. E. Luria. P1*c*, which forms clear plaques on *Shigella dysenteriae* Sh, was a stable mutant isolated in this laboratory. $\lambda + + \cdot K$ (a gift of M. Meselson) was used as a density reference standard (1.508 g/cm³) on CsCl gradients. T4D was used as an internal standard for magnification calibration in EM preparations.

Bacterial strains. The bacterial strains are described in Table 1.

Media. LB broth (1% tryptone, 0.5% yeast extract, 1% NaCl, 0.1% glucose; pH adjusted to about 7.4 with 1 N NaOH) was routinely used for growing bacteria. LC broth (LB broth brought to 2.5 mM CaCl₂) was used for adsorption mixtures in which Ca²⁺ is required for P1 adsorption (7). LC hard agar (LB broth with 1.0% agar; CaCl₂ added to a concentration of 2.5 mM just before pouring) was used for routine P1 assays and for the preparation of stocks from confluent lysis plates.

LC soft agar for the top layer of the plates consisted of LB broth with 0.6% agar containing 2.5 mM CaCl₂.

M14 buffer (1 liter of water containing 3.0 g of Na₂HPO₄, 4.0 g of NaCl, 22.5 g of K₂SO₄, and 1 ml of 1% gelatin; after autoclaving, 5 ml of 1.0 M MgSO₄ and 8 ml of 25 mM CaCl₂ were added) was used for dilutions and for storing P1.

Phosphate buffer (pH 7.2, 0.1 M) and tris (hydroxymethyl) aminomethane (Tris) buffer (pH 7.4, made by a 10-fold dilution of a 0.5 M stock solution containing 6% Tris, 5% NaCl, and 1% NH₄Cl dissolved in 0.5 N HCl) were also used.

It should be noted that another variation of Tris buffer (0.02 M Tris-hydrochloride, 0.01 M MgSO₄, 1% NH₄ acetate, pH 7.2) is currently being used in P1 experiments because even low titers of P1 show no loss after 2 years if stored in it at 4 to 8 C. MgSO₄ is the critical factor (Walker, *unpublished data*).

Phage assays. Sh was generally used as the host for P1 assays. Prior to plating, P1 was preadsorbed on Sh for 10 min at 37 C in the presence of Ca²⁺. Adsorption tubes were then transferred to 45 C, 3.0 ml of LC soft agar was added, and the mixtures were poured onto the surfaces of plates containing 45 to

TABLE 1. List of bacterial strains

Bacterial strain	Symbol used in text	Characteristic	Source	Source code
<i>Escherichia coli</i> B	B	Wild type	E. W. Six	B-06
<i>Escherichia coli</i> Bc	Bc	B cured of X ^a	E. W. Six	B-15
<i>Escherichia coli</i> K-12	K-12	Lysogenic for λ	A. D. Kaiser	B-40
<i>Escherichia coli</i> C600	C600	Cured of λ	M. Meselson	
<i>Escherichia coli</i> Cla	Cla	ATCC C122 adapted for better growth	G. Bertani	Cla
<i>Escherichia coli</i> Cla(P3)	Cla(P3)	Lysogenic for P3	E. W. Six	C-24
<i>Shigella dysenteriae</i> Sh16	Sh	Best P1 indicator	S. E. Luria	
<i>Salmonella typhimurium</i> LT-2/22	St/22	Does not adsorb P1	N. Yamamoto	

^a X is a defective prophage normally carried by *E. coli* B (12).

50 ml of LC hard agar warmed to 37 C. Plates were incubated right side up at 37 C, and plaques could easily be scored on the following day.

λ was assayed by the method of Adams (1) by using LB media containing 3 mM MgSO₄.

Preparation of high titer stocks of P1. P1 stocks of high titer were made from confluent lysis plates (40) with the following modifications. Phage was pre-adsorbed to host cells in LC broth at 37 C for 10 to 15 min. This adsorption mixture was added to LC soft agar and plated on fresh, nondried prewarmed LC hard agar plates, so that each plate received 4.5 to 5.0 ml of soft agar containing 5 to 10 $\times 10^8$ plaque-forming units (PFU) and about 10⁹ *Escherichia coli* host cells. Some LC broth was usually added to the LC soft agar at 45 C with the preincubation mixture so that the final concentration of agar was between 0.4 and 0.5%. After 8 to 12 hr of incubation at 37 C, when plaques were contiguous, the plates were tilted to allow the semisolid top layer to slide off into centrifuge tubes. The phage was then purified and concentrated by differential centrifugation. With C600 cells as the host bacteria, 5 $\times 10^{11}$ to 10 $\times 10^{11}$ PFU per confluent lysis plate could be obtained routinely.

Preparation of one-step broth lysates of P1. *E. coli* C600 cells were grown overnight in a reciprocal shaker-incubator (Lab-Line Instruments) at 37 C. The culture was then centrifuged, resuspended, and diluted in LC broth at 37 C to a concentration of 10⁸ to 2 $\times 10^8$ cells/ml. The culture was shaken vigorously in a cotton-plugged Erlenmeyer flask for 20 min, and P1 was added at a multiplicity of infection (MOI) of 3 to 4. At 25 min after infection, the C600 cells were superinfected with the same amount of P1. The turbidity of the culture leveled off after 35 min and subsequently decreased after 55 min, indicating lysis. At 100 min when the turbidity had stopped falling, the P1 in the lysate was purified and concentrated by differential centrifugation.

³²P-labeled P1. C600 cells were grown at 37 C for six to eight generations in LB broth containing 60 to 70 μ C of ³²PO₄ per ml (~22 μ C/mg of P) to label nucleotide precursor pools. LC broth was added to dilute the cells appropriately for P1 infection. P1 was added 10 to 15 min later at an MOI of 4. The culture was superinfected as described above, and the re-

sultant lysates were treated with deoxyribonuclease and ribonuclease before purification and concentration by differential centrifugation.

Assays of radioactivity. Counts of ³²P in labeled phage were carried out by putting 0.2-ml volumes directly onto glass-fiber filters, which were dried and then counted in a Packard Tri-Carb liquid scintillation spectrometer, with 10 ml per counting vial of toluene containing 4 g of 2,5-diphenyloxazole per liter and 100 mg of 1,4-bis-2-(5-phenyloxazolyl)-benzene per liter.

CsCl density gradient centrifugation. Methods generally used were those of Weigle et al. (43), Ting (42), and Vinograd and Hearst (44). CsCl was obtained from Penn Rare Metals, Inc., Maywood Chemical Co., and the American Potash and Chemical Co. (Trona). Stock solutions were made by dissolving CsCl in whatever solvent was being used (Tris buffer, phosphate buffer, LB broth, M14 buffer, or a 1:1 mixture of LB and M14). Densities of these solutions were in the range of 1.85 to 1.88 g/cm³ calculated from the index of refraction as determined with an Abbe refractometer.

CsCl stock solution and phage mixtures to be run as gradients were made up to 4.8 ml with average densities from 1.400 to 1.500 g/cm³, depending on which bands were being sought. The mixtures were placed in lusteroid tubes, covered with a thin layer of mineral oil, and run in an SW39 swinging-bucket head in a Spinco model L centrifuge. Uniformly mixed solutions were centrifuged at 20,000 to 39,000 rev/min (21,000 to 170,000 $\times g$) for 20 to 45 hr. At the end of the run, the tube was carefully removed, a photograph was taken when visible bands were present, and 30 to 100 fractions were collected for index of refraction readings, bioassays, and electron microscopy.

P1 is large enough to approach equilibrium within a few hours in preformed gradients. Such gradients were made either in a standard mixing chamber device or by layering three concentrations of CsCl in the centrifuge tube; they were run for 3 to 25 hr.

Electron microscopy. Phage preparations were "embedded" in a negative stain (pH 7), either phosphotungstate (PT; 9) or silicotungstate (ST; 4). A drop of the phage preparation, either from CsCl gradients or from concentrated lysates, was placed on a grid covered with a thin layer of carbon-coated

Formvar. The preparation was washed by touching the grid to the concave meniscus of a 1% ammonium acetate solution in a small test tube and stained by adding a loopful of 2% PT or ST, often containing

0.4% sucrose, to facilitate spreading. A piece of filter paper was then touched to the side of the drop and the excess was drawn off, leaving a thin film. The grid was placed in the grid holder and dried in the

FIG. 2. P1 particles present in the density range between 1.35 and 1.43 g/cm³ of a CsCl gradient. The particles were dialyzed against Tris buffer, washed on the EM grid with 1% ammonium acetate, and negatively stained with phosphotungstate. There are two PIB heads (one ghosted), nine PIS heads (one ghosted), and two PIM heads. Several other structural features should be noted. In addition to the four extended tail sheaths and four fully contracted sheaths, there are two sheaths (S) which are longer than the fully contracted sheaths and which still have baseplates and tail fibers attached. Note also the three sheaths (CE) which have both contracted and extended regions. It can be seen that the neckpieces (N) on these sheaths are slightly thicker than the hollow tail tubes. The head-neck connectors (HC) are annular structures, contiguous with the head capsid, and may well be specialized vertex capsomeres to which the neckpiece specifically attaches. Note also the rings of material attached to the bases of the neckpieces on the tails where the abnormal contraction has left the proximal portion of the tail tube exposed; these rings of material may represent specialized rows of subunits which normally stabilize the extended sheaths and anchor them to the tail tubes. (Photo taken on a Phillips 200 electron microscope at The Lister Institute, London, England.)

vacuum of a Siemens Elmiskop I electron microscope which was operated at 80 kv.

To estimate the relative numbers of the different kinds of particles in a sample, electron micrographs were taken at 10,000 to 20,000 $\times$. Scoring was done on photographic enlargements [11 by 14 inches (28 by 35 cm)] of whole fields. For high resolution studies of P1 structure, EM exposures were taken at magnifications of 25,000 to 50,000 $\times$.

The dimensions of the P1 morphological variants were determined by using phage T4D as an internal calibration standard in mixed EM fields of P1 and T4. The dimensions used for T4 were the extended tail sheath lengths (95 nm), the head width (85 nm), and the head length (115 nm) given by Kellenberger et al. (24). Within any given photograph, each of these parameters of T4 was measured several times to determine the dimensions of P1 in the same photograph. In fields containing only P1, the length of the extended tail sheath (216 nm) was used to determine the magnification.

RESULTS

Head sizes of P1. Examples of P1B, P1S, and P1M are shown in Fig. 1 to 3. It should be noted (Fig. 1, 2) that the tail structure attached to the variant heads, with the sheath either extended or contracted, appears the same as that attached to P1B heads. Figure 3 shows each of the three head types from the same EM field at two focal planes. The average side-to-side diameters of these heads measure 85 nm, 65 nm, and 45 nm, respectively. A summary of head size measurements from several micrographs is given in Table 2.

Another head size, intermediate between P1B and P1S, may also exist. Figure 4 shows one particle with a head which is intermediate in size between the P1B and P1S heads in the same photograph.

Structure of P1 heads. As already mentioned, when not flattened out, most P1 heads appear to have hexagonal outlines in negatively stained preparations. Occasionally, however, particles like those in Fig. 5b, c, d, and f are seen that appear pentagonal. Others in Fig. 5a, e, and f have threefold and twofold axes of symmetry. This strongly suggests that the head has the 5-3-2 symmetry of an icosahedron, which appears hexagonal when completely embedded and viewed either down a twofold or threefold axis. On the other hand, when completely embedded and viewed down a fivefold axis, an icosahedron would appear in silhouette as a regular ten-sided polygon. To appear pentagonal after negative staining, an icosahedron would have to be "floating" in the subunit with only about one quarter of its depth submerged and to be viewed down a fivefold axis as it sits on its vertex. In Fig. 5b, the tail appears to have been attached to such a vertex and to have broken off as it dried.

As with T4 (38), one occasionally observes particles with more than one tail. The particles seen in Fig. 6 have tails with included angles of 62° and 65° in agreement with the theoretical angle of 63° 26' for the attachment of two tails to adjacent vertices of an icosahedron.

High resolution micrographs of P1 heads are shown in Fig. 3. Some fine structure does appear but, as with T4 (25) and λ (17), the number and arrangement of capsomeres are not obvious.

Tail structure. As shown diagrammatically in Fig. 7, the P1 tail is a rigid structure consisting of a hollow tail tube surrounded by a contractile sheath. At the distal end of the sheath is an ill-defined baseplate to which several tail fibers, probably six, are attached. Proximally, as in T4, (13, 26, 32), the tail tube is anchored to the head by a distinct morphological entity which will be termed the "head-neck connector." This term allows for the possibility that the neck region may be functionally distinct from the tail tube proper.

The extended sheath often shows a pattern of 54 ± 2 striations running perpendicular to the long axis of the tail (Fig. 8 and 9b). The total length of the sheath is 216 nm, so the distance between neighboring striations is ~ 3.9 nm. [For comparison, the analogous distance is ~ 3.8 nm for the extended sheath of T4 (5, 29, 33, 34).] The width of the P1 sheath is 18 ± 1 nm.

As seen in Fig. 8, the contracted sheath has a length of 90.5 ± 2.0 nm and a width of 25.5 ± 1.5 nm, and has the appearance of a corncob. When cross-striations are apparent, the distance between them is ~ 3.4 nm. We can estimate that there are 27 ± 1 striations per contracted sheath. As in T4, this is one-half the number of striations in the extended sheath (5, 29, 33, 34).

After exposure to CsCl, the contracted sheath tended to break and sections can be found standing on end (Fig. 9a). The appearance is similar to cross sections of the T4 contracted sheath that show a "wormgear" with 12 "teeth." Each tooth has a central body with an arm extending away from the axis of the sheath.

After passing through the sheath, the neck region (Fig. 2, 9b) of the tail tube attaches to the head capsid by means of the head-neck connector that is visible on ghosts and on tails which have broken away from the head (Fig. 2, 8, and 9b). An analogous structure is seen in T4 (13, 26, 32). The hollow neck region, from the head capsid to the beginning of the sheath, is 13 to 14 nm long and shows no evidence of the collar that exists in T4 (3). Also, it may be slightly thicker than the tail tube itself. The tail tube is hollow, and its cylindrical hole is 2.5 to 3.0 nm in diameter.

The distal end of the extended sheath terminates with a baseplate ~ 3.5 nm in thickness and ~ 21

FIG. 3
770

TABLE 2. Summary of measurements of P1 head sizes^a

Head type	Total average diameter $\pm \sigma$ (nm)	Smallest individual head	Largest individual head	Total heads measured	No. of micrographs
P1B....	85.5 $\pm$ 1.8	80.5	91.5	100	1
P1S....	64.7 $\pm$ 1.5	58.0	67.5	134	3
P1M...	46.7 $\pm$ 2.0	43.0	51.0	32	26

^a Only heads showing hexagonal outlines were measured. The value for an individual head was taken to be the average of the three side-to-side widths of the hexagon.

nm in diameter (Fig. 8). To this structure are attached tail fibers 92 to 98 nm in length with a characteristic kink 17 to 18 nm from the base-plate (Fig. 10). These fibers are only 1 to 2 nm in thickness (distinctly thinner than T4 tail fibers) and can be seen best in micrographs, like those of Fig. 10, that are slightly underfocused to enhance phase contrast.

Attempted purification by cloning and by varying the host. Successive single-plaque isolations were made to exclude the possibility that the morphological variants might actually be different phages. If the variants bred true, one might hope to obtain populations of particles with only one head size. Using Sh as the host, isolated plaques were picked from plates containing less than 10 plaques to reduce contamination by diffusion of phage from one type of plaque to another. The progeny of each isolated plaque was replated, and the procedure was repeated. However, even after as many as four successive clonings, the resulting progeny always consisted of a mixed population of the variant types.

This was also found when P1 was grown on a series of different bacterial hosts and the progeny of single clones were examined in the electron microscope, the rationale being that the different variants might have different host ranges. However, as shown in Table 3, all P1 preparations contained all three morphological variants. Similar results were obtained with a purified preparation of P1B (99.9%). This indicates that P1B can give rise to the other two variants as well as to itself after infection of each of the hosts tested.

The possibility remains that P1B is supplying information, permitting the development and

FIG. 4. Micrograph of P1B, P1S, and one particle having a head of possible intermediate size (P1I) from the band 4 region of a CsCl gradient. The actual measurement of the P1I head is 74 by 77 by 80 nm. The arrow indicates a neckpiece attached to the tailless P1B head.

release of defective prophages contained in the host cells. Our results, however, suggest that the variants are produced by errors in the morphogenetic assembly of P1, rather than that all seven hosts carry the same defective prophages.

FIG. 3. Micrographs of the heads of P1B, P1S, and P1M particles found in a "light" band (ca. 1.400 g/cm³) isolated from a CsCl gradient of a P1c·C600 lysate. All three heads are from the same two EM plates in a focal series. A triangular facet is apparent on the P1B head capsid; its size is consistent with icosahedral symmetry. Arrows for the P1S and P1M heads point to possible capsomere "doughnuts." The distance between the centers of the two doughnuts indicated for P1M is ca. 11 nm. The difficulty in deducing subunit structure from these heads is multifold: (i) one is seeing the tops and bottoms of the capsids superimposed, (ii) the particles are not empty ghosts, and (iii) the "noise" level of the background is high.

FIG. 5. Several samples of P1 heads exhibiting axial symmetry. The symmetry is indicated by the number on each phage head. All were stained with silicotungstate, except (c) and (d) where 1% uranyl acetate was used. PIB heads are seen in (a), (d), (e), and (f). PIS heads are seen in (b) and (c). Those heads showing threefold axes of symmetry could be either icosahedra or octahedra, but those showing fivefold symmetry are probably icosahedra. The arrow in (b) points to what is probably a head-neck connector seen end-on; if this is true, the empty PIS capsid is actually sitting on a vertex rather than on a side or on an edge.

FIG. 6. *P1B* heads with two tails attached. They are rarely seen in lysates but can be found in *CsCl* gradients at $\rho = 1.454 \pm 0.004$ g/cm³.

Infectivity profiles of P1 in *CsCl* density gradients. The infectivity of the variants was first examined by taking advantage of the fact that they could be separated in *CsCl* density gradients

(6), evidently because they have different proportions of DNA and protein. In addition to bands of phage ghosts and bacterial debris, at least seven visible bands of P1 particles were obtained. Their

FIG. 7. Diagrammatic scale drawing of P1 morphological variants.

contents, as seen in the EM, are shown in Fig. 11 and summarized in Table 4. Band 5, which consists only of P1B particles, contained significantly more material than the other six bands. The relative turbidities of the other six bands varied considerably and did not seem to depend on either the strain of phage (P1 or P1c) used for making the initial lysates or the bacterial host on which the phage was grown. As will be discussed later, it is of particular interest that the P1B particles that persisted in band 2 after repurification in CsCl had only partially filled heads, as judged by the fact that many of them were only partially filled with negative stain.

The infectivity profiles of two successive CsCl density gradients are shown in Fig. 12. The fact that 99% of the PFU were found in band 5 after two successive runs establishes the infectivity of the P1B particles and strongly suggests that the P1S particles in bands 2 and 4 are not infective. The most likely explanation for the small peaks of infectivity observed in bands 2 and 4 (when the MOI is kept below 1 to avoid complementation between particles) is that a few infective P1B

particles became trapped in these bands as they formed during the run.

P1S particles were contained mainly in bands 2 and 4. After three successive runs, band 2 contained as much as 95% P1S. An equally pure preparation of P1S particles from band 4, however, was difficult to obtain, not only because of its proximity to band 5 but also because of the existence of "light P1B" particles which banded at or near the band 4 position (Walker, *unpublished data*).

Bands 6 and 7 contained heads of P1S and P1B, respectively. These tailless particles, as would be suspected, were noninfectious. As will be shown in the discussion, the density of the tailless big heads was useful in determining the relative mass fractions of DNA, head capsid protein, and tail protein.

Comparative infectivity of P1B and P1S from ^{32}P -labeling experiments. The noninfectivity of P1S particles suggested from CsCl gradient profiles was confirmed by an experiment showing that the ^{32}P -containing material in band 2 (P1S) formed very few plaques relative to the ^{32}P -containing material in band 5 (P1B). The fact that band 2 particles contained DNA was established both by Karamata (Certificate Report, Inst. of Mol. Biol., Univ. of Geneva, Switzerland, 1964) and by Ikeda and Tomizawa (20). Karamata identified phenol-extracted material as double-stranded DNA through diphenylamine and buoyant density analyses. Ikeda and Tomizawa further showed that the DNA isolated from band 2 and band 5 particles had the same buoyant density (1.706 g/cm^3) in CsCl gradients and that this density was lower than that of the *E. coli* host-cell DNA (1.710 g/cm^3). They concluded that the band 2 particles probably contained fragments of P1 genome.

P1 was grown in single-cycle broth lysates containing added $^{32}\text{PO}_4$. Bands 2 and 5 were isolated from the gradient and each rerun on second gradients. The corresponding fractions were collected and passed through G25 Sephadex columns to remove the CsCl. The fractions were then treated with deoxyribonuclease and ribonuclease and again passed through Sephadex to remove any ^{32}P -labeled material not contained within the phage heads.

The P1B and P1S fractions were then examined for homogeneity in the EM, assayed for PFU/ml, and counted for radioactivity. The results of two such experiments are summarized in Table 5. The values for PFU per counts per min indicated that the partially pure P1S fraction is only 0.67 to 4% as efficient in forming plaques as the pure P1B fraction. What little plaque-forming activity there was in band 2 can be attributed to infec-

FIG. 8. Concentrated lysate of *P1c·Cla*. In addition to the full *P1B* and *P1M* heads, the following structural features can be seen: extended sheaths showing horizontal striations terminated distally with baseplates (BP) and tail fibers (TF); contracted sheaths with hollow tail tubes exposed; free contracted sheaths showing stain penetration through the central channel; a head-neck connector (HC) with a pronounced hole indicating that it is an annular structure (see also Fig. 2, 5b).

tious, contaminating *P1B*. An additional and unexpected conclusion is that the majority of the *P1B* particles present in band 2 are also noninfectious.

Adsorption and injection by *P1B* and *P1S*. The preceding results provided good evidence that most, if not all, of the *P1S* particles were unable to form plaques, although the reason was not

clear. Therefore, we performed a Hershey-Chase (18) experiment to determine the extent to which *P1B* and *P1S* could adsorb to host cells and inject their ^{32}P -labeled DNA. In the control, *P1B* and *P1S* were mixed with St/22 which does not adsorb *P1*.

The results (Table 6) indicate that, after 20 min had been allowed for adsorption, 31 to 44% of the

FIG. 9. *P1* purified in CsCl gradients. (a) Part of a contracted *P1* sheath standing on end. The appearance is similar to an axial view of a wormgear having 12 projections. Part of the hollow tail tube can be seen. (b) Several head-neck connectors may be seen. Arrow points to a hollow tail tube where the contracted sheath has slid part way down the tube. Note also the many small doughnut structures in the background. These measure 11 ± 1 nm in diameter and are probably capsomeres. The reason for thinking so is that in the larger micrograph from which 9b was excerpted, the majority of the tails are intact whereas the heads are ghosted, and more than half of these head capsids are in a state of breakdown.

FIG. 10. Underfocused electron micrograph of *P1* particles showing tail fibers. The fibers have characteristic kinks 17 to 18 nm from the baseplate. Here phase contrast has made the fibers stand out more clearly than in the other micrographs.

TABLE 3. Proportions of P1S particles in P1 stocks grown on various hosts

Host	Proportion of P1S ^a		Total particles counted ^c
	Avg	Range ^b	
<i>Escherichia coli</i> B..	0.064	0.05 to 0.10	470 (3)
<i>E. coli</i> Bc.....	0.14	0.08 to 0.24	638 (4)
<i>E. coli</i> K-12 (λ)...	0.27	0.20 to 0.37	475 (2)
<i>E. coli</i> C600.....	0.25	0.20 to 0.44	2,950 (8)
<i>E. coli</i> Cla.....	0.14	0.08 to 0.22	700 (3)
<i>E. coli</i> Cla(P3)....	0.19		260 (1)
<i>Shigella dysenteriae</i> Sh.....	0.10	0.04 to 0.22	1,804 (7)

^a Each stock also contained smaller proportions of P1M varying from 0.002 to 0.036.

^b Range represents minimum and maximum proportions counted for different photographic fields (often from independently prepared lysates).

^c Numbers represent total particles counted on one or more photographic fields, the number of fields being given by the figures in parentheses.

radioactivity in the band 2 fraction adsorbed to Sh, compared to 56 to 59% of the radioactivity in the band 5 fraction. Assuming that each P1B particle in band 2 contained 1.5 times as much DNA as each P1S particle (*see below*), a maximum of 27% of the total counts of band 2 could be contained in P1B particles. Then, if the same proportion (57.5%) of P1B particles in band 2 adsorbed as in band 5, the P1B particles of band 2 could account for no more than half of the counts adsorbed from the band 2 fraction. The proportion of adsorbed counts which were injected was almost the same for band 2 and band 5 preparations, so we conclude that P1S particles can inject their DNA.

Killing by P1S particles. It was of interest to determine the ability of particles to kill host bacteria. We preadsorbed Sh cells with the band 2 preparation from experiment 2 of Table 5 under conditions such that one-half of the cells received an infectious MOI (PFU MOI) of 0.00034 and the other half received a PFU MOI of 0.034 (100-fold greater). In the first case, there were 100% bacterial survivors; in the second case, 52%. The killing MOI (kMOI) can be calculated from the fraction of surviving bacteria, S/S_0 , by the relation $S/S_0 = e^{-kMOI}$. For $S/S_0 = 0.52$, kMOI = 0.66. Since the ratio kMOI/PFU MOI = $0.66/0.034 = 19.5$, we could conclude that PFU particles accounted for only 5% of the killed bacteria. Most of the killing was evidently due to P1S particles.

Genetic marker rescue. If P1S particles contain fragments of the P1B genome, it should be possi-

ble to rescue genetic markers from the noninfectious particles by simultaneous infection with P1B particles. We infected Sh cells with P1B carrying a *c* (clear plaque) allele at an MOI of 2.5 and with band 2 noninfectious particles carrying the wild-type (turbid plaque) allele at an estimated particle MOI of 1.7. Preliminary experiments indicated that the interaction of the two alleles in the same infective center gives rise to many "mottled" plaques. Therefore, we plated the cells exposed to both P1Bc and P1S and compared the number of turbid and mottled plaques with those produced by cells plated after exposure to P1S alone at the same MOI. Plaques were scored from enlarged photographs of the plates on which the turbid, clear, and mottled types could easily be distinguished.

The results showed a 2.6-fold increase in turbid plus mottled plaques over the controls. A statistical analysis (null hypothesis) of the experiment indicated that there was less than 1 in 1,000 chance that marker rescue was not occurring. Therefore, we conclude that markers can be rescued from noninfectious band 2 particles, but we cannot tell whether the rescue is occurring from P1S, noninfectious P1B, or both.

P1M particles. Because P1M seldom accounts for more than 1% of the particle population in a given lysate, we have not attempted to obtain a purified preparation. When examining CsCl fractions in the EM, we have observed P1M particles in the buoyant density range of 1.370 to 1.400 g/cm³. Because its density was higher than that of P1 ghosts (1.327 g/cm³) and because the negative stain rarely penetrated the capsid, we assume that P1M contains DNA; however, the P1M head is so small that it could accommodate only about 10% of the P1 genome.

DISCUSSION

Possible triangulation numbers for P1 capsids. Applying the architectural principles elucidated by Caspar and Klug (10) for virus particles, we assume that different P1 capsids have different triangulation numbers, T , which characterize the distribution of structural elements on the surfaces of icosahedra. Then the length of an edge, E , of one of the 20 equilateral triangular facets of an icosahedron is given by the equation

$$E = dT^{1/2} \quad (1)$$

where d is the distance between centers of two dimensional hexagonal capsomeres on the equilateral triangular net depicted by Caspar and Klug (11). Furthermore,

$$E = 0.618 D_G \quad (2)$$

FIG. 11. Micrographs of particles from various bands formed during centrifugation of a concentrated lysate of P1 in CsCl. (a, b) P1S and PIB particles found in bands 2 and 4, respectively; (c) PIB particles in band 5; (d) P1S heads in band 6; (e) PIB heads found in band 7. The contents of the bands are further described in Table 4. Note that during preparation of the specimen many of the particles have lost the DNA once contained in their heads. The bar represents 0.5 μ m.

where D_G is the geometric diameter of such an icosahedron bounded by planes having no thickness (15). Combining equations 1 and 2 we obtain

$$d = 0.618 D_G T^{-1/2} \quad (3)$$

Now the capsomeres of P1 appear to be about 5 nm in thickness, so, as illustrated in Fig. 13, the geometric diameters on which our series of homologous capsids are constructed might well be about 5 nm less than the observed diameters, D_0 . The values of D_G would then be 80, 60, and 42 nm for PIB, P1S, and P1M, respectively. These ratios, 4:3:2, correspond closely to minimal

values for T of 16, 9, and 4 and for d of 12.4, 12.4, and 13.0 nm as shown in Table 7. The constancy of d suggests that the assignments of triangulation numbers are plausible for the P1 variants.

Table 7 also includes the putative intermediate head size of P1I (which would fit $T = 12$) and of the isometric head of P2 (which would fit well with $T = 7$ if d were the same for P1 and P2). Actually, capsomeres seen (Walker, *unpublished data*) on partially degraded capsids of P2 heads have surface diameters in the same range (10 to 12 nm) as those obtained for P1.

The only other phage for which morphological

TABLE 4. *Densities and contents of P1 bands in CsCl density gradients^a*

Band	Relative buoyant densities ^b (g/cm ³)	Per cent content as seen in electron microscope	
		Original ^c (after 1 run)	Purified (after 2 to 3 runs)
1	1.422 ± 0.002	85 P1B, 15 P1S	P1B goes to band 5 and/or 3
2	1.433 ± 0.002	40 P1B, 60 P1S	5 P1B, 95 P1S
3	1.450 ± 0.002	P1B > P1S	95 P1B, 5 P1S (often absent)
4	1.459 ± 0.002	P1B > P1S	60 P1B, 40 P1S
5	(1.473 ± 0.002)	99 P1B, 1 P1S	≥99.9 P1B, ≤0.1 P1S
6 ^d	1.490 ± 0.004		15 P1B heads, 85 P1S heads
7 ^d	1.505 ± 0.002	96 P1B, 4 P1S	99 P1B heads, 1 P1S head
(γ)	(1.508)	(γ)	

^a Similar results concerning bands 1 through 5 were found independently by Karamata (Certificate Report, Univ. of Geneva, 1964).

^b Density of band 5 was determined by using λ as a standard (43). The densities of the other bands were determined by using band 5 as a substandard.

^c More recent observations (Walker, *unpublished data*) indicate that bands 1 and 3 are often entirely absent if the gradients contain 0.005 M MgSO₄.

^d Bands 6 and 7 contain heads only; the tails are missing.

variants are known are the T-even phages. Infectious T4 particles have prolate-shaped heads but, as was mentioned above, in lysates of T4 one also observes particles with isometric heads which contain defective genomes. These heads, however, have the same diameter perpendicular to the long axis of the phage as normal T4, so it would appear that they result from a malfunction of the termination process of capsid elongation rather than from a basic change in triangulation number. By using the optical diffraction and image filtering techniques of Klug and his collaborators (27, 28), Yanagida, Klug, and DeRosier (*personal communication*) recently determined the lattice spacing (*d*) on two types of T4 polyheads to be 11.0 nm and 12.5 nm. Bayer and Remsen (6a) observed a value of about 10 nm for *d*.

Lysates of T4 also contain elongated and spherical τ particles which have antigens in common with those of T4 heads (Laemmli et al., (30a)). However, τ particles appear not to contain DNA, and they have no tails attached. They are probably analogous to the "petite" forms found in preparations of the isometric coliphages λ (23), φ80pt (37), and P2 and P4 (Walker, *unpublished data*) which appear essentially empty and never have tails attached. For this reason, we would not classify them as morphological variants in the sense that we have used the term in this paper. Their roles, if any, in phage morphogenesis are not clear.

Estimation of molecular weights of P1 tails, heads, and DNA contents. The molecular weights of the various components of the P1 variants can be estimated from their buoyant densities. We assume (i) that the tailless P1B particles of band 7 and the complete P1B particles in band 5 both

contain normal complements of P1 DNA and (ii) that the buoyant density of a particle reflects the relative mass of DNA and protein of that particle (43). Ignoring any internal head protein, we have calculated the mass of head protein (*M*_{hp}) and tail protein (*M*_{tp}) as was done for P22 by Israel et al. (22). By definition for complete particles in band 5

$$\begin{aligned}\rho_{P1B[5]} &= \frac{M_d + M_{hp} + M_{tp}}{V_d + V_{hp} + V_{tp}} \\ &= \frac{M_d + M_{hp} + M_{tp}}{M_d/\rho_d + M_{hp}/\rho_{hp} + M_{tp}/\rho_{tp}}\end{aligned}\quad (4)$$

and for the tailless particles in band 7

$$\rho_{P1B[7]} = (M_d + M_{hp}) / (M_d/\rho_d + M_{hp}/\rho_{hp}) \quad (5)$$

where *V*_d, *V*_{hp}, and *V*_{tp} are the volumes occupied by DNA, head protein, and tail protein with corresponding densities ρ_d, ρ_{hp}, and ρ_{tp}. Taking ρ_{P1B[5]} = 1.473 ± 0.002 g/cm³, ρ_{P1B[7]} = 1.505 ± 0.002 g/cm³, ρ_d = 1.706 g/cm³ (19), ρ_{hp} = ρ_{tp} = ρ_{ghosts} = 1.327 g/cm³ (our results), and *M*_d = 60 × 10⁶ daltons (19), we obtain *M*_{hp} = (52.7 ± 1.1) × 10⁶ daltons and *M*_{tp} = (21.8 ± 2.8) × 10⁶ daltons. As a check on the validity of these calculations, we can estimate a density of 1.451 g/cm³ for P1B capsids with two tails, and this is where we find them in density gradients.

If *T* = 16 for the P1B head capsid, the number of structural protein subunits = 60 *T* = 960 (10). The molecular weight for the subunit is then about 55,000 daltons. Dyson and Van Holde (16) reported that the head capsid subunit of λ has a molecular weight of about 55,000 daltons too.

We can now calculate the DNA content of

FIG. 12. *CsCl* density gradients on *P1*·*Sh*: correlation of peaks from PFU values and visible turbidity. The 4.8-ml gradient was run in Tris-phosphate buffer for 32 hr at 30,000 rev/min; 57 fractions were collected (upper figure). Estimated recovery of the initial 1.5×10^{12} PFU was 60%. A log scale is used simply to emphasize the peaks in the region of bands 1 to 4. Over 90% of the recovered PFU are contained in fractions 21 to 24. Fractions 25 to 44 were pooled and rerun in a volume of 4.8 ml for 32 hr at 28,000 rev/min (lower figure). Again, over 90% of the PFU were found in the band 5 position.

other particles. For example, the P1S head capsid ($T = 9$) would have 540 subunits, giving it a mass of $(29.7 \pm 0.6) \times 10^6$ daltons. Using the value of $\rho = 1.433 \pm 0.002$ g/cm³ for the P1S particles in band 2, we can estimate its DNA content to be $(25.7 \pm 1.7) \times 10^6$ daltons. This is in good agreement with the value of Ikeda and Tomizawa of 24×10^6 daltons obtained from the rate of sedimentation of P1 DNA (20). This represents about $42.8 \pm 2.7\%$ or roughly 40% of the normal P1B genome.

We can now estimate that the DNA in the P1S particles present in band 4 ($\rho = 1.459 \pm 0.002$ g/cm³) is $(35.4 \pm 2.3) \times 10^6$ daltons which is $58.9 \pm 3.8\%$ or roughly 60% of the P1 genome. It seems remarkable that the sum of the DNA contents of the P1S particles in bands 2 and 4 adds up to the equivalent of a complete P1B genome ($101.7 \pm 6.7\%$).

Several further calculations of this type can be made. One is of particular interest since it suggests that, like the P1S particles of band 4, the P1B particles found in the band 2 region contain about 60% of the DNA content of a complete P1B genome. Such defective P1B particles would band at $\rho = 1.430$ g/cm³ which is only 0.003 g/cm³ less dense than the P1S particles of band 2. In fact we have found that if the band 2 region is rerun on a very shallow *CsCl* gradient, it separates into two bands about 0.004 g/cm³ apart. The more dense part contains 98% P1S particles, whereas the less dense part contains up to 60% P1B particles (Walker, unpublished data).

Current theories of the packaging of DNA in phage heads (20, 36, 39) do not easily account for the fact that P1B and P1S heads can be partially

TABLE 5. Comparison of plaque-forming ability of band 2 material with band 5 material^a

Determination	Expt 1		Expt 2	
	Band 5	Band 2	Band 5	Band 2
P1 content (from EM counts).....	P1B (100%)	P1S (60%) + P1B (40%)	P1B (100%)	P1S (80%) + P1B (20%)
PFU/ml.....	6.4×10^{10}	1.0×10^8	4.9×10^{11}	8.7×10^8
Counts per min per ml.....	1.77×10^5	4.2×10^4	9.94×10^5	4.32×10^4
PFU per counts per min.....	3.6×10^5	2.4×10^3	4.9×10^5	2.0×10^4
PFU per counts per min relative to band 5.....	(1.0)	0.0067	(1.0)	0.040

^a Abbreviations: PFU, plaque-forming units; EM, electron microscopic.

TABLE 6. *Hershey-Chase experiment: host adsorption and DNA injection from ^{32}P -labeled P1B and P1S fractions^a*

Determination	Band 5 (100% P1B + bacteria)	Band 2 (80% P1S + 20% P1B + bacteria)
Counts in adsorption mixtures	(100)	(100)
Counts pelleted with Sh cells	56-59	31-44
Counts pelleted with St/22 cells (control)	7-18	7-16
Counts in resuspended pellets	(100)	(100)
Counts sedimented after blending Sh pellet	59-60	55-58
Counts sedimented after blending St/22 pellet (control)	10-13	

^a All figures are in relative per cent units. Both supernatants and resuspended pellets were counted at each step of the procedure; discrepancies are indicated by the ranges given. Duplicate samples were counted. Initial adsorption mixtures (28,000 counts/min) for band 5 contained 3×10^8 cells/ml with $\text{MOI} = 0.4$; for band 2 (12,000 counts/min in the adsorption mixture), the same number of cells was used and a similar MOI based on the assumption that P1S contains about one-half as much DNA per particle as P1B. (The band 2 preparation is the same as that of experiment 2 in Table 4.)

FIG. 13. *Exaggerated diagrammatic cross section through an isometric capsid. Symbols: c = capsid thickness = 5 nm; D_G = geometric diameter; D_O = outside diameter.*

filled with DNA. According to one theory (the "head full theory"), circularly permuted genomes like those of T4 and P22 (41) would simply fill all heads. P1 DNA doesn't fill all heads even though it is known to be circularly permuted in the P1B heads found in band 5 (21). It would be premature to discuss mechanisms of packaging DNA in the variants of P1 until we know more about the nature of the DNA fragments found in them.

TABLE 7. *Triangulation numbers (T), diameters (D_0), and calculated intercapsomere distances (d) of P1 variants*

T	$T^{1/2}$	Phage	Observed D_0 (nm)	Calculated d (nm)	Calculated D_0 for $d = 12.4$ nm
1	1.000				25.0
3	1.732				39.6
4	2.000	P1M	47	13.0	45.0
7	2.646	(P2)	(58)	12.4	57.9
9	3.000	P1S	65	12.4	65.0
12	3.464	P11	(~74)	12.3	74.3
13	3.606				77.1
16	4.000	P1B	85	12.4	85.0
19	4.359				92.2
21	4.583				96.6
25	5.000				105.0

ACKNOWLEDGMENTS

We are indebted to Dimitri Karamata for providing us with a copy of his certificate report. D. H. W. thanks the Lister Institute, London, England, where some of the writing was done, and G. G. Meynell for critical reading of parts of the manuscript.

This investigation was supported by Public Health Service Training Grant 5-TI GH-694-05 to the University of Pennsylvania, by National Science Foundation grants GB-982 and GB-4640, and by Public Health Service grant CA-06927 from the National Cancer Institute. An appropriation from the Commonwealth of Pennsylvania to The Institute for Cancer Research is also gratefully acknowledged.

LITERATURE CITED

- Adams, M. H. 1959. Bacteriophages. p. 450-451. Interscience, Publishers, Inc., New York.
- Anderson, T. F. 1953. The morphology and osmotic properties of bacteriophage systems. Cold Spring Harbor Symp. Quant. Biol. 18:197-203.
- Anderson, T. F. 1960. On the fine structures of the temperate bacteriophages P1, P2, and P22, p. 1008-1011. In Proc. Eur. Reg. Conf. Electron Microsc., Delft, 1960, vol. 2. Almquist and Wiksell, Uppsala.
- Anderson, T. F. 1962. Negative staining and its use in the study of viruses and their serological reactions, p. 251-262. In R. H. C. Harris (ed.), The interpretation of ultrastructure, vol. 1. Symp. Int. Soc. Cell Biol. Academic Press Inc., New York.
- Anderson, T. F., and S. Krimm. 1966. Diffraction of light from electron micrographs of helical structures in bacteriophage tail sheaths, p. 145-146. Proc. 6th Int. Cong. Electron Microsc., Kyoto, vol. 2.
- Anderson, T. F., and D. H. Walker, Jr. 1960. Morphological variants of the bacteriophage P1. Science 132:1488.
- Bayer, M. E., and C. C. Remsen. 1970. Bacteriophage T2 as seen with the freeze-etching technique. Virology 40: 703-718.
- Bertani, G. 1951. Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. J. Bacteriol. 62:293-300.
- Boy de la Tour, E., and E. Kellenberger. 1965. Aberrant forms of the T-even phage head. Virology 27:222-225.
- Brenner, S., and R. W. Horne. 1959. A negative staining method for high resolution electron microscopy of viruses. Biochim. Biophys. Acta 34:103-114.

10. Caspar, D. L. D., and A. Klug. 1962. Physical principles in the construction of regular viruses. Cold Spring Harbor Symp. Quant. Biol. 27:1-24.
11. Caspar, D. L. D., and A. Klug. 1963. Structure and assembly of regular virus particles, p. 27-39. In *Viruses, nucleic acids, and cancer*. Williams & Wilkins Co. Baltimore.
12. Cohen, D. 1959. A variant of phage P2 originating in *Escherichia coli*, strain B. Virology 7:112-126.
13. Cummings, D. J., V. A. Chapman, and S. S. DeLong. 1968. Disruption of T-even bacteriophages by dimethyl sulfoxide. J. Virol. 2:610-620.
14. Cummings, D. J., and L. M. Kozloff. 1962. Various properties of the head protein of T2 bacteriophage. J. Mol. Biol. 5:50-62.
15. Cundy, H. M., and A. P. Rollett. 1961. Mathematical models, p. 128. Oxford Univ. Press, London.
16. Dyson, R. D., and K. E. Van Holde. 1967. An investigation of bacteriophage λ , its protein ghosts and subunits. Virology 33:559-566.
17. Eiserling, F. A., and E. Boy de la Tour. 1965. Capsomeres and other structures observed on some bacteriophages. Pathol. Microbiol. 28:175-180.
18. Hershey, A. D., and M. Chase. 1952. Independent functions of viral protein and nucleic acid in growth of bacteriophage. J. Gen. Physiol. 36:39-56.
19. Ikeda, H., and J. Tomizawa. 1965. Transducing fragments in generalized transduction by phage P1. I. Molecular origin of the fragments. J. Mol. Biol. 14:85-109.
20. Ikeda, H., and J. Tomizawa. 1965. Transducing fragments in generalized transduction by phage P1. III. Studies with small phage particles. J. Mol. Biol. 14:120-129.
21. Ikeda, H., and J. Tomizawa. 1968. Prophage P1, an extra-chromosomal replication unit. Cold Spring Harbor Symp. Quant. Biol. 33:791-798.
22. Israel, J. V., T. F. Anderson, and M. Levine. 1967. *In vitro* morphogenesis of phage P22 from heads and base-plate parts. Proc. Nat. Acad. Sci. U. S. A. 57:284-291.
23. Karamata, D., E. Kellenberger, G. Kellenberger, and M. Terzi. 1962. Sur une particule accompagnant le développement du coliphage λ . Pathol. Microbiol. 25:575-585.
24. Kellenberger, E., A. Bolle, E. Boy de la Tour, R. H. Epstein, N. C. Franklin, N. K. Jerne, A. Reale-Scafati, J. Sechaud, I. Bendet, D. Goldstein, and M. A. Lauffer. 1965. Functions and properties related to the tail fibers of bacteriophage T4. Virology 26:419-440.
25. Kellenberger, E., and M. Boy de la Tour. 1965. Studies on the morphopoiesis of the head of phage T-even. II. Observations on the fine structure of polyheads. J. Ultrastruct. Res. 13:343-358.
26. King, J. 1968. Assembly of the tail of bacteriophage T4. J. Mol. Biol. 32:231-262.
27. Klug, A., and J. E. Berger. 1964. An optical method for the analysis of periodicities in electron micrographs, and some observations on the mechanism of negative staining. J. Mol. Biol. 10:565-569.
28. Klug, A., J. T. Finch, R. Leberman, and W. Longley. 1966. Design and structure of regular virus particles, p. 158-189. In G. E. W. Wolstenholme and M. O'Connor (ed.), Principles of biomolecular organization. J. and A. Churchill, Ltd., London.
29. Krimm, S., and T. F. Anderson. 1967. Structure of normal and contracted tail sheaths of T4 bacteriophage. J. Mol. Biol. 27:197-202.
30. Labaw, L. W., and V. M. Mosley. 1954. Demonstration of striated fibers in the capsule of the Lisbonne strain of lysogenic *Escherichia coli*. J. Bacteriol. 67:576-584.
- 30a. Laemmli, U. K., E. Mölbert, M. Showe, and E. Kellenberger. 1970. Form-determining function of the genes required for the assembly of the head of bacteriophage T4. J. Mol. Biol. 49:99-113.
31. Lennox, E. S. 1955. Transduction of linked genetic characters of the host by bacteriophage P1. Virology 1:190-206.
32. Moody, M. F. 1965. The shape of the T-even bacteriophage head. Virology 26:567-576.
33. Moody, M. F. 1967. Structure of the sheath of bacteriophage T4. I. Structure of the contracted sheath and polysheath. J. Mol. Biol. 25:167-200.
34. Moody, M. F. 1967. Structure of the sheath of bacteriophage T4. II. Rearrangement of the sheath subunits during contraction. J. Mol. Biol. 25:201-208.
35. Mosig, G. 1968. A map of distances along the DNA molecule of phage T4. Genetics 59:137-151.
36. Ozeki, H., and H. Ikeda. 1968. Transduction mechanisms. Annu. Rev. Genet. 2:245-278.
37. Shinagawa, H., Y. Hosaka, H. Yamagishi, and Y. Nishi. 1966. Electron microscope studies on $\phi 80$ and $\phi 80pt$ phage virions and their DNA. Biken. J. 9:135-148.
38. Smith, K. O., and M. Trousdale. 1965. Multiple-tailed T4 bacteriophage. J. Bacteriol. 90:796-802.
39. Streisinger, G., J. Emrich, and M. M. Stahl. 1967. Chromosome structure in phage T4. III. Terminal redundancy and length determination. Proc. Nat. Acad. Sci. U.S.A. 57:292-295.
40. Swannstrom, M., and M. H. Adams. 1951. Agar layer method for production of high titer phage stocks. Proc. Soc. Exp. Biol. Med. 78:372-375.
41. Thomas, C. A., Jr. 1967. The rule of the ring. J. Cell Physiol. 70(Suppl. 1):13-34.
42. Ting, R. C. 1962. The specific gravity of transducing particles of bacteriophage P1. Virology 16:115-121.
43. Weigle, J., M. Meselson, and K. Paigen. 1959. Density alterations associated with transducing ability in the bacteriophage lambda. J. Mol. Biol. 1:379-386.
44. Vinograd, J., and J. E. Hearst. 1962. Equilibrium sedimentation of macromolecules and viruses in a density gradient. Fortsch. Chem. Org. Naturst. 20:372-422.