Structure of *Bacillus subtilis* Bacteriophage $\phi 25$ and $\phi 25$ Deoxyribonucleic Acid

W. F. LILJEMARK AND D. L. ANDERSON

Department of Microbiology and School of Dentistry, University of Minnesota, Minneapolis, Minnesota 55455

Received for publication 17 February 1970

The structure of *Bacillus subtilis* bacteriophage $\phi 25$ and $\phi 25$ deoxyribonucleic acid (DNA) were studied by electron microscopy. The head of $\phi 25$ is a regular polyhedron measuring 75 nm in diameter. The uncontracted tail of $\phi 25$ is 130 nm in length and includes a large, complex tail plate. Phage $\phi 25$ DNA is double-stranded and has a molecular weight of approximately 100 million as determined by electron microscopic length measurements and analytical band sedimentation in CsCl. The complementary strands of $\phi 25$ DNA contain numerous random interruptions. Chemical analysis of $\phi 25$ DNA demonstrated that 5-hydroxymethyluracil replaces thymine and that the DNA has a mole per cent (guanine plus cytosine) of 42.

Bacillus subtilis bacteriophage $\phi 25$ is related morphologically and antigenically to a group of *B. subtilis* phages represented by SP8 (Liljemark and Anderson, Bacteriol. Proc., p. 164, 1968; B. E. Reilly, Ph.D. Thesis, Western Reserve University, 1965). In the present study, the structure of $\phi 25$ was elucidated, and the sedimentation coefficient, molecular length, and base composition of $\phi 25$ deoxyribonucleic acid (DNA) were determined.

MATERIALS AND METHODS

Phage production. Lysates of phage $\phi 25$ were prepared on *B. amyloliquefaciens* strain H as described by Reilly and Spizizen (15).

Phage purification. Lysates were clarified in a Sorvall RC-2 centrifuge with a GSA rotor by centrifugation for 20 min at $8,000 \times g$. Supernatant solutions were passed through a type O3 Sela filter, and the filtrate was then spun in a Spinco model L2-65 preparative ultracentrifuge, equipped with a no. 19 rotor, for 1 hr at $25,000 \times g$. Phage pellets were resuspended in a phase buffer containing 0.1 M NaCl, 0.001 M $MgSO_4$, and 0.005 M tris(hydroxymethyl)aminomethane (Tris)- PO_4 buffer (pH 7.5). Cesium chloride was added in an amount equal to 60% of the weight of the phage suspension, and the mixture was centrifuged for 15 to 16 hr at $70,000 \times g$ in the Spinco model L2-65 centrifuge equipped with an SW39L rotor. The phage band, located 27 to 28 mm from the meniscus, was collected, and the cesium chloride was removed by dialysis against 4 liters of phase buffer. Purified phage were stored at 4°C over chloroform.

Extraction of phage DNA. For electron microscopic studies, phage $\phi 25$ DNA was released with

sodium perchlorate by use of a modification of the procedure of Freifelder (10), as reported previously by Anderson and Mosharafa (3). Purified $\phi 25$ was mixed with 4 or 8 M sodium perchlorate to give a final concentration of 2 or 4 M and 7 M perchlorate, respectively, and the mixture was incubated at 37°C for 45 to 60 min.

For chromatographic analysis, phage $\phi 25$ DNA was prepared by the phenol extraction technique of Mandel and Hershey (14), as modified by Reilly and Spizizen (15).

Electron microscopy. Phages were negatively stained with a 2% (w/v) phosphotungstate (PTA) solution adjusted to pH 7 with 1 N KOH (6). Specimens were supported on fenestrated carbon films prepared by the technique of Huxley and Zubay (11).

Phage $\phi 25$ DNA molecules were prepared for electron microscopy by use of the protein film procedure of Kleinschmidt et al. (13), as modified by Anderson et al. (2). The lengths of the DNA molecules were measured as described previously (2, 3).

Electron micrographs were taken with an RCA EMU-3G electron microscope.

Denaturation of $\phi 25$ DNA. Phage $\phi 25$ DNA in 0.15 M NaCl-0.015 M sodium citrate (SSC) was denatured at a concentration of 100 $\mu g/ml$ or less in 0.1 M NaOH by the procedure of Studier (17), taking particular care to avoid shear. After 1 to 2 min at room temperature, the solution was adjusted to pH 8 with an appropriate volume of 1.1 M HCl-0.2 M Tris.

Sedimentation. The native, alkaline denatured and denatured neutralized forms of $\phi 25$ DNA were studied by analytical band sedimentation by using the method of Bruner and Vinograd (7), as modified by Anderson and Mosharafa (3). Calculation of

molecular weight was based on the equations of Studier (17).

Chromatography. Phage $\phi 25$ DNA was hydrolyzed with formic acid and analyzed by descending paper chromatography (4).

RESULTS

Size and morphology of $\phi 25$. The structure of $\phi 25$ is shown schematically in Fig. 1 and negatively stained in Fig. 2 to 5. The dimensions of $\phi 25$ from negatively stained particles are summarized in Table 1. The regular polyhedral head of $\phi 25$ measures 75 nm in diameter. The sheath and tail plate assembly of $\phi 25$ is 120 nm in length when uncontracted and 68 nm when contracted (Fig. 2-5). The uncontracted sheath exhibits a periodicity of about 3.8 nm with 30 such periods per phage tail (Fig. 3). The upper portion of the tail consists of a neck region and a collar 10 nm in diameter, located below the head and above the sheath (Fig. 5). The lower portion of the tail consists of a large, intricate tail plate, 70 nm in diameter. The detailed morphology of the tail plate is obscured on uncontracted tails but is more clearly revealed on contracted tails. The tail plate is apparently attached to the sheath by 12 tail processes, of which only 6 are visible on lateral view (Fig. 5). These

processes, seen only when the tail is contracted, are connected to each other some distance from the sheath to form the tail plate (Fig. 5). The tail spikes, 14 nm in length, are suspended from the tail plate (Fig. 2, 5) and are apparently used for attachment to the bacterial cell wall. In the contracted position, the tail core, measuring 130 nm in length and 6 nm in diameter, is seen protruding about 50 nm from the tail plate (Fig. 2).

Length of $\phi 25$ DNA. The lengths of $\phi 25$ DNA molecules extracted with 7 M sodium perchlorate were measured by electron microscopy (Fig. 6). Fifty-four molecules ranged in length from 45 to 55 nm, and the mean length was 50 nm (Fig. 7). Assuming that $\phi 25$ DNA is a double helix sodium salt in the B crystallographic form and that the base pairs are 0.34 nm apart, the molecular weight corresponding to the mean length of 50 nm would be 9.6×10^7 daltons.

Extraction of $\phi 25$ DNA at sodium perchlorate concentrations of 2 and 4 M gave a spectrum of fragments shorter than the molecules released in 7 M sodium perchlorate. The DNA was only partially released from the phage with 2 and 4 M perchlorate as indicated by protrusion of $\phi 25$ DNA from the phage tail in protein film preparations and by the presence of partially filled heads in negatively stained preparations. When 7 M sodium perchlorate was used, $\phi 25$ DNA was free from phage tails in the protein films, and the negatively stained preparations clearly showed that the $\phi 25$ heads had lysed or broken, effecting total release of the phage DNA.

Analytical band sedimentation of $\phi 25$ DNA. Phage $\phi 25$ DNA (0.6 μ g) was centrifuged in CsCl at a density of 1.54 g/cm³ and a speed of 35,600 rev/min. Ultraviolet absorption photographs were taken at 8-min intervals (Fig. 8). A standard sedimentation coefficient ($S_{20,w}$) of 55 was calculated for the sodium salt of $\phi 25$ DNA from the equations of Studier. The molecular weight from these data is 11.5×10^7 daltons, a value somewhat higher than that observed by electron microscopy. Repeated attempts to band the alkaline denatured and denatured neutralized forms of $\phi 25$ DNA failed. The diffuse distribution of the DNA in the tube indicated that the complementary strands of $\phi 25$ DNA contain numerous random interruptions.

Base composition of $\phi 25$ DNA. Phage $\phi 25$ DNA was hydrolyzed in formic acid, and the bases were separated by descending paper chromatography. Phage $\phi 25$ DNA contained 5-hydroxymethyluracil (HMU) rather than thy-

FIG. 1. Schematic diagram of phage $\phi 25$.

FIG. 2. Phage $\phi 25$ showing the regular polyhedral capsid with the tail in a normal and contracted position. $\times 240,000$.

FIG. 3. Phage $\phi 25$ showing the periodicity of the tail in its normal position. $\times 320,000$.

FIG. 4. Phage $\phi 25$ showing its tail collar. $\times 347,000$.

FIG. 5. Phage $\phi 25$ showing the organization of the tail plate with appendages. $\times 373,000$.

FIG. 6. Native phage $\phi 25$ DNA molecule released with 7 M sodium perchlorate. $\times 29,600$.

mine (Table 2). Spectrophotometric absorption ratios of 250 to 260 nm and 280 to 260 nm supported the identification.

Kallen et al. have shown partial degradation of HMU by formic acid hydrolysis (12). If the molar ratio of HMU to adenine were one in $\phi 25$ DNA, then the guanine plus cytosine (GC) content would be 42%.

DISCUSSION

Phage $\phi 25$ has an intricate tail plate, a long contractile tail, and a regular polyhedral head which is probably an icosahedron. Thus, $\phi 25$ is closely related morphologically to *B. subtilis* phage SP8 (8) and to the series of *B. subtilis* phages studied by Eiserling, including SP5-SP9 and SP13 (Ph.D. Thesis, University of California, Los Angeles, 1964).

The incomplete or partial release of $\phi 25$ DNA in 2 and 4 M sodium perchlorate is similar to that reported for phage SP50 under similar

conditions (5). DNA is expelled through the tail at these lower perchlorate concentrations, whereas 7 M perchlorate causes a rupture of the head and release of the complete genome.

Measurement of 54 $\phi 25$ DNA molecules released intact by 7 M sodium perchlorate revealed a mean length of 50 μm , which corresponds to a molecular weight of 96 million. In contrast, analytical band sedimentation of $\phi 25$ DNA in CsCl yielded molecular weight values of 115 to 120 million, by use of the empirically derived equations of Studier relating *S* and *M*

TABLE 1. Dimensions of phage $\phi 25$

Dimension	Size
<i>nm</i>	
<i>Normal</i>	
Head	
Diameter.....	75
Neck	
Length.....	9
Collar	
Diameter.....	10
Thickness.....	3
Sheath	
Diameter.....	13
Length (including plate).....	120
Band periodicity.....	3.8
Tail plate	
Diameter.....	70
<i>Contracted</i>	
Sheath	
Diameter.....	16
Length.....	54
Tail plate	
Diameter.....	70
Sheath plus tail plate	
Length.....	68
Core	
Diameter.....	6
Length.....	130

FIG. 7. Distribution of the lengths of $\phi 25$ DNA molecules extracted with 7 M sodium perchlorate. Micrographs were taken at magnifications of 5,450 $\times$ and enlarged 12 $\times$ to give a total magnification of $\times 65,400$.

TABLE 2. Chromatographic analysis of $\phi 25$ DNA

Bases ^a	Combined base standard			$\phi 25$ DNA				Calf thymus DNA			
	<i>R_F</i>	250/260 nm	280/260 nm	<i>R_F</i>	250/260 nm	280/260 nm	Amt (μmoles)	<i>R_F</i>	250/260 nm	280/260 nm	Amt (μmoles)
A	0.35	0.78	0.37	0.34	0.77	0.36	0.15	0.35	0.78	0.37	0.11
T	0.78	0.70	0.53					0.77	0.71	0.53	0.10
U	0.68	0.85	0.18								
HMU	0.61	0.77	0.37	0.62	0.78	0.35	0.04				
G	0.25	1.38	0.81	0.26	1.37	0.82	0.11 ^b	0.26	1.38	0.83	0.08 ^c
C	0.48	1.47	0.49	0.47	1.49	0.50	0.11 ^b	0.49	1.49	0.51	0.09 ^c

^a A, adenine; T, thymine; U, uracil; HMU, 5-hydroxymethyluracil; G, guanine; C, cytosine.
^b $\phi 25$ DNA % GC = 42%.
^c Calf thymus DNA % GC = 44%.

(17). This is about 20% higher than the molecular weight based on length measurements. Studier states that native T2 DNA gives a 10% higher $S_{20,w}$ value with his equations than would be predicted from published values of molecular weight based on other methods (17). An 8 to 10% reduction in the $S_{20,w}$ value obtained for native $\phi 25$ DNA would result in a molecular weight of 100 million, a value in good agreement with that obtained by electron microscopy.

The equations of Studier, however, did yield the anticipated value for the molecular weight of denatured T2 DNA. Presence of random single-stranded breaks in $\phi 25$ DNA prevented a test of Studier equations for denatured $\phi 25$ DNA. Anderson and Mosharrafa (3) demonstrated that, under the identical conditions used in this study, the molecular weight of alkaline-denatured DNA of *B. subtilis* phage $\phi 29$ was exactly half that of the native DNA, indicating no single-stranded interruptions. Davison et al. presented presumptive evidence for single-stranded interruptions in several phenol-extracted viral DNA species including SP8 which is closely related to $\phi 25$ (9).

Previous calculation of the mole fraction GC of $\phi 25$ DNA from equilibrium density gradient sedimentation data resulted in a value of 83%, and the GC calculated from thermal denaturation was 15% (B. E. Reilly, Ph.D. Thesis, Western Reserve University, 1965). DNA from *B. subtilis* phages SP8 and ϕe which exhibited these anomalies was shown to contain HMU instead of thymine (12, 16, 18). Chromatographic analysis of the $\phi 25$ DNA has shown that HMU replaces thymine and that $\phi 25$ DNA has a GC content of 42%.

ACKNOWLEDGMENTS

This investigation was supported by Public Health Service grants AI-08088 from the National Institute of Allergy and Infectious Diseases and 2 TO1 DE-00143-06 from the National Institute of Dental Research, and by grant GB-6695 from the National Science Foundation.

We are indebted to David E. Peterson for doing the ultracentrifugation.

FIG. 8. Analytical band sedimentation patterns of native $\phi 25$ DNA. Phage $\phi 25$ DNA extracted in 7 M sodium perchlorate was diluted in standard saline-citrate, and 0.6 μ g of DNA was sedimented in CsCl having a density of 1.54. The analytical band sedimentation patterns are from left to right. The intervals between the frames were 8 min, and the rotor speed was 35,600 rev/min.

LITERATURE CITED

1. Abelson, J., and C. A. Thomas, Jr. 1966. The anatomy of the T5 bacteriophage DNA molecule. *J. Mol. Biol.* 18:262-288.
2. Anderson, D. L., D. D. Hickman, and B. E. Reilly. 1966. Structure of *Bacillus subtilis* bacteriophage $\phi 29$ and the length of $\phi 29$ deoxyribonucleic acid. *J. Bacteriol.* 91:2081-2089.
3. Anderson, D. L., and E. T. Mosharafa. 1968. Physical and biological properties of phage $\phi 29$ deoxyribonucleic acid. *J. Virol.* 2:1185-1190.
4. Bendich, A. 1957. Methods for characterization of nucleic acids by base composition, p. 715-723. *In* S. P. Colowick and N. O. Kaplan (ed.), *Methods in enzymology*, vol. 3. Academic Press Inc., New York.
5. Biswal, N., A. K. Kleinschmidt, H. C. Spatz, and T. A. Trautner. 1967. Physical properties of the DNA of bacteriophage SP50. *Mol. Gen. Genet.* 100:39-55.
6. Brenner, S., and R. W. Horne. 1959. A negative staining method for high resolution electron microscopy of viruses. *Biochim. Biophys. Acta.* 34:103-110.
7. Brunner, R., and J. Vinograd. 1964. The evaluation of standard sedimentation coefficients of sodium RNA and sodium DNA from sedimentation velocity data in concentrated NaCl and CsCl solutions. *Biochim. Biophys. Acta* 108: 18-29.
8. Davison, P. F. 1963. The structure of bacteriophage SP8. *Virology* 21:146-151.
9. Davison, P. F., D. Freifelder, and B. W. Holloway. 1964. Interruptions in the polynucleotide strands in bacteriophage DNA. *J. Mol. Biol.* 8:1-10.
10. Freifelder, D. 1966. Effect of NaClO_4 on bacteriophage: release of DNA and evidence for population heterogeneity. *Virology* 28:742-751.
11. Huxley, H. E., and G. Zubay. 1960. Electron microscope observations on the structure of microsomal particles from *Escherichia coli*. *J. Mol. Biol.* 2:10-18.
12. Kallen, R. G., M. Simon, and J. Marmur. 1962. The occurrence of a new pyrimidine base replacing thymine in a bacteriophage DNA: 5-hydroxymethyluracil. *J. Mol. Biol.* 5:248-250.
13. Kleinschmidt, A. K., D. Lang, D. Jacherts, and R. K. Zahn. 1962. Darstellung und langennmessungen des gesamen desoxyribonucleinsaeure-inhaltes von T2-bakteriophagen. *Biochim. Biophys. Acta* 61:857-864.
14. Mandel, J. D., and A. D. Hershey. 1960. A fractionating column for analysis of nucleic acids. *Anal. Biochem.* 1: 66-77.
15. Reilly, B. E., and J. Spizizen. 1965. Bacteriophage deoxyribonucleate infection of competent *Bacillus subtilis*. *J. Bacteriol.* 89:782-790.
16. Roscoe, D. H., and R. G. Tucker. 1966. The biosynthesis of 5-hydroxymethyldeoxyuridylic acid in bacteriophage-infected *Bacillus subtilis*. *Virology* 29:157-166.
17. Studier, F. W. 1965. Sedimentation studies on the size and shape of DNA. *J. Mol. Biol.* 11:373-390.
18. Szybalski, W. 1968. Use of cesium sulfate for equilibrium density gradient centrifugation, p. 330-360. *In* L. Grossman and K. Moldave (ed.), *Methods in enzymology*, vol. 12, part B. Academic Press Inc., New York.