Morphology and Physiology of the Intracellular Development of *Bacillus subtilis* Bacteriophage $\phi 25$

W. F. LILJEMARK AND D. L. ANDERSON

Department of Microbiology and School of Dentistry, University of Minnesota, Minneapolis, Minnesota 55455

Received for publication 17 February 1970

The morphology of the intracellular development of bacteriophage $\phi 25$ in *Bacillus subtilis* 168M has been correlated with nucleic acid synthesis in infected cells. Host deoxyribonucleic acid (DNA) synthesis was shut off by a phage-induced enzyme within 5 min after infection, and another phage-mediated function extensively degraded host DNA at the time of cell lysis. Synthesis of phage DNA in infected cells began within 5 min and continued until late in the rise period. After phage DNA synthesis and coinciding with lysis, much of the unpackaged, newly synthesized phage DNA was degraded. Studies of thin sections of $\phi 25$ infected cells suggested that unfilled capsids may be precursors to filled capsids in the packaging process. To assess dependence of capsid formation on phage DNA replication, cells were either treated with mitomycin C and infected with normal phage or infected with ultraviolet-irradiated (99% killed) $\phi 25$. Only empty capsids were found in these cells, indicating that capsid production may be independent of the presence of newly synthesized viral DNA.

Our previous morphological studies on the intracellular development of bacteriophage $\phi 25$ in *Bacillus subtilis* 168M suggested that $\phi 25$ head capsids were formed and then filled with deoxyribonucleic acid (DNA) (Liljemark and Anderson, Bacteriol. Proc., p. 164, 1968). These results prompted us to correlate intracellular morphological changes with the kinetics of host and phage DNA synthesis to understand better the $\phi 25$ infection process in *B. subtilis* and to compare this infection with other *B. subtilis* phage systems. The present report provides morphological evidence that empty capsids are formed when DNA replication is inhibited.

MATERIALS AND METHODS

Phage and host. Phage $\phi 25$ was originally obtained from B. E. Reilly. *B. subtilis* 168M (*ind*⁻) was the host strain in these studies, and all experiments were conducted in Difco Antibiotic Medium No. 3. Phage $\phi 25$ has a latent period of 20 to 25 min and a rise period of 40 to 50 min under the conditions used in this study. The generation time of *B. subtilis* 168M was 30 min.

Electron microscopy. Infected and uninfected cells were fixed and dehydrated for electron microscopy by the method of Kellenberger et al. (4). The cells were embedded in Epon 812 by a modification of the procedure of Luft (6). Thin sections were cut on an LKB Ultratome, mounted on carbon-coated Formvar-covered grids, and doubly stained either with uranyl acetate and Millonig's lead hydroxide (9) or uranyl

acetate and Reynold's lead citrate (11). All electron micrographs were taken with an RCA EMU-3G electron microscope.

Cell fractionation and radioisotope assay procedures. ³H-thymidine incorporation was followed by placing 0.2 ml of culture containing the radioisotope in 3 ml of iced 10% trichloroacetic acid. After 60 min, the cold mixture was filtered through 25-mm membrane filters (Millipore Corp., Bedford, Mass.; pore size, 0.45 μ m) and washed with 10 ml of iced 10% trichloroacetic acid. The filters were allowed to dry at room temperature and assayed for radioactivity as described below.

Cells grown in the presence of ³²P-phosphate were fractionated to isolate the ribonucleic acid (RNA) and DNA components by the method of Schmidt and Thannhauser (13), as modified by Schachtele and Rogers (12). ³²PO₄-labeled RNA and DNA were assayed by placing 0.1-ml samples on Whatman no. 3 paper discs (22 mm) mounted on stick pins. After drying, the discs were placed in counting vials containing 5 ml of scintillation fluid, which consisted of 100 mg of 1,4-bis-2-(5-phenyloxazolyl)-benzene and 4 g of 2,5-diphenyloxazole per liter of toluene. Samples were counted with a Packard Tri-Carb liquid scintillation spectrometer.

RESULTS

Intracellular development of $\phi 25$. Samples of a culture of *B. subtilis* 168M infected with $\phi 25$ at a multiplicity of four particles per bacterium were taken at 10-min intervals from the time of phage

FIG. 1. *B. subtilis* 168M cell, less than 2 min after infection, with intact phage ϕ 25 attached by its tail plate appendages. $\times 106,000$.

addition (time zero) through 120 min and prepared for electron microscopy.

Attachment of ϕ 25 to the cell wall was accomplished by the tail plate and tail-plate appendages (Fig. 1). Adsorbed contracted phages exhibited a capsid with low electron density, consistent with discharge of the DNA into the cell.

Cells fixed 10 min after infection showed no change in intracellular morphology and resembled uninfected control cells (Fig. 2). In cells fixed at 20 min, many areas of very low electron density and a few structures with contents resembling the DNA-plasm of the host were observed, both corresponding to the size of ϕ 25 heads. The areas of lowest electron density were generally outside the DNA-plasm throughout the infection, whereas the "partially filled" structures were generally located in the DNA-plasm (Fig. 3-5). A distinct peripheral limiting structure, presumably the phage capsid or an immature form of the capsid, was observed around both the "empty" bodies and the "partially filled" structures (Fig. 6, 7). Electron-dense areas, interpreted as full capsids, were usually seen in the nuclear region. Changes in the nuclear area were confined to an increase in volume probably reflecting phage DNA synthesis, but no disruption or fragmentation of the nuclear region was observed (Fig. 4, 8). The number of electron-dense areas increased with time after

infection (Fig. 8), but at the point of lysis cells still contained many empty capsids (Fig. 9). The particular cell shown in Fig. 9 contains an unusually large number of empty capsids, and whether some of these capsids contained DNA at the time of fixation is unknown. Estimates of the number of complete versus incomplete phages made in ϕ 25-infected cells, based on electron microscopy of sections and the number of ghosts obtained from normal lysates, indicated that only about half of the capsids are completed.

Effect of ϕ 25 infection on host DNA synthesis. Phage ϕ 25 DNA contains hydroxymethyluracil, not thymine (5), and incorporation of ^3H -thymidine may be used to monitor host DNA synthesis in ϕ 25 infected cells. After ϕ 25 infection, there was an abrupt cessation of ^3H -thymidine uptake by *B. subtilis* 168 (Fig. 10). The addition of chloramphenicol (100 $\mu\text{g}/\text{ml}$) caused a reduction in the rate of DNA synthesis, but host DNA synthesis did occur in infected cells in the presence of the antibiotic.

Phage ϕ 25-induced degradation of host DNA. Morphological examination of ϕ 25 infected cells does not reveal the extensive disruption of the host DNA-plasm that characterizes infection by T2 virus (4). But the analysis of trichloroacetic acid-insoluble DNA in infected cells suggests degradation of host DNA. *B. subtilis* 168M was

grown for four generations in the presence of ^3H -thymidine and infected with phage $\phi 25$. At intervals, samples were treated with lysozyme (20 $\mu\text{g}/\text{ml}$) and trichloroacetic acid-insoluble radioactivity was assayed. DNA degradation was evident by 40 min (Fig. 11). Since the DNA of lysozyme-treated uninfected cells remained acid-insoluble for the duration of the experiment, it is not likely that host nuclease alone could account for the degradation.

RNA and phage DNA synthesis in infected cells.

A modification of the Schmidt-Thannhauser technique was used to study differential incorporation of $^{32}\text{PO}_4$ into RNA and phage DNA during phage $\phi 25$ infection (Fig. 12). RNA synthesis was stimulated in infected cells, but net synthesis ceased 10 min after infection. By this time, the maximal rate of phage DNA synthesis was attained. The extensive release of $^{32}\text{PO}_4$ late in the rise period suggests that unpackaged $\phi 25$ DNA as well as host DNA was degraded.

Capsid formation. If capsid formation is independent of the presence of newly synthesized viral DNA, then capsid formation should occur when DNA synthesis is inhibited. Ultraviolet-irradiated $\phi 25$ (>99% killed) was used to infect *B. subtilis* 168M at a multiplicity of infection (MOI) of 4, and $^{32}\text{PO}_4$ incorporation into RNA and DNA was measured. Cell samples were also prepared for electron microscopy. Net RNA synthesis continued for 50 min, and a low level of DNA synthesis was detected (Fig. 13). Viral capsids were observed, but no capsids containing electron-dense material were evident at any time during the experiment (Fig. 14, 15). Most of the sectioned cells contained from three to six empty capsids. Multiplying these average numbers per cell in sections by a factor of 20 (assuming an average section thickness of no more than 50 nm) would yield an estimate of as much as 120 empty capsids per cell.

In addition, cells were infected with phage $\phi 25$ (MOI of about 4) in the presence of mitomycin C (25 $\mu\text{g}/\text{ml}$). DNA synthesis was not detected, although RNA synthesis continued for 20 to 30 minutes (Fig. 13). The rate of DNA synthesis in untreated infected cells was far greater than in the uninfected control cells (Fig. 12). Electron micrographs of thin sections of treated cells again showed distinct, empty viral capsids (Fig. 16-18).

FIG. 2. Uninfected control cell of *B. subtilis* 168M. $\times 37,000$.

FIG. 3. *B. subtilis* 168M cells fixed 20 min after infection showing many areas of very low density (immature phage capsids) outside the DNA-plasm of the cell. $\times 27,000$.

FIG. 4. *B. subtilis* 168M cell fixed 20 min after infection showing areas of low electron density outside the nuclear region and some areas with cores on the periphery of the DNA-plasm. $\times 47,000$.

FIG. 5. *B. subtilis* 168M cells showing various developmental stages 30 min after infection. $\times 27,000$.

FIG. 6-7. *B. subtilis* 168M cells 30 min after infection showing various stages of phage development. $\times 107,000$; $120,000$.

FIG. 8. *B. subtilis* 168M cell fixed 40 min after infection. $\times 53,000$.

FIG. 9. *B. subtilis* 168M cell undergoing lysis. Note the large number of unfilled capsids. $\times 67,000$.

FIG. 10. Host DNA synthesis in $\phi 25$ -infected *B. subtilis* 168M. *B. subtilis* 168M was grown for one generation in the presence of ^3H -thymidine (16.5 c/mmole, 1.0 $\mu\text{C}/\text{ml}$). When the cells reached a concentration of $7 \times 10^7/\text{ml}$, chloramphenicol was added to two cultures. After 5 min, one of these cultures was infected with $\phi 25$ at a MOI of 4. Phage $\phi 25$ (MOI of 4) was added to a third culture without chloramphenicol. Cultures were assayed for radioactivity at 5-min intervals. Symbols: $\circ$, control cells; $\odot$, cells and chloramphenicol; $\bullet$, cells and chloramphenicol and $\phi 25$; $\bullet$, cells and $\phi 25$.

DISCUSSION

Empty, partially filled, and full capsids were observed in thin sections of $\phi 25$ -infected *B. subtilis* and the ratio of full to empty capsids increased

FIG. 11. Effect of $\phi 25$ infection on *B. subtilis* 168M DNA. Cells were grown for four generations in the presence of ^3H -thymidine (16.5 c/mmole, 1.0 $\mu\text{C}/\text{ml}$). They were collected after reaching a concentration of about $2 \times 10^8/\text{ml}$, and washed three times with unlabeled Penassay Broth. The culture was diluted slightly, allowed to equilibrate at 37 C for 5 to 10 min, and then divided into three parts. At zero time, one culture received the phage diluent, the second received $\phi 25$ at an MOI of 5, and the third lysozyme to give a final concentration of 20 $\mu\text{g}/\text{ml}$. Samples were assayed at intervals for radioactivity. Symbols: $\bullet$, control cells; Δ , cells and lysozyme; $\circ$, cells and phage $\phi 25$.

FIG. 12. Nucleic acid synthesis in $\phi 25$ -infected *B. subtilis* 168M. When exponentially growing *B. subtilis* 168M cells reached a concentration of $7.5 \times 10^7/\text{ml}$, $^{32}\text{PO}_4$ (4.0 $\mu\text{C}/\text{ml}$) was added, and the culture was divided into two parts. Phage $\phi 25$ was added to one of the flasks at an MOI of 2 to 3. Samples from both cultures were assayed at intervals for radioactivity. Symbols: $\bullet$, control cells; Δ , $\phi 25$ -infected cells.

FIG. 13. Phage and host nucleic acid synthesis in *B. subtilis* 168M (1) treated with mitomycin C and (2) infected with ultraviolet-irradiated $\phi 25$. Bacteria were grown to a concentration of 7.5×10^7 /ml, and the culture was divided into two parts. Carrier free $^{32}\text{PO}_4$ ($8.0 \mu\text{C}/\text{ml}$) was added to culture (1) and the culture was then subdivided into three parts. Mitomycin C, at a final concentration of $25 \mu\text{g}/\text{ml}$, and $\phi 25$ at an MOI of 4 were added to one part of this culture. Ultraviolet-irradiated $\phi 25$, 99% killed, was added at a MOI of 4 to the second portion. The third portion was left uninoculated. The cultures containing mitomycin C or ultraviolet-irradiated $\phi 25$ were kept in the dark throughout the experiment. Radioactivity was assayed at intervals. Culture (2) was divided and inoculated in the same manner as (1), and samples were fixed and embedded for electron microscopy. Symbols: $\circ$, uninfected control cells; $\bullet$, cells infected by ultraviolet-irradiated $\phi 25$; $\odot$, mitomycin C-treated cells infected by $\phi 25$.

with time after the infection. The micrographs suggest, but do not prove, that empty capsids are precursors of filled capsids. Unfilled capsids were observed in a phage-infected marine bacterium, but the authors did not interpret this as evidence for head filling (14). Other evidence for head filling was reported by Luftig (Abstr., Biophys. J. Soc., 9:A259, 1969) in coliphage T4.

Phage $\phi 25$ infection rapidly stops host DNA synthesis and also causes degradation of host DNA. Experiments based on $^{32}\text{PO}_4$ incorporation indicated that $\phi 25$ DNA synthesis commences within 5 min after infection, thus resembling the SP8 and 2C systems studied by Marmur et al. (8, 10). The $\phi 25$ system differs, however, in that unpackaged phage $\phi 25$ DNA is degraded about the time of lysis.

Two means of inhibiting DNA synthesis were investigated for effects on the morphology of $\phi 25$ development. The first, described in the *B. subtilis* phage 2C infection by Pene and Marmur (10), employed mitomycin C to block DNA synthesis.

Mitomycin C prevented the synthesis of lysozyme in the phage 2C infection but allowed formation of the "early" enzyme deoxycytidine monophosphate deaminase. The second means for inhibiting phage DNA synthesis utilized ultraviolet-irradiated phage which has been shown in the coliphage T2 and T4 systems (1, 2) to inhibit both synthesis of phage DNA and appearance of "late" proteins. In experiments with $\phi 25$, DNA synthesis was not detected in the mitomycin C-treated cells infected with normal phage and reduced to a low level in normal cells infected with ultraviolet-treated $\phi 25$. Electron microscopy demonstrated that capsids are formed in both of these experiments. Thus, capsid production is apparently independent of the presence of newly synthesized DNA. Other evidence for the expression of "late" functions in the absence of DNA synthesis was presented by Edlin, who used temperature-sensitive mutants of phage T4 blocked in DNA synthesis (3), and by Margaretten et al., who used T4-infected, hydroxyurea-treated *Escherichia coli* (7).

FIG. 14-15. Empty capsids in *B. subtilis* 168M cells 40 min after infection with ultraviolet-irradiated $\phi 25$.
 $\times 93,000$; $100,000$.

FIG. 16-18. Mitomycin C-treated *B. subtilis* 168M cells 20 min after infection with normal phage $\phi 25$ (Fig. 16, 17), showing a normal cellular morphology with numerous empty capsids. $\times 27,000$; 47,000. Fig. 18 shows a mitomycin C-treated *B. subtilis* 168M cell 40 min after infection with normal $\phi 25$. $\times 87,000$.

ACKNOWLEDGMENTS

This investigation was supported by Public Health Service grants AI-08088 from the National Institute of Allergy and Infectious Diseases and 2 TO1 DE-00143-06 from the National Institute of Dental Research, and GB-6695 from the National Science Foundation.

LITERATURE CITED

1. Dirksen, M.-L., J. S. Wiberg, J. F. Koerner, and J. M. Buchanan. 1960. Effect of ultraviolet irradiation of bacteriophage T2 on enzyme synthesis in host cells. *Proc. Nat. Acad. Sci. U.S.A.* 46:1425-1430.
2. Ebisuzaki, K. 1966. Ultraviolet sensitivity and functional capacity in bacteriophage T4. *J. Mol. Biol.* 20:545-558.
3. Edlin, G. 1965. Gene regulation during T4 development. I. Phenotypic reversion of T4 amber mutants by 5-fluorouracil. *J. Mol. Biol.* 12:363-374.
4. Kellenberger, E., A. Ryter, and J. Sechaud. 1958. Electron microscope study of DNA plasms, vegetative and mature phage DNA as compared with normal bacterial nucleoids in different physiological states. *J. Biophys. Biochem. Cytol.* 4:671-676.
5. Liljemark, W. F., and D. L. Anderson. 1970. Structure of *Bacillus subtilis* bacteriophage ϕ 25 and ϕ 25 deoxyribonucleic acid. *J. Virol.* 6:107-113.
6. Luft, J. H. 1961. Improvements in epoxy resin embedding methods. *J. Biophys. Biochem. Cytol.* 9:409-414.
7. Margaretten, W., C. Morgan, H. S. Rosenkranz, and H. M. Rose. 1966. Effect of hydroxyurea on virus development. I. Electron microscopic study of the effect on the development of bacteriophage T4. *J. Bacteriol.* 91:823-833.
8. Marmur, J., C. M. Greenspan, E. Palecek, F. M. Isahan, J. Levine, and M. Mandel. 1963. Specificity of the complementary RNA formed by *B. subtilis* infected with bacteriophage SP8. *Cold Spring Harbor Symp. Quant. Biol.* 28:191-199.
9. Millonig, G. 1961. A modified procedure for lead staining of thin sections. *J. Biophys. Biochem. Cytol.* 11:736-739.
10. Pene, J. J., and J. Marmur. 1967. Deoxyribonucleic acid replication and expression of early and late bacteriophage function in *Bacillus subtilis*. *J. Virol.* 1:86-91.
11. Reynolds, E. S. 1963. Lead citrate staining technique. *J. Cell Biol.* 17:208-213.
12. Schachtele, C. F., and P. Rogers. 1965. Canavanine death in *Escherichia coli*. *J. Mol. Biol.* 14:474-489.
13. Schmidt, G., and S. J. Thannhauser. 1945. A method for the determination of deoxyribonucleic acid, ribonucleic acid, and phosphoproteins in animal tissues. *J. Biol. Chem.* 161:83-89.
14. Valentine, A. F., and G. B. Chapman. 1966. Fine structure and host-virus relationship for a marine bacterium and its bacteriophage. *J. Bacteriol.* 92:1535-1554.